AUSTRIAN JOURNAL OF EARTH S



LARGER BENTHIC FORAMINIFERA FROM THE MIDDLE EDCENE TO OLIGOCENE OF TANZANIA

Laura J. COTTON" & Paul N. PEARSON

KEYWORDS

larger foraminifera Oligocene Tanzania Eocene MECO

School of Earth and Ocean Sciences, Cardiff University, Main Building, Park Place, Cardiff. CF10 3AT, UK; ⁹ Corresponding author, cottonlj@cardiff.ac.uk

ABSTRACT

Larger benthic foraminifera (LBF) are common and diverse throughout the Paleogene sediments of southern Tanzania, but have previously been little studied. A recent programme of onshore drilling known as the Tanzania Drilling Project has recovered large proportion of this succession for palaeoclimatic and palaeontological study. The sediment is largely a hemipelagic clay with secondary gravity sediment flow limestones and calcarenites. LBF occur concentrated in the secondary beds and are present in some clay horizons. Planktonic foraminiferal, nannofossil and, in some cases, stable isotope studies of the clays allow the larger benthic foraminiferal ranges to be tied to global stratigraphy. Here we use nine of these drill sites to examine the LBF from the Middle Eocene to Oligocene. Within this interval several global turnover events of long-ranging and widespread LBF genera are known to occur. However, problems with biostratigraphy mean the exact timing and therefore mechanisms remain uncertain. Our study shows that ranges of Tanzanian LBF genera are within known global ranges. Additionally, there is a change in the LBF assemblage with a number of local first and last occurrences of genera during the Bartonian, which may have potential links to the onset of the Mid Eocene Climatic Optimum.

1. INTRODUCTION

The coastal basins of southern Tanzania contain a thick succession of Paleogene marine sediments that have been the focus of much recent research on the palaeontological and palaeoclimatic history of the area (Pearson et al., 2004, 2006; Nicholas et al., 2006, 2007; Pearson et al., 2008; Wade and Pearson, 2008; Lear et al., 2008; Dunkley-Jones et al., 2008, 2009; Pearson et al., 2009). Rich assemblages of larger benthic foraminifera (LBF) occur throughout but have received relatively little attention. Blow and Banner (1962) gave a brief overview of LBF species found in much of the succession, but largely concentrated on the planktonic foraminifera. Here we focus on the Middle Eocene to Oligocene distribution of LBF taxa as recorded in drill cores and compare the record to other areas. This work builds on the study of Cotton and Pearson (2011) which focused on the Tanzanian LBF at the Eocene Oligocene transition (EOT), extending the record at lower resolution to a much longer time period.

The Kilwa and Lindi Districts of Tanzania contain a thick succession of clay sediment from Santonian to Oligocene age (Figure 1; Nicholas et al 2006; 2007). These sediments are formally defined as the Kilwa group and are split into four formations: the Nangurukuru (Santonian to Paleocene), Kivinje (Paleocene to lower Lutetian), Masoko (lower Lutetian to mid Bartonian) and Pande formations (mid Bartonian to the Rupelian; Nicholas et al., 2006). A large proportion of this succession has been recovered in a series of shallow drill cores by the Tanzania Drilling Project (TDP; see Nicholas et al., 2006 for review). The sediments of the Kilwa group are broadly homogeneous and consist of a succession of dark greenish grey clays and claystones to marls with limestones and calcarenites deposited as sediment gravity flows. LBF are abundant in

the limestone beds and they also occur in clay horizons in the succession. Specimens from the clays are generally better preserved than those from the limestone, but are smaller and more dispersed. Calcareous micro- and nannofossils are often exceptionally well-preserved (e.g. Pearson et al., 2008; Bown et al., 2008) and planktonic foraminiferal and nannofossil studies have been used to determine the stratigraphy of the succession (Pearson et al., 2004, 2006; Nicholas et al., 2006; Wade and Pearson, 2008, Dunkley Jones et al., 2008a,b, 2009). Stable isotope analysis has been carried out on three sites which span the Eocene - Oligocene transition enabling these sites to be correlated with the global isotope stratigraphy (Pearson et al., 2008).

Nine of the TDP sites are stratigraphically placed between the Upper Ypresian and Upper Rupelian (Pearson et al., 2004, 2006; Nicholas et al., 2006). During this interval LBF are known to have undergone several global turnover events (Hallock et al., 1991). Towards the late Middle Eocene there is a large turnover in nummulitids, followed by the extinction of *Assilina* and then the extinctions of *Alveolina* and large species of *Nummulites*. A further global extinction of LBF is then seen at the Eocene - Oligocene Transition, with the extinction of the orthophragmines, the pellatispirids and several species of *Nummulites* (Adams et al., 1986). All of these are long ranging and widespread groups of LBF.

However, biostratigraphy of LBF is often problematic. The mutually exclusive environments of planktonic microfossils and LBF mean that cross-correlation is often difficult. This, coupled with species endemism, has resulted in the use of regional zonations, such as the Shallow Benthic Zones in the Tethyan region (Cahuzac and Poignant, 1997; Serra Kiel et

Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania

al., 1998) and the East Indian Letter Classification in the Indo-Pacific region (Adams, 1970; Renema, 2007). Accurate correlations between LBF events and planktonic foraminiferal, nannofossil and climatic events are therefore often difficult to determine and mechanisms remain uncertain.

Here we use samples from the Middle Eocene to Oligocene TDP sites to give an overview of LBF genera present in southern Tanzania. The well-dated stratigraphy allows correlation of LBF ranges with global biostratigraphy and allows potential links between LBF extinctions and global climatic events to be explored.

2. MATERIALS AND METHODS

This study utilises nine TDP sites from the top of the Kivinje Formation to the top of the Pande Formation to examine LBF occurrences across the Middle Eocene to Oligocene interval. The locations (Figure 1), assigned planktonic foraminiferal Zones and ages of these cores are summarised in Table 1.

Clay samples were washed through a 63 μ m sieve and residues dried. Oriented sections of loose individual LBF from clays were made for identification. The secondary beds were studied in randomly oriented thin section and acetate peels (prepared according to Dickson, 1965, 1966).

3. RESULTS

The Eocene secondary beds are bioclastic pack-grainstones. LBF are the dominant bioclastic component, but also present are echinoid fragments, red algae, smaller foraminifera and serpulid worm tubes. The LBF occur in a matrix of finer carbonate fragments including smaller-sized foraminifera and quartz grains. Most LBF have suffered at least some abrasion to the outer test whorls, comparable with category 2 to 3 on the scale of Beavington-Penney (2004). Intraclasts are present in some samples, but are rare. Several large-sized LBF show evidence of boring and/or overgrowth by red algae. The Oligocene secondary beds are calcarenites with a high fine grained quartz content. Bioclasts including LBF are present but less abundant than in the Eocene beds. The samples from TDP Site 6



FIGURE 1: Location map showing Middle Eocene to Oligocene Tanzanian Drilling Project sites.

are distinctly different from the rest of the succession. These contain a large number of reworked carbonate clasts including LBF in a finer quartz-rich matrix.

The stratigraphic occurrences of the LBF genera are shown in Figure 2 with the levels of the TDP sites. The assemblage throughout the Lutetian (TDP Sites 2, 20 and 13) remains fairly constant and is generally dominated by *Nummulites* and *Alveolina* with orthophragmines and *Linderina*. Also present but less frequent are *Lockhartia*, *Assilina*, *Glomalveolina* and *Orbitolites*, although the latter two genera only occur in the lower half of the Lutetian. *Assilina* are rare in the core samples, but

Site	Location	UTM	Planktonic fora- miniferal Zone	Age
TDP 2	SW Kilwa prison (80m from TDP 20)	37L 555371 9013813	E7-E9	L. Ypresian to Mid Lutetian
TDP 20	SW Kilwa prison	37L 555457 9013846	E7-E9	L. Ypresian to Mid Lutetian
TDP 30	Roadside N of Mkazambo	37L; 558673 8975981	E9-E11	Middle to Upper Lutetian
TDP 18	Roadside N of Mkazambo	37L; 558640 8975370	E12 (lower part)	Bartonian
TDP 4	Ras Tipuli	37L; 578530 8900033	E12-E13	mid Bartonian
TDP 11	Close to the village of Stakishari	37L; 560250 8983211	E16-O1	Upper Priabonian to Lower Rupelian
TDP 12	Close to the village of Stakishari	37L; 560222 8981309	E16-O1	Upper Priabonian to Lower Rupelian
TDP 17	Close to the village of Stakishari	37L; 560539 8984483	E15-O1	Upper Priabonian to Lower Rupelian
TDP 6	W of Kilwa Masoko airstrip	37L; 555752 9014922	02-04	Lowe Oligocene

TABLE 1: Location and age data for the TDP sites used in this study.



FIGURE 2: Range chart showing the stratigraphic occurrence of LBF genera in the TDP succession. Stratigraphic levels of the TDP sites are shown in the left hand column. Crosses indicate the occurrence of a genus; black lines are used to indicate the occurrences of genera in TDP 11, 12 and 17 as a high resolution study has been carried out. Pale grey bars in the background show known global ranges of genera (Hallock et al., 1991; Renema, 2002; BouDagher-Fadel, 2008).

191

Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania

abundant in outcrops on the edge of Kilwa Creek, close to the drill sites of TDP 2 and TDP 20 and thought to be of similar age. Towards the top of TDP Site 13 the last Alveolina specimen is found, which has had an almost continuous pre-sence in the TDP samples to this point. In the Bartonian (TDP Site 18) there is a high diversity of LBF genera and there appears to have been a change in the LBF assemblage between the uppermost samples of TDP Site 13 and the lowermost samples of TDP Site 18. In TDP Site 18 there are the first occurrences of reticulate Nummulites, Operculina and Sphaerogypsina. Heterostegina also becomes more common, thus far represented by a single specimen towards the top of TDP 13. There is also the last occurrence of Lockhartia, and the last definite occurrence of Linderina. TDP Site 4 is still within the Bartonian but stratigraphically slightly higher than TDP Site 18. In these samples Nummulites are most common and occur with Discocyclina, Operculina and possible Linderina. There is then a gap in the TDP record and the next sites are Upper Priabonian to Lower Rupelian (TDP Sites 11, 12 and 17) and continuously span the EOT (Pearson et al., 2008). These samples also show a relatively high diversity of LBF. At these sites the majority of the genera present go extinct at the Eocene/Oligocene boundary, coincident with the extinction of the Hantkeninidae (see the more detailed study by Cotton and Pearson, 2011). Only Sphaerogypsina and some species of Nummulites pass through the boundary apparently unaffected. Several of the genera which do go extinct at the boundary, including Fabiania, Pellatispira and Palaeonummulites, are only present in these samples. In the Upper Rupelian (TDP Site 6) secondary beds there are a large number of intraclasts and the only LBF in them that do not appear to have been reworked are small Nummulites.

4. DISCUSSION

As LBF are known to inhabit the shallow carbonate environment (Beavington-Penney and Racey, 2004) all specimens occurring in the hemipelagic clay succession of the Kilwa group have been transported. Despite this, they can still be used to give ranges of LBF as the majority appear to have been deposited penecontemporaneously with the clay sediments. Intraclasts are rare and the foraminifera appear to have been deposited as uncemented grains, evidenced by tests that have been abraded during transport. The high levels of abrasion suggest high energy transport and rapid deposition (Beavington-Penney, 2004). There are occasional examples of largesized LBF which have been bored and so were dead tests when re-deposited, the borings are infilled by the surrounding matrix indicating they were still uncemented when re-deposited. The exception to this penecontemporaneous sedimentation are the secondary beds from site TDP Site 6 which contain a very large amount of intraclasts often with biostratigraphically older LBF. The only LBF in this site that do not appear to be reworked are small Nummulites.

The ranges of the LBF genera found in the TDP sites from Tanzania all fall within global ranges (Adams 1970; Adams et

al., 1986; Hallock et al., 1991; Serra Kiel et al., 1998; Renema, 2002, Renema, 2007). One of the largest changes in the Tanzanian LBF assemblage takes place between the last sample of TDP Site 13 and the first sample of TDP Site 18. These two sites were drilled less than 1 km from each other, making it unlikely that the assemblage change is due to geographic location or varying sediment source. The succession in TDP Site 18 has been assigned to the lower part of planktonic foraminiferal Zone E12 (Nicholas et al., 2006) and is therefore close to a global climatic warming event known as the Mid Eocene Climatic Optimum (MECO). The MECO was a global interval of rapid warming from ~40 Ma to ~40.8 Ma which interrupted the overall cooling trend of the Eocene (Bohaty et al., 2009; Edgar et al., 2010). The MECO is relatively little studied with regards to both planktonic foraminifera and LBF response. Some overturning within the planktonic acaraninids and morozovelloidids is associated with this period and there is the occurrence of a short ranging 'excursion taxon', Orbulinoides beckmanni, which spans the MECO (Edgar et al., 2010). In LBF, overturning within Nummulites and the first occurrences of reticulate Nummulites have been noted to occur around this level (Hallock et al., 1991) although have not been directly associated with the MECO event.

Studies of planktonic biostratigraphy from TDP Site 18 show that the *Orbulinoides beckmanni* present are apparently early forms (Nicholas et al., 2006), suggesting that the succession is from the very early stages of, or even slightly before, the MECO. Therefore the faunal change in Tanzania occurred before the peak of the MECO. Additionally TDP Site 18 samples contain reticulate *Nummulites* from the *N. fabianii* group and therefore their evolution pre-dates the main phase of the MECO. Preceding the MECO is an interval of cooling from ~40.6-41.6 Ma which has also been recognised in δ^{18} O records (Bohaty et al., 2009). A cooling trend followed by the onset of rapid warming may have been responsible for at least local changes in the faunal distribution.

The global extinctions of *Alveolina* and large species of *Num-mulites* are known to occur in the late Middle Eocene, slightly preceded by the extinction of *Assilina* (Hallock et al., 1991). Although attempts have been made by the TDP to locate and drill this stratigraphy, thus far these have proved unsuccessful.

Assilina is rare in the TDP samples, and little light is shed on its extinction level. No examples of Assilina were seen above the Lutetian, however the abundance of them in outcrop suggests that this may be due to the small area sampled by the TDP cores and the fact that LBF assemblages vary across the shelf. Further outcrop samples may therefore be able to help with this issue. The disappearance of large sized species of *Nummulites* requires a more detailed species-level study. However, *Nummulites* specimens > 10 mm diameter are present in the clays of TDP Site 18 and *Nummulites* > 25 mm belonging to the *N. perforatus* group were found in outcrop close to the drill site. These large *Nummulites* are not found at TDP Sites 11, 12 and 17. This suggests that they became extinct within the un-sampled interval. *Alveolina* has a lower last occurrence at the top of TDP Site 13 and is found in almost all TDP samples until this point. This indicates that the mechanism causing the assemblage changes between the successions in TDP Sites 13 and 18 may have also caused a local "early" extinction of *Alveolina*.

The Eocene/Oligocene boundary shows a rapid extinction of a number of genera shown to have long ranges in the Tanzanian record. The Priabonian samples are the most diverse, but this could be in part due to the higher-resolution study carried out on them. The Tanzanian sites have allowed accurate correlation between the LBF extinctions and plankton extinctions in the Eocene-Oligocene transition (Cotton and Pearson, 2011). The extinction was found to be closely coincident with the extinction of the planktonic foraminiferal Family Hantkeninidae which defines the Eocene/Oligocene boundary. This precedes the global sea level drop known to occur in the transition by ~200,000 years, indicating this was not the cause of the LBF extinction as previously suggested.

5. CONCLUSION

This study gives an insight into the range of LBF genera present in a previously little-studied region. The penecontemporaneous redeposition of the LBF into the hemipelagic environment enables LBF ranges to be tied to global nannofossil and planktonic foraminiferal biostratigraphy. With further work, there is therefore a high potential for using the TDP data to help improve global LBF stratigraphy and provide links between LBF and climatic events. The Eocene-Oligocene transition cores have previously enabled the exact level of the LBF extinction to be determined relative to global isotope stratigraphy. Here we have shown faunal changes take place on the Tanzanian platform possibly linked to the MECO and preceding cooling interval.

ACKNOWLEDGEMENTS

The work was supported by the Tanzania Commission for Science and Technology and the Tanzania Petroleum Development Corporation work. E. Ozcan, C. Papazzoni and W. Renema are thanked for their help and advice with identifications. We also thank the reviewers M. Wagreich and G. Less for their constructive comments which helped to improve the manuscript.

REFERENCES

Adams, C.G., 1970. A reconsideration of the East Indian letter classification of the Tertiary. Bulletin of the British Museum (Natural History) Geology, 19, 1-137.

Adams, C.G., Butterlin, J. and Samanta, B.K., 1986. Larger foraminifera and events at the Eocene / Oligocene boundary in the Indo-West Pacific region. In: C. Pomerol and I. Premoli Silva (eds.), Terminal Eocene events. Elsevier, Amsterdam, pp. 237-252. Beavington-Penney, S.J., 2004. Analysis of the effects of abrasion on the test of *Palaeonummulites venosus*: implications for the origin of nummulithoclastic sediments. Palaios, 19, 143-155.

Beavington-Penney, S.J. and Racey, A., 2004. Ecology of extant nummulitids and other LBF: applications in palaeoenvironmental analysis. Earth-Science Reviews, 67, 219-265.

Berggren, W.A., Kent, D.V., Swisher, I.C.C. and Aubry, M.P., 1995. A revised Cenozoic geochronology and chronostratigraphy. In: W.A Berggren., D.A. Kent and J. Hardenbol (eds.), Geochronology, time scales and global stratigraphic correlation: a unified temporal framework for an historical geology. SEPM (Society for Sedimentary Geology) Special Publications, pp. 129-212.

Blow, W. H. and Banner, F.T., 1962. Part 2: The Mid Tertiary (Upper Eocene to Aquitanian) Globigerinaceae. In: F.E. Eames, F.T. Banner, W.H. Blow, and W.J. Clarke (eds), Fundamentals of Mid-Tertiary stratigraphical correlation. Cambridge University Press, pp. 61-154.

Bohaty, S.M., Zachos, J. C., Florindo, F. and Delaney, M. L., 2009. Coupled greenhouse warming and deep-sea acidification in the middle Eocene. Paleoceanography, 24, PA2207 doi:10.1029/2008PA001676.

Bown, P.R., Dunkley Jones, T., Lees, J.A., Randell, R.D., Mizzi, J.A., Pearson, P.N., Coxall, H.K., Young, J.R., Nicholas, C.J., Karega, A., Singano, J. and Wade, B.S., 2008. A Paleogene calcareous microfossil Konservat-Lagerstätte from the Kilwa group of coastal Tanzania. Geological Society of America Bulletin, 120, 3-12.

BouDagher-Fadel, M.K., 2008. Evolution and geological significance of larger benthic foraminifera. Developments in Palaeontology and Stratigraphy 21, Elsevier, Amsterdam. 540 pp.

Cahuzac, B. and Poignant, A., 1997. Essai de biozonation de l'Oligo—Miocène dans les bassins européens à l'aide des grands foraminifères néritiques. Bulletin de la Société Géologique de France, 168, 155-169.

Cotton, L.J. and Pearson, P.N., 2011. Extinction of larger benthic foraminifera at the Eocene/Oligocene boundary. Palaeogeography, Palaeoclimatology, Palaeoecology, doi:10.1016/j. palaeo.2011.09.008.

Dickson, J.A.D., 1965. A modified staining technique for carbonates in thin section. Nature, 4971, 587.

Dickson, J.A.D., 1966. Carbonate identification and genesis as revealed by staining. Journal of Sedimentary Petrology, 36, 491-505.

Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania

Dunkley Jones, T., Bown, P.R. and Pearson, P.N., 2009. Exceptionally well preserved upper Eocene to lower Oligocene calcareous nannofossils (Prymnesiophycidae) from the Pande Formation (Kilwa group) Tanzania. Journal of Systematic Palaeontology, 4, 359-411.

Dunkley Jones, T., Bown, P.R., Pearson, P.N., Wade, B.S., Coxall, H.K. and Lear, C.H., 2008. Major shifts in calcareous phytoplankton assemblages through the Eocene-Oligocene transition of Tanzania and their implications for low-latitude primary production. Paleoceanography, 23, PA4204 doi:10. 1029/2008PA001640.

Edgar, K. M., Wilson, P.A., Sexton, P. F., Gibbs, S.J., Roberts, A. P. and Norris, R.D., 2010. New biostratigraphic, magnetostratigraphic and isotopic insights into the Middle Eocene Climatic Optimum in low latitudes. Palaeogeography, Palaeoclimatology, Palaeoecology, 297, 670-682.

Hallock, P., Premoli Silva, I. and Boersma, A., 1991. Similarities between planktonic and larger foraminiferal evolutionary trends through Paleogene paleoceanographic changes. Palaeogeography, Palaeoclimatology, Palaeoecology, 83, 49-64.

Lear, C.H., Bailey, T.R., Pearson, P.N., Coxall, H.K. and Rosenthal, Y., 2008. Cooling and ice growth across the Eocene— Oligocene transition. Geology, 36, 251-254.

Nicholas, C.J., Pearson, P.N., Bown, P.R., Dunkley Jones, T., Huber, B.T., Karega, A., Lees, J.A., McMillan, I.K., O'Halloran, A., Singano, J.M. and Wade, B.S., 2006. Stratigraphy and sedimentology of the Upper Cretaceous to Paleogene Kilwa Group, southern coastal Tanzania. Journal of African Earth Sciences, 45, 431-466.

Nicholas, C.J., Pearson, P.N., McMillan, I.K., Ditchfield, P.J. and Singano, J.M., 2007. Structural evolution of southern coastal Tanzania since the Jurassic. Journal of African Earth Sciences, 48, 273-297.

Pearson, P.N., Foster, G.L. and Wade, B.S., 2009. Atmospheric carbon dioxide through the Eocene–Oligocene climate transition. Nature, 461, 1110-1113.

Pearson, P.N., McMillan, I.K., Wade, B.S., Dunkley Jones, T., Coxall, H.K. Bown, P.R. and Lear, C.H., 2008. Extinction and environmental change across the Eocene-Oligocene boundary in Tanzania. Geology, 36, 179-182.

Pearson, P.N, Nicholas, C.J., Singano, J.M, Bown, P.R., Coxall, H.K., van Dongen, B.E., Huber, B.T., Karega, A., Lees, J.A., MacLeod, K., McMillan, I.K., Pancost, R.D., Pearson, M. and Msaky, E., 2006. Further Paleogene and Cretaceous sediment cores from the Kilwa area of coastal Tanzania: Tanzania Drilling Project Sites 6-10. Journal of African Earth Sciences, 45, 279-317. Pearson, P.N., Nicholas, C.J, Singano, J.M., Bown, P.R., Coxall, H.K., van Dongen, B.E., Huber, B.T., Karega, A., Lees, J.A., Msaky, E., Pancost, R.D., Pearson, M. and Roberts, A. P., 2004. Paleogene and Cretaceous sediment cores from the Kilwa and Lindi areas of coastal Tanzania: Tanzania Drilling Project Sites 1–5. Journal of African Earth Sciences, 39, 25-62.

Renema, W., 2002. Larger foraminifera as marine environmental indicators. Scripta Geologica, 124, 1-260.

Renema, W., 2007. Fauna Development of Larger Benthic Foraminifera in the Cenozoic of Southeast Asia. In: W. Renema (ed.), Biogeography, time and place: distributions, barriers and islands. Springer, Dordrecht, pp. 179-215.

Serra-Kiel, J., Hottinger, L., Caus, E., Drobne, K., Ferrandez, C., Jauhri, A.K., Less, G., Pavlovec, R., Pignatti, J., Samso, J.M., Schaub, H., Sirel, E., Strougo, A., Tambareau, Y., Tosquella, J. and Zakrevskaya, E., 1998. Larger foraminiferal biostratigraphy of the Tethyan Paleocene and Eocene. Bulletin de la Société Géologique de France, 169, 281-299.

Wade, B. S. and Pearson, P.N., 2008. Planktonic foraminiferal turnover, diversity fluctuations and geochemical signals across the Eocene / Oligocene boundary in Tanzania. Marine Micropaleontology, 68, 244-255.

Wade, B.S., Pearson, P.N., Berggren, W.A. and Pälike, H., 2011. Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the Geomagnetic Polarity and Astonomical Time Scale. Earth Science Reviews, 104, 111-142.

Received: 24 October 2011 Accepted: 16 March 2012

Laura J. COTTON" & Paul N. PEARSON

School of Earth and Ocean Sciences, Cardiff University, Main Building, Park Place, Cardiff. CF10 3AT, UK;

" Corresponding author, cottonlj@cardiff.ac.uk

© Österreichische Geologische Gesellschaft/Austria; download unter www.geol-ges.at/ und www.biologiezentrum.at

Laura J. COTTON & Paul N. PEARSON

© Österreichische Geologische Gesellschaft/Austria; download unter www.geol-ges.at/ und www.biologiezentrum.at

Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania

PLATE 1:

LBF in petrological thin sections from secondary limestones. FIGURE A: Nummulites sp. (TDP 20) FIGURE B: Nummulites sp. (TDP 18) FIGURE C: Nummulites sp. (TDP 4) FIGURE D: Asterocyclina sp. (TDP 20) FIGURE E: Discocyclina sp. (TDP 13) FIGURE F: Heterostegina sp. (TDP 18) FIGURE G: Linderina sp. (TDP 18) FIGURE H: Lockhartia sp. (TDP 20)

Laura J. COTTON & Paul N. PEARSON



© Österreichische Geologische Gesellschaft/Austria; download unter www.geol-ges.at/ und www.biologiezentrum.at

Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania

PLATE 2:

a-d) LBF in petrological thin sections from secondary limestones; e-h) Oriented individual LBF thin sections of specimens from clays.
FIGURE A: Alveolina sp. (TDP 2)
FIGURE B: Alveolina sp. (TDP 2)
FIGURE C: Glomalveolina sp. (TDP 2)
FIGURE D: Orbitolites sp. (TDP 2)
FIGURE E: Reticulate Nummulites of the N. fabianii group (TDP 18)
FIGURE F: Asterocyclina sp. (TDP 18)
FIGURE G: Operculina gomezi (TDP 18)

FIGURE H: Orbitoclypeus sp. (TDP 18)

Laura J. COTTON & Paul N. PEARSON



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Austrian Journal of Earth Sciences

Jahr/Year: 2012

Band/Volume: 105_1

Autor(en)/Author(s): Cotton Laura J., Pearson Paul

Artikel/Article: Larger benthic foraminifera from the Middle Eocene to Oligocene of Tanzania. 189-199