#### 2012

VIENNA

### QUANTITATIVE CHARACTERISATION OF CATACLASITES USING A STATISTICAL APPROACH (ANALYSIS OF VARIANCE)

Norbert KOHLMAYER<sup>1</sup> & Bernhard GRASEMANN

#### KEYWORDS Analysis of variance

Brittle deformation Cataclasites Cyclades Tectonic

D-Value

Aegaen

Department of Geodynamics and Sedimentology, University of Vienna, Althanstraße 14, Vienna, Austria; <sup>9</sup> Corresponding author, norbert.kohlmayer@aon.at

#### ABSTRACT

Cataclastic rocks can be described by both a range of shape parameters and the particle size distribution. In this work, we apply statistical methods to investigate these variables in samples with different degrees of maturity from high- and low-angle normal faults within marbles in the W-Cyclades (Greece). Four shape parameters (circularity, elliptical PARIS factor, solidity and the aspect ratio) were used to describe the particles. Using a statistical analysis, the variance of the cataclastic particles of a particular shape is partitioned into fractions attributable to different sources of variation. In addition to the shape parameters, the particle size distribution was included, to test whether there is a relationship between particle size and shape.

The investigations demonstrate that the analysis of variance is a statistical method ideally suited for quantitatively studying the grain-size and shape parameters in cataclasites. Our results indicate that although the particle size distributions of the test samples are very similar, three shape parameters (solidity, circularity and the elliptical PARIS factor) can discriminate between the samples. Ideally, numbers of the shape parameter discretized in individual classes of equivalent diameter of the particle components should be used to quantitatively describe the different samples in order to derive information about the deformation mechanisms in the fault rock. Some of the investigated shape parameters record a clear dependence on the grain-size. In the investigated samples, smaller particles record a higher circularity and a lower elliptical PARIS Factor than larger particles. However, no relationship between the solidity or aspect ratio and the grain size has been observed. This suggests that abrasion and comminution are the dominant deformation mechanism in the finer grained particles but fracturing and cracking are prevalent in the more angular, coarser grained particles.

Kataklasite können mit einzelnen Formparametern der Komponenten respektive mit Verteilungen der Korngrößen der kataklastischen Komponenten beschrieben werden. In dieser Arbeit werden verschiedene Formparameter der Komponenten zusammen mit ihrer Größenverteilung statistisch mit der Varianzanalyse untersucht. Die Methoden wurden an kataklastischen Störungsgesteinen mit unterschiedlichem Reifegrad von steilen und flachwinkeligen Störungen in Marmoren aus den W-Kykladen (Griechenland) durchgeführt. Als unabhängige Variablen wurden neben den unterschiedlichen Kataklasiten aus steilen und flachwinkeligen Abschiebungen noch die Variable Korngröße in fünf Klassen berücksichtigt. Als abhängige Variable fungierten die Kornformparameter Zirkularität, ein elliptischer PARIS Faktor, Solidität sowie das Achsenverhältnis d.h. das Längen-, Breitenverhältnis der Komponenten. Es sollte die Frage geklärt werden, ob in den verwendeten Proben ein Zusammenhang zwischen Korngröße und Kornform gegeben ist bzw. ob sich die Proben hinsichtlich der Kornformparameter signifikant unterscheiden.

Generell kann gesagt werden, dass die Varianzanalyse eine äußerst geeignete Methode darstellt um Kataklasite quantitative zu beschreiben. Die Untersuchungen an den Testproben erbrachten im wesentlichen folgendes Ergebnis. Obwohl die untersuchten Proben alle eine sehr ähnliche Korngrößenverteilung aufweisen, können sie über die Formparameter Zirkularität, elliptischer PARIS Faktor und Solidität signifikant differenziert werden. Die genannten Kornformparameter diskretisiert in unterschiedliche Korngrößenklassen wurden verwendet um die verschiedenen Proben quantitativ zu beschreiben bzw. daraus Informationen auf die zugrundeliegenden Deformationsmechanismen in den Störungsgesteinen abzuleiten. Ein wichtiger Zusammenhang besteht zwischen den Kornformparametern und der Korngröße da kleinere Komponenten eine höhere Zirkularität und einen geringeren elliptischen PARIS Faktor zeigen als größere Komponenten. Bezüglich Solidität und Achsenverhältnis konnte kein Zusammenhang zur Korngröße festgestellt werden. Zusammenfassend kann für die untersuchten Proben gesagt werden, dass die runderen, kleinen Korngrößen durch Rollen und Abreiben deformiert worden sind, während bei den eckigen größeren Komponenten die Deformation durch Zerbrechen dominiert hat.

#### 1. INTRODUCTION

Cataclasites are fault rocks that form by the mechanical fragmentation of rocks due to microcracking and frictonal processes, such as sliding, grinding, and rotation of the fragments (Passchier and Trouw, 2005 and references cited therein). During cataclastic flow, which may be associated with a dilational component of strain, voids are created that may be filled with

material precipitated from fluids. This material may be subsequently involved in the cataclastic processes (cf. Hausegger et al. 2010). Cataclastic flow usually occurs at non- to low-grade metamorphic conditions and at relatively high strain rates. High fluid-pressures promote cataclastic flow and are responsible for the common occurrence of veins in cataclasites and tectonic breccias (Blenkinsop, 2000 and references cited therein).

All of these processes lead to a particle size distribution (PSD) which is characteristically fractal (e.g., Blenkinsop, 1991). A fractal distribution of particle sizes d can be described by the relation:

$$N(d) \sim d^{-D}$$
 Ep. 1

where N(d) is the number of particles greater than size d, and D is the fractal dimension. Since the D-value is used to describe the relationship between grain size and frequency, it is useful to specify the range over which the relationship has a good fit to the data. Truncation, sampling bias and low image resolution effects may lead to the deviation of the large and the small grain sizes from this relationship (Blenkinsop, 1991).

During translation and rotation in cataclastic flow, initially angular fragments may be rounded by fracturing (including grinding and abrasion). Thus an analysis of the progressive shape changes and the spatial rearrangement of the particles during shearing is of great importance (Mair et al., 2002; Storti et al., 2003). Since increasing roundness is a sign of increasing wear, by increasing deformation or displacement, the shape of the fragments can be used to distinguish mature cataclasites (i.e. gouges which have accommodated large displacements) from newly fragmented rocks (Cladouhos, 1999; Storti et al., 2003). It has been suggested that the D-value, in conjunction with shape parameters, can be used to describe cataclastic rocks (Heilbronner and Keulen, 2006) and to correlate this description with the conditions of formation. The present study expands on these ideas and suggests an approach based on a statistical analysis of variance to combine the shape parameters for a quantitative description of cataclastic gouge rocks.

#### 2. SHAPE AND GRAIN SIZE DESCRIPTION OF CA-TACLASITES

A standard method to describe the frequencies of particle grain sizes in a sample has been described by Blenkinsop (1991); this demonstrated that multiple fracturing of basalts produced a PSD with a higher fractal dimension than a single fracturing event. Grady and Kipp (1987) generalized from experimental data to show that single tensile fragmentation leads to particle size distributions with D-values less than 2, in contrast to shearing and comminution which results in D-values between 2 and 2.4. High D-values, between 2.60 and 2.82, were derived from fault gouges in the Lopes and Witwatersrand faults (Blenkinsop 1991), where PSDs were influenced by lithology (mineralogical composition), fragmentation process, initial size distribution, number of fracturing events and energy input (Arbiter and Harris, 1965; Hartman, 1969). Some of the earliest experiments on gouge formation (Engelder 1974) showed that the proportion of finer fragments increased with displacement and confining pressure. A reasonably linear reduction in median grain size with increasing confining pressure was recognized for a natural fault gouge in a granodiorite from the San Andreas fault (Sammis et al. 1986). Heilbronner and Keulen (2006) used the D-value to distinguish between cracked fragments from gouge material. Both types of fault rocks exhibit two slopes: for grain size <  $2\mu$ m D ~ 1.0 for both fault rock types; for grain size >  $2\mu$ m, cracked material shows D ~ 1.6 while gouge recorded D > 2.0.

Image analysis programs were used to automatically derive parameters which describe the size and shapes of the particles (Heilbronner and Keulen, 2006). Two important parameters are the aspect ratio and the PARIS factor. The aspect ratio ( $\Delta R$ ) is defined as the ratio between the longest L and the shortest S projection of a particle. The PARIS factor (Panazzo and Hurlimann, 1983) is a measure of the irregularity of grain boundaries and is defined as the ratio of the perimeter length divided by the outline of the convex hull that envelopes the grain. A smooth grain, such as a circular one or any grain with a convex outline will have a PARIS factor of 1. The value increases as the grain boundary becomes irregular and lobate. Other parameters frequently used are the area, A, and the perimeter, P, of the grains, the circularity C (C =  $4 \pi A/P^2$ ; a value of 1.0 indicates a perfect circle), the solidity S (S = A/cA), where cA is the convex area) and the equivalent diameter EquiD (EquiD = 2  $(A/\pi)^{1/2}$ ).

#### 3. SAMPLE DESCRIPTION

The two samples examined (Tab. 1) are cataclasites from lowand high-angle normal faults on Kea (W-Cyclades, Greece). The fault system belongs to the West Cycladic Detachment System (Grasemann et al., 2011; Iglseder et al., 2011); this accommodated Miocene extension of the Aegean crust caused by the roll-back of the Hellenic subduction zone (for a regional geological description see the review of Jolivet and Brun 2010). The samples are from cataclastic faults that localized



**FIGURE 1:** Synoptic block diagram showing the interaction of highand low-angle normal faults (LANF) from the island Kea (Greece). The LANF has a top-to-the SW directed shear sense. Structural location of the two investigated samples HF1 and LF1.

Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)

sample	Sample number	GPS coordinates UTM 35	lithology	shear-strain
low-angle fault	LF1	4158684.6/261132.0	marble	high
high-angle fault	HF1	4156750.9/260721.7	marble	low
low-angle fault	LF2	4158684.6/261132.0	marble	high
clast in low-angle fault	CLF2	4158684.6/261132.0	marble	low

TABLE 1: Sample description and sample locations. "high" and "low" shear strains refers to decimeter thick fault rocks, which create an offset in the order of hundreds of meters and a few meters respectively.

in marble host rocks (Fig. 1).

The low-angle faults (analysed image LF1, LF2 and CLF2) represents a brittle segment of a ductile/brittle detachment fault that accommodated displacements in the order of tens of kilometers in total. Although the exact shear-strain taken up by the brittle deformation is unknown, data from other parts of the West Cycladic Detachment System, where a syn-tectonic granodiorite intrusion can be used as a strain marker, suggest that the brittle fault segments accommodated displacements

high-angle faults (analysed image HF1) are mechanically linked with the low-angle faults and accommodate extensional strain in the hanging wall of the detachments by bookshelf faulting. The shear strain in such faults is kinematically limited (e.g., Ramsay and Huber, 1987) and marker horizons suggest a few meters to tens of meters of offset. Since we have no better control on the shear strain values, we assume that the shear-strain in the samples from the low-angle fault was "high"

in the order of 1.5 km (Tschegg and Grasemann, 2009). The





(i.e. in the order of few 100 m to 1 km) and in the high-angle fault was "low" (i.e. in the order 1 m - 10 m). Figure 2 shows two thin-sections of the investigated samples, from which the analysed images were taken.

Low angle fault rocks (LF1, LF2 and CLF2): The ultra-cataclasites from the low-angle normal fault record a random distribution of particles with a size of less than 10 µm to up to 1 cm (Fig. 2a). The calcitic matrix has a dark brown colour containing angular to rounded fragments that consists of either aggregates of dolomite crystals, single dolomite and calcite crystals or dolomite cataclasites. Generally, the dolomite cataclasite fragments are more rounded than the dolomite aggregates, suggesting reworking of fault rocks within the catclasites by abrasive wear. The dolomite aggregates break either along the grain boundaries or along the cleavage of the dolomite crystals. Locally, a spaced pressure solution cleavage and micro-veins, filled with calcite, indicate dissolutionprecipitation processes. Some of the calcite veins record strong overprinting by deformation twinning and by frictional deformation, suggesting that the calcite fragments in the cataclasites are a reworked filling of the veins. In general, the microstructures suggest alternating periods of cataclastic flow dominated by fracturing, translation and grinding of particles (velocity weakening) and periods of dissolution precipitation creep (velocity hardening). LF1 and LF2 are from the ultracataclasites and CLF2 is a cataclastic component in LF2.

High angle fault (HF1): The investigated thin section records two different parts: (i) a protocataclastic zone (central and right part in Fig. 2b) and (ii) an ultracataclastic zone (left part in Fig. 2b). The protocataclasites are characterized by cm size dolomite host-rock components with extensive microfractures. The much finer grained ultracataclastic material with a dark brown calcite matrix and with a random fabric was injected into fractures and voids that developed between the dolomitic host rock particles, especially in the transition zone between the ultracataclasite and the protocataclasite. In the ultracataclasite zone, the largest components are less than 2 mm in size and are angular to partly rounded. The particles consist of dolomite single crystals and aggregates similar to the larger protocataclastic components. No counterparts of fractured particles have been observed. No cataclasite particles occur in the ultracataclasite and no evidence for dissolution precitpitation creep is observed. Our interpretation is that these fault rocks formed by fluidization processes; this would explain the random fabric, the injection structures and the missing counterpart particles (Monzawa and Otsuki, 2003). HF1 is from the ultracataclastic zone.

#### 4. DATA ACQUISITION

Electron microprobe backscattered electron (BSE) images were acquired on a Cameca SX-100 electron microprobe (Department of Lithospheric Research, University of Vienna). Methods and measurement conditions are described in Tschegg and Grasemann (2009). Representative parts of the BSE images of samples LF1, LF2, HF1 and CLF2 were analysed with the program ImageJ (rsbweb.nih.gov/ij). Due to the different grey shadings of the dolomite clasts and the calcitic matrix in the BSE images, the particles were automatically detected using a color coded filter (Fig. 3a).

All particles smaller than 19  $\mu$ m were then deleted because of resolution effects influencing further statistical analysis (Fig. 3b).

For each remaining particle, the following parameters were automatically calculated (length and area parameter are measured in µm and µm<sup>2</sup> respectively): Area *A*, Perimeter *P*, aspect ratio *AR*, circularity *C*, solidity *S* and equivalent diameter *EquiD*. In contrast to the PARIS factor of Panazzo and Hurlimann (1983), we calculated an elliptical PARIS factor *EP*, which is the difference between the perimeter of a particle and the perimeter of a fit-ellipse envelop  $P_{ell}$  of the particle  $(EP = (P-P_{ell})/P_{ell})$ .

For further statistical data analysis, we used the multivariate statistics software package SPSS (www.ibm.com/software/at/ analytics/spss/products/statistics). Before further processing, 5 classes of equivalent diameter were generated for samples LF1, LF2, HF1 and CLF2. In particular, the particles of the samples were divided in 5 classes of diameter with minimum and maximum diameter and number of particles (Tab. 2; see also box-plots Appendix A in the electronic supplement). Based on this, SPSS tables including all parameters derived by the automatic image analysis were compiled. A template of such an SPSS table is given in Appendix C in the electronic supplements.

# 5 METHOD AND IMPLEMENTATION OF DATA ANA-

As outlined above, the investigated samples clearly had a different tectonic history in terms of deformation mechanism and finite strain. Using a quantitative statistical approach, the following questions could be addressed:



FIGURE 3: a: Automatic detected particles from the BSE images using the program ImagJ. b: Selection of particles with a grain size > 19 µm and exclusion of particles, which are cut by the border of the image.

Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)

			Sample											
			LF1			HF1			LF2			CLF2		
			EquiD			EquiD		EquiD		EquiD				
		Minimum	Maximum	Count										
ter	1	19.09	24.54	40	19.21	25.22	34	19.59	24.84	35	19.04	25.07	80	
of imet	2	25.51	32.54	35	25.32	32.61	44	25.32	32.61	52	25.26	32.69	57	
ses ( t dia	3	32.76	42.73	41	32.76	42.73	52	33.06	42.56	44	32.84	42.81	52	
lass alen	4	43.02	65.26	46	43.91	65.60	58	43.07	66.33	46	43.26	65.34	38	
uiva C	5	66.92	590.75	45	66.81	663.82	56	66.48	298.30	44	66.66	652.72	43	
ed	Total	19.09	590.75	207	19.21	663.82	244	19.59	298.30	221	19.04	652.72	270	

TABLE 2: Classification of the equivalent diameter for particles > 19 µm in samples LF1, LF2, HF1 and CLF2.

- Is there a significant difference in the shape parameters (circularity, aspect ratio, solidity and elliptical PARIS factor) between samples of different tectonic history?
- Is there a significant difference between the five equivalent diameter classes in all specific questions above?
- Are there significant interactions between tectonic history and classes to equivalent diameter?

#### 5.1 THEORY OF THE VARIANCE ANALYSIS:

"...In investigations in which many factors are involved, it will not be economic or efficient to investigate the effect of one factor at a time on the particular result under investigation. Such a procedure gives no information about the possible interactions which may exist between relevant factors." (Huitson, 1966).

In this work, the variance analysis is applied using two factors, the factor tectonic (i.e. samples from the high- and lowangle normal fault deformed by different deformation mechanisms) and the factor grain-size. For the factor grain-size, five classes are used, based on which model equations can be formulated (Backhaus et al., 1996; Field, 2005). For example the equation for the dependent variable  $y_{abb}$ :

$$y_{ghk} = \mu + \alpha_g + \beta_h + (\alpha\beta)_{gh} + e_{ghk}$$
 Eq. 2

Where the index *g* represents the samples (1-4), *h* is the number of grain-size classes (1-5) and *k* represents the number of investigated particles in each section. This equation states that, for example, the circularity of a particle  $y_{ghk}$ , is caused by an overall mean value  $\mu$ , plus the effect of the  $g^{th}$  level of factor A (e.g. tectonics) plus the effect of the  $h^{th}$  level of factor B (e.g. grain-size), plus the  $gh^{th}$  interaction between factor A and factor B (e.g. tectonics with grain size). The interaction ( $\alpha\beta$ ) indicates an effect that acts in addition to the isolated main effects  $\alpha$  and  $\beta$  (see Bortz, 1999). The term  $e_{ghk}$  takes into account all those factors which have not been included in the study. The partition of one observation into different cases of influence can be assigned to the total variability of a set of data

$$SS_t = SS_A + SS_B + SS_{A \times B} + SS_W$$
 Eq. 3

where  $SS_A$  is the sum of squares of factor *A* (e.g. tectonics),  $SS_B$  is the sum of squares of factor *B* (e.g. grain-size),  $SS_{AxB}$  is the sum of squares cause by the interaction of factor *A* and *B* and  $SS_w$  is the residual sum of squares.

To find the total amount of variation within the data, we calculate the difference between each observed data point  $y_{ghk}$ and the grand mean  $\bar{Y}$ . The squared differences were added resulting in the total sum of squares (SS<sub>t</sub>):

$$SS_t = \sum_{g=1}^G \sum_{h=1}^H \sum_{k=1}^K (y_{ghk} - \bar{Y})^2$$
 Eq. 4

where G is the number of levels of factor A, H is the number of levels (classes) of factor B and K is the number of observations combining factor A and B.

The model sum of squares  $SS_M$  unites three components:

$$SS_M = SS_A + SS_B + SS_{A \times B}$$
 Eq. 5

 $SS_M$  is calculated from the data with the difference between each group mean and the overall mean.

$$SS_M = K \sum_{g=1}^G \sum_{h=1}^H \left( \bar{y}_{gh} - \bar{Y} \right)^2 \qquad \qquad \text{Eq. 6}$$

 $SS_A$  represents the variation of the data controlled by the contrasting tectonic history.

$$SS_A = H.K.\sum_{g=1}^{G} \left(\bar{y}_g - \bar{Y}\right)^2$$
 Eq. 7

 $SS_{B}$  describes the variation in data controlled by the different grain sizes.

$$SS_B = G.K.\sum_{h=1}^{H} (\bar{y}_h - \bar{Y})^2$$
 Eq. 8

Finally, the interaction effect  $SS_{AxB}$  represents the variance controlled by interaction of the two variables (tectonic history and grain size). This can be calculated very easily, because  $SS_{M} = SS_{A} + SS_{B} + SS_{AxB}$  (see above).

$$SS_{A \times B} = SS_M - SS_A - SS_B$$
 Eq. 9

The residual sum of squares  $SS_w$  represents the effects of

Osterreichische Geologische Gesellschaft/Austria; download unter www.geol-ges.at/ und www.biologiezentrum.at

Norbert KOHLMAYER & Bernhard GRASEMANN

variations, which cannot be explained by the model.

$$SS_W = SS_t - SS_M$$
 Eq. 10

To find the average amount of variation explained by the model,  $SS_A$ ,  $SS_B$ ,  $SS_{AxB}$  and  $SS_w$  must be divided by the degrees of freedom (Backhaus et al., 1996). With these values, the Fratio can then be calculated (Field, 2005).

The F-ratio is a measure of the ratio of the variation explained by the model and the variation explained by unsystematic factors. If this value is less than 1 then it represents a nonsignificant effect (i.e. more unsystematic than systematic variance). If the F-value is greater than 1 it means that the systematic effect is greater than the unsystematic effect, but it is not constrained whether this is a chance result. In order to test this, the F-value is compared with the maximum random value in an F-distribution with the same degrees of freedom.

#### 5.2 DATA PROCESSING WITH SPSS

The univariate analysis of variance was calculated with the software SPSS; this firstly gathers descriptive statistical information from the sample data and secondly tests the validity of the calculation and the precondition (for the assumptions and a SPSS output and example datafile see Appendices B, C and D in the electronic supplement). The full output file of the SPSS calculations is provided in the electronic supplements. In this, the mean results for every dependent variable are separately shown in several tables as well as graphically. Table 3 shows an example for the Test of *Between-Subjects Effects* for the circularity. The first and the last columns are

important in this table. Columns 2 to 5 show the sum of squares, the degree of freedom *df*, the mean square and the F-value. The last column *Sig* shows the probability that the observed results come about only under the condition chance. Generally, probabilities smaller than 0.05 are considered to be significant; that is, the observed results have only a 5% probability of being random.

#### 6. RESULTS OF THE ANALY-SIS OF VARIANCE

#### 6.1 CIRCULARITY, SAMPLE AND EQUIVALENT DIAMETER

Table 3 shows in the column "source" and "Sig" that for Model, Sample and EquiDclasses, the probability for a random result is smaller than 0.000. The first row with the entry Model indicates that the calculation with the used dependent and independent variables throughout is significant with 0.000. The subsequent rows, with the entry Sample and EquiDclasses, mean that the possibility for the differences in the circularity between the factor levels of the samples and the factor levels of the diameter classes under the condition chance are also very low. In contrast, the interaction "Sample \* EquiDclasses" is under the condition chance possible in 40% percent of the cases.

The effect size of the model is the ratio of model sum of square (Corrected Model) to total sum of square (Corrected Total). It is 40.1% for the variable Circularity. That means 40% of the total sum of square (Corrected Total) is explained by the model. This percentage is divided equally between the sample and grain size. The 60% of the variance that is not explained was caused by factors which were not considered here (Backhaus et al., 1996).

The subsequent calculated *Post Hoc Test* (Table 7) highlights those samples that differ significantly from the others. In this case, the samples are grouped by their circularity. It can be seen that samples HF1 and CLF2 belong to the same group because the circularities of these samples are very similar. However, all the other samples are classified in separate groups.

The grouping of the samples in terms of the measured circularity can be visualized in a plot of the estimated means of the circularity for each sample versus the class of equivalent diameter (Fig. 4). This clearly shows the close relationship between samples HF1 and CLF2. In contrast, samples LF1 and LF2 represent separate classes with generally higher circularities. However, the plot also demonstrates that for all samples there is a clear trend of the circularity to decrease

Tests of Between-Subjects Effects										
Dependent Variable:Circ.										
Source	Type III Sum of Squares <sup>1</sup>	df	Mean Square	F	Sig.					
Corrected Model	11.145ª	19	.587	32.533	.000					
Intercept <sup>2</sup>	335.915	1	335.915	18630.158	.000					
Sample	6.034	3	2.011	111.545	.000					
EquiDclasses	5.026	4	1.257	69.691	.000					
Sample * EquiDclasses	.226	12	.019	1.045	.405					
Error	16.624	922	.018							
Total	373.184	942								
Corrected Total	27.770	941								

a. R Squared = .401 (Adjusted R Squared = .389)

TABLE 3: Test of Between-Subjects Effects for the dependent variable circularity

<sup>2</sup> The value of the intercept is for the interpretation not important, reason for this entry is that SPSS performs the calculation with a regressions analysis.

<sup>&</sup>lt;sup>1</sup> The column "Type III Sum of Squares" means that the sum of squares were adapted to the fact that not all cells of the model are equal full. In the 4x5 cells (4 samples and 5 Equidclasses) the amount of components are not equal.

Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)



FIGURE 4: Diagram of the circularity versus the classes of equivalent diameters for the four investigated sample.

with higher equivalent diameter classes.

# 6.2. ASPECT RATIO, SAMPLE AND EQUIVALENT DIAMETER

Table 4 shows that with respect to the dependent variable aspect ratio, only the samples record significant differences but the classes of diameter do not differ significantly. The interaction between the two factors sample and equivalent diameter classes with a value of 0.059 is also not significant. The post hoc test (Tab. 7) demonstrates that the significance of the sample value in Table 4 is dominated by the aspect ratio value of LF2. The value of this sample is the lowest in all classes of equivalent diameters and is therefore classified into a single group. The three other samples form a second group.

Tests of Between-Subjects Effects											
Dependent Variable:AR	Dependent Variable:AR										
Source	Type III Sum of Squares	df	Mean Square	F	Sig.						
Corrected Model	19.243ª	19	1.013	2.792	.000						
Intercept	2825.163	1	2825.163	7786.803	.000						
Sample	10.076	3	3.359	9.257	.000						
EquiDclasses	1.852	4	.463	1.276	.278						
Sample * EquiDclasses	7.469	12	.622	1.716	.059						
Error	334.515	922	.363								
Total	3342.419	942									
Corrected Total	353.758	941									

a. R Squared = .054 (Adjusted R Squared = .035)

TABLE 4: Test of Between-Subjects Effects for the dependent variable aspect ratio AR.

The plot of the aspect ratio versus the classes of equivalent diameters for the four investigated samples (Fig. 5) shows that LF2 differs from the other three samples, which record higher aspect ratios. Whereas HF has higher aspect ratios of the particles in the coarser classes of equivalent diameters, LF1 and CLF2 have unsystematically scattered aspect ratios. Therefore, the effect size for the variance of the aspect ratio is only 5% of the total variance.

#### 6.3 SOLIDITY, SAMPLE AND EQUIVALENT DIA-METER

The parameter solidity (the ratio of area to convex area), shows significance only dependent on the sample factor (Tab. 5), but as highlighted by the post hoc test, there are great differences between the samples (Tab. 7). In fact, the results of the post hoc test suggest that all samples strongly differ from each other and that all the samples have to be classified into separate groups. This can be visualized in a plot of the solidity versus the classes of equivalent diameters for the four samples (Fig. 6). The solidity of the particles in all samples records a minor variability and unsystematically scatters in the different classes of equivalent diameter. However, the four samples can be clearly distinguished using the solidity of their particles, with particles from LF2 recording the highest solidity.

The model effect for the dependent variable solidity is 19% which is only influenced by the factor sample. The factor grain size effects only 0.8% of the variance.

#### 6.4 ELLIPTICAL PARISFACTOR, SAMPLE AND EQUI-VALENT DIAMETER

The statistical test Between-Subject Effects clearly demonstrates that all factors as well as the interaction between the factors are significant (Tab. 6). Similarly, the post hoc test (Tab. 7) allows a clear grouping of the samples where HF1

> and CLF2 record the closest relationships (i.e. they belong to the same group). All other samples are classified in individual groups.

> The plot in Figure 7 shows the elliptical PARIS factor versus the classes of equivalent diameters. It can be seen that in all four investigated samples the elliptical PARIS factor of the particles increases with the classes of equivalent diameter. The larger the grain size, the more irregular the particle boundaries become. CLF2 and HF1 form the group with the greater values in the elliptical PARIS Factor and have more irregular particle boundaries compared to smoother particles of LF1 and Lf2.

> The model effect for the Elliptical PARIS Factor is 36%, which is influenced by sample and grain size

Norbert KOHLMAYER & Bernhard GRASEMANN

in approximately equal amounts.

#### 7. DISCUSSION

The particle size distribution in cataclasites can be quantified using the D-value (Eq. 1), which provides information about the frequency of particle sizes for a particular particle size range. Many processes that lead to the formation of cataclastic fault rocks, such as mechanical fragmentation, microcracking, sliding, grinding or spalling of the fragments, result in different particle size distributions, which are further complicated by mineralogical phases, chemomechanical feedback processes and the finite strain (e.g. Blenkinsop and Rutter, 1986; Sammis et al., 1986; Marone and Scholz, 1989; An and Sammis, 1994; Monzawa and Otsuki, 2003; Rawling and Goodwin, 2003; Billi 2005; Keulen et al., 2007; Stünitz et al., 2010). However, in most studies, the dominant process of cataclasis under lithostatic compaction or shear deformation has been observed to be transgranular fracturing initiated by loading at grain-to-grain contacts abrasion with additional flaking or spalling from grain edges. Therefore, cataclastic fault rocks generated in the laboratory, preferentially record D-values at about 2.5-2.6 (Sammis et al., 1986; Biegel et al., 1989; Marone and Scholz, 1989; Billi and Storti, 2004) suggesting scale and time invariance of the fragmentation processes. However, other studies present evidences indicating that D-values systematically increase from immature to mature cataclastic rocks (Marone and Scholz, 1989; Hattori and Yamamoto, 1999; Storti et al., 2003; Billi and Storti, 2004). Relating the maturity of cataclasites with finite strain in the brittle fault zone, the observed systematic variation in D-values suggests that the dominant particle fragmentation mechanism changes and the contribution of chipping and surface abrasion increases with progressing particle interaction (Hattori and Yamamoto, 1999; Storti et al., 2003).

The PSD of our samples (Fig. 8) records D-values that are in good agreement with other published Dvalues for natural and experimental fault rocks (e.g., Heilbronner and Keulen, 2006). Although we are aware of the problems of this technique because of the sensitivity to cut-off effects at large sizes, there is little variation in the D-values between the investigated samples (LF1: 1.7; LF2: 1.8; CLF2: 1.7; HF1: 1.5, see Fig. 8), in agreement with the proposed scale and time invariance of the fragmentation processes. A detailed discussion about the D-value and its validity is given in Hergarten (2002). A number of other studies focused on the shape parameters of particles, to unravel processes during catclastic deformation. It has



FIGURE 5: Diagram of the aspect ratio versus the classes of equivalent diameters for the four investigated sample.

been shown, for example, that a microstructural analysis of the fragment shapes shows a dependency on the amount of slip and may record some information on slip magnitude. With increasing displacement, grain shapes evolve towards more rounded and less serrated grains, while the grain size distribution remains constant (Mair and Marone, 1999; Heilbronner and Keulen, 2006; Storti et al., 2007; Stünitz et al., 2010).

However, in most of these studies the particle shape parameters were mainly evaluated individually or compared with the statistic averages of each individual sample. A quantitative interpretation of the shape values is difficult since without statistical tests it is impossible to establish which trends are significant and which can be regarded as coincidental.

Tests of Between-Subjects Effects										
Dependent Variable:Solidity										
Source	Type III Sum of Squares	df	Mean Square	F	Sig.					
Corrected Model	1.013ª	19	.053	11.670	.000					
Intercept	637.609	1	637.609	139572.707	.000					
Sample	.928	3	.309	67.735	.000					
EquiDclasses	.042	4	.011	2.302	.057					
Sample * EquiDclasses	.048	12	.004	.874	.573					
Error	4.212	922	.005							
Total	666.055	942								
Corrected Total	5.225	941								

a. R Squared = .194 (Adjusted R Squared = .177)

TABLE 5: Test of Between-Subjects Effects for the dependent variable solidity.

Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)



**FIGURE 6:** Diagram of the solidity versus the classes of equivalent diameters for the four investigated sample.

The presented analysis of variance provides more robust information than the simple statistical analysis of data series. The main advantage of the analysis of variance presented here is that, firstly, the indicators can be tested for random influences, and, secondly, multiple parameters (e.g. measured shape factors) enter in the analysis simultaneously and, therefore, mutual influences and known interactions can be detected. The statistical methods used here investigate the sample and the effect of grain-size in relation to the particle shape parameters and therefore it can be quantified whether or not a particular sample in a given particle size shows random or significant trends in relation to grain-shape parameters.

The analysis of variance clearly demonstrated significant differences between the distribution of some particle sizes in re-

Tests of Between-Subjects Effects										
Dependent Variable:Elliptic	Dependent Variable:Elliptical PARIS Factor									
Source	Type III Sum of Squares	df	Mean Square	F	Sig.					
Corrected Model	12.984ª	19	.683	27.613	.000					
Intercept	54.632	1	54.632	2207.560	.000					
Sample	5.815	3	1.938	78.325	.000					
EquiDclasses	6.285	4	1.571	63.491	.000					
Sample * EquiDclasses	1.108	12	.092	3.732	.000					
Error	22.818	922	.025							
Total	93.445	942								
Corrected Total	35.801	941								

a. R Squared = .363 (Adjusted R Squared = .350)

TABLE 6: Test of Between-Subjects Effects for the dependent variable elliptical PARIS factor.



FIGURE 7: Diagram of the elliptical PARIS Factor versus the classes of equivalent diameters for the four investigated sample.

lation to the various shape parameters, in the investigated samples and between the samples. The statistical analysis presented clearly provides additional information that can help to quantitatively discriminate between similar and dissimilar cataclastic fault rocks. Further, it can also provide information about the deformation mechanisms that controlled the formation of the cataclasites.

In the following, we present a short and cautious interpretation of the PSD in combination with the statistically investigated shape parameters of the studied samples:

The circularity and elliptical PARIS factor are well suited for distinguishing between the different samples and show a clear trend in the investigated grain-sizes. In all samples, there is a decrease in circularity with increasing grain-size classes. Simi-

> larly, there is an increase in the elliptical PARIS factor with increasing grain-sizes. HF1 and CLF2 show very similar trends and record less circular particles (and higher elliptical Paris factor) than the samples LF1 and LF2; this probably reflects the higher maturity (higher finite strain) of the samples from the low-angle faults. In the early stages of brittle fault development, particle fragmentation is the dominant mechanism resulting in coarse and angular particles. With progressing deformation, particle interaction by rolling, sliding and rotation favours surface abrasion and chipping (i.e. grinding) that eventually become the dominant deformation mechanisms resulting in more rounded grains (Biegel et al., 1989; Blenkinsop, 1991; Hattori and

Yamamoto, 1999). Additionally, the efficiency of particle cracking decreases with decreasing grain-size and therefore smaller particles get more rounded by continuous abrasion (Blenkinsop, 1991). Rounded particles accommodate shear mainly by rolling resulting in fault weakening, i.e. the progressive decrease in friction with increasing fault displacement (Mair et al., 2002; Guo and Morgan, 2004; Anthony and Marone, 2005). Interestingly, Storti et al. (2007) presented data from carbonate cataclasites that show that particle angularity systematically decreases with increasing particle size and with increasing fractal dimension. The evidence that smaller particles are more angular than larger ones was interpreted to indicate a deformation process where particle fragmentation dominates in the early evolutionary stages of the fault, while comminution and spalling control cataclasis at higher

of fault displacements.

Generally, fracturing of grains produces higher aspect ratios, which is further controlled by the mineralogical composition (Heilbronner and Keulen, 2006), although the shape anisotropy of the particles may be also an inherited feature, associated with the orthorhombic symmetry of joint patterns, which enhances fracturing of lithons perpendicular to their long symmetry axes (e.g. Engelder, 1987; Ramsay and Lisle, 2000; Billi et al., 2004). The aspect ratio of particles in cataclasites is significantly influenced (increased) by dissolution-precipitation processes (Babaie et al., 1991; Stünitz et al., 2010; Rowe et al., 2011). In the investigated samples, the aspect ratio shows no systematic trends and is independent on the classes of equivalent diameter.

The solidity, which has rarely been used as a shape parameter in quantifying particle structures in cataclasites, gives information about the roughness or smoothness of particle edges. The solidity in the investigated particle shapes clearly separates the samples into four different groups. The order of separation is the same as that derived by the circularity and elliptical PARIS factor, highlighting the relationship between roundness and smoothness of particle borders. In contrast to the circularity and the elliptical PARIS factor, the solidity is nearly insensitive to the classes of equivalent diameters. To our knowledge, the present study is the first application of the well established statistical method of variance analysis to quantitatively investigate the grain-size and shape parameters of cataclasites. To test this method, we selected samples from marble cataclasites, where only the influence of the grainsize on the particle shape parameters and their interactions were investigated. The method can be easily extended in future to fault rocks that consist of different mineralogical particles, which may have a significant influence of grain-size and particle shape parameters.

#### 8. CONCLUSIONS

1) The analysis of variance is a powerful statistical method that can be applied to quantitatively investigate the grain-



FIGURE 8: Log-Log diagram of particle size versus number of particles. The absolute value of the slope of the regression line corresponds to the D-value. a) LF1; b) HF1; c) LF2; d) CLF2

Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)

Resuts: Post Hoc Tests		SAM	PLE	
	CLF2	HF1	LF1	LF2
Circularity 1	0.55	0.54		
Circularity 2			0.61	
Circularity 3				0.74
AR 1				1.61
AR 2	1.89	1.81	1.77	
Solitity 1	0.79			
Solidity 2		0.82		
Solidity 3			0.85	
Solidity 4				0.88
EII. PARIS-f 3	0.31	0.30		
Ell. PARIS-f 2			0.23	
EII. PARIS-f 1				0.13

**TABLE 7:** Post Hoc Tests. The test after the analysis of variance in order to group the samples according to their values of circularity, aspect ratio (AR), solidity and elliptical PARIS factor (Ell. PARIS-f).

size and particle shape parameters of cataclasites.

- 2) Four test samples from low- and high-angle faults in the Western Cyclades representing fault rocks that were formed by different deformation mechanisms and accommodated different finite strains, record partly very different particle shape parameters but similar particle-size distributions.
- 3) In the test samples, circularity and the elliptical PARIS factor show systematic trends in the classes of equivalent diameters. The finer grained particles are more rounded suggesting that abrasion and comminution was the dominant deformation mechanism in comparison to the more angular, coarser-grained particles, where fracturing prevailed.
- 4) Solidity, which is similar to circularity and the elliptical PA-RIS factor, which is also a function of the roundness of the particles, shows no dependence on the classes of equivalent diameters, emphasizing the need to analyzing a broad variation of particle shape parameters.
- 5) The present study identified the circularity, the elliptical PARIS factor and the solidity as useful shape parameters for discriminating between the different samples.

#### ACKNOWLEDGEMENTS

We thank Christoph Iglseder for providing the investigated samples from Kea including the thin sections and the BSE images and for discussions about the geology of Kea and the Aegean region. Two reviews by S. Hergarten and W. Kurz and handling of the manuscript by W. Kurz, Associate Editor of the Austrian Journal of Earth Sciences, are gratefully acknowledged. Hugh Rice significantly improved the language and style of the manuscript.

#### REFERENCES

An, L.-J., Sammis, C.G., 1994. Particle size distribution of cataclastic fault materials from Southern California: A 3-D study. Pure and Applied Geophysics, 143, 203-227.

Anthony, J.L. and Marone, C., 2005. Influence of particle characteristics on granular friction. Journal of Geophysical Research, 110, B08409.

Arbiter, N. and Harris, C.C., 1965. Particle size distribution time relationships in comminution. Britisch Chemical Engineering, 10, 240-247.

Babaie, H.A., Babaei, A. and Hadizadeh, J., 1991. Initiation of cataclastic flow and development of cataclastic foliation in non-porous quartzites from a natural fault zone. Tectonophysics, 200, 67-77.

Backhaus, K., Erichson, B., Plinke, W. and Weiber, R., 1996. Multivariate Analysemethoden. Springer-Verlag, Berlin Heidelberg New York, 591 pp.

Billi, A. and Storti, F., 2004. Fractal distribution of particle size in carbonate cataclastic rocks from the core of a regional strike-slip fault zone. Tectonophysics, 384, 115-128.

Billi, A., 2005. Grain size distribution and thickness of breccia and gouge zones from thin (<1 m) strike-slip fault cores in limestone. Journal of Structural Geology, 27, 1823-1837.

Biegel, R.L., Sammis, C.G. and Dieterich, J.H., 1989. The frictional properties of a simulated gouge having a fractal particle distribution. Journal of Structural Geology, 11, 827-846.

Bjork, T.E., Mair, K. and Austrheim, H., 2009. Quantifying granular material and deformation: Advantages of combining grain size, shape, and mineral phase recognition analysis. Journal of Structural Geology, 31, 637-653.

Blenkinsop, T.G. and Rutter, E.H., 1986. Cataclastic deformation of quartzite in the moine thrust zone. Journal of Structural Geology, 8, 669-681.

Blenkinsop, T., 1991. Cataclasis and processes of particle size reduction. Pure and Applied Geophysics, 136, 59-86.

Blenkinsop, T., 2000. Deformation Microstructures and Mechanisms in Minerals and Rocks. Kluwer Academic Publishers, Dordrecht Boston London, 150 pp.

Bortz, J., 1999. Statistik für Sozialwissenschaftler. Springer-Verlag, Berlin, Heidelberg, New York, 836 pp.

Norbert KOHLMAYER & Bernhard GRASEMANN

Cladouhos, T.T., 1999. Shape preferred orientation of survivor grains in fault gouge. Journal of Structural Geology, 21, 419-436.

Dowdy, S. and Wearden, S., 1991. Statistics for Research. John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore, 629 pp.

Engelder, J.T., 1974. Cataclasis and the Generation of Fault Gouge. GSA Bulletin, 85, 1515-1522.

Engelder, T., 1987. Joints and some fractures in rocks. In: Atkinson, B.K. (Ed.), Fracture Mechanics of Rock. Academic Press, pp. 27-69.

Field, A., 2005. Discovering Statistics Using SPSS. SAGE Publicatons, London Thousand Oaks New Delhi, 779 pp.

Grady, D.E. and Kipp, M.E., 1987. Dynamic rock fragmentation. Academic Press Inc., London, 1987, 420-475.

Grasemann, B., Schneider, D.A., Stöckli, D.F. and Iglseder, C., 2011. Miocene bivergent crustal extension in the Aegean: Evidence from the western Cyclades (Greece). Lithosphere, 4, doi:10.1130/L164.1.

Guo, Y., and J. K. Morgan, 2004. Influence of normal stress and grain shape on granular friction: Results of discrete element simulations. Journal of Geophysical Research, 109, doi: 10.1029/2004JB003044.

Hartman, W. K., 1969. Terrestrial, lunar and interplanetary rock fragmentation. Icarus, 10, 201-213.

Hausegger, S., Kurz, W., Rabitsch, R., Kiechel E. and Brosch F.-J., 2010. Analysis of the internal structure of a carbonate damage zone: Implications for the mechanisms of fault breccia formation and fluid flow. Journal of Structural Geology, 32, 1349-1362.

Hattori, I. and Yamamoto, H., 1999. Rock Fragmentation and Particle Size in Crushed Zones by Faulting. The Journal of Geology, 107, 209-222.

Heilbronner, R. and Keulen, N., 2006. Grain size and grain shape analysis of fault rocks. Tectonophysics, 427, 199-216.

Hergarten, S., 2002. Self-Organized Criticality in Earth Systems. Springer-Verlag, Berlin, Heidelberg, New Yourk, 272 pp.

Huitson, A., 1966. The Analysis of Variance, Number eighteen of Griffin's Statistical Monographs & Courses. Charles Griffin & Company Limited, London, 83 pp.

Iglseder, C., Grasemann, B., Rice, A.H.N., Petrakakis, K. and Schneider, D.A., 2011. Miocene south directed low-angle normal fault evolution on Kea Island (West Cycladic Detachment System, Greece). Tectonics, 30, 10.1029/2010tc002802. Jolivet, L. and Brun, J.-P., 2010. Cenozoic geodynamic evolution of the Aegean. International Journal of Earth Sciences, 99, 109-138.

Keulen, N., Heilbronner, R., Stunitz, H., Boullier, A.-M. and Ito, H., 2007. Grain size distributions of fault rocks: A comparison between experimentally and naturally deformed granitoids. Journal of Structural Geology, 29, 1282-1300.

Mair, K., Marone, C., 1999. Friction of simulated fault gouge for a wide range of velocities and normal stresses. Journal of Geophysical Research, 104, 28899-28914.

Mair, K., Frye, K.M. and Marone, C., 2002. Influence of grain characteristics on the friction of granular shear zones. Journal of Geophysical Research, 107, 10.1029/2001jb000516.

Marone, C. and Scholz, C.H., 1989. Particle-size distribution and microstructures within simulated fault gouge. Journal of Structural Geology, 11, 799-814.

Monzawa, N. and Otsuki, K., 2003. Comminution and fluidization of granular fault materials: implications for fault slip behavior. Tectonophysics, 367, 127-143.

Panazzo, R. and Hurlimann, H., 1983. A simple method for the quantitative discrimination of convex-concave lines. Microscopica Acta 87, 169-176.

Passchier, C.W. and Trouw, R.A.J., 1996. Microtectonics. Springer-Verlag, Berlin, Heidelberg, New York, 289 pp.

Ramsay, J.G. and Huber, I.M., 1987. The Techniques of Modern Structural Geology. Volume 2: Folds and Fractures. Academic Press Inc. Ltd, London, 391 pp.

Ramsay, J.G. and Lisle, R.J., 2000. The Techniques of Modern Structural Geology. Volume 3: Applications of Continuum Mechanics in Structural Geology. Academic Press Inc. Ltd, London, 360 pp.

Rawling, G.C. and Goodwin, L.B., 2003. Cataclasis and particulate flow in faulted, poorly lithified sediments. Journal of Structural Geology, 25, 317-331.

Rowe, C.D., Meneghini, F. and Moore, J.C., 2011. Textural record of the seismic cycle: strain-rate variation in an ancient subduction thrust. Geological Society, London, Special Publications, 359, 77-95.

Sammis, C.G., Osborne, R.H., Anderson, J.L., Badert, M. and White, P., 1986. Self-similar Cataclasis in the Formation of Fault Gouge. Pure and Applied Geophysics, 124, 53-78.

Storti, F., Billi, A. and Salvini, F., 2003. Particle size distribution in natural carbonate fault rocks; insights for non-self-similar cataclasis. Earth and Planetary Science Letters, 206 (1-2), 173-186. Quantitative characterisation of cataclasites using a statistical approach (analysis of variance)

Storti, F., Balsamo, F. and Salvini, F., 2007. Particle shape evolution in natural carbonate granular wear material. Terra Nova, 19, 344-352.

Stünitz, H., Keulen, N., Hirose, T. and Heilbronner, R., 2010. Grain size distribution and microstructures of experimentally sheared granitoid gouge at coseismic slip rates - Criteria to distinguish seismic and aseismic faults? Journal of Structural Geology, 32, 59-69.

Tschegg, C. and Grasemann, B., 2009. Deformation and alteration of a granodiorite during low-angle normal faulting (Serifos, Greece). Lithosphere, 1, 139-154.

Wang, W. and Scholz, C.H., 1994. Wear processes during frictional sliding of a rock: A theoretical and experimental study. Journal of Geophysical Research, 99, 6789-6799.

> Received: 17 January 2012 Accepted: 29 March 2012

#### Norbert KOHLMAYER<sup>9</sup> & Bernhard GRASEMANN

Department of Geodynamics and Sedimentology, University of Vienna, Althanstraße 14, Vienna, Austria;

\*) Corresponding author, norbert.kohlmayer@aon.at

Electronic supplement: APPENDIX A: Boxplots of the used Data.

The statistical outlier are marked with the number of the case.





### Electronic supplement: APPENDIX B:

Assumptions of Analysis of Variance (see Bortz, 1999).

Regarding the variance decomposition no prerequisites are necessary. Assuptions for the validity of significance tests are normal distribution, independence of the data and homogeneity of variances. In the present work, the independence of the data is given by different samples. The normal distribution assumption is reviewed by Kolmogorov-Smirnov Test but the homogeneity of variances is not given in the data. According to Bortz inhomogeneous variances have a negative effect only for small samples and unequal cell occupation on the F-test. Buehl and Zöffel (1999) suggest in the case of inhomogeneous variances to increase the significance level of 0.05 to 0.01. In this work the results are significant at 0.000 level and therefore confident. To further confirm the results, a rank analysis of variance with low requirements (Kruskal-Wallis test) was performed, these tests affirm the results of the main effects, but the interactions could not be tested with this method.

Electronic supplement: APPENDIX C: Outputfile of the SPSS program.

### NPar Tests Sample = LF1, Classes of equivalent diameter = 1

One-Sample Kolmogorov-Smirnov Test <sup>a</sup>							
		Circ.	AR	Solidity	Elliptical PARIS		
					Factor		
Ν		40	40	40	40		
Normal Parameters <sup>b,c</sup>	Mean	.71580	1.87765	.85180	.1106		
Normal Farameters	Std. Deviation	.136087	.621977	.061558	.07950		
	Absolute	.111	.141	.252	.163		
Most Extreme Differences	Positive	.111	.141	.106	.163		
	Negative	111	117	252	111		
Kolmogorov-Smirnov Z		.704	.891	1.595	1.030		
Asymp. Sig. (2-tailed)		.705	.406	.012	.239		

a. Sample = LF1, Classes of equivalent diameter = 1

b. Test distribution is Normal.

c. Calculated from data.

### Sample = LF1, Classes of equivalent diameter = 2

		Circ.	AR	Solidity	Elliptical PARIS		
					Factor		
Ν		35	35	35	35		
Normal Davage stars <sup>b,C</sup>	Mean	.65091	1.60394	.84589	.2084		
Normal Parameters	Std. Deviation	.145648	.459964	.058788	.14328		
	Absolute	.119	.136	.186	.193		
Most Extreme Differences	Positive	.097	.136	.125	.193		
	Negative	119	126	186	128		
Kolmogorov-Smirnov Z		.706	.805	1.103	1.142		
Asymp. Sig. (2-tailed)		.701	.535	.175	.147		

#### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = LF1, Classes of equivalent diameter = 2

b. Test distribution is Normal.

Sample = LF1	, Classes of	equivalent	diameter = 3
--------------	--------------	------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS Factor
Ν		41	41	41	41
Normal Daramatara <sup>b,c</sup>	Mean	.63073	1.77085	.85351	.1980
Normal Parameters	Std. Deviation	.121613	.592804	.056827	.11644
	Absolute	.157	.150	.190	.159
Most Extreme Differences	Positive	.079	.150	.121	.159
	Negative	157	119	190	117
Kolmogorov-Smirnov Z		1.007	.958	1.216	1.019
Asymp. Sig. (2-tailed)		.262	.317	.104	.250

a. Sample = LF1, Classes of equivalent diameter = 3

b. Test distribution is Normal.

c. Calculated from data.

### Sample = LF1, Classes of equivalent diameter = 4

		Circ.	AR	Solidity	Elliptical PARIS	
					Factor	
Ν		46	46	46	46	
Name al Danama (anab.C	Mean	.56650	1.78063	.83613	.2769	
Normal Parameters	Std. Deviation	.145394	.693153	.067450	.16240	
	Absolute	.078	.189	.167	.153	
Most Extreme Differences	Positive	.072	.189	.078	.153	
	Negative	078	151	167	104	
Kolmogorov-Smirnov Z		.526	1.284	1.134	1.041	
Asymp. Sig. (2-tailed)		.945	.074	.153	.229	

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = LF1, Classes of equivalent diameter = 4

b. Test distribution is Normal.

Sample = LF1;	<b>Classes of</b>	equivalent	diameter = 5
---------------	-------------------	------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS Factor
Ν		45	45	45	45
Normal Daramatara <sup>b,c</sup>	Mean	.51013	1.83349	.85256	.3316
Normal Parameters	Std. Deviation	.122487	.623240	.053095	.15429
	Absolute	.081	.133	.103	.116
Most Extreme Differences	Positive	.081	.133	.070	.116
	Negative	076	099	103	106
Kolmogorov-Smirnov Z		.543	.895	.691	.777
Asymp. Sig. (2-tailed)		.930	.399	.726	.582

a. Sample = LF1, Classes of equivalent diameter = 5

b. Test distribution is Normal.

c. Calculated from data.

### Sample = HF1, Classes of equivalent diameter = 1

		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		34	34	34	34
Normal Davage stars <sup>b,C</sup>	Mean	.64159	1.66065	.82841	.2031
Normal Parameters	Std. Deviation	.129304	.432699	.053161	.12224
	Absolute	.145	.114	.158	.186
Most Extreme Differences	Positive	.089	.114	.069	.186
	Negative	145	114	158	090
Kolmogorov-Smirnov Z		.848	.666	.919	1.086
Asymp. Sig. (2-tailed)		.468	.766	.367	.189

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = HF1, Classes of equivalent diameter = 1

b. Test distribution is Normal.

Sample = HF1	, Classes of	equivalent	diameter :	= 2
--------------	--------------	------------	------------	-----

		- <b>J</b>			
		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		44	44	44	44
Normal Parameters <sup>b,c</sup>	Mean	.59893	1.73745	.82839	.2400
Normal Parameters	Std. Deviation	.131730	.520849	.060409	.13144
	Absolute	.104	.150	.146	.115
Most Extreme Differences	Positive	.087	.150	.080	.115
	Negative	104	104	146	111
Kolmogorov-Smirnov Z		.688	.995	.970	.762
Asymp. Sig. (2-tailed)		.731	.276	.303	.607

a. Sample = HF1, Classes of equivalent diameter = 2

b. Test distribution is Normal.

c. Calculated from data.

### Sample = HF1, Classes of equivalent diameter = 3

One-Sample Kolmogorov-Siminov Test					
		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		52	52	52	52
Normal Davamatara <sup>b,c</sup>	Mean	.55979	1.80952	.81965	.2782
Normal Parameters	Std. Deviation	.131435	.690511	.066184	.13567
	Absolute	.073	.195	.154	.108
Most Extreme Differences	Positive	.057	.195	.124	.108
	Negative	073	170	154	065
Kolmogorov-Smirnov Z		.525	1.409	1.107	.776
Asymp. Sig. (2-tailed)		.946	.038	.172	.584

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = HF1, Classes of equivalent diameter = 3

b. Test distribution is Normal.

Sample = HF1	, Classes of	equivalent	diameter = 4
--------------	--------------	------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		58	58	58	58
Normal Parameters <sup>b,c</sup>	Mean	.49843	1.88648	.81514	.3504
	Std. Deviation	.135766	.699767	.071742	.17787
	Absolute	.130	.165	.141	.161
Most Extreme Differences	Positive	.130	.165	.080	.161
	Negative	100	109	141	098
Kolmogorov-Smirnov Z		.993	1.260	1.072	1.229
Asymp. Sig. (2-tailed)		.277	.084	.201	.097

a. Sample = HF1, Classes of equivalent diameter = 4

b. Test distribution is Normal.

c. Calculated from data.

### Sample = HF1, Classes of equivalent diameter = 5

		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		56	56	56	56
Normal Davage stars <sup>b,C</sup>	Mean	.47798	1.87584	.84964	.3809
Normal Parameters	Std. Deviation	.128257	.935205	.062529	.16175
	Absolute	.066	.202	.148	.091
Most Extreme Differences	Positive	.049	.202	.084	.091
	Negative	066	185	148	086
Kolmogorov-Smirnov Z		.492	1.514	1.106	.682
Asymp. Sig. (2-tailed)		.969	.020	.173	.740

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = HF1, Classes of equivalent diameter = 5

b. Test distribution is Normal.

Sample = L	_F2, Classes	of equivalent	diameter = 1
------------	--------------	---------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS Factor
Ν		35	35	35	35
Normal Daramatara <sup>b,c</sup>	Mean	.86266	1.42591	.89811	.0500
Normal Parameters	Std. Deviation	.066481	.313177	.026234	.03178
	Absolute	.124	.218	.114	.112
Most Extreme Differences	Positive	.077	.218	.114	.112
	Negative	124	109	062	083
Kolmogorov-Smirnov Z		.736	1.290	.673	.661
Asymp. Sig. (2-tailed)		.651	.072	.755	.775

a. Sample = LF2, Classes of equivalent diameter = 1

b. Test distribution is Normal.

c. Calculated from data.

### Sample = LF2, Classes of equivalent diameter = 2

		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		52	52	52	52
Normal Davage stars <sup>b,C</sup>	Mean	.80838	1.50617	.89663	.0775
Normal Parameters	Std. Deviation	.088369	.407827	.034484	.04718
	Absolute	.111	.176	.135	.210
Most Extreme Differences	Positive	.075	.176	.115	.210
	Negative	111	118	135	161
Kolmogorov-Smirnov Z		.804	1.268	.973	1.514
Asymp. Sig. (2-tailed)		.538	.080	.300	.020

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = LF2, Classes of equivalent diameter = 2

b. Test distribution is Normal.

Sample = LF2,	Classes of	equivalent	diameter = 3
---------------	------------	------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS Factor
Ν		44	44	44	44
Normal Daramatara <sup>b,c</sup>	Mean	.70361	1.74343	.87155	.1366
Normal Farameters	Std. Deviation	.123749	.515081	.058942	.09008
	Absolute	.137	.148	.169	.156
Most Extreme Differences	Positive	.079	.148	.107	.156
	Negative	137	099	169	128
Kolmogorov-Smirnov Z		.907	.981	1.121	1.035
Asymp. Sig. (2-tailed)		.383	.290	.162	.234

a. Sample = LF2, Classes of equivalent diameter = 3

b. Test distribution is Normal.

c. Calculated from data.

### Sample = LF2, Classes of equivalent diameter = 4

		Circ.	AR	Solidity	Elliptical PARIS	
					Factor	
Ν		46	46	46	46	
Normal Davage stars <sup>b,C</sup>	Mean	.70163	1.70704	.88143	.1412	
Normal Parameters	Std. Deviation	.122934	.488858	.061401	.08083	
	Absolute	.142	.133	.201	.179	
Most Extreme Differences	Positive	.074	.133	.123	.179	
	Negative	142	101	201	120	
Kolmogorov-Smirnov Z		.963	.900	1.365	1.217	
Asymp. Sig. (2-tailed)		.312	.392	.048	.103	

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = LF2, Classes of equivalent diameter = 4

b. Test distribution is Normal.

Sample = LF2, Classes	of equivalent diameter = 5
-----------------------	----------------------------

	· · ·	Circ.	AR	Solidity	Elliptical PARIS Factor
Ν		44	44	44	44
Normal Developmentary b.c	Mean	.62132	1.64441	.87532	.2324
Normal Parameters	Std. Deviation	.134559	.394860	.064711	.14679
	Absolute	.181	.130	.141	.211
Most Extreme Differences	Positive	.077	.130	.118	.211
	Negative	181	090	141	137
Kolmogorov-Smirnov Z		1.201	.865	.936	1.398
Asymp. Sig. (2-tailed)		.112	.443	.345	.040

a. Sample = LF2, Classes of equivalent diameter = 5

b. Test distribution is Normal.

c. Calculated from data.

### Sample = CLF2, Classes of equivalent diameter = 1

		Circ.	AR	Solidity	Elliptical PARIS	
					Factor	
Ν		80	80	80	80	
Normal Davage stars <sup>b,C</sup>	Mean	.66086	1.88786	.80902	.1689	
Normal Parameters	Std. Deviation	.157431	.598619	.082004	.16008	
	Absolute	.113	.088	.136	.177	
Most Extreme Differences	Positive	.069	.088	.135	.177	
	Negative	113	078	136	175	
Kolmogorov-Smirnov Z		1.012	.789	1.214	1.579	
Asymp. Sig. (2-tailed)		.257	.562	.105	.014	

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = CLF2, Classes of equivalent diameter = 1

b. Test distribution is Normal.

Sample = CLF2,	Classes of	equivalent	diameter = 2
----------------	------------	------------	--------------

0					
		Circ.	AR	Solidity	Elliptical PARIS
					Factor
Ν		57	57	57	57
Normal Parameters <sup>b,c</sup>	Mean	.58163	2.04582	.80577	.2198
	Std. Deviation	.132716	.666345	.070170	.14881
	Absolute	.057	.088	.077	.146
Most Extreme Differences	Positive	.044	.088	.064	.146
	Negative	057	075	077	114
Kolmogorov-Smirnov Z		.430	.661	.582	1.101
Asymp. Sig. (2-tailed)		.993	.774	.888	.177

a. Sample = CLF2, Classes of equivalent diameter = 2

b. Test distribution is Normal.

c. Calculated from data.

### Sample = CLF2, Classes of equivalent diameter = 3

		Circ.	AR	Solidity	Elliptical PARIS	
					Factor	
Ν		52	52	52	52	
Name I Davamatava <sup>b,c</sup>	Mean	.51567	1.77394	.79387	.3444	
Normal Parameters	Std. Deviation	.136759	.593905	.070336	.19297	
	Absolute	.115	.158	.091	.147	
Most Extreme Differences	Positive	.088	.158	.062	.147	
	Negative	115	142	091	086	
Kolmogorov-Smirnov Z		.826	1.136	.655	1.060	
Asymp. Sig. (2-tailed)		.502	.151	.785	.211	

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = CLF2, Classes of equivalent diameter = 3

b. Test distribution is Normal.

Sample = CLF2,	Classes of	of equivalent	diameter = 4
----------------	------------	---------------	--------------

		Circ.	AR	Solidity	Elliptical PARIS
					1 40101
N		38	38	38	38
Normal Parameters <sup>b,c</sup>	Mean	.47379	1.97813	.78208	.4153
Normal Farameters	Std. Deviation	.187825	.657537	.113770	.27189
	Absolute	.142	.174	.166	.168
Most Extreme Differences	Positive	.142	.174	.118	.168
	Negative	103	093	166	112
Kolmogorov-Smirnov Z		.872	1.072	1.026	1.037
Asymp. Sig. (2-tailed)		.432	.200	.243	.232

a. Sample = CLF2, Classes of equivalent diameter = 4

b. Test distribution is Normal.

c. Calculated from data.

### Sample = CLF2, Classes of equivalent diameter = 5

		Circ.	AR	Solidity	Elliptical PARIS	
					Factor	
Ν		43	43	43	43	
Name al Danama (anab.C	Mean	.40786	1.79735	.79712	.5509	
Normal Parameters	Std. Deviation	.152310	.565814	.095791	.31103	
	Absolute	.112	.160	.177	.129	
Most Extreme Differences	Positive	.112	.160	.086	.129	
	Negative	087	117	177	113	
Kolmogorov-Smirnov Z		.731	1.051	1.158	.847	
Asymp. Sig. (2-tailed)		.659	.219	.137	.470	

### One-Sample Kolmogorov-Smirnov Test<sup>a</sup>

a. Sample = CLF2, Classes of equivalent diameter = 5

b. Test distribution is Normal.

```
UNIANOVA Circ BY Sample EquiDclasses
/METHOD=SSTYPE(3)
/INTERCEPT=INCLUDE
/POSTHOC=Sample(SCHEFFE)
/PLOT=PROFILE(EquiDclasses*Sample)
/PRINT=HOMOGENEITY
/CRITERIA=ALPHA(.05)
/DESIGN=Sample EquiDclasses Sample*EquiDclasses.
```

### **Univariate Analysis of Variance**

		Value Label	N
	1	LF1	207
Sample	2	HF1	244
	3	LF2	221
	4	CLF2	270
	1	1	189
	2	2	188
classes of equivalent diameter	3	3	189
	4	4	188
	5	5	188

#### **Between-Subjects Factors**

#### Levene's Test of Equality of Error Variances<sup>a</sup>

Dependent Variable: Circ.

F	df1	df2	Sig.
4.439	19	922	.000

Tests the null hypothesis that the error variance of the dependent variable is equal across groups.<sup>a</sup> a. Design: Intercept + Sample + EquiDclasses + Sample \* EquiDclasses

#### **Tests of Between-Subjects Effects**

Dependent Variable: Circ.					
Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	11.145 <sup>a</sup>	19	.587	32.533	.000
Intercept	335.915	1	335.915	18630.158	.000
Sample	6.034	3	2.011	111.545	.000
EquiDclasses	5.026	4	1.257	69.691	.000
Sample * EquiDclasses	.226	12	.019	1.045	.405
Error	16.624	922	.018		
Total	373.184	942			
Corrected Total	27.770	941			

a. R Squared = .401 (Adjusted R Squared = .389)

### **Post Hoc Tests**

### Sample

#### **Multiple Comparisons**

Dependent Variable: Circ.

Scheffe

(I) Sample	(J) Sample	Mean Difference	Std. Error	Sig.	95% Confide	ence Interval
		(I-J)			Lower Bound	Upper Bound
	HF1	.06521 <sup>*</sup>	.012689	.000	.02967	.10074
LF1	LF2	12656 <sup>*</sup>	.012988	.000	16294	09019
	CLF2	.06054 <sup>*</sup>	.012405	.000	.02580	.09528
	LF1	06521 <sup>*</sup>	.012689	.000	10074	02967
HF1	LF2	19177 <sup>*</sup>	.012469	.000	22669	15685
	CLF2	00467	.011861	.985	03788	.02855
	LF1	.12656 <sup>*</sup>	.012988	.000	.09019	.16294
LF2	HF1	.19177 <sup>*</sup>	.012469	.000	.15685	.22669
	CLF2	.18710 <sup>*</sup>	.012181	.000	.15299	.22122
	LF1	06054 <sup>*</sup>	.012405	.000	09528	02580
CLF2	HF1	.00467	.011861	.985	02855	.03788
	LF2	18710 <sup>*</sup>	.012181	.000	22122	15299

Based on observed means.

The error term is Mean Square(Error) = .018.

\*. The mean difference is significant at the .05 level.

### **Homogeneous Subsets**

Circ.

Scheffe					
Sample	N	Subset			
		1 2 3			
HF1	244	.54489			
CLF2	270	.54955			
LF1	207		.61009		
LF2	221			.73666	
Sig.		.987	1.000	1.000	

Means for groups in homogeneous subsets are displayed. Based on observed means.

The error term is Mean Square(Error) = .018.

a. Uses Harmonic Mean Sample Size = 233.129.

b. The group sizes are unequal. The harmonic mean of the group sizes is used. Type I error levels are not guaranteed.

c. Alpha = .05.

### **Profile Plots**



```
UNIANOVA AR BY Sample EquiDclasses
/METHOD=SSTYPE(3)
/INTERCEPT=INCLUDE
/POSTHOC=Sample(SCHEFFE)
/PLOT=PROFILE(EquiDclasses*Sample)
/PRINT=HOMOGENEITY
/CRITERIA=ALPHA(.05)
/DESIGN=Sample EquiDclasses Sample*EquiDclasses.
```

### **Univariate Analysis of Variance**

		Value Label	Ν	
	1	LF1	207	
Sample	2	HF1	244	
	3	LF2	221	
	4	CLF2	270	
	1	1	189	
	2	2	188	
classes of equivalent diameter	3	3	189	
	4	4	188	
	5	5	188	

#### **Between-Subjects Factors**

#### Levene's Test of Equality of Error Variances<sup>a</sup>

Dependent Variable: AR

F	df1	df2	Sig.
2.229	19	922	.002

Tests the null hypothesis that the error variance of the dependent variable is equal across groups.<sup>a</sup> a. Design: Intercept + Sample + EquiDclasses + Sample \* EquiDclasses

#### **Tests of Between-Subjects Effects**

#### Dependent Variable: AR

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	19.243 <sup>a</sup>	19	1.013	2.792	.000
Intercept	2825.163	1	2825.163	7786.803	.000
Sample	10.076	3	3.359	9.257	.000
EquiDclasses	1.852	4	.463	1.276	.278
Sample * EquiDclasses	7.469	12	.622	1.716	.059
Error	334.515	922	.363		
Total	3342.419	942			
Corrected Total	353.758	941			

a. R Squared = .054 (Adjusted R Squared = .035)

### **Post Hoc Tests**

### Sample

**Multiple Comparisons** 

Dependent Variable: AR

#### Scheffe

(I) Sample	(J) Sample	Mean Difference	Std. Error	Sig.	95% Confide	ence Interval
		(I-J)			Lower Bound	Upper Bound
	HF1	03024	.056918	.963	18965	.12917
LF1	LF2	.16903 <sup>*</sup>	.058262	.039	.00586	.33220
	CLF2	11850	.055646	.210	27435	.03734
	LF1	.03024	.056918	.963	12917	.18965
HF1	LF2	.19926 <sup>*</sup>	.055934	.006	.04261	.35592
	CLF2	08826	.053204	.432	23727	.06074
	LF1	16903 <sup>*</sup>	.058262	.039	33220	00586
LF2	HF1	19926 <sup>*</sup>	.055934	.006	35592	04261
	CLF2	28753 <sup>*</sup>	.054639	.000	44055	13450
	LF1	.11850	.055646	.210	03734	.27435
CLF2	HF1	.08826	.053204	.432	06074	.23727
	LF2	.28753 <sup>*</sup>	.054639	.000	.13450	.44055

Based on observed means.

The error term is Mean Square(Error) = .363.

\*. The mean difference is significant at the .05 level.

### **Homogeneous Subsets**

AR

Scheffe				
Sample	N	Subset		
		1	2	
LF2	221	1.61003		
LF1	207		1.77906	
HF1	244		1.80930	
CLF2	270		1.89756	
Sig.		1.000	.212	

Means for groups in homogeneous subsets are displayed.

Based on observed means.

The error term is Mean Square(Error) = .363.

a. Uses Harmonic Mean Sample Size = 233.129.

b. The group sizes are unequal. The harmonic

mean of the group sizes is used. Type I error

levels are not guaranteed.

c. Alpha = .05.

### **Profile Plots**





```
UNIANOVA Solidity BY Sample EquiDclasses
/METHOD=SSTYPE(3)
/INTERCEPT=INCLUDE
/POSTHOC=Sample(SCHEFFE)
/PLOT=PROFILE(EquiDclasses*Sample)
/PRINT=HOMOGENEITY
/CRITERIA=ALPHA(.05)
/DESIGN=Sample EquiDclasses Sample*EquiDclasses.
```

### **Univariate Analysis of Variance**

		Value Label	N
	1	LF1	207
Sample	2	HF1	244
	3	LF2	221
	4	CLF2	270
	1	1	189
	2	2	188
classes of equivalent diameter	3	3	189
	4	4	188
	5	5	188

**Between-Subjects Factors** 

#### Levene's Test of Equality of Error Variances<sup>a</sup>

Dependent Variable: Solidity

F	df1	df2	Sig.	
6.905	19	922	.000	

Tests the null hypothesis that the error variance of the dependent variable is equal across groups.<sup>a</sup> a. Design: Intercept + Sample + EquiDclasses + Sample \* EquiDclasses

#### **Tests of Between-Subjects Effects**

#### Dependent Variable: Solidity

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	1.013 <sup>a</sup>	19	.053	11.670	.000
Intercept	637.609	1	637.609	139572.707	.000
Sample	.928	3	.309	67.735	.000
EquiDclasses	.042	4	.011	2.302	.057
Sample * EquiDclasses	.048	12	.004	.874	.573
Error	4.212	922	.005		
Total	666.055	942			
Corrected Total	5.225	941			

a. R Squared = .194 (Adjusted R Squared = .177)

### Post Hoc Tests Sample

#### **Multiple Comparisons**

Dependent Variable: Solidity

#### Scheffe

(I) Sample	(J) Sample	Mean Difference	Std. Error	Sig.	95% Confidence Interval	
		(I-J)			Lower Bound	Upper Bound
	HF1	.01956 <sup>*</sup>	.006387	.025	.00168	.03745
LF1	LF2	03664*	.006538	.000	05495	01834
	CLF2	.04809 <sup>*</sup>	.006244	.000	.03060	.06558
	LF1	01956 <sup>*</sup>	.006387	.025	03745	00168
HF1	LF2	05621 <sup>*</sup>	.006276	.000	07379	03863
	CLF2	.02853 <sup>*</sup>	.005970	.000	.01181	.04525
	LF1	.03664 <sup>*</sup>	.006538	.000	.01834	.05495
LF2	HF1	.05621 <sup>*</sup>	.006276	.000	.03863	.07379
	CLF2	.08474 <sup>*</sup>	.006131	.000	.06757	.10191
	LF1	04809*	.006244	.000	06558	03060
CLF2	HF1	02853*	.005970	.000	04525	01181
	LF2	08474 <sup>*</sup>	.006131	.000	10191	06757

Based on observed means.

The error term is Mean Square(Error) = .005.

\*. The mean difference is significant at the .05 level.

### **Homogeneous Subsets**

#### Solidity

Scheffe							
Sample	N		Subset				
		1	2	3	4		
CLF2	270	.79973					
HF1	244		.82826				
LF1	207			.84782			
LF2	221				.88447		
Sig.		1.000	1.000	1.000	1.000		

Means for groups in homogeneous subsets are displayed.

Based on observed means.

The error term is Mean Square(Error) = .005.

a. Uses Harmonic Mean Sample Size = 233.129.

b. The group sizes are unequal. The harmonic mean of the group sizes is

used. Type I error levels are not guaranteed.

c. Alpha = .05.

### **Profile Plots**



```
UNIANOVA Ellparisfaktor BY Sample EquiDclasses
  /METHOD=SSTYPE(3)
  /INTERCEPT=INCLUDE
  /POSTHOC=Sample(SCHEFFE)
  /PLOT=PROFILE(EquiDclasses*Sample)
  /PRINT=HOMOGENEITY
  /CRITERIA=ALPHA(.05)
  /DESIGN=Sample EquiDclasses Sample*EquiDclasses.
```

### **Univariate Analysis of Variance**

Detween-Oubjeets Factors			
		Value Label	N
	1	LF1	207
0	2	HF1	244
Sample	3	LF2	221
	4	CLF2	270
	1	1	189
	2	2	188
Classes of equivalent diameter	3	3	189
	4	4	188
	5	5	188

### **Between-Subjects Factors**

#### Levene's Test of Equality of Error Variances<sup>a</sup>

Dependent Variable: Elliptical PARIS Factor

F	df1	df2	Sig.
13.690	19	922	.000

Tests the null hypothesis that the error variance of the dependent variable is equal across groups.<sup>a</sup> a. Design: Intercept + Sample + EquiDclasses + Sample \* EquiDclasses

Dependent Variable: Elliptical PARIS Factor						
Source	Type III Sum of	df	Mean Square	F	Sig.	
	Squares					
Corrected Model	12.984 <sup>a</sup>	19	.683	27.613	.000	
Intercept	54.632	1	54.632	2207.560	.000	
Sample	5.815	3	1.938	78.325	.000	
EquiDclasses	6.285	4	1.571	63.491	.000	
Sample * EquiDclasses	1.108	12	.092	3.732	.000	
Error	22.818	922	.025			
Total	93.445	942				
Corrected Total	35.801	941				

#### **Tests of Between-Subjects Effects**

a. R Squared = .363 (Adjusted R Squared = .350)

### **Post Hoc Tests**

### Sample

#### **Multiple Comparisons**

Dependent Variable: Elliptical PARIS Factor

#### Scheffe

(I) Sample	(J) Sample	Mean Difference	Std. Error	Sig.	95% Confidence Interval	
		(I-J)			Lower Bound	Upper Bound
	HF1	0722*	.01487	.000	1138	0305
LF1	LF2	.1004 <sup>*</sup>	.01522	.000	.0578	.1431
	CLF2	0795 <sup>*</sup>	.01453	.000	1202	0388
	LF1	.0722*	.01487	.000	.0305	.1138
HF1	LF2	.1726 <sup>*</sup>	.01461	.000	.1317	.2135
	CLF2	0074	.01390	.963	0463	.0315
	LF1	1004 <sup>*</sup>	.01522	.000	1431	0578
LF2	HF1	1726 <sup>*</sup>	.01461	.000	2135	1317
	CLF2	1800 <sup>*</sup>	.01427	.000	2199	1400
	LF1	.0795 <sup>*</sup>	.01453	.000	.0388	.1202
CLF2	HF1	.0074	.01390	.963	0315	.0463
	LF2	.1800 <sup>*</sup>	.01427	.000	.1400	.2199

Based on observed means.

The error term is Mean Square(Error) = .025.

\*. The mean difference is significant at the .05 level.

### **Homogeneous Subsets**

Scheffe					
Sample	N	Subset			
		1	2	3	
LF2	221	.1290			
LF1	207		.2295		
HF1	244			.3016	
CLF2	270			.3090	
Sig.		1.000	1.000	.968	

#### **Elliptical PARIS Factor**

Means for groups in homogeneous subsets are displayed.

Based on observed means.

The error term is Mean Square(Error) = .025.

a. Uses Harmonic Mean Sample Size = 233.129.

b. The group sizes are unequal. The harmonic mean of the group sizes is used. Type I error levels are not guaranteed.c. Alpha = .05.

### **Profile Plots**



Classes of equivalent diameter

### Kruskal-Wallis Test

Ranks				
	Sample	N	Mean Rank	
	LF1	207	473.93	
	HF1	244	369.44	
Circ.	LF2	221	682.63	
	CLF2	270	389.04	
	Total	942		
	LF1	207	464.96	
	HF1	244	477.41	
AR	LF2	221	396.97	
	CLF2	270	532.17	
	Total	942		
	LF1	207	497.58	
	HF1	244	411.87	
Solidity	LF2	221	675.30	
	CLF2	270	338.58	
	Total	942		
	LF1	207	468.23	
	HF1	244	591.38	
Elliptical PARIS Factor	LF2	221	265.97	
	CLF2	270	533.90	
	Total	942		

Te	st Statistics <sup>a,b</sup>	

	Circ.	AR	Solidity	Elliptical PARIS
				Factor
Chi-Square	192.231	30.245	202.067	187.713
df	3	3	3	3
Asymp. Sig.	.000	.000	.000	.000

a. Kruskal Wallis Test

b. Grouping Variable: Sample

### Kruskal-Wallis Test

Ranks					
	Classes of equivalent	Ν	Mean Rank		
	diameter				
	1	189	631.41		
	2	188	557.29		
Circ	3	189	453.32		
Circ.	4	188	402.82		
	5	188	311.90		
	Total	942			
	1	189	467.50		
	2	188	452.15		
	3	189	472.77		
AK	4	188	491.67		
	5	188	473.43		
	Total	942			
	1	189	469.76		
	2	188	490.25		
Solidity	3	189	440.80		
Solidity	4	188	458.74		
	5	188	498.13		
	Total	942			
	1	189	284.26		
	2	188	376.07		
Elliptical DADIS Eastor	3	189	495.69		
	4	188	545.96		
	5	188	656.38		
	Total	942			

### Test Statistics<sup>a,b</sup>

	Circ.	AR	Solidity	Elliptical PARIS							
				Factor							
Chi-Square	161.494	2.039	5.521	215.017							
df	4	4	4	4							
Asymp. Sig.	.000	.729	.238	.000							

a. Kruskal Wallis Test

b. Grouping Variable: Classes of equivalent diameter

## Electronic supplement: APPENDIX D: SPSS example datafile

Sample	Nr	Area	Perim	Fitellips-Major	Fitellips-Minor	Circ	AR	Solidity	EquiD	Ellparisfaktc
1	1	328.045	77.73	22.465	18.592	0.682	1.208	0.843	20.44	0.20
1	2	331.859	73.82	28.392	14.882	0.765	1.908	0.921	20.56	0.06
1	3	1'224.446	154.79	45.376	34.357	0.642	1.321	0.881	39.48	0.23
1	4	1'136.713	136.13	39.377	36,755	0.771	1.071	0.909	38.04	0.14
1	5	1'945 382	206 60	55 170	44 897	0.573	1 229	0.818	49 77	0.31
1	6	1'651 667	163.08	52 829	39 807	0 780	1 327	0.905	45.86	0.12
1	7	324 230	71.06	22 827	18 085	0.807	1 262	0.885	20.32	0.10
1	8	381 117	83.25	30.823	15 757	0.607	1.202	0.000	22.02	0.10
1	0	942 000	125.65	50.025	10.157	0.032	2 250	0.077	22.04	0.11
1	9	042.999	135.05	29.131	10.152	0.576	3.230	0.090	32.70	0.05
I	10	306.972	00.49	21.554	10.209	0.070	1.179	0.076	19.05	0.06
•	•	•		•	•	•	•			•
•	•		•	•	•	•	•	•	•	
1	207	637 017	101 45	33 325	24 338	0 778	1 369	0 895	28 48	0 11
2	1	1'399 912	184.31	52 681	33 834	0.518	1.557	0.826	42 22	0.34
2	2	1'178 672	172 31	44 203	33 051	0.400	1 302	0.707	38.74	0.01
2	2	442.470	90.02	22 252	16 042	0.433	1.002	0.737	22.74	0.40
2	3	442.479	121.25	20.202	21 002	0.000	1.903	0.875	23.74	0.11
2	4	901.247	131.35	39.304	31.092	0.700	1.200	0.070	54.90	0.16
2	5	2063.630	216.23	01.751	42.550	0.555	1.451	0.834	51.20	0.31
2	6	5023.662	337.20	92.671	69.022	0.555	1.343	0.831	79.98	0.32
2	1	15 292.225	587.99	206.634	94.228	0.556	2.193	0.906	139.54	0.20
2	8	945.989	152.98	48.149	25.015	0.508	1.925	0.813	34.71	0.30
2	9	1'724.142	188.41	51.519	42.610	0.610	1.209	0.850	46.85	0.27
2	10	3'318.592	271.66	85.027	49.695	0.565	1.711	0.878	65.00	0.26
	•		:				:		:	
2	244	1'411.355	152.03	52.032	34.537	0.767	1.507	0.896	42.39	0.11
3	1	3'795.401	249.29	79.543	60.753	0.767	1.309	0.915	69.52	0.13
3	2	869.700	128.79	52.643	21.035	0.659	2.503	0.889	33.28	0.07
3	3	1'499.088	156.41	48.743	39.158	0.770	1.245	0.891	43.69	0.13
3	4	3'330.036	313.96	101.003	41.978	0.425	2.406	0.667	65.11	0.34
3	5	831.555	108.78	38.417	27.560	0.883	1.394	0.926	32.54	0.04
3	6	808.668	107.64	33.968	30.312	0.877	1.121	0.920	32.09	0.07
3	7	1'819.504	172.70	57.392	40.366	0.767	1.422	0.931	48.13	0.12
3	8	5'397.480	293.68	98.974	69.435	0.786	1.425	0.944	82.90	0.10
3	9	6'900.383	419.03	127.511	68.902	0.494	1.851	0.773	93.73	0.33
3	10	740 008	100.97	33 705	27 955	0.912	1 206	0.922	30.70	0.04
3	221	7/3 822	104.88	38.604	24 533	0.850	1 574		30.77	
1	1	611 442	122.25	12 251	17.059	0.000	2 414	0.322	27.00	0.04
4	1	011.442	133.30	43.331	17.900	0.432	2.414	0.754	27.90	0.33
4	2	341.800	111.58	35.468	12.270	0.345	2.891	0.062	20.86	0.41
4	3	1412.774	163.39	49.478	30.300	0.000	1.301	0.849	42.41	0.20
4	4	1165.918	210.71	41.581	35.701	0.330	1.165	0.760	38.53	0.73
4	5	341.800	81.73	26.866	16.199	0.643	1.659	0.793	20.86	0.19
4	6	463.329	96.18	24.742	23.843	0.629	1.038	0.795	24.29	0.26
4	7	349.396	75.28	25.128	17.704	0.775	1.419	0.876	21.09	0.11
4	8	3'490.158	253.11	77.141	57.606	0.685	1.339	0.879	66.66	0.19
4	9	3'304.067	404.22	80.811	52.058	0.254	1.552	0.660	64.86	0.91
4	10	35'744.691	911.31	287.685	158.199	0.541	1.819	0.912	213.33	0.27
•	•		•		•	•	•	•	•	•
4	270	349.396	77.84	27.221	16.343	0.725	1.666	0.807	21.09	0.12

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Austrian Journal of Earth Sciences

Jahr/Year: 2012

Band/Volume: 105\_3

Autor(en)/Author(s): Kohlmayer Norbert, Grasemann Bernhard

Artikel/Article: <u>Quantitative characterisation of cataclasites using a statistical approach</u> (analysis of variance) 48-60