

The sharpshooters of Panguana (Peru) (Auchenorrhyncha, Cicadellidae, Cicadellinae)

KLAUS SCHÖNITZER & WIELAND FEUERABENDT

Abstract

In this communication we list 38 species of Cicadellini and 14 species of Proconiini (Cicadellinae) from Panguana in the Amazon lowland of Peru, as well as some other interesting records of sharpshooters from South America. The following species are new for Peru: *Cardioscarta electa* MELICHAR, 1932, *Jakrama taeniata* YOUNG, 1977, *Oragua elegantula* YOUNG, 1977, *Omagua fitchii* (SIGNORET, 1855), *Propetes schmidti* MELICHAR, 1925, *Propetes triquetra* (FABRICIUS, 1803). New records for the countries given in brackets are: *Soosiulus interpolis* YOUNG, 1977 (Peru and Bolivia), *Dilobopterus lucentis* YOUNG, 1977 (Brazil), *Juliacaca scalarum* YOUNG, 1977 (Brazil), *Paromenia isabellina* (FOWLER, 1899) (Ecuador), *Zyzzogeton viridipennis* (LATREILLE, 1811) (Ecuador), *Pawiloma rubra* CAVICHIOLI, 1998 (Guatemala) and *Aulacizes quadripunctata* (GERMAR, 1821) (Paraguay).

Introduction

The ecological field station Panguana in the Amazon lowland of Peru is a research area predominantly for zoology. It is frequently called „hotspot of biodiversity“ (e.g. DILLER & BURMEISTER 2007). For over 40 years the station has been used for research, since the foundation of Panguana by Hans-Wilhelm and Maria KOEPCKE. Within the last years the research station was expanded and is run by Juliane DILLER.

Panguana is located at about 260 m altitude in the departamento de Huánuco, its geographical position is 9° 36' 49,3" S, 74° 56' 8,2" W (see map in this issue in HORNBURG 2014, page 27, Fig. 68, point 8). There are many different habitats, partly dense rainforests, low hills, little creeks, swamps with palm trees and pastures. The river Llullapichis (= Yuyapichis) which originates from the Sira mountains passes through the area of Panguana. This river is usually only about one or two meters deep and some 40 m wide but it can become a rapid stream of six to seven meters depth within a few hours. The mean air temperature in Panguana is about 24,5 °C, but in the dry season (October to April) maximum temperatures of about 40 °C are quite common. The annual precipitations are up to 3.000 mm with 180 days of rain per year. The humidity of the air is about 90 % within the forest (DILLER & BURMEISTER 2006, HENZE 2011). Since 2011 the area of Panguana is a private natural reserve by appointment of the Peruvian authorities (SPDA, weblink).

In this communication we provide data on the sharpshooters collected in Panguana within the last years. A first, preliminary account on Cicadellidae from Panguana was published by LOZADA et al. (2007). In addition, we list some new records from other South American countries based on specimens housed in the ZSM collections.

Material and Methods

The material was collected in Panguana by members of the Bavarian State Collection of Zoology (ZSM) and accompanying zoologists. The material is deposited in ZSM and in the Museo de Historia Natural Javier Prado of the University San Marcos in Lima (Universidad Nacional Mayor de San Marcos).

Collection times and collectors. To avoid unnecessary repetitions the main collectors and the collecting times are listed here in alphabetical order, single collectors are listed below.

Apr. 06 - Apr. 17/2003: T. KOTHE, E.-G., H. and J. BURMEISTER, C. J. ZWAKHALS

Sept. 21 - Oct. 04/2004: M. BREITSAMETER, T. KOTHE, K. SCHÖNITZER

Sept. 19 - Oct. 02/2005: M. BREITSAMETER, E.-G. BURMEISTER, E. DILLER, O. GRULER, T. KOTHE

Sept. 20 - Oct. 07/2007: E. DILLER

Nov. 23 - Dec. 11/2008: E. DILLER, F. GLAW, K. SCHÖNITZER, F. WACHTEL

Oct. 02 - Oct. 18/2009: E.-G. BURMEISTER

Sept. 20 - Oct. 06/2013: E. DILLER

The identification of the sharpshooters was predominantly done with the help of the excellent website by WILSON et al. (2009). Furthermore the publications by MELICHAR (1951), YOUNG (1968, 1977) and McKAMEY (2007) were used. Previously determined material in the collection, especially determined by D. A. YOUNG and P. W. LOZADA, could be used for comparison or could be included in this communication. These two colleagues frequently made genitalia preparations. The distribution as mentioned follows WILSON et al (2009) if not otherwise stated. In several cases the pictures in WILSON et al. (2009) provide additional locality information.

Results

Tribus Cicadellini

Alocha rossi (YOUNG, 1977)

Number of specimens: 1; leg.: April 8th 2003, E. DILLER.

Distribution: Peru, Brazil.

Determined by P. LOZADA, labelled as *Paromenia rossi* (i.e. synonym).

Amblyscarta trinotata YOUNG, 1977 (Fig. 1)

Specimens: 8; leg.: 2004, 2007.

Distribution: Peru, Brazil.

Only known from Amazon lowland of Peru and Brazil (YOUNG 1977). McKAMEY (2007) listed for Peru only.

Caldwelliola solimoeana YOUNG, 1977 (Fig. 2)

Specimens: 1; leg.: 2004 (T. KOTHE).

Distribution: Peru, Brazil.

Only found in Amazon lowland (YOUNG 1977, ZAMOL & MENEZES 1982).

Cardioscarta electa MELICHAR, 1932 (Fig. 3)

Specimens: 2; leg.: 2003.

Distribution: Ecuador, Bolivia, **new record for Peru**.

Determined by P. LOZADA.

Cardioscarta quadrifasciata (LINNAEUS, 1758)

Specimens: 1; leg.: 2003 (T. KOTHE).

Distribution: Colombia, Guiana, French Guiana, Peru, Bolivia, Brazil.

Determined by P. LOZADA

Catagonalia conjunctula (OSBORN, 1926) (Fig. 4)

Specimens: 3; leg.: 2003, 2008.

Distribution: Colombia, Bolivia, Brazil, Ecuador, Peru, Paraguay.

One of the specimens looks rather like *Graphocephala lemnisca*, but the latter species is known from Mexico only. In Peru found in Madre de Dios (LOZADA 2010).

Catagonalia lunata (SIGORET, 1854) (Fig. 5)

Specimens: 18; leg.: 2003, 2004, 2007, 2008.

Distribution: Ecuador, Peru, Bolivia, Brazil, Colombia.

Partly determined by P. LOZADA and listed by LOZADA et al. (2007). According to LOZADA & ARELLANO (2008) this species is very common in the lowland of Peru, especially on open, secondary vegetation.

Diedrocephala bimaculata (GMELIN, 1789) (Fig. 6)

Specimens: 4; leg.: 2004, 2005, 2008 (one specimen collected with hand-net).

Distribution: Mexico, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Guiana, Suriname, French Guiana, Brazil, Paraguay, Argentina, Peru, Bolivia (DELLAPÉ et al. 2013).


This species is of economic importance as vector for bacteria *Xylella fastidiosa* in citrus orchards (DELLAPÉ et al. 2013), it is also common in the Satipo province of Peru (LOZADA & ARELLANO 2008). Species with similar appearance are *Tipuana chirensis*, *Platygonia angrana* and *Lebaja mediana*.

Dilobopterus adocetus YOUNG, 1977

Specimens: 1; leg.: 2003 (E.-G. BURMEISTER.).

Distribution: Peru, Brazil, Bolivia.

A rather small specimen (size = 10,4 mm), possibly female (YOUNG, 1977, had males only).



Figs 1-9: 1. *Amblyscarta trinotata* (Length: 10,1 mm); 2. *Caldwelliola solimoeana* (6,0 mm); 3. *Cardioscarta electa* (11,0 mm); 4. *Catagonalia conjunctula* (7,6 mm); 5. *Catagonalia lunata* (9,1 mm); 6. *Diedocephala bimaculata* (6,9 mm); 7. *Dilobopterus gaudens* (16,0 mm); 8. *Dilobopterus vicinus* (12,9 mm); 9. *Hortlesia similis* (4,9 mm).

***Dilobopterus demissus* (FABRICIUS, 1803)**

Specimens: 3; leg.: 2003.

Distribution: Panama, Colombia, French Guiana, Guiana, Ecuador, Peru, Bolivia, Brazil, Venezuela.

***Dilobopterus gaudens* (WALKER, 1851) (Fig. 7)**

Specimens: 3; leg.: 2009, 2013.

Distribution: Peru, Brazil.

Rather large species (size = 16,0 mm), with conspicuous coloration.

***Dilobopterus jemima* (DISTANT, 1908)**

Specimen: 1; leg.: 2004.

Distribution: Ecuador, Peru, Bolivia, Brazil.

Determined by P. LOZADA. Also found in rather high altitude in the sierra of Peru (up to 2.400 m; LOZADA & ARELLANO 2008).

***Dilobopterus manni* YOUNG, 1977**

Specimens: 4; leg.: 2003.

Distribution: Bolivia, Peru.

***Dilobopterus obliquatulus* (JACOBI, 1905)**

Specimens: 2; leg.: 2003, 2004.

Distribution: Colombia, Ecuador, Peru, Bolivia, Brazil.

One specimen (size = 9,3 mm) determined by P. LOZADA; listed for the region of Cusco (LOZADA & ARELLANO 2008).

***Dilobopterus syrphoidulus* (JACOBI, 1905)**

Specimens: 2; leg.: 2003, 2004.

Distribution: Ecuador, Peru, Bolivia, Brazil, Colombia.

One specimen determined by P. LOZADA.

***Dilobopterus vicinus* (SIGNORET, 1853) (Fig. 8)**

Specimens: 2; leg.: 2005, 2009.

Distribution: Colombia, Bolivia, Brazil, Paraguay, Ecuador, Peru.

***Draeculacephala clypeata* OSBORN, 1926**

Specimens: 2; leg.: 2003, T. KOTHE.

Distribution: Mexico to Colombia, Guiana, Argentina, Peru.

Determined by P. LOZADA.

***Fusigonalia optata* YOUNG, 1977**

Specimens: 1; leg.: 2003, T. KOTHE.

Distribution: Bolivia, Peru.

Determined by P. LOZADA. This species was found frequently in coffee plantations (LOZADA & ARELLANO 2008).

***Hortlesia similis* (WALKER, 1851) (Fig. 9)**

Specimens > 200; leg.: 2003, 2004 (a large series of 130 individuals was collected by hand-net on pasture near Panguana, 17.4.2003 by E. DILLER and T. KOTHE).

Distribution: widely distributed in North, Middle and South America.

This species is frequently found on agricultural land. In Peru it is found in the coastal areas, in the mountain ranges (COSTA & LOZADA 2010) and in the Amazon basin (LOZADA 1993, LOZADA & ARELLANO 2008, DIAZ & ZAMORA 2001). In an orchard in Brazil it was found to be frequent throughout all the year (OTT et al. 2006). In our investigation this species was very abundant in April 2003, but was found only rarely in other years, when collection trips were performed in September and October. Thus it seems that this species might be more frequent in April than September and October.

***Iragua montana* YOUNG, 1977**

Specimens: 3; leg.: 2004, 2006 (G. RIEDEL).

Distribution: Peru, Bolivia.

Determined by P. LOZADA. This species is also reported from the sierra of Peru (COSTA & LOZADA 2010). Furthermore LOZADA et al. (2007) mentioned *Iragua diversa* (SIGNORET, 1855), but material of this species could not be found (possibly a mistake, or material passed to Lima).

***Jakrama krameri* YOUNG, 1977 (Fig. 10)**

Specimens: 3; leg.: 2003.

Distribution: Venezuela, Guiana, Brazil, Peru, Colombia.

Very similar to *J. eureta*, one sample was previously determined as *J. eureta* by P. LOZADA.

***Jakrama taeniata* YOUNG, 1977**

Specimens: 3; leg.: 2004, one specimen collected in malaise trap.

Distribution: Bolivia, Brazil, Ecuador, Colombia, **new record for Peru**.

This species may be distinguished by the broader dark stripes and a generally lighter coloration, fine silvery shade. One sample was previously determined as *J. eureta* by LOZADA and listed by LOZADA et al. (2007). According to our results, however, the latter species was not recorded from Panguana.

***Julia cinctella* (MELICHAR, 1951) (Fig. 11)**

Specimens: 15; leg.: 2003, 2004, (mostly collected with hand net on pasture, 17.4.2003 by E. DILLER and T. KOTHE).

Distribution: Peru.

Rather small species (size = 4,5 mm), with distinctive color pattern. By LOZADA determined to genus. A rare species, up to now only known from the original description from Pachitea (MELICHAR 1951, p. 97). Panguana and Pachitea are both in the department Huánuco.

***Ladoffa obscurana* YOUNG, 1977 (Fig. 12)**

Specimens: 1; leg.: 2004 (A. OEHlke, malaise trap).

Distribution: Peru, Ecuador, Bolivia, Brazil, Colombia (FREYTAG & SHARKEY 2002).

***Lissoscarta nipata* YOUNG, 1977**

Specimens: 4; leg.: 2003, 2004.

Distribution: Peru.

Determined by P. LOZADA and listed by LOZADA et al. (2007). This species was previously only reported from the original description from Cusco country, Hacienda María (YOUNG, 1977), listed also in COSTA & LOZADA (2010). These cicadas probably imitate wasps.

***Macugonalia moesta* (FABRICIUS, 1803)**

Specimens: 48; leg.: 2003, 2004, 2008 (31 specimens collected with hand-net, on pasture near Panguana, 17.4.2003 E. DILLER, T. KOTHE, one specimen collected in Malaise trap).

Distribution: Guyana, Suriname, French Guiana, Brazil, Ecuador, Peru, Bolivia, Colombia.

Mostly determined by P. LOZADA. Common in plantations of papaya and avocado trees (LOZADA & ARELLANO 2008).

***Microgoniella pudica* (FABRICIUS, 1803)**

Specimens: 4, leg.: 2003.

Distribution: Colombia, Brazil, Bolivia, Peru.

Determined by P. LOZADA. All year round to be found in plantations in Peru in Chanchomayo and Satipo (LOZADA & ARELLANO 2008), also found in Ucayali province (DIAZ & ZAMORA 2001).

***Oragua elegantula* YOUNG, 1977 (Fig. 13)**

Specimens: 3; leg.: 2004, one specimen collected by malaise trap.

Distribution: Brazil, **new record for Peru**.

Very similar to *O. alvarengai*. By LOZADA labelled as „*Oragua* spec. nov.”

***Oragua nusinasa* YOUNG, 1977 (Fig. 14)**

Specimens: 5; leg.: 2003, 2007.

Distribution: Peru.


This species is only known from Peru, it was reported from Cusco and Arequipa regions (Callanga), interestingly from higher elevations (YOUNG 1977, COSTA & LOZADA 2010), but also found in the northern lowland, San Martin region (WILSON et al. 2009).

***Oragua partitula* (JACOBI, 1905) (Fig. 15)**

Specimens: 1; leg.: 2007 (E. DILLER).

Distribution: Peru, Bolivia.

According to LOZADA & ARELLANO (2008) a very common species in tropical forests of Peru, also reported from the department Cusco (COSTA & LOZADA, 2010), listed by LOZADA et al. (2007).



Figs 10-18: **10.** *Jakrama krameri* (Length: 13,3 mm); **11.** *Juliaca cinctella* (4,5 mm); **12.** *Ladoffe obscurana* 6,3 mm); **13.** *Oragua elegantula* (7,8 mm); **14.** *Oragua nusinasa* (9,7 mm); **15.** *Oragua partitula* (9,9 mm); **16.** *Pachitea jezima* (9,4 mm); **17.** *Poeciloscarta aurorula* (11,7 mm); **18.** *Ramosulus corrugipennis* (4,6 mm).

***Pachitea jezima* YOUNG, 1977 (Fig. 16)**

Specimens: 1; leg.: 2009 (E.-G. BURMEISTER).

Distribution: Peru.

The black surface of the wings has a bluish tinge. Only known from Amazon lowland in Peru, Tingo Maria, Madre de Dios (YOUNG 1977, p. 333).

***Plesiomma mollicella* FOWLER, 1900**

Specimens: 23; leg.: 2003, 2004, partly by hand-net.

Distribution: Mexico, Belize, Guatemala, Costa Rica, Panama, Colombia, Venezuela, French Guiana, Ecuador, Bolivia, Paraguay, Brazil, Argentina, Peru.

According to LOZADA (1993) and LOZADA & ARELLANO (2008) a very common species in tropical Peru (although Peru is not listed in WILSON et al. 2009).

Determined by P. LOZADA.

***Poeciloscarta aurorula* (BREDDIN, 1901) (Fig. 17)**

Specimens: 1; leg.: April 2012, E. DILLER.

Distribution: Colombia, Peru, Bolivia, Brazil.

Coloration may vary considerably (WILSON et al. 2009).

***Ramosulus corrugipennis* (OSBORN, 1926) (Fig. 18)**

Specimens: 7; leg.: 2003, 2004 (hand net, one specimen from pasture near Panguana).

Distribution: Brazil, Ecuador, Peru, Bolivia, Colombia.

According to LOZADA & ARELLANO (2008) a species which is common in disturbed areas of the tropical forest, also reported from the department Cusco (COSTA & LOZADA, 2010). The species is similar to *Julia pergrata* and *J. cruenta*.

***Rotigonalia concedula* (MELICHAR, 1926)**Specimens: 1; leg.: Apr. 17th 2003, E. DILLER & T. KOTHE, hand-net on pasture near Panguana.

Distribution: Peru, Bolivia, Brazil (CAVICHIOLI & TAKIYA 2012).

Determined by P. LOZADA.

***Scopogonalia subolivacea* (STÅL, 1862)**

Specimens: 3; leg.: 2003 (T. KOTHE), 2006 (G. RIEDEL).

Distribution: Peru, Bolivia, Brazil, Paraguay, Argentina.

Determined by P. LOZADA. According to LOZADA & ARELLANO (2008) a species which is common in disturbed areas of the tropical forest. In Peru also known from the Chanchamayo province (LOZADA et al. 2007, WILSON et al. 2009).

***Soosiulus interpolis* YOUNG, 1977 (Fig. 19)**

Specimens: 2; leg.: 2003, 2004 (T. KOTHE) hand net.

Distribution: Brazil, Colombia, Guiana (BASSET et al. 2001), new record for Peru and Bolivia.

This species feeds on leaves in the canopy (BASSET et al. 2001). Additional material in ZSM: Bolivia, Chapare, La Jungla, Chipiriri near Villa Tunari 9.11.2002; two specimens; leg. B. BEMBÉ.

***Tylozygus fasciatus* (WALKER, 1851)**

Specimens 3; leg.: 2004.

Distribution: US, Mexico, Guatemala, Belize, El Salvador, Costa Rica, Panama, Haiti, Dominica, Grenada, Trinidad, Jamaica, Venezuela, Colombia, Ecuador, Peru, Brazil, Cuba.

Determined by P. LOZADA, listed by LOZADA et al. (2007). This very widespread and colorful species was found only rarely in Panguana.

Additional species

Several specimens could only be identified to genus level within this study:

Amblyscarta sp. (possibly *A. opulenta* ?, LOZADA et al. 2007), *Lautereria* sp., *Lissocarta* sp., *Mesogonia* sp., *Dilobopterus* sp., and *Oragua* sp..

The genus *Soosiulus* contains several rather similar species, one specimen was determined as *S. servulus* (MELICHAR, 1932) by P. LOZADA, but this is doubtful.

Of the genus *Microgoniella* one specimen was determined by LOZADA as *Microgoniella* sp. n., cf. *apicula* (OSBORN, 1926) and another one as *Microgoniella* sp. n.. These identifications require confirmation.

Tribus Proconiini

***Acrogonia terminalis* YOUNG, 1968**

Specimens 2; leg.: 2003.

Distribution: Peru, Brazil, Bolivia.

Determined by P. LOZADA. This species occurs from February to August on leaves of papaya trees (LOZADA & ARELLANO 2008); this and the following species are known to be vectors for plant diseases in Brazil (AZEVEDO FILHO & CARVALHO 2004).

***Acrogonia virescens* (METCALF, 1949)**

Specimens 2; leg.: VII.1982, 2009 (E.-G. BURMEISTER).

Distribution: Peru, Guiana, Brazil, Paraguay.

LOZADA & ARELLANO (2008) reported this species from papaya leaves.

***Desamera intersecta* (GERMAR, 1821) (Fig. 20)**

Specimens 3; leg.: 2005, 2009.

Distribution: Peru, Brazil (CARVALHO et al. 2011, ZAMOL & MENEZES 1982).

***Diestostemma huallagana* YOUNG, 1968**

Specimens 1; leg.: 2004.

Distribution: Peru, Brazil, Bolivia (DELLAPE et al. 2011).

Determined by P. LOZADA; in Peru reported from several locations, mostly from department of Huánuco (YOUNG 1968, p.44, WILSON et al. 2009).

***Ichthyobelus bellicosus* MELICHAR, 1925**

Specimens 1; leg.: 2005.

Distribution: Ecuador, Peru, Bolivia.

Also reported from the department Cusco (COSTA & LOZADA, 2010) and listed by LOZADA et al. (2007).

***Molomea consorta* (MELICHAR, 1925)**

Specimens: 2; leg.: 2005, 2009 (E.-G. BURMEISTER).

Distribution: Brazil, Peru.

Reported for the department Cusco (COSTA & LOZADA, 2010) and listed by LOZADA et al. (2007); Peru is not mentioned, however, in WILSON et al. (2009).

***Omagua fitchii* (SIGNORET, 1855) (Fig. 21)**

Specimens 2; leg.: 2005, 2007.

Distribution: French Guiana, Guiana, Ecuador (WILSON et al. 2009), **new record for Peru**.

According to YOUNG (1968) also reported from „Amazonas”, which probably means Brazil, listed in ZAMOL & MENEZES (1982).

***Proconia marmorata* (FABRICIUS, 1803)**

Specimens 6; leg.: 2003 (T. KOTHE).

Distribution: Colombia, Bolivia, Brazil, Ecuador, Guiana, Peru.

In Peru previously reported from the department Cusco (COSTA & LOZADA, 2010) and Huanuco (WILSON et al. 2009), listed by LOZADA et al. (2007). Determined by P. LOZADA.

***Propetes schmidti* MELICHAR, 1925 (Figs 22, 23)**

Specimens 1♂; leg.: 2003 (T. KOTHE).

Distribution: Paraguay, Brazil, **new record for Peru**.

This species is imitating wasps (TAKIYA et al. 1999). Males and females mimic different (morpho-)species of wasps.

***Propetes triquetra* (FABRICIUS, 1803)**

Specimens 2; leg.: 2005.

Distribution: Guiana, Brazil, **new record for Peru**.

Also this species apparently mimics wasp species. The fore tibiae are remarkably broadened. For taxonomy of this species see TAKIYA et al. (2006).

***Pseudometopia amblardii* (SIGNORET, 1855) (Fig 24)**

Specimens 28; leg.: 2003, 2004, 2006, 2008, 2009, 2013.

Distribution: Bolivia, Mexico, Brazil, Colombia, Ecuador, Peru, Paraguay, Argentina.

Remarkably often found in Panguana, with variable coloration, but nevertheless with a characteristic color pattern. Part of the material was determined P. LOZADA. Very common species in plantations in Chanchamayo province (LOZADA & ARELLANO 2008), also known from Tingo Maria (WILSON et al. 2009).

***Pseudometopia phalaesia* (DISTANT, 1908)**

Specimens 2; leg.: 2003.

Distribution: Colombia, Peru, Bolivia.

Determined by P. LOZADA. In Peru also frequent in plantations in Chanchamayo province (LOZADA & ARELLANO 2008) and known from Tingo Maria (WILSON et al. 2009).

***Raphirhinus phosphoreus* (LINNAEUS, 1758) (Fig. 25)**

Specimens 14; leg.: 2003 (T. KOTHE, C. ZWAKHALS), 2004, 2005, 2009.

Distribution: Ecuador, Bolivia, Peru, Colombia, Venezuela, Trinidad, Guiana, Suriname, French Guiana, Brazil.

Part of the material determined by P. LOZADA and listed by LOZADA et al. (2007). In Peru also known from the department Cusco (COSTA & LOZADA, 2010) and Tingo Maria (WILSON et al. 2009).

***Tretogonia cibrata* MELICHAR, 1926**

Specimens 253; leg.: 2003, 2004.

Distribution: Brazil, Argentina (WILSON et al. 2009), French Guiana, Bolivia, Peru (LOZADA & ARELLANO 2008).

Determined by P. LOZADA, listed by LOZADA et al. (2007). Frequently found in Chanchamayo province (LOZADA & ARELLANO 2008). For morphology and distribution in Brazil see MEJDALANI (1998).

Additional species

Of the following genera some specimens could not be determined to species level:

Acrogonia sp., *Hygonia* sp., *Dichrophleps* sp. and *Proconia* sp.. This material should be investigated in more detail.

Additional material from ZSM

Cicadellini

***Dilobopterus lucentis* YOUNG, 1977**

Number of specimens: 1; Brazil, Tapuruquara am Rio Negro / Amazonas; 4.II.1963, leg.: C. LINDEMANN.

Distribution: Guiana, French Guiana (WILSON et al. 2009), **new record for Brazil**.

***Juliacaca scalarum* YOUNG, 1977**

Specimens: 1♀; St. Catharina, Brazil, leg.: M. WIRTNER.

Distribution: Colombia, Peru, Venezuela (WILSON et al. 2009), **new record for Brazil**.

Determined by YOUNG „*Juliacaca* sp.” (genital preparation) otherwise not distinguishable from *Microgoniella pudica*.

***Paromenia isabellina* (FOWLER, 1899)**

Specimens: 4; Ecuador, Prov. Napo, Lumbaqui, 900 m, (Ölpumpstation) 6.1.1976; leg.: W. SCHACHT.

Distribution: Mexico, Guatemala, Costa Rica, Panama, Colombia, Bolivia, Nicaragua, Venezuela, **new record for Ecuador**.

***Pawiloma rubra* CAVICHIOLI, 1998**

Specimens: 2; Guatemala, Verapaz, San Cristobal, Quixal, 23.IV.80; leg.: H. & L. FREUNDE.

Distribution: Belize, **new record for Guatemala**.

Proconiini

***Aulacizes quadripunctata* (GERMAR, 1821)**

Specimens: 1; Paraguay, Independencia, 10.X.1951 (name of collector not indicated).

Distribution: Brazil, Argentina, **new record for Paraguay**.

See MEJDALANI et al. (2006) for further details, taxonomy and images.

***Zyzzogeton viridipennis* (LATREILLE, 1811) (Fig. 26)**

Specimens: 1; Ecuador, Pr. Imbabura, Cojapi bei Carolina, 900 m, 18.-20.12.1975, leg.: W. SCHACHT.

Distribution: Colombia and Peru (FREYTAG & SHARKEY 2002, WILSON et al. 2009), **new record for Ecuador**.



19



20



21



22



23



24



25



26

Figs 19-26: 19. *Soosiulus interpolis* (Length: 6,8 mm); 20. *Desamera intersecta* (14,3 mm); 21. *Omagua fitchii* (16,5 mm); 22. and 23. *Propetes schmidti* (13,4 mm); 24. *Pseudometopia amblardii* (10,9 mm); 25. *Raphirhinus phosphoreus* (18,6 mm).; 26. *Zyzzogeton viridipennis* (head, frontal).

Discussion

This list of species is a contribution to the exploration of the fauna of Panguana. Our contribution lists 65 species of Cicadellinae, with 52 species identified to species level, but certainly there are more species to be found. On the one hand we could not yet determine all material and on the other hand more species may be collected in the future. It is quite obvious that also some new species may be expected there, but we are not able to describe new species of Cicadellinae since this has to be done by experts.

An interesting point is that a good deal of species (8 Cicadellini, 3 Proconiini) are represented by single specimens, indicating that several species occur at very low densities. Common species are predominantly those inhabiting pastures and are also common on other places on crops, e.g. *Catagonalia lunata*, *Hortlesia similis*, *Macugonalia moesta*, and *Tretogonia cibraria*. Other species which are reported to be common in agricultural areas e.g. *Scopogonalia sublivacea*, *Oragua partitula* and *Acrogonia terminalis* are only rarely found in Panguana, possibly because collectors have been active mainly in forests and less on pastures.

Some of the species reported here are obviously rather rare, e.g. *Amblyscarta trinotata*, *Julia cinctella*, *Propetes schmidti*, or *Lissoscarta nipata*. These species have been mainly known from their original description and are now reported from Panguana.

Most species were collected during field trips in 2003 and 2004. This is probably due to the efforts of Tanja KOTHE who collection manager of ZSM Hemiptera section at that time.

Acknowledgements

We are very grateful to Dr. Juliane DILLER (ZSM), who is the head of the field station Panguana, organized all field trips and runs the station. We thank all colleagues who collected cicadas and the family MODENA who took care of the scientists in Panguana. Thanks are due to the colleagues and authorities in Peru, Museo de Historia Natural de la Universidad National Mayor de San Marcos, Lima, and Dirección General Forestal y de Fauna Silvestre (DGFFS, formerly INRENA) for cooperation and permissions. We are also thankful to Margaretha and Siegfried STOCKER, Hofpfisterei (Munich), for financial support of the field station Panguana. We thank Mike WILSON for his invaluable website and help. Special thanks are due to Pedro W. LOZADA for his valuable cooperation.

Literature

- BASSET, Y., CHARLES, E., HAMMOND, D. S. & V. K. BROWN 2001: Short-term effects of canopy openness on insect herbivores in a rain forest in Guyana – Journal of Applied Ecology **38**, 1045-1058.
- CARVALHO, R. A., MEJDALANI, G. & D. M. TAKIYA 2011: Phylogenetic placement and taxonomy of the Neotropical sharpshooter genus *Desamera* YOUNG, with description of its sister group, *Ciccamera* gen. nov. (Hemiptera: Cicadellidae: Cicadellinae) – Systematics and Biodiversity **9**: 59-75.
- CAVICHIOLI R. R. & D. M. TAKIYA 2012: Description of a new species of *Wolfiania* and new records of *Rotigonalia* (Hemiptera: Cicadellidae: Cicadellinae) from the state of Amazonas, Brazil. – Zoologia **29** (1): 85-88.
- COSTA, J. F. & P. W. LOZADA 2010: Lista de Cigarritas (Hemiptera: Cicadellidae) de Cusco, Perú. – Revista Peruana de Biología **17** (3), 303-316.
- DELLAPE, G., LOGARZO, G. A., VIRLA, E. G., & S. L. PARADELL 2011: New Records on the Geographical Distribution of South American Sharpshooters (Cicadellidae: Cicadellinae: Proconiini) and their Potential as Vectors of *Xylella fastidiosa* – Florida Entomologist **94**, 364-366.
- DELLAPE, G., J. P. BOUVET & S. L. PARADELL 2013: Diversity of Cicadomorpha (Hemiptera: Auchenorrhyncha) in Citrus Orchards in Northeastern Argentina – Florida Entomologist **96**: 1125-1134.
- DIAZ, W. & J. ZAMORA 2001: Insectos del Algodonero de los Departamentos de San Martín y Ucayaly. – Área de Entomología. SENSA 1-12.
- DILLER, J. & E.-G. BURMEISTER 2007: Panguana – ein peruanischer Vogelname für eine bayerische Urwaldforschungsstation. – Aviso **1**/2007: 46-49.
- FREYTAG, P. H. & M. J. SHARKEY 2002: A preliminary list of the leafhoppers (Homoptera: Cicadellidae) of Colombia. – Bioita Colombiana **3** (2) 235-283.
- HENZE, G. 2011: Paradies Panguana. – Bild der Wissenschaft, 2011 (8), 36-39.
- HORNBURG, M. 2014: Taxonomic synopsis of the genera *Euhylaeogena* HESPENHEIDE, 2014 and *Hedwigella* OBENBERGER, 1941 (Coleoptera: Buprestidae) for South America with the description of eight new species. – Mitteilungen der Münchener Entomologischen Gesellschaft **104**, 5-29.

- LOZADA, P. W. 1993: Notas sobre Cicadellidae (Homoptera) en plantas forrajeras de Loreto, Perú. – Revista peruana de Entomología **35**: 24-26.
- LOZADA, P. W. 2010: Neotropical Cicadellinae (Hemiptera: Cicadellidae) in the Staatliches Museum für Naturkunde Stuttgart. – Stuttgarter Beiträge zur Naturkunde, A, Neue Serie **3**: 103-110.
- LOZADA, P. W., SCHÖNITZER, K. & T. KOTHE 2007: Las especies de „Cigarritas” (Hemiptera: Cicadellidae) de Panguana, Peru. – XLIX Convenión Nacional de Entomología, Lambayeque - Peru, p. 16.
- LOZADA, P. W. & G. A. ARELLANO CRUZ 2008: Lista preliminar comentada de las „Cigarritas” (Insecta: Hemiptera: Cicadellidae) de Chanchamayo y Satipo, Peru – Ecología Aplicada **7**: 117-122.
- MCKAMEY, S.H. 2007: Taxonomic Catalogue of the Leafhoppers (Membracoidea). Part 1. Cicadellinae. – Memoirs of the American Entomological Institute **78**: 1-394.
- MEJDALANI G. 1998: Morfologia externa dos Cicadellinae (Homoptera, Cicadellidae): Comparação entre *Versigonalla ruficauda* (WALKER) (Cicadellini) e *Tretogonia cibrata*. MELLCHAR (Proconiini), com notas sobre outras espécies e análise da terminologia – Revista Brasileira de Zoologia **15** (2): 451-544.
- MEJDALANI, G., TAKIYA, D. M. & R. A. CARVALHO 2006: Notes on Neotropical Proconiini (Hemiptera: Cicadellidae: Cicadellinae), IV: Lectotype designations of *Aulacizes* AMYOT & AUDINET-SERVILLE species described by GERMAR and revalidation of *A. reythrocephala* (GERMAR, 1821). – Arthropoed Systematics & Phylogeny **64**, 105-111.
- MELICHAR, L. 1951: Monographie der Cicadellinen. V. – Annales Historico-Naturales musei Nationalis Hungarici **42** (N.S. 1): 72-112.
- OTT, A. P., AZEVEDO-FILHO, W. S., FERRARI, A. & G. S. CARVALHO 2006: Abundância e sazonalidade de cigarrinhas (Hemiptera, Cicadellidae, Cicadellinae) em vegetação herbácea de pomar de laranja doce, no município de Montenegro, Estado do Rio Grande do Sul, Brasil. – Iheringia, Série Zoologia, **96**: 425-429.
- TAKIYA, D. M., MEJDALANI G. & M. FELIX 1999: Dual-mimicry of Wasps by the Neotropical Leafhopper *Propetes schmidti* MELICHAR with a Description of its Female (Hemiptera: Cicadellidae: Cicadellinae) – Proceedings of the Entomological Society of Washington **101**: 722-728.
- TAKIYA, D.M., MCKAMEY, S. H. & R. R. CAVICHIOLI 2006: Validity of *Homalodisca* and of *H. vitripennis* as the Name for Glassy-Winged Sharpshooter (Hemiptera: Cicadellidae: Cicadellinae) – Annals of the Entomological Society of America, **99**: 648-655.
- WILSON M. R., TURNER, J. A. & MCKAMEY, S. H. 2009: Sharpshooter Leafhoppers of the World (Hemiptera: Cicadellidae subfamily Cicadellinae). Amgueddfa Cymru – National Museum Wales (link see below).
- YOUNG, D. A. 1968: Taxonomic study of the Cicadellinae (Homoptera: Cicadellidae). Part 1. Proconiini. – United States National Museum Bulletin **261**, 287 pp.
- YOUNG, D. A. 1977: Taxonomic study of the Cicadellinae (Homoptera: Cicadellidae). Part 2. New World Cicadellini and the genus Cicadella. – North Carolina Agricultural Experiment Station Technical Bulletin **239**, 1135 pp.
- ZAMOL, K. M. R. & M. de MENEZES 1982: Lista preliminar dos cicadelídeos (Homoptera, Cicadellidae) do Brasil. – Inheringia Série Zoologia (**61**), 8-65.

Weblinks:

- SPDA: <http://www.conservamospornaturaleza.org/area/panguana/> (accessed September 8, 2014).
- Sharpshooter Leafhoppers of the World: <http://naturalhistory.museumwales.ac.uk/Sharpshooters> (accessed: August 22, 2014).

Addresses of the authors:

- | | |
|--|---|
| Klaus SCHÖNITZER
Zoologische Staatssammlung München
Münchhausenstr. 21
81247 München
Germany
schoenitzer@zsm.mwn.de | Wieland FEUERABENDT
Penzinger Straße 12
86940 Schwifting
Germany
wieland@posteo.de |
|--|---|

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Mitteilungen der Münchener Entomologischen Gesellschaft](#)

Jahr/Year: 2014

Band/Volume: [104](#)

Autor(en)/Author(s): Schönitzer Klaus, Feuerabendt Wieland

Artikel/Article: [The sharpshooters of Panguana \(Peru\) \(Auchenorrhyncha, Cicadellidae, Cicadellinae\) 121-132](#)