

## Abstract\*

**Interspecies hierarchy and social control in ant communities (Hymenoptera: Formicidae)**

Zhanna REZNIKOVA

Prof. Dr. Zhanna Reznikova, Institute for Animal Systematics and Ecology, Siberian Branch, Russian Academy of Sciences, Frunze 11 ul., 630091, Novosibirsk, Russia. E-mail: zhanna@reznikova.net

Myrmecol. News 10: 119

Interspecific hierarchy in ant communities was originally described by KACZMAREK (1953) who coined the terms "dominants", "subdominants" and "influent". However, his concept was not supported experimentally, but rather was based on records achieved by catching with an entomological net. This idea was developed by PISARSKI (1973) and GALLÉ (1975, 1978). The first experimental demonstration of interspecific hierarchy was made by REZNIKOVA (1980), based on the method of baiting observations. The role of behavioural plasticity in sophisticated inter-relations between dominating *Formica* and subdominating *Serviformica* species was firstly studied experimentally by STEBAEV & REZNIKOVA (1972) and REZNIKOVA (1975, 1982) by manipulations with limited food resources. The hierarchical structure of ant communities was later investigated by many myrmecologists (VEPSÄLÄINEN & PISARSKI 1982, ROSENGREN 1986, SAVOLAINEN & al. 1989, PALMER & al. 2000, SANDERS & GORDON 2003, GIBB & HOCHULI 2004, and others). Here, I summarize long-term data on the concept of "interspecific social control" in hierarchically organized ant assemblages that was first developed on the basis of transplantation experiments (REZNIKOVA 1994, 1999).

The original data on the zonal and biotopical distribution of 70 ant species in the southern Western Siberian Plain and adjacent regions were analysed. The results showed that, against a background of a considerable diversity of ant fauna, model communities dominated by *Formica pratensis* RETZIUS, 1783 retain constant species composition and the quantitative ratio of nests over the area from the northern forest steppe to the southern boundary of the steppe zone. In such coadaptive complexes (sensu: DLUSSKYI 1981), stable interspecies relationships and characteristic management of density regulation develop. We conducted field transplantation experiments in which some ants' nests were removed or replaced. In other series of experiments, we manipulated with population densities and sizes of ant colonies by means of adopted pupae. Both series of experiments were carried out in different landscapes where the ant communities were dominated by different species: *Formica polyctena* FÖRSTER, 1850, *F. pratensis* or *F. uralensis* RUZSKY, 1895. A new form of interspecies interaction in ant communities has been revealed and named "interspecies social control": ants of dominant species actively regulate the level of dynamic density of subdominant species and exterminate "superfluous" individuals. Comparative experimental data revealed the threshold values of the dynamic density of subordinate species at which their coexistence with dominants is still possible. Supplemental laboratory experiments enabled us to suggest that the fine management of the biotic relationship in ant communities is based on the ability of the dominating species to estimate numbers of encoun-

ters with individuals of subordinate species fairly accurately. – Elaborated within the frame of a HAS-RAS Academic Exchange Project and supported by the RFBR (05-04-48104).

**References**

- DLUSSKYI, G. 1981: Ants of deserts. – Nauka, Moscow, 230 pp. (in Russian)
- GALLÉ, L. 1975: Factors stabilizing the ant populations (Hymenoptera, Formicidae) in the grass associations of the Tisza Basin. – Tiscia 10: 61-66.
- GALLÉ, L. 1978: Dispersion of the nests of ant species (Hymenoptera, Formicidae). – Acta Biologica Szeged 24: 105-109.
- GIBB, H. & HOCHULI, D.F. 2004: Removal experiment reveals limited effects of a behaviourally dominant species on ant assemblages. – Ecology 85: 648-657.
- KACZMAREK, W. 1953: Badania nad zespolani mrowek lesnych. – Ekologia Polska 1: 69-96.
- PALMER, T.M., YOUNG, T.P., STANTON, M.L. & WENK, E. 2000: Short-term dynamics of an acacia-ant community in Laikipia, Kenya. – Oecologia 123: 425-435.
- PISARSKI, B. 1973: Les principes d'organisation des colonies polycaliques des fourmis. – Proceedings of the VII Congress of the IUSSE: 311-316.
- REZNIKOVA, J. 1982: Interspecific communication among ants. – Behaviour 80: 84-95.
- REZNIKOVA, ZH. 1975: Non-antagonistic relations between ant species occupying similar ecological niches. – Zoologicheskii Zhurnal 54: 1020-1031. (In Russian with English summary)
- REZNIKOVA, ZH. 1980: Interspecific hierarchy in steppe ants. – Zoologicheskii Zhurnal 59: 1168-1176. (In Russian with English summary)
- REZNIKOVA, ZH. 1994: The original pattern of management in ant communities: Interspecies social control. – Progress to meet the Challenge of Environmental Change. Manchester, UK, p. 27.
- REZNIKOVA, ZH. 1999: Ethological mechanisms of population density control in coadaptive complexes of ants. – Russian Journal of Ecology 30: 187-192.
- ROSENGREN, R. 1986: Competition and coexistence in an insular ant community – a manipulation experiment (Hymenoptera: Formicidae). – Annales Zoologici Fennici 23: 297-302.
- SANDERS, N.J. & GORDON, D.M. 2003: Resource-dependent interactions and the organization of desert ant communities. – Ecology 84: 1024-1031.
- SAVOLAINEN, R., VEPSÄLÄINEN, K. & WUORENINNE, H. 1989: Ant assemblages in the taiga biome: testing the role of territorial wood ants. – Oecologia 81: 481-486.
- STEBAEV, I.V. & REZNIKOVA, J. 1972: Two interaction types of ants living in steppe ecosystems in South Siberia. – Ekologia Polska 20: 103-109.
- VEPSÄLÄINEN, K. & PISARSKI, B. 1982: Assembly of island ant communities. – Annales Zoologici Fennici 19: 327-335.

\* Based on a presentation at the 2<sup>nd</sup> Central European Workshop of Myrmecology at Szeged, Hungary, 17 - 19 May 2007