

## Worldwide spread of the longhorn crazy ant, *Paratrechina longicornis* (Hymenoptera: Formicidae)

James K. WETTERER



### Abstract

The longhorn crazy ant, *Paratrechina longicornis* (LATREILLE, 1802), is a ubiquitous agricultural and household pest throughout much of the tropics and subtropics and a pervasive indoor pest in temperate areas. I compiled specimen records of *P. longicornis* from > 2100 sites worldwide and analyzed historical, ecological, and evolutionary information to evaluate its known distribution, geographic origin, and potential future spread. I documented the earliest known *P. longicornis* records for 181 geographic areas (countries, major islands, island groups, US states, and Canadian provinces), including many for which I found no previously published records, e.g., Anguilla, Argentina, Austral Islands, Bahrain, British Indian Ocean Territory, Cayman Islands, Comoro Islands, Dominica, East Timor, El Salvador, Gambia, Ghana, Gibraltar, Guatemala, Iles Eparses, Kenya, Maldives, Mali, Montserrat, Nepal, Nevis, Pakistan, St. Kitts, St. Lucia, Turks and Caicos Islands, Tuvalu, Maryland, Missouri, and Virginia. *Paratrechina longicornis* is arguably the most broadly dispersed of all ant species, distributed widely across the Old World and New World in both the northern and southern hemispheres. The successful worldwide spread of *P. longicornis* relates to its ability to flourish in highly disturbed and artificial environments, including ships at sea. Whereas *P. longicornis* records are most frequent near major commercial waterways, I expect that with the increasing importance of air transport, *P. longicornis* will spread to more inland areas as well. Many *P. longicornis* records, including most if not all records from temperate latitudes, come from inside or near buildings. Because *P. longicornis* can live indoors anywhere that humans live, there is no limit to the latitude where it could thrive. The worldwide distribution records of *P. longicornis* and two species-specific symbionts provide ambiguous clues to the original native range of *P. longicornis*. However, the distributions of three closely related *Paratrechina* species offer good evidence that *P. longicornis* is native to Southeast Asia and Melanesia.

**Key words:** Biogeography, biological invasions, worldwide distribution, exotic species, Formicidae, invasive species, tramp ants.

Myrmecol. News 11: 137-149 (online 29 June 2008)

ISSN 1994-4136 (print), ISSN 1997-3500 (online)

Received 17 April 2008; revision received 30 April 2008; accepted 5 May 2008

Prof. Dr. James K. Wetterer, Wilkes Honors College, Florida Atlantic University, 5353 Parkside Drive, Jupiter, FL 33458, USA. E-mail: [wetterer@fau.edu](mailto:wetterer@fau.edu)

### Introduction

Tramp ants are species that commonly are spread by human commerce and associate with human disturbance. Perhaps the most widespread of all tramp ants is the longhorn crazy ant, *Paratrechina longicornis* (LATREILLE, 1802). *Paratrechina longicornis* is a ubiquitous household and garden pest throughout much of the tropics and subtropics and is a pervasive indoor pest in temperate areas. In addition, *P. longicornis* can be a significant agricultural pest, enhancing populations of phloem-feeding Hemiptera, such as mealybugs, scale insects, and aphids. Hemiptera cause damage by sapping plants of nutrients and increasing the occurrence of diseases, including viral and fungal infections. Although *P. longicornis* is usually rare or absent in intact natural environments, it can be very common in disturbed and semi-natural environments. For example, FORYS & ALLEN (2005) found that *P. longicornis* was by far the most widespread exotic ant in the lower Florida Keys. *Paratrechina longicornis* has long been known as an important greenhouse pest, in both temperate and tropical regions (e.g., France: NYLANDER 1856, Sri Lanka: MOTSCHOUJSKY

1863), where it can reach enormously high population levels. *Paratrechina longicornis* has even invaded Biosphere 2, a 1.3-hectare greenhouse structure in the Arizona desert used as a microcosm for studying ecological interactions and global change. WETTERER & al. (1999) found that virtually all ants (> 99.9 %) coming to bait in Biosphere 2 were *P. longicornis*. WETTERER & al. (1999) observed *P. longicornis* foragers feeding almost exclusively on the sugary excretions (honeydew) produced by vast numbers of Hemiptera found on many plants. The only invertebrates thriving in Biosphere 2, besides *P. longicornis* and phloem-feeding Hemiptera, were either species with effective defenses against ants (well-armored isopods and millipedes) or tiny subterranean species that can escape ant predation (mites, thief ants, and springtails). A convergent pattern of biodiversity occurs in disturbed tropical and subtropical ecosystems dominated by tramp ants (e.g., COLE & al. 1992). Here, I document the known distribution of *P. longicornis*, and evaluate where it may have originated and where it may spread in the future.

Determining the native and exotic ranges of pest species can be useful in management strategies, such as searching for appropriate biocontrol agents. When evaluating the native and exotic ranges of a species, researchers may consider a spectrum of distributional, historical, evolutionary, ecological, and genetic information (e.g., see EMERY 1893, WASMANN 1905a, b, CHAPMAN & CARLTON 1991). Evidence considered indicative of a species' native range includes older records largely confined to a single continuous geographic region, occurrence in inland native communities, high genetic diversity, co-occurrence of species-specific symbionts, and proximity to the ranges of closely related species. In contrast, evidence indicative of a species' exotic range includes the sudden appearance and spread of the species through an area discontinuous with other known populations, occurrence exclusively in coastal and highly disturbed environments, low genetic diversity due to a founder effect, absence of species-specific symbionts, and geographic isolation from closely related species. In addition, the absence of natural enemies in a species' exotic range may result in population explosions. Finally, some tramp ant species (e.g., *Linepithema humile* (MAYR, 1868) and *Solenopsis invicta* BUREN, 1972) show extensive unicoloniality within their exotic range, where colonies spread over a broad geographic area show no intraspecific aggression, behaving as if they all belonged to a single colony (e.g., see WETTERER & WETTERER 2006).

EMERY (1893) evaluated the geographic origins of several widespread ant species, but was unable to discern an original homeland of *P. longicornis*, because it was already very broadly dispersed in tropical areas around the world. WASMANN (1905b) speculated on the geographic origin of *P. longicornis* and two myrmecophiles that live in *P. longicornis* nests: *Coluocera maderae* WOLLASTON, 1854 (Coleoptera: Lathridiidae) and *Myrmecophilus americanus* SAUSURE, 1877 (Orthoptera: Myrmecophilidae). WASMANN (1905b) noted that *P. longicornis* and its symbionts were widespread in India and neighboring parts of Southeast Asia, whereas in other parts of the world *P. longicornis* was known only from coastal areas along major trade routes used by Portuguese and Spanish ships and the two symbionts were rarely encountered. From this, WASMANN (1905b) concluded that *P. longicornis* and its symbionts probably originated in the Indian region. WHEELER (1910) wrote that WASMANN (1905b) offered "good reasons for believing that the original home of this ant is India, and that it has been carried to all parts of the tropics in ships." Later, WHEELER (1919) wrote with apparent confidence that *P. longicornis* "is of Indomalaysian origin." WILSON & TAYLOR (1967) wrote that *P. longicornis* "probably originated within the Old World tropics, perhaps specifically in southeastern Asia or Melanesia." BERNARD (1968) and DLUSSKY (1994) also judged *P. longicornis* to be of Asian origin, and in earlier papers I have considered it an Asian species (e.g., WETTERER 2002, WETTERER & VARGO 2003). In contrast, FOWLER & al. (1994) listed *P. longicornis* as originating in tropical Africa and HOLWAY & al. (2002) also called this species African. KENNE & al. (2005) premised their study of the predatory behavior of *P. longicornis* in Cameroon on the assumption that *P. longicornis* was native to West Africa. Thus, whereas there is general agreement that *P. longicornis* originated in the Old World tropics, there is disagreement as to the subregion. I there-

fore evaluated multiple lines of evidence in an effort to discern the native range of *P. longicornis*.

### Common names

By far the most-used common English name for *P. longicornis* is the "crazy ant," due to its rapid, seemingly erratic movements. Unfortunately, this common name is also often used for many other ant species, including all *Paratrechina* species, many of which are also widespread tramp species, e.g., *P. bourbonica* (FOREL, 1886), *P. flavipes* (SMITH, 1874), *P. fulva* (MAYR, 1862), *P. guatemalensis* (FOREL, 1885), *P. jaegerskioeldi* (MAYR, 1904), *P. pubens* (FOREL, 1893), and *P. vividula* (NYLANDER, 1846). Recently, some authors have begun to use the name "black crazy ant" to distinguish *P. longicornis* from another widespread pest species often called the "crazy ant" or the "yellow crazy ant," *Anoplolepis gracilipes* (SMITH, 1857) (e.g., HOFFMANN & O'CONNOR 2004). Unfortunately, the common name "black crazy ant," does not refer to a unique character and may better describe other *Paratrechina* species, e.g., *P. bourbonica*, which is darker in color than *P. longicornis*. Additional common names for *P. longicornis* include the long-horned ant (NEUMANN 1993 in HARRIS & BERRY 2005), the hairy ant (CHIN 1998), and the slender crazy ant (DEYRUP & al. 2000). I propose that "longhorn crazy ant" is a more descriptive common name, derived from a primary distinguishing characteristic and its Latin name (*longicornis* = longhorn). This name is unlikely to be confused with the common name of any other ant species.

### Methods

I documented the worldwide range of *Paratrechina longicornis* using both published and unpublished records. Unlike many other major pest ant species, *P. longicornis* is very easy to identify correctly, and therefore identifications in the literature are highly reliable. I obtained unpublished site records from museum specimens in the collections of the American Museum of Natural History (AMNH), the Archbold Biological Station (ABS), the British Natural History Museum (BMNH), the Museum of Comparative Zoology (MCZ), the National Museums Liverpool (NML), the Oxford University Natural History Museum (ONHM), the Smithsonian Institute's National Museum of Natural History (SI), Tel Aviv University (TAU), and the University of Guam Museum (UGM). I also examined specimens at the MCZ on loan from Museum Paris (MP). In addition, I used on-line databases with collection data of specimens in the Australian National Insect Collection (ANIC), the Essig Museum at UC Berkeley (UCB), and the California Academy of Sciences (CAS). I received unpublished site records and specimens from many sources: O. Bourquin (Mariana Islands), H. Cagniant (Algeria, Morocco), R. Clouse (Micronesia), M. Deyrup (Florida), G. Dlussky (Samoa), J. Endeman (Antigua, Ascension Island, Barbados, British Indian Ocean Territory, Colombia, East Timor, Grenada, Indonesia, the Maldives, Peru, St Helena), X. Espadaler (Brazil, Canary Islands, India, Malaysia, Nepal, Spain, Thailand), J. Fellowes (China), A. Francoeur (Canada), M. Haji (Pakistan), S. Hedges (Bahamas), G. Heller (Canary Islands, Germany), W. Jaitrong (Thailand), M.A. Kabir (Bangladesh), G. Knight (Cape Verde), J. Lewallen (Vietnam), M. Lush (Gambia), J. MacGown (Florida, Louisiana, Mississippi), P.D. Rajan (India), A. San Juan (Mex-

ico), M. Vonshak (Israel), D. Ward (Fiji), and J. Woinarski (Australia). Finally, I collected *P. longicornis* specimens on numerous Pacific, Atlantic, and Caribbean islands (e.g., WETTERER 1997, 1998a, 2002, WETTERER & VARGO 2003, WETTERER & WETTERER 2004, WETTERER & al. 2004, 2006), in Costa Rica (e.g., WETTERER 1998b), and Florida (e.g., WETTERER & O'HARA 2002, WETTERER & MOORE 2005).

I obtained geographic coordinates for sites from published references, from specimen labels, or I looked up the coordinates. For older references and specimens, many site names, particularly in Asia, were no longer in use or are now spelled differently and I searched, not always successfully, to determine current names.

Many authors reported that *P. longicornis* was found in an area, but presented no specific data. For example, BINGHAM (1903) wrote that *P. longicornis* in India, Sri Lanka, and Burma was found "throughout our limits and spread over all tropical countries," but did not provide a single site record. If a record listed a geographic region rather than a "point locale," and I had no other record for this region, I used the coordinates of the largest town within the region or, in the case of small islands and natural areas, the center of the region. If one source had many sites less than 10 - 20 km apart (e.g., HUDDLESTON & FLUKER 1968; my own collections on Pacific and Caribbean islands), I often did not plot every site. I did not map records of *P. longicornis* intercepted in transit by quarantine inspectors.

I documented the range of two symbionts that inhabit *P. longicornis* nests and three closely related *Paratrechina* species (*Paratrechina pallida* DONISTHORPE, 1947, *Paratrechina otome* TERAYAMA, 1999, and *Paratrechina yambaru* TERAYAMA, 1999) using published and unpublished records. In addition, in May - July 2007, while surveying ants on seven West Indian islands (Antigua, Aruba, Barbuda, Montserrat, Nevis, St. Kitts, and St. Martin), I looked for symbionts in every large *P. longicornis* nest I encountered.

## Results

I plotted records of *P. longicornis* from > 2100 sites around the world, based on information in > 400 published articles, from thousands of museum specimens, and specimens I collected from hundreds of sites (Fig. 1). In total, I found *P. longicornis* records for 181 geographic areas (i.e., countries, island groups, US states, and Canadian provinces), including many from geographic areas for which I found no previously published records, e.g., Anguilla, Argentina, Austral Islands, Bahrain, British Indian Ocean Territory (BIOT), Cayman Islands, Comoro Islands, Dominica, East Timor, El Salvador, Gambia, Ghana, Gibraltar, Guatemala, Iles Eparses, Kenya, Maldives, Mali, Montserrat, Nepal, Nevis, Pakistan, St. Kitts, St. Lucia, Turks and Caicos Islands, Tuvalu, Maryland, Missouri, and Virginia (Tabs. 1 - 7). *Paratrechina longicornis* has not yet been recorded from only a limited number of tropical maritime countries (Africa: Angola, Benin, Djibouti, Gabon, Gambia, Guinea-Bissau, Ivory Coast, Liberia, Mauritania, Mozambique, Namibia, Togo, and Western Sahara; Asia: Brunei and Cambodia; Oceania: Nauru; Neotropics: none). By 1900, *P. longicornis* was already known from a diversity of locales worldwide (Tabs. 1 - 7).

In all parts of its range, *P. longicornis* occurs primarily in highly disturbed coastal and urban environments. For

Tab. 1: Earliest known records for *Paratrechina longicornis* from Asia, the Middle East, and neighboring islands. Unpublished records include collector, museum source, and site. BIOT = British Indian Ocean Territory. BMNH = Natural History Museum, London. SI = Smithsonian Institution. MCZ = Museum of Comparative Zoology. + = no known published records.

|                      | Earliest record                              |
|----------------------|--|
| India                | ≤ 1851 (JERDON 1851)                         |
| Sri Lanka            | 1858 (MAYR 1862)                             |
| Indonesia            | ≤ 1866 (MAYR 1867)                           |
| Palestine/Israel     | ≤ 1880 (ANDRÉ 1881)                          |
| Syria                | ≤ 1880 (ANDRÉ 1881)                          |
| Yemen                | ≤ 1880 (EMERY 1881)                          |
| Burma/Myanmar        | ≤ 1887 (EMERY 1889)                          |
| Papua New Guinea     | ≤ 1901 (FOREL 1901b)                         |
| Taiwan               | ≤ 1909 (WHEELER 1909)                        |
| Philippines          | 1913 (STITZ 1925)                            |
| Malaysia             | ≤ 1914 (VIEHMEYER 1915)                      |
| Iraq                 | 1917 (DONISTHORPE 1918)                      |
| Vietnam              | ≤ 1920 (SANTSCHI 1920a)                      |
| China                | ≤ 1921 (WHEELER 1921)                        |
| Japan                | ≤ 1924 (TERANISHI 1924 in ONOYAMA 1980)      |
| Thailand             | 1928 (T.D. Cockerell, SI): Nan               |
| Singapore            | 1931 (collector unknown, BMNH): site unknown |
| Christmas Island     | 1933 (DONISTHORPE 1935)                      |
| + Bahrain            | 1951 (N.A. Weber, MCZ): site unknown         |
| + Nepal              | 1956 (E.I. Coher, MCZ): Amlekhganj           |
| Lebanon              | ≤ 1970 (CAGNIANT 1970)                       |
| + BIOT               | 1971 (Hutson, BMNH): Diego Garcia Island     |
| Saudi Arabia         | 1979 (COLLINGWOOD 1985)                      |
| Oman                 | ≤ 1985 (COLLINGWOOD 1985)                    |
| United Arab Emirates | 1995 (COLLINGWOOD & al. 1997)                |
| Iran                 | ≤ 2000 (ALIPANAH 2000 in PAKNIA & KAMI 2007) |
| + Maldives           | 2004 (J. Endeman, MCZ): Hulhule Island       |
| Bangladesh           | 2005 (HANNAN 2007)                           |
| + East Timor         | 2005 (J. Endeman, MCZ): Dili                 |
| + Pakistan           | 2007 (S. & Z. Valliani, MCZ): Karachi        |

Tab. 2: Earliest known records for *Paratrechina longicornis* from Africa and neighboring islands. Abbreviations as in Tab. 1, and MP = Museum Paris. NML = National Museums Liverpool.

|                      | Earliest record                           |
|----------------------|---|
| Senegal              | ≤ 1802 (LATREILLE 1802)                   |
| Madeira              | ≤ 1859 (ROGER 1859)                       |
| Guinea               | ≤ 1876 (MAYR 1876)                        |
| Rodrigues Island     | ≤ 1879 (SMITH 1879)                       |
| Egypt                | ≤ 1880 (ANDRÉ 1881, EMERY 1881)           |
| Madagascar           | ≤ 1891 (FOREL 1891)                       |
| Canary Islands       | ≤ 1893 (FOREL 1893a)                      |
| Réunion              | ≤ 1895 (FOREL 1895)                       |
| Seychelles           | ≤ 1897 (FOREL 1897a)                      |
| Congo (Zaire)        | ≤ 1901 (FOREL 1901a)                      |
| Eritrea              | 1906 (FOREL 1910)                         |
| Sudan                | ≤ 1911 (KARAVAIIEV 1911)                  |
| Central African Rep. | 1911 (STITZ 1916)                         |
| Tanzania             | 1912 (SANTSCHI 1914b)                     |
| Somalia              | 1913 (MENOZZI 1926)                       |
| Nigeria              | ≤ 1914 (SANTSCHI 1914a)                   |
| Cameroon             | 1916 (G. Schwab, MCZ): Batanga            |
| Sao Tome & Principe  | ≤ 1920 (SANTSCHI 1920b)                   |
| South Africa         | ≤ 1922 (ARNOLD 1922)                      |
| Morocco              | ≤ 1929 (SANTSCHI 1929)                    |
| Ethiopia             | ≤ 1933 (SANTSCHI 1935)                    |
| Mauritius            | 1942 (R. Mamet, BMNH): Beau Bassin        |
| Equatorial Guinea    | ≤ 1940 (MENOZZI 1942)                     |
| Cape Verde           | 1953 (O. Wellenius, NML): Ribeira Juliao  |
| Algeria              | 1962 (H. Cagniant, pers comm.): Algiers   |
| + Iles Eparses       | 1964 (R. Legendre, MP): Europa Island     |
| + Ghana              | 1968 (C. Collingwood, BMNH): Accra        |
| Malawi               | < 1970 (SWEENEY 1970 in MORRIS 2004)      |
| + Mali               | 1970 (P. Room, BMNH): Gao                 |
| Sierra Leone         | 1976 (GRIFFITHS 1980)                     |
| + Comoro Islands     | 1994 (Roger, MCZ): Mutasamudu             |
| + Kenya              | < 2004 (collector unknown, BMNH): Nairobi |
| + Gambia             | 2007 (M. Lush, pers. comm.): Kololi       |

Tab. 3: Earliest known records for *Paratrechina longicornis* from Australia and Pacific Ocean islands. Abbreviations as in Tab. 1, and ONHM = Oxford University Natural History Museum. UCB = Essig Museum at UC Berkeley.

|                   | Earliest record                         |
|-------------------|---|
| Samoa             | ≤ 1870 (MAYR 1870)                      |
| Australia         | ≤ 1886 (EMERY 1886)                     |
| Hawaii            | ≤ 1887 (BLACKBURN & CAMERON 1887)       |
| Galapagos         | 1906 (WHEELER 1919)                     |
| Society Islands   | 1907 (WHEELER 1908b)                    |
| Mariana Islands   | 1911 (WHEELER 1912)                     |
| New Caledonia     | ≤ 1914 (EMERY 1914)                     |
| Banaba            | 1914 ("DWP", ONHM): Ocean Island        |
| Solomon Islands   | 1916 (MANN 1919)                        |
| Fiji              | 1920 (MANN 1921)                        |
| Tonga             | 1923 (WILSON & TAYLOR 1967)             |
| Tokelau Islands   | 1924 (WILSON & TAYLOR 1967)             |
| Cook Islands      | 1925 (WILSON & TAYLOR 1967)             |
| Marquesas Isl.    | 1925 CHEESMAN & CRAWLEY 1928)           |
| New Zealand       | ≤ 1926 (GRIMMETT 1926)                  |
| Vanuatu           | 1930 (L.E. Cheesman, BMNH): Malekula    |
| Pitcairn Islands  | 1934 (WHEELER 1936)                     |
| Line Islands      | 1935 (WILSON & TAYLOR 1967)             |
| FS Micronesia     | 1935 (CLOUSE 2007)                      |
| Marshall Islands  | 1937 (CLOUSE 2007)                      |
| Phoenix Islands   | 1940-41 (VAN ZWALUWENBURG 1943)         |
| Palau             | 1952 (CLOUSE 2007)                      |
| Wake Island       | 1957 (NLH Krauss, SI): site unknown     |
| Gilbert Islands   | 1957 (NLH Krauss, MCZ): Butaritari      |
| Wallis & Futuna   | 1965 (WILSON & HUNT 1967)               |
| Niue              | ≤ 1967 (TAYLOR 1967)                    |
| Easter Island     | 1971 (WILSON 1973)                      |
| + Tuvalu          | 1976 (P. Maddison, BMNH): Vaitupu       |
| Tuamotu Islands   | 1996 (MORRISON 1997)                    |
| + Austral Islands | 2006 (P.D. Krushelnycky; UCB): Rimatara |

example, on islands in the Pacific, the Atlantic, the Caribbean, and in Costa Rica and Florida, I usually found *P. longicornis* in weedy areas and in trees at most waterfront and urban sites. A few records of *P. longicornis*, however, came from inland natural areas, including national parks and wildlife sanctuaries. For example, in Thailand, *P. longicornis* has been documented at diverse sites throughout all provinces in the country, including more than 30 national parks and wildlife sanctuaries (JAITRONG & NAB-

Tab. 4: Earliest known records for *Paratrechina longicornis* from South and Central America. Abbreviations as in Tab. 1.

|               | Earliest record                                 |
|---------------|---|
| Mexico        | ≤ 1859 (ROGER 1859)                             |
| Chile         | 1859 (MAYR 1862)                                |
| French Guiana | 1868 (RADOSZKOWSKY 1884)                        |
| Brazil        | ≤ 1876 (MAYR 1876)                              |
| Colombia      | ≤ 1876 (MAYR 1876)                              |
| Nicaragua     | 1897 (Baker, SI): Chinandega                    |
| Costa Rica    | ≤ 1899 (FOREL 1899)                             |
| Panama        | ≤ 1899 (FOREL 1899)                             |
| Belize        | 1906 (WHEELER 1907)                             |
| + Guatemala   | 1911 (W.M. Wheeler, MCZ): Zacapa                |
| Guyana        | 1911 (WHEELER 1916)                             |
| + Argentina   | 1915 (collector unknown, SI): La Plata          |
| Honduras      | 1916 (F.J. Dyer, SI): La Ceiba                  |
| Surinam       | ≤ 1932 (BÜNZLI 1935)                            |
| Venezuela     | 1935 (N.A. Weber, MCZ): Orinoco Delta           |
| Peru          | 1939 (W. Weyrauch, MCZ): Lima                   |
| + El Salvador | 1958 (L.J. Bottimer, SI): Volcan de Conchagua   |
| Ecuador       | ≤ 1969 (GUTIERREZ 1969 in WOJCIK & PORTER 2005) |
| Paraguay      | 1997 (A. Wild, pers. comm.): Asunción           |

HITABHATA 2005; W. Jaitrong, pers. comm.). *Paratrechina longicornis* has also been reported from national parks in Australia (HOFFMANN & O'CONNOR 2004; ANIC database), Florida (FERSTER & PRUSAK 1994; ABS database), Hawaii (WETTERER 1998a), India (P.D. Rajan; pers. comm.), and Vietnam (YAMANE & al. 2002, EGUCHI & al. 2004).

Many records of *P. longicornis* come from inside or near buildings, particularly greenhouses, including most, if not all records from more temperate parts of Europe and North America. The highest latitude record comes from inside the Orchid House of the Dorpat Botanical Gardens in what is now Tartu, Estonia (latitude = 58.4° N, MÜHLEN 1887). At the Smithsonian, there were *P. longicornis* specimens from Göteborg, Sweden (57.7° N, Tab. 6) labeled "Växthus" (= greenhouse). DONISTHORPE (1927) listed records from 16 sites around England (50.7 - 55.9° N), primarily from indoors. The British Museum has a number of unpublished records of *P. longicornis* collected in England, including specimens from a kitchen in Cheltenham (51.9° N, 1945, H. Donisthorpe), a bakery in Chatham (51.4° N, 1949, collector unknown), and from Redhill (51.4° N, 1906, G.E. Frisby). Other high latitude records come from greenhouses in Hamburg, Germany (53.6° N, JACOBSON 1939), the Vancouver Aquarium in British Columbia (49.3° N, NAUMANN 1994), Paris Botanical Garden greenhouses (48.9° N, NYLANDER 1856), a Zurich, Switzerland airport building (47.5° N, FREITAG & al. 2000), the basement of

Tab. 5: Earliest known records for *Paratrechina longicornis* from the West Indies. Abbreviations as in Tab. 1.

|                    | Earliest record                                   |
|--------------------|---|
| Cuba               | ≤ 1876 (MAYR 1876)                                |
| Virgin Islands     | 1878 (FOREL 1881)                                 |
| Saint Vincent      | ≤ 1892 (FOREL 1893b)                              |
| Trinidad & Tobago  | 1895 (WASMANN 1905b)                              |
| Grenada            | ≤ 1897 (FOREL 1897b)                              |
| Bahamas            | 1904 (WHEELER 1905)                               |
| Puerto Rico        | 1906 (WHEELER 1906a)                              |
| Jamaica            | 1906 (WHEELER 1911)                               |
| Haiti              | ≤ 1907 (FOREL 1907)                               |
| Martinique         | ≤ 1913 (WHEELER 1913b)                            |
| Dominican Republic | 1915 (collector unknown, MCZ): Sanchez            |
| Antigua & Barbuda  | 1918 (WHEELER 1923)                               |
| Barbados           | 1918 (WHEELER 1923)                               |
| + Montserrat       | 1934 (N.A. Weber, MCZ): site unknown              |
| + Saint Kitts      | 1934 (N.A. Weber, MCZ): site unknown              |
| + Nevis            | 1934 (N.A. Weber, MCZ): site unknown              |
| + Dominica         | no date (N.A. Weber, MCZ): site unknown           |
| + Saint Lucia      | 1935 (N.E. Box, MCZ): site unknown                |
| Dutch Leeward Is.  | 1936 (WEBER 1948)                                 |
| + Turks & Caicos   | 1953 (E.B. Haydenin, AMNH): South Caicos          |
| Guadeloupe         | ≤ 1990 (JAFJE & al. 1990)                         |
| Aruba              | ≤ 1994 (JAFJE & LATTKE 1994)                      |
| Saint Martin       | ≤ 1994 (JAFJE & LATTKE 1994)                      |
| + Anguilla         | 2006 (J.K. Wetterer, MCZ): Meads Bay              |
| + Cayman Islands   | 2008 (J.K. Wetterer, MCZ): Grand Cayman; Websters |

a convent in Sillery, Quebec (46.8° N, A. Francoeur, pers. comm.), and the Toronto Zoo in Ontario (43.7° N, CUTTEN & al. 1979). The northernmost records in the US come from Buffalo, New York (42.9° N, 1926, collector unknown, CZ), buildings in Boston (42.4° N, MARLATT 1928), and buildings in New York City (40.7° N, WHEELER 1906b). The highest latitude *P. longicornis* records in the southern hemisphere come from New Zealand, primarily from port areas (36.9 - 41.3° S, PASCOE 2002, HARRIS & BERRY 2005, HEMBRY 2005). GRIMMETT (1926) wrote, concerning Hymenoptera in the forest of Days Bay, New Zealand (41.3° S), that *P. longicornis* "is the species most commonly

Tab. 6: Earliest known records for *Paratrechina longicornis* from Europe, the Mediterranean, and Atlantic Islands. Abbreviations as in Tab. 1.

|                  | Earliest record                                    |
|------------------|--|
| France           | ≤ 1856 (NYLANDER 1856)                             |
| England          | ≤ 1859 (ROGER 1859)                                |
| Estonia          | ≤ 1887 (MÜHLEN 1887)                               |
| Netherlands      | 1909 (B. Vierbergen, pers. comm.): site unknown    |
| Azores           | 1929 (DONISTHORPE 1936)                            |
| Czech Republic   | 1937 (NOVÁK 1947)                                  |
| Germany          | ≤ 1939 (JACOBSON 1939)                             |
| + Gibraltar      | 1956 (N.L.H. Krauss, SI): site unknown             |
| Sweden           | 1957 (B. Hanson, SI): Göteborg                     |
| Ascension Isl.   | 1958 (DUFFEY 1964)                                 |
| Malta            | 1975 (SCHEMBRI & COLLINGWOOD 1981)                 |
| Italy            | ≤ 1981 (SCHEMBRI & COLLINGWOOD 1981)               |
| Greece           | ≤ 1988 (KUGLER 1988)                               |
| Bermuda          | 1990 (WETTERER & WETTERER 2004)                    |
| Saint Helena     | 1994 (ASHMOLE & ASHMOLE 2000)                      |
| Denmark          | 1996 (JENSEN & BILLE 1997 in WOJCIK & PORTER 2005) |
| Spain            | 1998 (TINAUT & AÑO 2000)                           |
| Switzerland      | 1999 (FREITAG & al. 2000)                          |
| Balearic Islands | 2004 (GÓMEZ & ESPADALER 2006)                      |

found, and in some samples constituted an overwhelming majority of the fauna," though this may have been a transient population.

### Nest symbionts

The two myrmecophilic insect species known to live almost exclusively in *P. longicornis* nests have site records from around the world: *Coluocera maderae* and *Myrmecophilus americanus*. WOLLASTON (1854) described the beetle *Coluocera maderae* collected from the nest of an unidentified ant on the Atlantic island of Madeira. *Coluocera maderae* has been subsequently reported from *P. longicornis* nests in Brazil, Burma, Cuba, Galapagos, Haiti, Hawaii, India, Taiwan, and Trinidad (WASMANN 1905b, ASSMUTH 1907, MANN 1915, 1918, ILLINGWORTH 1931, KRAUSS 1944, KISTNER 1982, PECK 1993, TARI 2007). In India, *C. maderae* has also been reported from the nests of *Paratrechina indica* (FOREL, 1894) and an unidentified *Pheidole* species (WASMANN 1905b).

SAUSSURE (1877) described the ant cricket *Myrmecophilus americanus* from an unspecified locale in Colombia, with no indication of the host ant species. Published site records of *M. americanus* come from around the world, including India, the western Indian Ocean (Madagascar, Seychelles, Réunion), Oceania (Hawaii), and the Neotro-

Tab. 7: Earliest known records for *Paratrechina longicornis* from the United States and Canada. Abbreviations as in Tab. 1.

|                  | Earliest record                                    |
|------------------|--|
| New York         | 1886 (C.V. Riley, SI): site unknown                |
| Washington DC    | 1886 (C.V. Riley, SI): site unknown                |
| + Missouri       | 1901 (collector unknown, MCZ): Shaw Botanic Garden |
| Texas            | 1905 (F.C. Pratt, SI): Dallas                      |
| Florida          | ≤ 1906 (WHEELER 1906b)                             |
| Pennsylvania     | 1906 (collector unknown, MCZ): Philadelphia        |
| Alabama          | 1910 (W.D. Pierce, SI): Mobile                     |
| Georgia          | ≤ 1913 (WHEELER 1913a)                             |
| + Maryland       | 1913 (R.L. Ball, SI): Baltimore                    |
| + Virginia       | 1915 (D.E. Finck, SI): Norfolk Navy Yard           |
| Indiana          | ≤ 1921 (DIETZ 1921)                                |
| Mississippi      | 1922 (J. MacGown, pers. comm.): Oktibbeha          |
| Massachusetts    | ≤ 1928 (MARLATT 1928)                              |
| South Carolina   | ≤ 1934 (SMITH 1934)                                |
| Oklahoma         | ≤ 1935 (SMITH 1935)                                |
| North Carolina   | ≤ 1937 (POWELL 1937 in WOJCIK & PORTER 2005)       |
| Louisiana        | 1943 (W. Buren, SI): Alexandria                    |
| Quebec           | 1963 (A. Francoeur, pers. comm.): Sillery          |
| California       | 1967 (F. Yaruss, SI): San Diego                    |
| Ontario          | 1978 (CUTTEN & al. 1979)                           |
| Illinois         | ≤ 1988 (DUBOIS & LABERGE 1988)                     |
| Arizona          | 1993 (WETTERER & al. 1999)                         |
| British Columbia | 1994 (NAUMANN 1994)                                |
| Ohio             | ≤ 1998 (HEDGES 1998)                               |
| New Mexico       | ≤ 2000 (HEDGES 2000)                               |

pics (Brazil, Colombia, Cuba, Haiti). Additional records of *M. americanus* come from Sudan, Libya, Egypt, and Israel, though some or all of these may be *Myrmecophilus cottami* CHOPARD, 1922, if this species proves to be valid (see WETTERER & HUGEL in press). In all recorded cases except one, the ant host of *M. americanus* was the longhorn crazy ant, *Paratrechina longicornis*. Finally, WETTERER & HUGEL (in press) collected *M. americanus* specimens from *P. longicornis* nests at 13 sites on six West Indian islands: Antigua, Aruba, Montserrat, Nevis, St. Kitts, and St. Martin.

### Related *Paratrechina* species

WILSON & TAYLOR (1967) proposed that *P. longicornis* is most closely related to *Paratrechina pallida*, a species known only from the lowlands of New Guinea, primarily




Fig. 1: Worldwide records of *Paratrechina longicornis*.

from intact rainforest (DONISTHORPE 1947, WILSON 1959, ROOM 1975, SNELLING 2000, unpublished records in the MCZ). TRAGER (1984) similarly concluded that: "*longicornis* and *pallida* appear to form a real species-group."

Recently, TERAYAMA (1999) described two *Paratrechina* species from grasslands and forest margins in the far south of Japan that closely resemble *P. longicornis*: *Paratrechina otome* from two of the Yaeyama Islands and *Paratrechina yambaru* from northern Okinawa Island.

### Discussion

*Paratrechina longicornis* is arguably the most broadly dispersed of any ant species, with records widely distributed across the Old World and New World in both the northern and southern hemispheres (Fig. 1). DELABIE & BLARD (2002) proposed that the ant *Hypoponera punctatissima* (ROGER, 1859) "has the widest distribution within Formicidae, being present on many oceanic and continental islands and all continental regions except Antarctica and the Orient." However, in compiling worldwide site records for > 40 major tramp ants, my analyses indicate that several ant species have broader distributions than *H. punctatissima*. Although *H. punctatissima* is widespread in Europe and on Pacific Islands, it has a very limited distribution in other parts of the world, particularly in South America (two records), Asia (one record), and Australia (no continental records). In addition, ant specimens that have been identified as *H. punctatissima* represent more than one species (S. Cover, pers. comm.; SEIFERT 2003).

I found that the only ant species that might be considered as widely distributed as *P. longicornis* is the Pharaoh ant, *Monomorium pharaonis* (LINNAEUS, 1758), a species thought to be from Africa or Asia that has long been known as an important house pest in Europe and North America. WEBER (1939) wrote that *P. longicornis* and *Monomorium pharaonis* "are probably the most common ship ants and may be expected on any vessels visiting ports in warmer

regions." In much of its range, however, *M. pharaonis* has been collected only inside buildings.

*Paratrechina longicornis* has become so widespread, in part, because it can thrive in even the most artificial environments, including ships at sea. For example, the British Museum collection included a *P. longicornis* specimen that E.S. Brown collected on a boat traveling from Mombasa, Kenya to the Seychelles Islands, and J. Endeman sent me *P. longicornis* specimens she collected on a tour boat visiting Komodo Island, Indonesia (for additional records from ships, see WHEELER 1908a, 1916, MYERS 1934, WEBER 1939, 1940, MILLER 1994). Its habit of hitchhiking in cargo has allowed *P. longicornis* to spread quickly around the world. HEDGES (1998) reported *P. longicornis* from a car driven from Florida to Indiana, and noted "this species' propensity for nesting in the soil of potted tropical plants enables it to be shipped literally anywhere in the United States." Internationally, *P. longicornis* is one of the most common ant species intercepted in cargo by quarantine officials (e.g., see LESTER 2005).

There are only a few tropical maritime nations from which *P. longicornis* has not yet been recorded. All these countries, however, have been poorly sampled for ants. For example, I found no published records for any ants at all from Nauru, the only tropical Pacific nation without *P. longicornis* records. It seems likely that even a cursory ant survey in Nauru or any tropical port city would find *P. longicornis* already present. In fact, TRAGER (1984) speculated that *P. longicornis* was so ubiquitous that: "perhaps no large city in the tropics is without this ant."

Currently, *P. longicornis* records are uncommon in many inland tropical areas, except along major waterways, such as the Nile, Congo, and Amazon Rivers (Fig. 1). With the increasing prevalence of air commerce, I expect that in the future *P. longicornis* will spread to other inland areas as well. Because *P. longicornis* can live indoors anywhere that humans live, there is no limit to the latitude where this spe-

cies could thrive, joining humans and German cockroaches (*Blattella germanica* LINNAEUS, 1767) (Blattodea: Blattellidae), as one of the most widespread and populous animal species on Earth.

### Native and exotic range

The worldwide distribution of all known *P. longicornis* site records (Fig. 1) offers ambiguous clues to where the species is native. The greatest saturation of *P. longicornis* site records comes from around the Caribbean, but this pattern is almost certainly a result of higher sampling intensity in this region, including my own collecting. In the Old World, the greatest saturation of *P. longicornis* site records is in Southeast Asia and to a lesser extent the north coast of Australia, but again, this may be an artifact of sampling intensity.

In 1900, when most of the world's ant fauna was still very poorly documented, *P. longicornis* was already known from a diversity of locales scattered across Asia, Africa, the Middle East, the Indian Ocean, Central America, South America, the West Indies, and Atlantic islands, as well as a few records in Europe and the US (Tabs 1 - 7). Although WASMANN (1905b) wrote that *P. longicornis* was more widespread in India and neighboring areas than in other parts of the world, he provided scant evidence to support this claim in terms of site records. In many parts of the world, the earliest collection date for *P. longicornis* does not appear to indicate the earliest arrival of the ant, but often corresponds to the date of the first thorough ant surveys. Thus, reconstructing the spread of *P. longicornis* out from its original native range is not possible using a chronology of historical specimen records.

The occurrence of *P. longicornis* in inland natural areas appears to offer some useful evidence about the native range of the species. The greatest number of records of *P. longicornis* in national parks and wildlife refuges come from Southeast Asia. For example, YAMANE & al. (2002) and EGUCHI & al. (2004) reported *P. longicornis* from two inland national parks in Vietnam. In Thailand, *P. longicornis* was found even in extremely remote forest reserves (JAITRONG & NABHITABHATA 2005, W. Jaitrong, pers. comm.), suggesting that it is native to the area. In contrast, in natural areas elsewhere in the world, *P. longicornis* may invade only human-impacted areas. For example, I found *P. longicornis* only once in Hawaii Volcanoes National Park, in the cooking shelter at a campground (WETTERER 1998a), a site not indicative of a native species (and, in fact, no ants are thought to be native to Hawaii). FERSTER & PRUSAK (1994) and HOFFMANN & O'CONNOR (2004) considered *P. longicornis* to be exotic to Everglades National Park, USA and Kakadu National Park, Australia, respectively.

In some ant species, intraspecific aggression is common in native populations, but largely absent in exotic populations (e.g., see WETTERER & WETTERER 2006). LIM & al. (2003), in the only published study of intraspecific aggression in *P. longicornis*, found that colonies from five sites on Panang Island, Malaysia all showed fierce intercolonial aggression. But this intercolonial aggression does not appear to provide strong evidence that *P. longicornis* is native to this region: a preliminary study of intraspecific aggression among *P. longicornis* colonies in Florida, where *P. longicornis* is certainly exotic, also found no evidence of unicoloniality (A. Corsaro & J. Wetterer, unpubl.). It is un-

clear how widespread unicoloniality is among invasive ant species.

The two myrmecophilic insects (*C. maderae* and *M. americanus*) found almost exclusively in *P. longicornis* nests have been recorded from locales scattered around the world. Thus, the simple presence of these species-specific symbionts is not indicative of the ants' native range. WASMANN (1905b) concluded that the widespread occurrence of these nest associates in India and Southeast Asia indicated that *P. longicornis* originated in this region. In contrast, in other parts of the world, the myrmecophiles seem to be common at only a few scattered locales. For example, MANN (1915) found *P. longicornis* at many sites in Haiti, but collected *C. maderae* and *M. americanus* at only a single site. Similarly, I found *M. americanus* only sporadically in *P. longicornis* colonies in the Caribbean.

The distributions of three *Paratrechina* species believed to be closely related to *P. longicornis* provide the most unambiguous evidence that *P. longicornis* is native to Papuaia (Southeast Asia and Melanesia) as proposed by WILSON & TAYLOR (1967): *P. pallida* known only from New Guinea, and *P. otome* and *P. yambaru* known only from southern Japanese islands. The limits of the native range of *P. longicornis* within Asia and Melanesia, however, remain ambiguous.

Outside of Asia, there are many *P. longicornis* populations that are almost certainly exotic. For example, populations in the New World and on mid-Atlantic islands are all very distant from the range of any closely related *Paratrechina* species. In North America and Europe, *P. longicornis* populations suddenly appeared in isolated urban areas with very extensive earliest sampling for ants, with most records coming from inside buildings. On most or all isolated islands of the eastern Pacific, *P. longicornis* is also probably exotic. In fact, WILSON & TAYLOR (1967) concluded that: "few if any ant species are native to the islands east of Rotuma, Samoa, Tonga, and New Zealand."

I found scant evidence suggesting that *P. longicornis* originated in or is native to Africa. I found relatively few records of *P. longicornis* and few (possibly no) records of its species-specific nest symbionts from continental Africa, a region distant from the range of closely related *Paratrechina* species in East Asia and New Guinea. No African *Paratrechina* species even remotely resemble *P. longicornis* (S. Cover, pers. comm.). KENNE & al. (2005) apparently considered *P. longicornis* native to West Africa simply because DEJEAN & al. (1996) found a *P. longicornis* colony in a Cameroon rainforest, though I do not find this single record from an abandoned termite nest to be compelling evidence. FOWLER & al. (1994) and HOLWAY & al. (2002) may have reported *P. longicornis* as native to Africa simply because LATREILLE (1802) first described the species from Senegal in the far western tip of Africa.

Future research on the phylogeny of the genera *Paratrechina*, *Coluocera*, and *Myrmecophilus*, and on the genetic diversity of *P. longicornis* populations in different parts of the world should help elucidate further the native and exotic ranges of *P. longicornis*.

### Acknowledgements

I thank A. Wetterer, M. Wetterer, P. Lester, R. Clouse, J. Berry, and J. Trager for comments on this manuscript; S. Cover for help, encouragement, and ant identification; J.



Woinarski, D. Ward, M. Vonshak, A. San Juan, P.D Rajan, J. MacGown, M. Lush, J. Lewallen, G. Knight, M.A. Kabir, W. Jaitrong, G. Heller, S. Hedges, M. Haji, A. Francoeur, J. Fellowes, X. Espadaler, G. Dlussky, M. Deyrup, R. Clouse, H. Cagniant, and O. Bourquin for providing unpublished records; B. Taylor, C.Y. Lee, and S. Hedges for additional information; B. Bolton (BMNH), J. Carpenter (AMNH), S. Cover (MCZ), M. Deyrup (ABS), J. Hogan (ONHM), G. Knight (NML), and T. Schultz (SI) for help with their respective ant collections; W. Haines for compiling data from the Smithsonian; J. Endeman for collecting ants around the world; S. Hugel for valuable information and advice concerning *M. americanus*, W. O'Brien for GIS help; D. P. Wojcik and S.D. Porter for compiling their valuable FORMIS bibliography; C. Scheid and R. Pasos of the FAU library for processing so many interlibrary loans; Florida Atlantic University, the National Science Foundation, and the National Geographic Society for financial support.

### Zusammenfassung

Die Ameisenart *Paratrechina longicornis* (LATREILLE, 1802), auch "longhorn crazy ant" genannt, ist ein ubiquitärer Landwirtschafts- und Haushaltslästling in großen Teilen der Tropen und Subtropen und eine Plage in Gebäuden temperater Gebiete. Ich habe Nachweise von *P. longicornis* von > 2100 Fundorten weltweit zusammengetragen und anhand historischer, ökologischer und evolutionsbiologischer Befunde die bekannte Verbreitung, den geographischen Ursprung und die potentielle zukünftige Ausbreitung evaluiert. Ich habe die frühesten bekannten Nachweise für 181 geographische Gebiete (Länder, große Inseln, Inselgruppen, US-Bundesstaaten und Kanadische Provinzen) dokumentiert, einschließlich vieler, für die meines Wissens bisher keine Nachweise veröffentlicht worden sind, beispielsweise Anguilla, Argentinien, Austral-Inseln, Bahrain, British Indian Ocean Territory, Cayman Inseln, Comoro Inseln, Dominica, El Salvador, Gambia, Ghana, Gibraltar, Guatemala, Iles Eparses, Kenya, Malediven, Mali, Montserrat, Nepal, Nevis, Ost Timor, Pakistan, St. Kitts, St. Lucia, Turks und Caicos Inseln, Tuvalu, Maryland, Missouri und Virginia. *Paratrechina longicornis* ist somit die nachweislich am weitesten verbreitete aller Ameisenarten, mit Vorkommen in der Alten und Neuen Welt, sowohl auf der Nord- als auch der Südhalbkugel. Die erfolgreiche weltweite Ausbreitung von *P. longicornis* fußt auf der Fähigkeit, vitale Populationen in hochgradig gestörten und künstlichen Lebensräumen aufzubauen, sogar in Schiffen auf See. Nachweise von *P. longicornis* stammen meist aus der Nähe großer kommerzieller Wasserwege, aber ich erwarte, dass die Art mit der zunehmenden Bedeutung des Lufttransports auch Gebiete im Landesinneren verstärkt besiedeln wird. Viele Nachweise, insbesondere die meisten, wenn nicht sogar alle, aus temperaten Breiten, stammen von innerhalb oder aus der Nähe von Gebäuden. Weil *P. longicornis* in Gebäuden überall dort leben kann, wo Menschen leben, ist der Ausbreitung in hohe Breitengrade keine Grenze gesetzt. Die weltweiten Nachweise von *P. longicornis* und von zwei artspezifischen Symbionten geben wenig eindeutige Hinweise auf den geographischen Ursprung von *P. longicornis*. Die Areale von drei nahe verwandten *Paratrechina*-Arten jedoch machen Südostasien und Melanesien als natives Verbreitungsgebiet von *P. longicornis* glaubhaft.

### References

- ANDRÉ, E. 1881: Catalogue raisonné des Formicides provenant du voyage en Orient de M. Abeille de Perrin et description des espèces nouvelles. – Annales de la Société Entomologique de France (6)1: 53-78.
- ARNOLD, G. 1922: A monograph of the Formicidae of South Africa (Myrmicinae). – Annals of the South African Museum 14: 579-674.
- ASHMOLE, P. & ASHMOLE, M. 2000: St Helena and Ascension Island: a natural history. – Anthony Nelson, Shropshire, 528 pp.
- ASSMUTH, J. 1907: Einige Notizen über *Prenolepis longicornis* LATR. – Zeitschrift für Wissenschaftliche Insektenbiologie 3: 301-309, 328-334, 357-368.
- BERNARD, F. 1968: Faune de l'Europe et du Bassin Méditerranéen. 3. Les fourmis (Hymenoptera Formicidae) d'Europe occidentale et septentrionale. – Masson, Paris, 411 pp.
- BINGHAM, C.T. 1903: The fauna of British India, including Ceylon and Burma. Hymenoptera, vol. II. ants and cuckoo-wasps. – Taylor & Francis, London, 507 pp.
- BLACKBURN, T. & CAMERON, P. 1887: On the Hymenoptera of the Hawaiian Islands. – Memoirs of the Manchester Literary and Philosophical Society 10: 194-244.
- BÜNZLI, G.H. 1935: Untersuchungen über coccidophile Ameisen aus den Kaffeefeldern von Surinam. – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 16: 453-593.
- CAGNIANT, H. 1970: Deuxième liste de fourmis d'Algérie récoltées principalement en forêt (Deuxième partie). – Bulletin de la Société d'Histoire Naturelle de Toulouse 106: 28-40.
- CHAPMAN, J.W. & CARLTON, J.T. 1991: A test of criteria for introduced species: the global invasion by the isopod *Synidotea laevidorsalis* (MIERS 1881). – Journal of Crustacean Biology 11: 386-400.
- CHEESMAN, L.E. & CRAWLEY, W.C. 1928: A contribution towards the insect fauna of French Oceania. - Part III. Formicidae. – Annals and Magazine of Natural History (10)2: 514-525.
- CHIN, D. 1998: Ants in the household and backyard. – Agnote 156: 1-6.
- CLOUSE, R.M. 2007: The ants of Micronesia (Hymenoptera: Formicidae). – Micronesica 39: 171-295.
- COLE, F.R., MEDEIROS, A.C., LOOPE, L.L. & ZUEHLKE, W.W. 1992: Effects of the Argentine ant on arthropod fauna of Hawaiian high-elevation shrubland. – Ecology 73: 1313-1322.
- COLLINGWOOD, C.A. 1985: Hymenoptera: Fam. Formicidae of Saudi Arabia. – Fauna of Saudi Arabia 7: 230-302.
- COLLINGWOOD, C.A., TIGAR, B.J. & AGOSTI, D. 1997: Introduced ants in the United Arab Emirates. – Journal of Arid Environments 37: 505-512.
- CUTTEN, G.M., RAPLEY, W.A. & CHAN, A. 1979: Solving problems in the Toronto Zoo. – Pest Control Technology 7(10): 14-15.
- DEJEAN, A., DURAND, J.L. & BOLTON, B. 1996: Ants inhabiting *Cubitermes* termitaries in African rain forest. – Biotropica 28: 701-713.
- DELABIE, J.H.C. & BLARD, F. 2002: The tramp ant *Hypoconer punctatissima* (ROGER) (Hymenoptera: Formicidae: Ponerinae): new records from the southern hemisphere. – Neotropical Entomology 31: 149-151.
- DEYRUP, M., DAVIS, L. & COVER, S. 2000: Exotic ants in Florida. – Transactions of the American Entomological Society 126: 293-326.
- DIETZ, H.F. 1921: Some problems in greenhouse inspection work in Indiana. – Journal of Economic Entomology 14: 188-194.

- DLUSSKY, G.M. 1994: Zoogeography of southwestern Oceania (in Russian). In: PUZATCHENKO, Y.G., GOLOVATCH, S.I., DLUSSKY, G.M., DIAKONOV, K.N., ZAKHAROV, A.A. & KORGANOVA, G.A. (Eds.): Animal population of the islands of Southwestern Oceania (ecogeographic studies). – Nauka Publishers, Moscow, pp. 48-93.
- DONISTHORPE, H. 1918: A list of ants from Mesopotamia; with a description of a new species and a new variety. – Entomologist's Record and Journal of Variation 30: 165-168.
- DONISTHORPE, H. 1927: British ants, their life-history and classification, 2nd edn. – George Routledge and Sons, London, 436 pp.
- DONISTHORPE, H. 1935: The ants of Christmas Island. – Annals and Magazine of Natural History (10)15: 629-635.
- DONISTHORPE, H. 1936: The ants of the Azores. – Entomologists' Monthly Magazine 72: 130-133.
- DONISTHORPE, H. 1947: Some new ants from New Guinea. – Annals and Magazine of Natural History (11)14: 183-197.
- DUBOIS, M.B. & LABERGE, W.E. 1988: Annotated list of ants in Illinois (Hymenoptera: Formicidae) In: TRAGER, J.C. (Ed.): Advances in myrmecology. – E.J. Brill, Leiden, Netherlands, pp. 133-156.
- DUFFEY, E. 1964: The terrestrial ecology of Ascension Island. – Journal of Applied Ecology 1: 219-251.
- EGUCHI, K., BUI, T.V., YAMANE, S.K., OKIDO, H. & OGATA, K. 2004: Ant faunas of Ba Vi and Tam Dao, North Vietnam (Insecta: Hymenoptera: Formicidae). – Bulletin of the Institute of Tropical Agriculture Kyushu University 27: 77-98.
- EMERY, C. 1881: Viaggio ad Assab nel Mar Rosso dei Signori G. Doria ed O. Beccari con il R. Avviso "Esploratore" dal 16 novembre 1879 al 26 febbraio 1880. I. Formiche. – Annali del Museo Civico di Storia Naturale 16: 525-535.
- EMERY, C. 1886: Catalogo delle formiche esistenti nelle collezioni del Museo Civico di Genova. Parte terza. Formiche della regione Indo-Malese e dell' Australia. [part b] – Annali del Museo Civico di Storia Naturale 24: 209-258.
- EMERY, C. 1889: Formiche di Birmania e del Tenasserim raccolte da Leonardo Fea (1885-87). – Annali del Museo Civico di Storia Naturale 27: 485-520.
- EMERY, C. 1893: Über die Herkunft der Pharao-Ameise. – Biologisches Centralblatt 13: 435-436.
- EMERY, C. 1914: Les fourmis de la Nouvelle-Calédonie et des îles Loyalty. In: SARASIN, F. & ROUX, J. (Eds.): Nova Caledonia. A. zoologie. – C.W. Kreidels Verlag, Wiesbaden, pp. 393-437.
- FERSTER, B. & PRUSAK, Z. 1994: A preliminary checklist of the ants (Hymenoptera: Formicidae) of Everglades National Park. – Florida Entomologist 77: 508-512.
- FOREL, A. 1881: Die Ameisen der Antille St. Thomas. – Mitteilungen des Münchner Entomologischen Vereins 5: 1-16.
- FOREL, A. 1891: Histoire physique, naturelle, et politique de Madagascar. Histoire naturelle des hyménoptères. Deuxième partie: les formicides. – Hachette et Cie, Paris, 237 pp.
- FOREL, A. 1893a: Nouvelles fourmis d'Australie et des Canaries. – Annales de la Société Entomologique de Belgique 37: 454-466.
- FOREL, A. 1893b: Formicides de l'Antille St. Vincent, récoltées par Mons. H. H. Smith. – Transactions of the Entomological Society of London 1893: 333-418.
- FOREL, A. 1895: Nouvelles fourmis de diverses provenances, surtout d'Australie. – Annales de la Société Entomologique de Belgique 39: 41-49.
- FOREL, A. 1897a: Ameisen aus Nossi-Bé, Majunga, Juan de Nova (Madagaskar), den Aldabra-Inseln und Sansibar, gesammelt von Herrn Dr. A. Voeltzkow aus Berlin. Mit einem Anhang über die von Herrn Privatdocenten Dr. A. Brauer in Marburg auf den Seychellen und von Herrn Perrot auf Ste. Marie (Madagaskar) gesammelten Ameisen. – Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft 21: 185-208.
- FOREL, A. 1897b: Quelques formicides de l'Antille de Grenada récoltés par M. H. H. Smith. – Transactions of the Entomological Society of London 1897: 297-300.
- FOREL, A. 1899: Biologia Centrali-Americana; or, contributions to the knowledge of the fauna and flora of Mexico and Central America. Insecta. Hymenoptera. 3. Formicidae. – R.H. Porter, London, 160 pp.
- FOREL, A. 1901a: Formiciden des Naturhistorischen Museums zu Hamburg. Neue *Calypatomyrmex*-, *Dacryon*-, *Podomyrma*- und *Echinopla*-Arten. – Mitteilungen aus dem Naturhistorischen Museum in Hamburg 18: 43-82.
- FOREL, A. 1901b: Formiciden aus dem Bismarck-Archipel, auf Grundlage des von Prof. Dr. F. Dahl gesammelten Materials. – Mitteilungen aus dem Zoologischen Museum in Berlin 2: 4-37.
- FOREL, A. 1907: Formiciden aus dem Naturhistorischen Museum in Hamburg. – Mitteilungen aus dem Naturhistorischen Museum in Hamburg 24: 1-20.
- FOREL, A. 1910: Ameisen aus der Kolonie Erythräa. Gesammelt von Prof. Dr. K. Escherich (nebst einigen in West-Abessinien von Herrn A. Ilg gesammelten Ameisen). – Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 29: 243-274.
- FORYS, E.A. & ALLEN, C.R. 2005: The impacts of sprawl on biodiversity: the ant fauna of the lower Florida Keys. – Ecology & Society 10 (1): 25 [online].
- FOWLER, H.G., SCHLINDWEIN, M.N. & MEDEIROS, M.A. 1994: Exotic ants and community simplification in Brazil: a review of the impact of exotic ants on native ant assemblages. In: WILLIAMS, D.F. (Ed.): Exotic ants: biology, impact, and control of introduced species. – Westview Press, Boulder, pp. 151-162.
- FREITAG, A., DORN, K. & CHERIX, D. 2000: First occurrence of the crazy ant *Paratrechina longicornis* (LATREILLE) (Hym. Formicidae: Formicinae) in Switzerland. – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 73: 301-303.
- GÓMEZ, K. & ESPADALER, X. 2006: Exotic ants (Hymenoptera: Formicidae) in the Balearic Islands. – Myrmecologische Nachrichten 8: 225-233.
- GRIFFITHS, D. 1980: The feeding biology of ant-lion larvae: prey capture, handling and utilization. – Journal of Animal Ecology 49: 99-125.
- GRIMMETT, R.E.R. 1926: Forest-floor covering and its life. – Transactions and Proceedings of the New Zealand Institute 56: 423-440.
- HANNAN, M.A. 2007: Ant records from Savar, Dhaka, Bangladesh. – Asian Myrmecology 1: 101-104.
- HARRIS, R. & BERRY, J. 2005: *Paratrechina longicornis* (LATREILLE). Invasive ant threat information sheet 20. – Landcare Research contract report to Biosecurity New Zealand, Lincoln, New Zealand, 61 pp.
- HEDGES, S.A. 1998: Field guide for the management of structure-infesting ants. 2<sup>nd</sup> edn. – GIE Inc., Cleveland, 304 pp.
- HEDGES, S.A. 2000: Coast to coast. – PCT Online 77 (10 December 2000), <<http://pctonline.com/articles/article.asp?ID=1273&IssueID=77>>, retrieved on 1 February 2008.
- HEMBRY, O. 2005: "Standing army" ready to keep NZ clean, green. – New Zealand Herald 7 November 2005.
- HOFFMANN, B.D. & O'CONNOR, S. 2004: Eradication of two exotic ants from Kakadu National Park. – Ecological Management & Restoration 5: 98-105.

- HOLWAY, D.A., LACH, L., SUAREZ, A.V., YSUTSUI, N.D. & CASE, T.J. 2002: The causes and consequences of ant invasions. – *Annual Review of Ecology & Systematics* 33: 181-233.
- HUDDLESTON, E.W. & FLUKER, S.S. 1968: Distribution of ant species of Hawaii. – *Proceedings of the Hawaiian Entomological Society* 20: 45-69.
- ILLINGWORTH, J.F. 1931: *Coluocera maderae* WOLLASTON. – *Proceedings of the Hawaiian Entomological Society* 7: 378.
- JACOBSON, H. 1939: Die Ameisenfauna des ostbaltischen Gebietes. – *Zoomorphologie* 35: 389-454.
- JAFFE, K. & LATTKE, J.E. 1994: Ant fauna of the French and Venezuelan islands in the Caribbean. In: WILLIAMS, D.F. (Ed.): *Exotic ants: biology, impact, and control of introduced species*. – Westview Press, Boulder, pp. 181-190.
- JAFFE, K., MAULEON, H. & KERMARREC, A. 1990: Predatory ants of *Diaprepes abbreviatus* (Coleoptera: Curculionidae) in citrus groves in Martinique and Guadeloupe, F.W.I. – *Florida Entomologist* 73: 684-687.
- JAITRONG, W. & NABHITABHATA, J. 2005: A list of known ant species of Thailand (Formicidae: Hymenoptera). – *The Thailand Natural History Museum Journal* 1: 9-54.
- JERDON, T.C. 1851: A catalogue of the species of ants found in Southern India. – *Madras Journal of Literature and Science* 17: 103-127.
- KARAVAEV, V. 1911: Murav'i, sobrannye v Egipte i Sudane. – *Russkoye Entomologicheskoye Obozreniye* 11: 1-12.
- KENNE, M., MONY, R., TINDO, M., NJALEU, L.C.K., ORIVEL, J. & DEJEAN, A. 2005: The predatory behavior of a tramp ant species in its native range. – *Comptes Rendus Biologies* 328: 1025-1030.
- KISTNER, D.H. 1982: 1. The social insects' bestiary. In: HERMANN, H.R. (Ed.): *Social insects*. – Academic Press, New York, pp. 1-244.
- KRAUSS, N.L.H. 1944: Notes on insects and other arthropods from the islands of Molokai and Maui, Hawaii. – *Proceedings of the Hawaiian Entomological Society* 12: 81-94.
- KUGLER, J. 1988: The zoogeography of Israel. 9. The zoogeography of social insects of Israel and Sinai. – *Monographiae Biologicae* 62: 251-275.
- LATREILLE, P.A. 1802: *Histoire naturelle des fourmis, et recueil de memoires et d'observations sur les abeilles, les araignees, les faucheurs, et autres insectes*. – Imprimerie de Crapelet (chez T. Barrois), Paris, 445 pp.
- LESTER, P.J. 2005: Determinants for the successful establishment of exotic ants in New Zealand. – *Diversity & Distributions* 11: 279-288.
- LIM, S.P., CHONG, A.S.C. & LEE, C.Y. 2003: Nestmate recognition and intercolonial aggression in the crazy ant, *Paratrechina longicornis* (Hymenoptera: Formicidae). – *Sociobiology* 41: 295-305.
- MANN, W.M. 1915: Some myrmecophilous insects from Hayti. – *Psyche* 22: 161-166.
- MANN, W.M. 1918: Myrmecophilous insects from Cuba. – *Psyche* 25: 104-106.
- MANN, W.M. 1919: The ants of the British Solomon Islands. – *Bulletin of the Museum of Comparative Zoology* 63: 273-391.
- MANN, W.M. 1921: The ants of the Fiji Islands. – *Bulletin of the Museum of Comparative Zoology* 64: 401-499.
- MARLATT, C.L. 1928: House ants, kinds and methods of control. Revised. – *United States Department of Agriculture Farmers' Bulletin* 740: 1-12.
- MAYR, G. 1862: *Myrmecologische Studien*. – *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 12: 649-776.
- MAYR, G. 1867: *Adnotationes in monographiam formicidarum Indo-Neerlandicarum*. – *Tijdschrift voor Entomologie* 10: 33-117.
- MAYR, G. 1870: *Neue Formiciden*. – *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 20: 939-996.
- MAYR, G. 1876: *Die australischen Ameisen*. – *Journal des Museums Godeffroy Hamburg* 12: 56-115.
- MENOZZI, C. 1926: *Formiche dell'Africa centrale*. – *Bollettino della Società Entomologica Italiana* 58: 36-41.
- MENOZZI, C. 1942: *Formiche dell'isola Fernando Poo e del territorio del Rio Muni (Guinea Spagnol)*. 24. Beitrag zu den Wissenschaftlichen Ergebnissen der Forschungsreise H. Eidmann nach Spanisch-Guinea 1939 bis 1940. – *Zoologischer Anzeiger* 140: 164-182.
- MILLER, S.E. 1994: Dispersal of plant pests into the Virgin Islands. – *Florida Entomologist* 77: 520-521.
- MORRIS, B. 2004: *Insects and human life*. – Berg Publishers, Oxford, 320 pp.
- MORRISON, L.W. 1997: Polynesian ant (Hymenoptera: Formicidae) species richness and distribution: a regional survey. – *Acta Oecologica* 18: 685-695.
- MOTSCHOULSKY, V. 1863: *Essai d'un catalogue des insectes de l'île Ceylan*. – *Bulletin de la Société Impériale des Naturalistes de Moscou* 36: 1-153.
- MÜHLEN, V.Z. 1887: *Hiesige Formiciden*. – *Sitzungsberichte der Naturforscher-Gesellschaft bei der Universität Dorpat* 8: 327-333.
- MYERS, J.G. 1934: The Arthropod fauna of a rice-ship, trading from Burma to the West Indies. – *Journal of Animal Ecology* 3: 146-149.
- NAUMANN, K. 1994: An occurrence of two exotic ant (Formicidae) species in British Columbia. – *Journal of the Entomological Society of British Columbia* 91: 69-70.
- NOVÁK, V. 1947: *Exotické mravenci ve sklenících Pražské botanické zahrady*. – *Casopis České Společnosti Entomologické* 44: 144-146.
- NYLANDER, W. 1856: *Synopsis des formicides de France et d'Algérie*. – *Annales des Sciences Naturelles, Zoologie* (4)5: 51-109.
- ONOYAMA, K. 1980: *An introduction to the ant fauna of Japan, with a check list (Hymenoptera, Formicidae)*. – *Kontyû* 48: 193-212.
- PAKNIA, O. & KAMI, H.G. 2007: New and additional record for Formicid (Hymenoptera: Insecta) fauna of Iran. – *Zoology in the Middle East* 40: 85-90.
- PASCOE, A. 2002: Crazy ant species found. – *Stowaways* 2: 12.
- PECK, S.B. 1993: New beetle records from the Galapagos-Islands, Ecuador (Coleoptera). – *Coleopterists Bulletin* 47: 151-157.
- RADOSZKOWSKY, O. 1884: *Fourmis de Cayenne Française*. – *Trudy Russkago Entomologicheskago Obshchestva* 18: 30-39.
- ROGER, J. 1859: *Beiträge zur Kenntniss der Ameisenfauna der Mittelmeerländer*. I. – *Berliner Entomologische Zeitschrift* 3: 225-259.
- ROOM, P.M. 1975: Diversity and organization of the ground foraging ant faunas of forest, grassland and tree crops in Papua New Guinea. – *Australian Journal of Zoology* 23: 71-89.
- SANTSCHI, F. 1914a: *Formicides de l'Afrique occidentale et australe du voyage de Mr. le Professeur F. Silvestri*. – *Bollettino del Laboratorio di Zoologia Generale ed Agraria, Portici* 8: 309-385.

- SANTSCHI, F. 1914b: Formicidae. – Voyage de Ch. Alluaud et R. Jeannel en Afrique Orientale, 1911-1912. Résultats scientifiques. – Insectes Hyménoptères 2: 41-148.
- SANTSCHI, F. 1920a: Fourmis d'Indo-Chine. – Annales de la Société Entomologique de Belgique 60: 158-176.
- SANTSCHI, F. 1920b: Études sur les maladies et les parasites du cacaoyer et d'autres plantes cultivées à S. Thomé. X. Fourmis de S. Thomé. – Société Portugaise des Sciences Naturelles. Librairie Ferin, Lisbon, 4 pp.
- SANTSCHI, F. 1929: Fourmis du Maroc, d'Algérie et de Tunisie. – Annales de la Société Entomologique de Belgique 69: 138-165.
- SANTSCHI, F. 1935: Hymenoptera. I. Formicidae. – Mission scientifique de l'Omo. Vol. 2 Zoologie 15: 255-277.
- SAUSSURE, H. 1877: Gryllides. – Mémoires de la Société de Physique et d'Histoire naturelle de Genève 25: 111-341.
- SCHEMBRI, S.P. & COLLINGWOOD, C.A. 1981: A revision of the myrmecofauna of the Maltese Islands (Hymenoptera, Formicidae). – Annali del Museo Civico di Storia Naturale "Giacomo Doria" 83: 417-442.
- SEIFERT, B. 2003: *Hypoponera punctatissima* (ROGER) and *H. schauinslandi* (EMERY) – two morphologically and biologically distinct species (Hymenoptera: Formicidae). – Abhandlungen und Berichte des Naturkundemuseums Görlitz 75: 61-81.
- SMITH, F. 1879: The collections from Rodriguez. Hymenoptera, Diptera, and Neuroptera. – Philosophical Transactions of the Royal Society of London 168: 534-540.
- SMITH, M.R. 1934: A list of the ants of South Carolina. – Journal of the New York Entomological Society 42: 353-361.
- SMITH, M.R. 1935: A list of the ants of Oklahoma (Hymen.: Formicidae). – Entomological News 46: 261-264.
- SNELLING, R. 2000: Ants recorded at two camps in the Wapoga area, Irian Jaya, Indonesia. – RAP Bulletin of Biological Assessment 14: 95-100.
- STITZ, H. 1916: Formiciden. – Ergebnisse der Zweiten Deutschen Zentral-Afrika Expedition 1: 369-405.
- STITZ, H. 1925: Ameisen von den Philippinen, den malayischen und ozeanischen Inseln. – Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1923: 110-136.
- TARI 2007: Insect and Mite Collection of Taiwan Agricultural Research Institute (Taichung County). – <<http://timdm.tari.gov.tw>>, retrieved on 1 February 2008.
- TAYLOR, R.W. 1967: Entomological survey of the Cook Islands and Niue. 1 – Hymenoptera-Formicidae. – New Zealand Journal of Science 10: 1092-1095.
- TERAYAMA, M. 1999: Taxonomic studies of the Japanese Formicidae, Part 5. Genus *Paratrechina* MOTSCHOUJSKY. – Memoirs of the Myrmecological Society of Japan 1: 49-64.
- TINAUT, A. & AÑO, J.L. 2000: *Paratrechina longicornis*, a new record for the Iberian Peninsula (Hymenoptera, Formicidae). – Boletín de la Asociación Española de Entomología 24: 253-254.
- TRAGER, J.C. 1984: A revision of the genus *Paratrechina* (Hymenoptera: Formicidae) of the continental United States. – Sociobiology 9: 49-162.
- VAN ZWALUWENBERG, R.H. 1943: The insects of Canton Island. – Proceedings of the Hawaiian Entomological Society 11: 300-312.
- VIHMEYER, H. 1915: Ameisen von Singapore. Beobachtet und gesammelt von H. Overbeck. – Archiv für Naturgeschichte 81: 108-168.
- WASMANN, E. 1905a: Zur Lebensweise einiger in- und ausländischer Ameisengäste (148. Beitrag zur Kenntnis der Myrmecophilen und Termitophilen). – Zeitschrift für Wissenschaftliche Insektenbiologie 10: 329-336.
- WASMANN, E. 1905b: Zur Lebensweise einiger in- und ausländischer Ameisengäste (148. Beitrag zur Kenntnis der Myrmecophilen und Termitophilen). – Zeitschrift für Wissenschaftliche Insektenbiologie 11: 384-390.
- WEBER, N.A. 1939: Tourist ants. – Ecology 20: 442-446.
- WEBER, N.A. 1940: Ants on a Nile River steamer. – Ecology 21: 292-293.
- WEBER, N.A. 1948: Studies on the fauna of Curaçao, Aruba, Bonaire and the Venezuelan islands: No. 14. Ants from the Leeward Group and some other Caribbean localities. – Natuurwetenschappelijke Studiekring voor Suriname en de Nederlandse Antillen 5: 78-86.
- WETTERER, J.K. 1997: Ants on *Cecropia* in Hawaii. – Biotropica 29: 128-132.
- WETTERER, J.K. 1998a: Nonindigenous ants associated with geothermal and human disturbance in Hawai'i Volcanoes National Park. – Pacific Science 52: 40-50.
- WETTERER, J.K. 1998b: Ants on *Cecropia* trees in urban San José, Costa Rica. – Florida Entomologist 81: 118-121.
- WETTERER, J.K. 2002: Ants of Tonga. – Pacific Science 56: 125-135.
- WETTERER, J.K., ESPADALER, X., WETTERER, A.L., AGUIN-POMBO, D. & FRANQUINHO-AGUIAR, A.M. 2006: Ants (Hymenoptera: Formicidae) of the Madeiran Archipelago. – Sociobiology 49: 265-297.
- WETTERER, J.K., ESPADALER, X., WETTERER, A.L. & CABRAL, S.G.M. 2004: Native and exotic ants of the Azores (Hymenoptera: Formicidae). – Sociobiology 44: 1-20.
- WETTERER, J.K. & HUGEL S. in press: Worldwide spread of the ant cricket *Myrmecophilus americanus*, a symbiont of the longhorn crazy ant, *Paratrechina longicornis*. – Sociobiology.
- WETTERER, J.K., MILLER, S.E., WHEELER, D.E., OLSON, C.A., POLHEMUS, D.A., PITTS, M., ASHTON, I.W., HIMLER, A.G., YOSPIN, M., HELMS, K.R., HARKEN, E.L., GALLAHER, J., DUNNING, C.E., NELSON, M., LITSINGER, J., SOUTHERN, A. & BURGESS, T.L. 1999: Ecological dominance by *Paratrechina longicornis* (Hymenoptera: Formicidae), an invasive tramp ant, in Biosphere 2. – Florida Entomologist 82: 381-388.
- WETTERER, J.K. & MOORE, J.A. 2005: Red imported fire ants (Hymenoptera: Formicidae) at gopher tortoise (Testudines: Testudinidae) burrows. – Florida Entomologist 88: 349-354.
- WETTERER, J.K. & O'HARA, B.C. 2002: Ants of the Dry Tortugas, the outermost Florida Keys. – Florida Entomologist 85: 303-307.
- WETTERER, J.K. & VARGO, D.L. 2003: Ants (Hymenoptera: Formicidae) of Samoa. – Pacific Science 57: 409-419.
- WETTERER, J.K. & WETTERER, A.L. 2004: Ants (Hymenoptera: Formicidae) of Bermuda. – Florida Entomologist 87: 212-221.
- WETTERER, J.K. & WETTERER, A.L. 2006: A disjunct Argentine ant metacolon in Macaronesia and southwestern Europe. – Biological Invasions 8: 1123-1129.
- WHEELER, W.M. 1905: The ants of the Bahamas, with a list of the known West Indian species. – Bulletin of the American Museum of Natural History 21: 79-135.
- WHEELER, W.M. 1906a: A new wingless fly (*Puliciphora borinquenensis*) from Porto Rico. – Bulletin of the American Museum of Natural History 22(14): 267-271.
- WHEELER, W.M. 1906b: On certain tropical ants introduced into the United States. – Entomological News 17: 23-26.
- WHEELER, W.M. 1907: A collection of ants from British Honduras. – Bulletin of the American Museum of Natural History 23: 271-277.

- WHEELER, W.M. 1908a: The ants of Porto Rico and the Virgin Islands. – *Bulletin of the American Museum of Natural History* 24: 117-158.
- WHEELER, W.M. 1908b: Ants from Moorea, Society Islands. – *Bulletin of the American Museum of Natural History* 24: 165-167.
- WHEELER, W.M. 1909: Ants of Formosa and the Philippines. – *Bulletin of the American Museum of Natural History* 26: 333-345.
- WHEELER, W.M. 1910: Ants: their structure, development and behavior. – Columbia University Press, New York, 663 pp.
- WHEELER, W.M. 1911: Additions to the ant fauna of Jamaica. – *Bulletin of the American Museum of Natural History* 30: 21-29.
- WHEELER, W.M. 1912: The ants of Guam. – *Journal of the New York Entomological Society* 20: 44-48.
- WHEELER, W.M. 1913a: Ants collected in Georgia by Dr. J. C. Bradley and Mr. W. T. Davis. – *Psyche* 20: 112-117.
- WHEELER, W.M. 1913b: Ants collected in the West Indies. – *Bulletin of the American Museum of Natural History* 32: 239-244.
- WHEELER, W.M. 1916: Ants collected in British Guiana by the expedition of the American Museum of Natural History during 1911. – *Bulletin of the American Museum of Natural History* 35: 1-14.
- WHEELER, W.M. 1919: Expedition of the California Academy of Sciences to the Galapagos Islands, 1905-1906. Part 14. The ants of the Galapagos Islands. – *Proceedings of the California Academy of Sciences* (4)2(2): 259-297.
- WHEELER, W.M. 1921: Chinese ants collected by Prof. C. W. Howard. – *Psyche* 28: 110-115.
- WHEELER, W.M. 1923: Report on the ants collected by the Barbados-Antigua Expedition from the University of Iowa in 1918. – *Studies in Natural History, Iowa University* 10: 3-9.
- WHEELER, W.M. 1936: Ants from the Society, Austral, Tuamotu and Mangareva Islands. – *Occasional Papers of the Bernice Pauahi Bishop Museum* 12(18): 3-17.
- WILSON, E.O. 1959: Some ecological characteristics of ants in New Guinea rain forests. – *Ecology* 40: 437-447.
- WILSON, E.O. 1973: The ants of Easter Island and Juan Fernández. – *Pacific Insects* 15: 285-287.
- WILSON, E.O. & HUNT, G.L. 1967: Ant fauna of Futuna and Wallis Islands, stepping stones to Polynesia. – *Pacific Insects* 9: 563-584.
- WILSON, E.O. & TAYLOR, R.W. 1967: Ants of Polynesia. – *Pacific Insects Monographs* 14: 1-109.
- WOJCIK, D.P. & PORTER, S.D. (compilers) 2005: FORMIS: a master bibliography of ant literature. – USDA-ARS, CMAVE, Gainesville, Florida.
- WOLLASTON, T.V. 1854: *Insecta Maderensia*; being an account of the insects of the islands of the Madeiran group. – Van Voorst, London, 634 pp.
- YAMANE, S.K., VIET, B.T., OGATA, K., OKIDO, H. & EGUCHI, K. 2002: Ant fauna of Cuc Phuong National Park, North Vietnam (Hymenoptera: Formicidae). – *Bulletin of the Institute of Tropical Agriculture Kyushu University* 25: 51-62.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Myrmecological News = Myrmecologische Nachrichten](#)

Jahr/Year: 2008

Band/Volume: [011](#)

Autor(en)/Author(s): Wetterer James K.

Artikel/Article: [Worldwide spread of the longhorn crazy ant, \*Paratrechina longicornis\* \(Hymenoptera: Formicidae\). 137-149](#)