

## The effect of ants on soil properties and processes (Hymenoptera: Formicidae)

Jan FROUZ & Veronika JILKOVÁ



### Abstract

Ants are ecosystem engineers, greatly affecting physical, chemical, and biological properties of the soil. The effects on physical soil properties are connected with the building of corridors and galleries, which increase soil porosity and may cause separation of soil particles according to their size. Ant-mediated chemical changes of soil are represented mainly by a shift of pH towards neutral and an increase in nutrient content (mostly nitrogen and phosphorus) in ant nest-affected soil. These effects correspond with accumulation of food in the nests and the effect on biological processes, such as acceleration of decomposition rate. Effects on biological soil properties may be connected with increased or decreased microbial activity, which is affected by accumulation of organic matter and internal nest temperature and especially moisture. Effects on the soil vary between ant species; substantial variation can be found in the same species living in different conditions.

**Key words:** Ants, soil, nutrient cycling, porosity, organic matter, moisture, microbial activity, review.

Myrmecol. News 11: 191-199 (online 25 July 2008)

ISSN 1994-4136 (print), ISSN 1997-3500 (online)

Received 25 March 2008; revision received 20 June 2008; accepted 22 June 2008

Assoc. Prof. Dr. Jan Frouz (contact author), Veronika Jilková, Institute of Soil Biology BC ASCR, Na sádkách 7; Faculty of Sciences, South Bohemian University, Branišovská 31, České Budějovice, CZ37005, Czech Republic. E-mail: frouz@upb.cas.cz

### Introduction

The role of ants in pest management has been emphasized several times (NIEMELA & LAINE 1986, WAY & KHOO 1992), but much less attention has been paid to the effects of ants on soil properties (CHERIX 1991). In comparison, modification of soil by termites and earthworms is much better understood (LEE 1982, LAVELLE & al. 1997) as is their role in nutrient cycling in ecosystems (JONES 1990). In this review, we will summarize the main mechanisms by which ants affect the soil environment and soil processes.

Ants may build nests, which are used for different periods of time, from several months to decades. Many ants build their nests completely or partly in the soil. The building of such a permanent structure as a nest will affect the properties of the surrounding soil (PETAL 1978, LOBRY DE BRUYN & CONACHER 1990, FOLGARAIT 1998). Ants accumulate a large amount of food in their nests and deposit excreta or food residues inside the nest or in close vicinity, which may alter the nutrient status of the soil. At the same time, food or nest building material is removed from the surrounding ecosystem, and such removal can result in complex interactions. Moreover, ants can directly or indirectly affect other organisms such as aphids, plants, and fungal decomposers, which may lead to complex multitrophic interactions and may also affect soil conditions.

### Types of soil nests and bioturbation

Ants build several types of nests in the soil. Soil nests that consist of chambers and corridors can be covered by a stone, dead wood, or some other natural structure on the soil surface. These nests may be surrounded by a crater of the excavated soil, which may also be used to build an above-ground structure, in some species containing both soil and

organic material brought from the surroundings. The annual input of plant remains (i.e., coniferous litter) to a *Formica polyctena* nest constitutes 12 - 37 % of the nest volume (POKARZHEVSKIJ 1981). It is generally assumed that ants build aboveground structures for thermoregulation of the nest. Nests built of mineral particles trap solar radiation, which can create large variability in temperatures. In response to this variation, ants can move into that part of the nest that offers the best microclimatic conditions. Ants that build organic anthill nests, especially several species of the subgenus *Formica* s.str., can maintain a relatively constant temperature inside the nest (about 10 °C higher than the ambient temperature) because organic material provides much greater insulation than mineral soil.

Belowground nests typically consist of vertical tunnels that connect horizontal chambers (MIKHEYEV & TSCHINKEL 2004). The descending angle of the tunnels may vary, forming a zig-zag or helix structure (TSCHINKEL 2004, MIKHEYEV & TSCHINKEL 2004). Chambers are built as horizontal appendages or enlargements of tunnels (TSCHINKEL 2004). Chambers may be similar throughout the nest and their size or number may be proportional to the number of workers, as shown by MIKHEYEV & TSCHINKEL (2004) for *Formica pallidefulva*, or more complex rules may apply. *Pogonomyrmex badius* construct top-heavy nests with larger chambers in the shallower parts, most likely due to age division of labor and differences in depth distribution of workers of different age. Older workers are more engaged in digging than younger workers and occur more frequently in the upper parts of a nest (TSCHINKEL 2004). Tunnels or chambers excavated belowground are coated with fine material in some cases but not in others

Tab. 1: Amount of soil excavated per year and ha by various ant species.

Species	Habitat / Country	Amount of soil	Reference
<i>Formica cf. picea</i>	forest / Russia	30 m <sup>3</sup>	DMITRIENKO & PETRENKO (1976)
Whole community	prairie / Russia	500 kg	DMITRIENKO & PETRENKO (1976)
<i>Chelaner</i> sp.	prairie / Australia	5 - 6 kg	BRIESE (1982)
<i>Pheidole</i> sp.	prairie / Australia	75 - 90 kg	BRIESE (1982)
<i>Iridomyrmex</i> sp.	prairie / Australia	150 - 180 kg	BRIESE (1982)
Whole community	prairie / Australia	350 - 420 kg	BRIESE (1982)
<i>Pogonomyrmex occidentalis</i>	prairie / USA	2.8 - 8 kg	ROGERS (1972)
<i>Lasius niger</i>	fallow / USA	855 kg	TALBOT (1953)
Whole community	podzol / USA	600 kg	LYFORD (1963)
<i>Myrmica</i> sp.	meadow / Poland	230 - 9,700 kg	PETAL & al. (1977)
<i>Lasius</i> sp.	meadow / Poland	230 - 9,700 kg	PETAL & al. (1977)
<i>Lasius flavus</i>	meadow / Russia	3,000 - 13,000 kg	DLUSSKIJ (1981)

(WANG & al. 1995, COSARINSKY 2006, COSARINSKY & ROCES 2007). Walls of aboveground structures are formed by pellets of excavated material, which are glued together by their edges, leaving star-shaped voids between the pellets (COSARINSKY & ROCES 2007). Sometimes pellets of excavated material may keep their original microstructure, but soil transport may cause separation of materials based on grain size (COSARINSKY 2006). This most likely depends on the preferred load size carried by the ants and the grain size composition of a particular soil.

Two major processes of nest building can be important for soil modification: bioturbation, which involves the mixing and accumulation of soils from different sources and horizons (NKEM & al. 2000), and transport of organic material from the surroundings into the nests as food or building material. Ants can excavate a significant amount of soil from deeper layers and deposit it on the soil surface. For *Formica cinerea*, about 85 % of aboveground nest material came from illuvial soil horizons (BAXTER & HOLE 1967). The overall amount of excavated soil in an area also depends on the number of nests, which may vary substantially. For example, the number of *Formica* s.str. nests in eastern Finland is 2.7 - 2.9 / ha (DOMISCH & al. 2006); for meadows in Denmark, NIELSEN (1986) reported that about 15 % of the area was covered by anthills; finally, nests of *Camponotus punctulatus* in northeastern Argentina may reach a density of 1,800 / ha, and individual nests may be up to 2 m in diameter (GOROSITO & al. 2006), which means that about half of the soil surface is affected by anthills. Macropore production may also be important, depending on the number and size of the nests and on the frequency at which nests are abandoned or the old structures replaced by new ones. In some species, a nest is used for only a few months. For example, MIKHEYEV & TSCHINKEL (2004) expect that nests of *Formica pallidefulva* are replaced each half year. On the other hand, some nests of wood ants, *Formica* s.str., may be used for decades (DLUSSKIJ 1967).

Nest lifespan is important particularly when nest density is high because empty space is immediately occupied by new nests (NIELSEN 1986). Bioturbation is likely to be more intensive when ants are forming new nests because the formation of new nests requires the building of many new structures. Even with existing nests, however, bioturbation continues because ants repair eroded chambers, fill abandoned chambers with soil, and dig new ones as a replacement (DLUSSKIJ 1981). Estimates of amount of the soil excavated by ants in various ecosystems are given in Table 1.

Bioturbation can also affect the surrounding vegetation. Continuous heaping of soil may support the persistence of some annual plants that would otherwise suffer from competition in dense meadow vegetation (DOSTAL 2007). The ant environment may also support species with fast root growth or long rhizomes (KOVAROVA & al. 2001, DOSTAL & al. 2005) or even support selection of strains of clonal plants with faster root growth and longer rhizomes (ROTHANZL & al. 2007). In some cases, these bioturbations can substantially change the environment for plant growth. For example, *Formica podzolica* forms hummocks in peatland soils, which have much better aeration than the surrounding peat and serve as a habitat for diverse plant species (LESICA & KANOWSKI 1998).

#### Physical properties of soil

Building of tunnels and chambers both above- and belowground increases soil macroporosity (MCCAHERN & LOCKWOOD 1990) and reduces bulk density. For example, bulk density in nests of *Pogonomyrmex occidentalis* was 1.47 g / cm<sup>3</sup> compared with 1.54 in the surrounding soil (ROGERS 1972). Reduced bulk density may increase soil aeration and permeability of soil for water (DLUSSKIJ 1967, ELDRIDGE 1993, 1994). The effect on water infiltration, however, can be complex. Nests increase not only soil macroporosity but also organic matter content, which may increase water repellency at low soil moistures. Thus, ant




Fig. 1: Relationships between soil pH and the nest-soil difference in pH. Filled circles based on data from *Lasius niger*, according to FROUZ & al. (2003). Empty squares with numbers: 1 - *Formica rufibarbis* according to DLUSSKIJ (1967), 2 - *Formica lugubris* according to MALOZEMOVA & KORUMA (1973), 3 - *Formica polyctena* according to MALOZEMOVA & KORUMA (1973), 4 - *Formica pratensis* according to MALOZEMOVA & KORUMA (1973), 5 - *Formica obscuripes* according to MCCAHOON & LOCKWOOD (1990), 6 - *Formica canadensis* according to CULVER & BEATTIE (1983), 7 - *Formica cunicularia* according to CZERWINSKI & al. (1971), 8 - *Lasius niger* according to DLUSSKIJ (1967), 9 - *Lasius flavus* according to DLUSSKIJ (1967), 10 - *Lasius flavus* according to GASPAR (1972), 11 - *Myrmica* sp. according to DLUSSKIJ (1967), 12 - *Myrmica* sp. according to CZERWINSKI & al. (1971), 13 - *Lasius flavus* according to CZERWINSKI & al. (1971), 14 - *Lasius niger* according to PETAL (1978), 15 - *Myrmica* sp. according to PETAL & al. (1977). Solid line represents a trend line for *L. niger* ( $r = 0.517$ ), dotted line represents a trend line for multispecies data ( $r = 0.668$ ); both significant at  $p < 0.01$ .

Ant nests increased water infiltration in moist or wet conditions but decreased water infiltration in dry conditions (CAMMERAAAT & al. 2002). GREEN & al. (1999) found that macroporosity in nests of imported fire ants can increase drainage, quickly bringing water to deeper soil layers and ensuring higher moisture in soil below the nest while reducing moisture in the nest compared with that in the surrounding soil. *Camponotus punctulatus* nests are surrounded by a peripheral ditch where water accumulates, producing a constantly-wetted zone inside the anthill. Nest moisture is often significantly different from the moisture of surrounding soil and can be lower (MCCAHOON & LOCKWOOD 1990) or higher (COENEN-STASS & al. 1980); it varies even within the same species (FROUZ 2000). For example, *Formica polyctena* can have wet and dry nests, which differ in temperature regime and also in the location and intensity of microbial activity, which is related to temperature and moisture content (FROUZ 2000, FROUZ & FINER 2007).

Temperature is another physical factor that is altered in an ant nest, and regulation of internal nest temperature has been mostly described for nests with anthills (DLUSSKIJ 1967, BRIAN 1978, HÖLLDOBLER & WILSON 1990). The most advanced thermoregulation has been described in wood ants, *Formica* s.str. (STEINER 1924, RAIGNIER 1948, DLUSSKIJ 1967, GALLÉ 1973, HORSTMANN 1983, 1990, ROSENGREN & al. 1987, FROUZ 1996, 2000, FROUZ & FINER 2007). Environmental factors influencing nest temperature are primarily air temperature, nest size, and nest moisture

(FROUZ & FINER 2007). Maintenance of higher internal nest temperatures is possibly due to a combination of the insulation provided by the nest (GALLÉ 1973, BRANDT 1980, FROUZ 1996, 2000), the trapping of solar radiation by the nest and by ant bodies (FROUZ 2000), and the production of metabolic heat by the ants (HORSTMANN 1987, 1990, HORSTMANN & SCHMID 1986) and by the microorganisms associated with the nest material (COENEN-STASS & al. 1980, FROUZ 2000).

### Chemical properties of soil

Many studies have reported significant differences in chemical soil properties between ant nests and the surrounding soil. In general, ants shift nest pH toward a neutral value (Fig. 1), i.e., ants increase pH in acidic soils and decrease it in basic soils (DLUSSKIJ 1967, FROUZ & al. 2003). Some interspecific differences were noted in this context; for example, MALOZEMOVA & KORUMA (1973) found that *Formica polyctena* and *Formica lugubris* shift pH from acidic to neutral values more intensively than *Formica pratensis*. The effect of ants on soil pH and other chemical properties can increase as the nest ages, and the effect is greatest near the nest periphery (ZACHAROV & al. 1981). The mechanism behind the influence of ants on nest pH is not clear, but an increase in pH may result from an increase in basic cations, whereas a decrease in pH may result from an accumulation of organic matter (PETAL 1978, FROUZ & al. 2003).

Changes in N and P content in ant nests have often been reported (FROUZ & KALCIK 1996, LAFLEUR & al. 2002, SNYDER & al. 2002, PETAL & al. 2003, WAGNER & JONES 2004, FROUZ & al. 2003, 2005, KILPELAINEN & al. 2007) (Tab. 2). There are interspecific differences in the accumulation of macronutrients in the nests as seen from measurements of several species in one locality (MALOZEMOVA & KORUMA 1973, BRIESE 1982, JONES & WAGNER 2006) (Tab. 2). On the other hand, nutrient accumulation in a particular species is affected by properties of the surrounding soil and the material used for building the nest. For example, FROUZ & al. (2003) showed that the phosphorus content of *Lasius niger* nests was greater in soils already rich in phosphorus. Beside affecting the total content of nutrients in nests, ants also affect the availability of nutrients; studies of *L. niger* and *Formica* s.str. showed that the increase in total P content in the nest was accompanied by a substantial increase in the available forms of P (FROUZ & al. 2003, 2005). In addition to accumulating easily decomposable substances in the nest (Fig. 2), ants can also increase the availability of P. A shift in nest pH may be also important; as mentioned earlier, ants shift pH towards neutral values, and P availability is highest with near neutral pH (BRADY & WEILL 1999). Increases in total C and organic C (BRIAN 1978) as well as humus in the nest (DMITRIENKO & PETRENKO 1976) have often been reported.

A shift in the content of basic cations ( $\text{Ca}^{2+}$ ,  $\text{K}^{+}$ , and  $\text{Mg}^{2+}$ ) has often been noted in ant nests, including those of *Lasius* spp. and *Myrmica* spp. (CZERWINSKI & al. 1971, FROUZ & al. 2003, STERNBERG & al. 2007) as well as those of wood ants, *Formica* s.str. (MALOZEMOVA & KORUMA 1973, ZACHAROV & al. 1981). A decrease in N content was detected in nests in soil highly contaminated by nitrogen (PETAL 1978; Tab. 2). This was explained by the increased numbers of soil bacteria that bound N in their biomass. Similarly, there was a lower salt content in nests than in the

Tab. 2: Changes (%) in content of total nitrogen and phosphorus in ant nests compared to the surrounding soil. Content in soil is assumed to be 100 %; – data not available.

Species	N	P	Habitat / Country	Reference
<i>Formica polyctena</i>	167	131	forest / Russia	DMITRIENKO & PETRENKO (1976)
<i>Formica rufa</i>	139	–	forest / Russia	GRIMALSKIJ (1960)
<i>Formica pratensis</i>	–	210	forest / Russia	MALOZEMOVA & KORUMA (1973)
<i>Formica polyctena</i>	–	516	forest / Russia	MALOZEMOVA & KORUMA (1973)
<i>Formica lugubris</i>	–	607	forest / Russia	MALOZEMOVA & KORUMA (1973)
<i>Formica aquilona</i>	278	239	forest / Russia	ZACHAROV & al. (1981)
<i>Formica canadensis</i>	158	139	meadow / USA	CULVER & BEATTIE (1983)
<i>Myrmica</i> sp.	–	129	meadow / Poland	PETAL & al. (1977)
<i>Lasius</i> sp.	–	129	meadow / Poland	PETAL & al. (1977)
<i>Myrmica</i> sp.	65	–	sand pit / Poland	PETAL (1978)
<i>Lasius niger</i>	65	–	sand pit / Poland	PETAL (1978)
<i>Pheidole</i> sp.	153	534	prairie / Australia	BRIESE (1982)
<i>Chelaner</i> sp.	110	1,266	prairie / Australia	BRIESE (1982)
<i>Iridomyrmex</i> sp.	62	100	prairie / Australia	BRIESE (1982)
<i>Solenopsis invicta</i>	100	100	prairie / USA	COX & al. (1992)

surrounding soil in soil with a high content of salt (COX & al. 1992).

For all chemical effects of ant nests, the strongest was observed in the center of the aboveground part of the nest (MALOZEMOVA & KORUMA 1973, ZACHAROV & al. 1981, FROUZ & al. 2003, 2005). In addition, nutrient concentrations and changes in pH increase with nest age (ZACHAROV & al. 1981, WAGNER & al. 2004) and may persist for several years after nest abandonment (JAKUBCZYK & al. 1972, KRISTIANSEN & AMELUNG 2001, KRISTIANSEN & al. 2001, FROUZ & al. 2003).

A question arises – do ants really alter soil chemical conditions or rather do they select soil spots with certain chemical conditions to build their nests? The above-mentioned studies indicated that ants do alter soil properties: nutrient contents follow a consistent pattern inside the nests and increase over time. Moreover, an experiment with an artificial nest (established in the laboratory) showed that ants do change the chemistry of the substrate and deposit specific substances in the nest (CHEN 2005, 2007, LAFLEUR & al. 2005).

Several factors affect the changes in the nutrient content of ant nests in comparison with the surrounding soil. The role of bioturbation is evident if the content of a given element changes significantly with depth of the soil profile. For example, FROUZ & al. (2003) showed that changes in the C content in nests of *Lasius niger* were the most pronounced in forest soil with a shallow but highly organic topsoil. In such conditions, ants substantially reduced the C content in the nest by bringing up mineral soil from deeper layers.




Fig. 2: The effect of the wood ant *Formica polyctena* on P availability (g of P per ha per year) in a forest ecosystem. The upper parts of the scheme represent the flow of total P. The bottom parts summarize the annual input of available P in individual compartments. Based on data from FROUZ & al. (1997).

Ants also collect large amounts of food, and food residues and excreta are important sources of nutrient increase in the nest (PETAL & al. 1977, KLOFT 1978, FROUZ & al. 1997). FROUZ & al. (1997) quantified the flow of P into a nest in the form of food (Fig. 2); their results showed that the amount of P brought annually into the ant nests in 1 ha of forest exceeded the amount of P supplied by annual lit-

ter fall. In nests built in organic soil or from organic material, a higher content of available nutrients may be caused by faster mineralisation of organic matter (FROUZ & al. 1997, PETAL 1998, STADLER & al. 2006, WAGNER & JONES 2006). Rapid mineralisation can also explain the higher concentrations of macronutrients in the refuse chamber in *Atta* nests than in the leaf material (GUERRA & al. 2007).

How much of these nutrients that accumulate in the nests can be used by plants? Using  $^{15}\text{N}$ , STERNBERG & al. (2007) showed that trees near *Atta* nests used more nest N and contained a higher concentration of Ca than trees more distant from the nests. Similarly, *Acacia constricta* trees near *Formica perpilosa* nests produced more seeds than trees distant from nests (WAGNER 1997). Also, seedlings grew faster and produced more biomass in soil from nests of various ant species, such as *Formica* or *Solenopsis*, than in soil from the surrounding area (FROUZ & al. 2005, LAFLEUR & al. 2005). On the other hand, some field surveys have not shown better establishment of seedlings on ant nests (LAFLEUR & al. 2002). The effect of ants on tree growth may be complex. There are fewer roots below the wood ant (*Formica*) nest than in the surrounding soil but the density of fine roots and macronutrient content of these roots is higher than in the surrounding soil (OHASHI & al. 2007a). This indicates that trees near the nest may use some nutrients stored in the nest. Trees adjacent to a wood ant (*Formica polycytena*) nest grew faster than those 5 - 50 m away but slower than those in a neighboring forest (200 m apart) unaffected by ants (FROUZ & al. 2008). If a substantial portion of the nutrients brought into the nest comes from honeydew (FROUZ & al. 2005, 2008), honeydew depletion may slow the growth of nearby trees (ROSENGREN & SUNDSTRÖM 1991, FROUZ & al. 2008). There are also specific plant species associated with ant nests that usually differ from species growing in adjacent areas (CULVER & BEATTIE 1983). These species must be capable of overcoming the negative effects of the mounds (e.g., the mixing of material) or may benefit from such conditions because of lower competition (DOSTAL 2007).

### **Biological processes and decomposer communities inside the nest**

Ant nests are hot spots of CO<sub>2</sub> production and thus are also hot spots of metabolic activity in an ecosystem, as has been repeatedly shown for wood ants *Formica* s.str. (OHASHI & al. 2005, 2007b, RISCH & al. 2005, DOMISCH & al. 2006). CO<sub>2</sub> emitted from the ant nest probably originates mainly from ant respiration, but in some cases microbial respiration may also be important. Microbial activity may be substantially higher in the nest than in the surrounding soil because of a surplus of available nutrients and because of suitable moisture and temperature conditions (FROUZ & al. 1997, FROUZ 2000). Microbial biomass is mostly concentrated in the upper layer of the nest where temperature and moisture are highest (FROUZ & al. 1997, FROUZ 2000). Higher numbers of bacteria, including N<sub>2</sub> fixers, have been found in wood ant nests (GORNÝ 1976, FROUZ & al. 1997) than in the surrounding soil. In nests of the meadow ants *Lasius* spp. and *Myrmica* spp., counts of bacteria were 14-fold greater and those of fungi 10-fold greater than in the surrounding soil, whereas counts of Actinomycetales were six-fold less than in the surrounding soil (CZERWINSKI & al. 1971). Microbial biomass was also higher in ant nests

(DAUBER & al. 2001). In some cases, however, microbial activity in an ant nest may be lower than in the surrounding soil, mainly because of lower moisture in the nest (HOLEC & FROUZ 2006).

Ant nests of *Lasius* and *Pogonomyrmex* have been reported to promote root colonization by arbuscular mycorrhizal fungi (AMF) (FRIESE & ALLEN 1993, DAUBER & al. 2008). This may be due to soil mixing, AMF spore transport, and a more suitable environment (decreased moisture and increased temperature) in the nest (DAUBER & al. 2008).

Other groups of soil biota, including protozoa, were also more abundant in ant nests than in surrounding soil (ZARAGOZA & al. 2007), but the differences are sometimes small (KORGANOVA & RAKHLEEVA 2006). Ant nests may also serve as hot spots for soil invertebrates (LAAKSO & SETALA 1997, 1998).

### **The effect of ants on soil outside the nest**

The effect of ants on the soil surrounding nests is much less understood than their effect on the nest itself, although we can assume that ants can alter soil conditions outside the nest through their foraging activities. The size of the foraging area varies between species and is also affected by colony size, food supply in the habitat, size of workers, and characteristics of the terrain (STRADLING 1978, PETAL 1980). The frequent use of foraging tracks by ants could alter soil properties because ants use such tracks to remove food and because the tracks are often near sites of ant bioturbation (NKEM & al. 2000). Ants may affect soil invertebrates in their territory and, in some cases, may also affect the input of organic matter to the soil. Ants can be important non-specific predators of soil invertebrates, thereby substantially reducing the densities of soil invertebrates, as shown by VINSON (1991) for *Solenopsis invicta* or by HORSTMANN (1970) for *Formica polycytena*. *Formica* ants can collect a significant amount of needles and other plant material and can thus slow litter input to soil. Wood ants also collect a significant amount of honeydew, which may reach up to 719 kg per nest per year (OEKLAND 1930, WELLENSTEIN 1952, HOLT 1955, HORSTMANN 1974, GOEBEL 1988). It seems likely that collection of so much honeydew can alter litter input into the soil. Moreover, ants can also alter litter input to the soil by reducing foliar herbivory (NIE-MELA & LAINE 1986).

### **Conclusion**

Because ants affect many soil properties, they are justifiably considered as ecosystem engineers (JONES & al. 1994, JOUQUET & al. 2006). Depending on ant species and soil conditions, ants can either increase or decrease certain soil parameters.

Recent literature provides many examples of differences between ant nests and surrounding soil in physical, chemical, and biological parameters. These examples also show that ants may affect the same parameter in different ways under different conditions. Although the pattern is sometimes clear and consistent (e.g., ants increase pH in acidic soil and decrease pH in alkaline soil), a better understanding of factors that determine how ants influence soil is needed. Most studies have thus far provided descriptive information, and even if the studies presented hypotheses about how the ants alter individual soil properties, they

have seldom tested these hypotheses. Thus, future research should test hypotheses so that we increase our understanding of mechanisms underlying how ants affect individual soil properties.

The strongest ant effect has been observed in the ant nest, which represents only a small area in the ecosystem. The effect on the area surrounding the nest is much less understood. This surrounding area is less conspicuous and less defined than the nest, but may include a much larger area. It follows that the impact of ants may be greater in the area surrounding the nest than in the nest itself.

An important research area that has yet to be explored concerns the effect of ants at various spatio-temporal scales of the ecosystem. For example, we know that ants increase available nutrients in their nests. Is the acceleration in nutrient turnover only a local effect? Or do ants deplete the surrounding ecosystem of available nutrients? How do ants affect the spatio-temporal use of nutrients by plants in the surrounding ecosystem? These and similar questions should be addressed by future research. In other words, more research should be focussed on how ants affect processes on the ecosystem level rather than on the level of the individual ant nest.

Finally, ants participate in many multitrophic interactions, such as predation (NIEMELA 1986), protection of aphids (GOEBEL 1988), and seed removal and dispersion (HOWE & SMALLWOOD 1982, ZHOU & al. 2007). How these interactions relate to the effect of ants on soil properties also requires study.

#### Acknowledgements

This study was supported by the research plan of ISB BC ASCR (Z6066911). Dr. Keith Edwards is thanked for linguistic improvements. Dr. J. Dauber and an anonymous referee are thanked for their helpful comments. Bruce Jaffee (JaffeeRevises, USA) is thanked for reading the manuscript.

#### Zusammenfassung

Ameisen sind Ingenieure des Ökosystems, mit enormem Einfluss auf physikalische, chemische und biologische Eigenschaften des Bodens. Der Einfluss auf die physikalischen Eigenschaften des Bodens steht in Zusammenhang mit dem Bau von Gängen und Gallerien, die die Porosität des Bodens erhöhen und die Auftrennung von Bodenpartikeln nach ihrer Größe bedingen können. Durch Ameisen bedingte Änderungen der Bodenchemie betreffen vor allem eine Verschiebung des pH in den neutralen Bereich und eine Steigerung des Nährstoffgehalts (hauptsächlich Stickstoff und Phosphor). Diese Effekte korrespondieren mit der Anhäufung von Nahrung in den Nestern und der Beeinflussung biologischer Prozesse, beispielsweise der Beschleunigung der Abbaurate. Der Einfluss auf biologische Eigenschaften des Bodens kann mit gesteigerter oder verringerter mikrobieller Aktivität zusammenhängen, welche durch die Anhäufung organischer Substanzen, die Temperatur und insbesondere die Feuchtigkeit im Nestinneren beeinflusst wird. Der Einfluss auf den Boden variiert zwischen den verschiedenen Ameisenarten; eine starke Variation findet sich aber auch innerhalb der Arten, in Abhängigkeit vom Lebensraum.

#### References

- BAXTER, F.P. & HOLE, F.D. 1967: Ant (*Formica cinerea*) pedoturbation in a prairie soil. – Soil Science Society of America Proceedings 31: 425-428.
- BRADY, N.C. & WEILL, R.R. 1999: The nature and properties of soils. – Prentice Hall, Upper Saddle River, 960 pp.
- BRANDT, D.C. 1980: The thermal diffusivity of the organic material of a mound of *Formica polyctena* FOERST. in relation to the thermoregulation of the brood (Hymenoptera, Formicidae). – Netherlands Journal of Zoology 30: 326-344.
- BRIAN, A.M. 1978: Production ecology of ants and termites. – Cambridge University Press, London, 409 pp.
- BRIESE, D.T. 1982: The effect of ants on the soil of semi-arid salt-bush habitat. – Insectes Sociaux 29: 375-382.
- CAMMERAAT, L.H., WILLOTT, S.J., COMPTON, S.G. & INCOLL, L.D. 2002: The effects of ants' nests on the physical, chemical and hydrological properties of a rangeland soil in semi-arid Spain. – Geoderma 105: 1-20.
- CHEN, J. 2005: Excretion of phosphoric acid by red imported fire ants, *Solenopsis invicta* BUREN (Hymenoptera, Formicidae). – Environmental Entomology 34: 1009-1012.
- CHEN, J. 2007: Qualitative analysis of red imported fire ant nests constructed in silica gel. – Journal of Chemical Ecology 33: 631-642.
- CHERIX, D. 1991: Red wood ants. – Ethology Ecology & Evolution Special issue 1: 165.
- COENEN-STASS, D., SCHAARSCHMIDT, B. & LAMPRECHT, I. 1980: Temperature distribution and calorimetric determination of heat production in the nest of the wood ants *Formica polyctena* (Hymenoptera, Formicidae). – Ecology 61: 238-244.
- COSARINSKY, M.I. 2006: Nest micromorphology of the neotropical mound building ants *Camponotus punctulatus* and *Solenopsis* sp. – Sociobiology 47: 329-344.
- COSARINSKY, M.I. & ROCES, F. 2007: Neighboring leaf-cutting ants and mound-building termites: Comparative nest micromorphology. – Geoderma 141: 224-234.
- COX, G.W., MILLS, J.M. & ELLIS, B.A. 1992: Fire ants (Hymenoptera, Formicidae) as major agents of landscape development. – Environmental Entomology 21: 281-286.
- CULVER, D.C. & BEATTIE, A.J. 1983: Effects of ant mounds on soil chemistry and vegetation patterns in a Colorado montane meadow. – Ecology 64: 485-492.
- CZERWINSKI, Z., JAKUBCZYK, H. & PETAL, J. 1971: Influence of ant hills on meadow soils. – Pedobiologia 11: 277-285.
- DAUBER, J., NIECHOJ, R., BALTRUSCHAT, H. & WOLTERS, V. 2008: Soil engineering ants increase grass root arbuscular mycorrhizal colonization. – Biology and Fertility of Soils DOI 10.1007/s00374-008-0285-5.
- DAUBER, J., SCHROETER, D. & WOLTERS, V. 2001: Species specific effects of ants on microbial activity and N-availability in the soil of an old-field. – European Journal of Soil Biology 37: 259-261.
- DLUSSKIJ, G.M. 1967: Muravji roda *Formica*. – Nauka, Moskva, 236 pp.
- DLUSSKIJ, G.M. 1981: Nester von *Lasius flavus* (Hymenoptera, Formicidae). – Pedobiologia 21: 81-99.
- DMITRIENKO, V.K. & PETRENKO, E.S. 1976: Muravji tazenych biocenosis sibiry. – Nauka, Novosibirsk, 220 pp.
- DOMISCH, T., FINER, L., OHASHI, M., RISCH, A.C., SUNDSTROM, L., NIEMELA, P. & JURGENSEN, M.F. 2006: Contribution of red

- wood ant mounds to forest floor CO<sub>2</sub> efflux in boreal coniferous forests. – *Soil Biology and Biochemistry* 38: 2425-2433.
- DOSTAL, P. 2007: Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. – *Journal of Vegetation Science* 18: 91-102.
- DOSTAL, P., BREYNIKOVA, M., KOYLICKOVA, V., HERBEN, T. & KOVAR, P. 2005: Ant-induced soil modification and its effect on plant below-ground biomass. – *Pedobiologia* 49: 127-137.
- ELDRIDGE, D.J. 1993: Effect of ants on sandy soils in semiarid eastern Australia – local distribution of nest entrances and their effect on infiltration of water. – *Australian Journal of Soil Research* 31: 509-518.
- ELDRIDGE, D.J. 1994: Nests of ants and termites influence infiltration in a semiarid woodland. – *Pedobiologia* 38: 481-492.
- FOLGARAIT, P.J. 1998: Ant biodiversity and its relationship to ecosystem functioning: a review. – *Biodiversity and Conservation* 7: 1221-1244.
- FRIESE, C.F. & ALLEN, M.F. 1993: The interaction of harvester ants and vesicular-arbuscular mycorrhizal fungi in a patchy semi-arid environment: the effect of mound structure on fungal dispersion and establishment. – *Functional Ecology* 7: 13-21.
- FROUZ, J. 1996: The role of nest moisture in thermoregulation of ant (*Formica polyctena*, Hymenoptera, Formicidae) nests. – *Biologia, Bratislava* 51: 541-547.
- FROUZ, J. 2000: The effect of nest moisture on daily temperature regime in the nest of *Formica polyctena* wood ants. – *Insectes Sociaux* 47: 229-235.
- FROUZ, J. & FINER, L. 2007: Diurnal and seasonal fluctuations in wood ant (*Formica polyctena*) nest temperature in two geographically distant populations along a south - north gradient. – *Insectes Sociaux* 54: 251-259.
- FROUZ, J., HOLEC, M. & KALCIK, J. 2003: The effect of *Lasius niger* (Hymenoptera, Formicidae) ant nests on selected soil chemical properties. – *Pedobiologia* 47: 205-212.
- FROUZ, J. & KALCIK, J. 1996: The role of ants for availability of phosphorus in spruce plantation. – *Proceedings of the National Conference about study of phosphorus and other elements, České Budějovice*: 75-80.
- FROUZ, J., KALCIK, J. & CUDLIN, P. 2005: Accumulation of phosphorus in nests of wood ants *Formica* s. str. – *Annales Zoologici Fennici* 42: 269-275.
- FROUZ, J., RYBNICEK, M., CUDLIN, P. & CHMELIKOVA, E. 2008: Influence of the wood ant, *Formica polyctena*, on soil nutrient and the spruce tree growth. – *Journal of Applied Entomology* 132: 281-284.
- FROUZ, J., SANTRUCKOVA, H. & KALCIK, J. 1997: The effect of wood ants (*Formica polyctena* FOERST) on the transformation of phosphorus in a spruce plantation. – *Pedobiologia* 4: 437-447.
- GALLÉ, L. 1973: Thermoregulation in the nest of *Formica pratensis* RETZ. (Hymenoptera, Formicidae). – *Acta Biologica Szeged* 19: 139-142.
- GASPAR, C. 1972: Action des fourmis du genre *Lasius* dans l'écosystème prairie. – *Ekologia Polska* 20: 145-152.
- GOEBEL, S. 1988: Freilanduntersuchungen an Waldameisenkolonien in Eichen- und Fichtenbeständen im Hinblick auf die Populationsdynamik von Rindenläusen (Aphidina) und deren Honigproduktion sowie die Auswirkungen auf die Insektenfauna. – PhD thesis, Albert-Ludwigs Universität, Freiburg, 272 pp.
- GORNY, H. 1976: Einige pedoökologische Probleme der Wirkung von industrieller Immission auf Waldstandorte. – *Pedobiologia* 16: 27-35.
- GOROSITO, N.B., CURMI, P., HALLAIRE, V., FOLGARAIT, P.J. & LAVELLE, P.M. 2006: Morphological changes in *Camponotus punctulatus* (MAYR) anthills of different ages. – *Geoderma* 132: 249-260.
- GREEN, W.P., PETTRY, D.E. & SWITZER, R.E. 1999: Structure and hydrology of mounds of the imported fire ants in the southeastern United States. – *Geoderma* 93: 1-17.
- GRIMALSKIJ, V.I.O. 1960: O roly rživich lesnich muravjev *Formica rufa* L. v lesnich biocenozach na Levoberežnom polesie Ukrainy. – *Zoologicheskij Zhurnal* 39: 394-398.
- GUERRA, M.B.B., SCHAEFER, C.E.G.R. & SOUSA-SOUTO, L. 2007: Chemical characteristics of nest refuse of *Atta sexdens rubropilosa* (Hymenoptera, Formicidae) reared with different substrates. – *Revista Brasileira de Ciencia do Solo* 31: 1185-1189.
- HOLEC, M. & FROUZ, J. 2006: The effect of two ant species *Lasius niger* and *Lasius flavus* on soil properties in two contrasting habitats. – *European Journal of Soil Biology* 42: 213-217.
- HÖLLDOBLER, B. & WILSON, E.O. 1990: The ants. – Springer Verlag, Berlin, 732 pp.
- HOLT, S.J. 1955: On the foraging activity of the wood ant. – *Journal of Animal Ecology* 24: 1-34.
- HORSTMANN, K. 1970: Untersuchungen über den Nahrungserwerb der Waldameisen (*Formica polyctena* FOERSTER) im Eichenwald. – *Oekologia* 5: 138-157.
- HORSTMANN, K. 1974: Untersuchungen über den Nahrungserwerb der Waldameisen (*Formica polyctena* FOERSTER) im Eichenwald. III. Jahresbilanz. – *Oecologia* 15: 187-204.
- HORSTMANN, K. 1983: Regulation der Temperatur in Waldameisen-Nestern (*Formica polyctena* FOERSTER). – *Zeitschrift für Naturforschung* 38: 508-510.
- HORSTMANN, K. 1987: Über die Reaktion von Waldameisen-Nestern (*Formica polyctena* FOERSTER) auf das Erscheinen von Glänzendschwarzen Holzameisen (*Lasius fuliginosus* LATREILLE) in ihren Nestern. – *Waldhygiene* 17: 23-28.
- HORSTMANN, K. 1990: Zur Entstehung des Wärmezentrums in Waldameisennestern (*Formica polyctena* FOERSTER, Hymenoptera, Formicidae). – *Zoologische Beiträge N.F.* 33: 105-124.
- HORSTMANN, K. & SCHMID, H. 1986: Temperature regulation in nests of the wood ant, *Formica polyctena* (Hymenoptera, Formicidae). – *Entomologia Generalis* 11: 229-236.
- HOWE, F.H. & SMALLWOOD, J. 1982: Ecology of seed dispersal. – *Annual Review of Ecology and Systematics* 13: 201-228.
- JAKUBCZYK, H., CZERWINSKI, Z. & PETAL, J. 1972: Ants as agents of the soil habitat changes. – *Ekologia Polska* 20: 153-163.
- JONES, C.G., LAWTON, J.H. & SHACHAK, M. 1994: Organisms as ecosystem engineers. – *Oikos* 69: 373-386.
- JONES, J.A. 1990: Termites, soil fertility and carbon cycling in dry tropical Africa: a hypothesis. – *Journal of Tropical Ecology* 6: 291-305.
- JONES, J.B. & WAGNER, D. 2006: Microhabitat-specific controls on soil respiration and denitrification in the Mojave Desert: The role of harvester ant nests and vegetation. – *Western North American Naturalist* 66: 426-433.
- JOUQUET, P., DAUBER, J., LAGERLÖF, J., LAVELLE, P. & LEPAGE, M. 2006: Soil invertebrates as ecosystem engineers: Intended and accidental effects on soil and feedback loops. – *Applied Soil Ecology* 32: 153-164.
- KILPELAINEN, J., FINER, L., NIEMELA, P., DOMISCH, T., NEUVONEN, S., OHASHI, M., RISCH, A.C. & SUNDSTROM, L. 2007: Carbon, nitrogen and phosphorus dynamics of ant mounds (*Formica rufa* group) in managed boreal forests of different successional stages. – *Applied Soil Ecology* 36: 156-163.
- KLOFT, W.J. 1978: Ökologie der Tiere. – UTB Ulmer, Stuttgart, 304 pp.

- KORGANOVA, G.A. & RAKHLEEVA, A.A. 2006: Shell amoebae (Testacea) in ant hills of *Formica lugubris*: Diversity and population structure. – Zoologichesky Zhurnal 85: 1281-1293.
- KOVAROVA, M., DOSTAL, P. & HERBEN, T. 2001: Vegetation of ant-hills in a mountain grassland: effects of mound history and of dominant ant species. – Plant Ecology 156: 215-227.
- KRISTIANSEN, S.M. & AMELUNG, W. 2001: Abandoned anthills of *Formica polyctena* and soil heterogeneity in a temperate deciduous forest: morphology and organic matter composition. – European Journal of Soil Science 52: 355-363.
- KRISTIANSEN, S.M., AMELUNG, W. & ZECH, W. 2001: Phosphorus forms as affected by abandoned anthills (*Formica polyctena* FOERSTER) in forest soils: sequential extraction and liquid-state P-31-NMR spectroscopy. – Journal of Plant Nutrition and Soil Science 164: 49-55.
- LAAKSO, J. & SETALA, H. 1997: Nest mounds of red wood ants (*Formica aquilonia*): hot spots for litter-dwelling earthworms. – Oecologia 111: 565-569.
- LAAKSO, J. & SETALA, H. 1998: Composition and trophic structure of detrital food webs in ant nest mounds of *Formica aquilonia* and in the surrounding forest soil. – Oikos 81: 266-278.
- LAFLEUR, B., BRADLEY, R.L. & FRANCOEUR, A. 2002: Soil modifications created by ants along a post-fire chronosequence in lichen-spruce woodland. – Ecoscience 9: 63-73.
- LAFLEUR, B., HOOPER-BUI, L.M., MUMMA, E.P. & GEAGHAN, J.P. 2005: Soil fertility and plant growth in soils from pine forests and plantations: effect of invasive red imported fire ants *Solenopsis invicta* BUREN. – Pedobiologia 49: 415-423.
- LAVELLE, P., BIGNELL, D., LEPAGE, M., WOLTERS, V., ROGERS, P., INESON, P., HEAL, O.W. & DHILLON, S. 1997: Soil function in a changing world: the role of invertebrate ecosystem engineers. – European Journal of Soil Biology 33: 159-193.
- LEE, K.E. 1982: The influence of earthworms and termites on soil nitrogen cycling. In: LEBRUN, P., ANDRE, H.M., DEMEDS, A., GREGORIE-WIBO, C. & WAUTHY, G. (Eds.): Proceedings of the VIII International Colloquium of Soil Zoology. Louvial La Neuve August 30 - September 2: 35-48.
- LESICA, P. & KANOWSKI, P.B. 1998: Ants create hummocks and alter structure and vegetation of a Montana Fen. – American Midland Naturalist 139: 58-68.
- LOBRY DE BRUYN, L.A. & CONACHER, A.J. 1990: The role of termites and ants in soil modification: a review. – Australian Journal of Soil Research 28: 55-93.
- LYFORD, W.H. 1963: Importance of ants to brown podzolic soil genesis in New England. – Harvard Forest Paper, Harvard University, Cambridge, MA, 7: 1-18.
- MALOZEMOVA, L.A. & KORUMA, N.P. 1973: O vlnaniji muravjev na počvu. – Ekologia 4: 98-101.
- MCCAHERN, T.J. & LOCKWOOD, J.A. 1990: Nest architecture and pedoturbation of *Formica obscuripes* FOREL (Hymenoptera, Formicidae). – Pan-Pacific Entomologist 66: 147-156.
- MIKHEYEV, A.S. & TSCHINKEL, W.R. 2004: Nest architecture of the ant *Formica pallidefulva*: structure, costs and rules of excavation. – Insectes Sociaux 51: 30-36.
- NIELSEN, M.G. 1986: Ant nests on tidal meadows in Denmark. – Entomologia Generalis 11: 191-195.
- NIEMELA, P. & LAINE, K.J. 1986: Green islands – predation not nutrition. – Oecologia 68: 476-478.
- NKEM, J.N., DE BRUYN, L.A.L., GRANT, C.D. & HULUGALLE, N.R. 2000: The impact of ant bioturbation and foraging activities on surrounding soil properties. – Pedobiologia 44: 609-621.
- OEKLAND, F. 1930: Wieviel Blattlauszucker verbraucht die Rote Waldameise. – Biologisches Zentralblatt 50: 449-459.
- OHASHI, M., FINER, L., DOMISCH, T., RISCH, A.C. & JURGENSEN, M.F. 2005: CO<sub>2</sub> efflux from a red wood ant mound in a boreal forest. – Agricultural and Forest Meteorology 130: 131-136.
- OHASHI, M., KILPELAINEN, J., FINER, L., RISCH, A.C., DOMISCH, T., NEUVONEN, S. & NIEMELA, P. 2007a: The effect of red wood ant (*Formica rufa* group) mounds on root biomass, density, and nutrient concentrations in boreal managed forests. – Journal of Forest Research 12: 113-119.
- OHASHI, M., FINER, L., DOMISCH, T., RISCH, A.C., JURGENSEN, M.F. & NIEMELA, P. 2007b: Seasonal and diurnal CO<sub>2</sub> efflux from red wood ant (*Formica aquilonia*) mounds in boreal coniferous forests. – Soil Biology & Biochemistry 39: 1504-1511.
- PETAL, J. 1978: The role of ants in ecosystems. In: BRIAN, M.V. (Ed.): Production ecology of ants and termites. – Cambridge University Press, London, pp. 293-325.
- PETAL, J. 1980: Ant populations, their regulation and effect on soil in meadows. – Ekologia Polska 28: 297-326.
- PETAL, J. 1998: The influence of ants on carbon and nitrogen mineralization in drained fen soils. – Applied Soil Ecology 9: 271-275.
- PETAL, J., CHMIELEWSKI, K., KUSINKA, A., KACZOROWSKA, R., STACHURSKY, A. & ZIMKA, J. 2003: Biological and chemical properties of fen soils affected by anthills of *Myrmica* spp. – Polish Journal of Ecology 51: 67-78.
- PETAL, J., NOWAK, E., JAKUBCZYK, H. & CZERWINSKI, Z. 1977: Effect of ants and earthworms on soil habitat modification. – Ecological Bulletins NFR Statens Naturvetenskapliga Forskningsrad 25: 501-503.
- POKARYHEVSKIJ, A.D. 1981: The distribution and accumulation of nutrients in nests of ant *Formica polyctena* (Hymenoptera, Formicidae). – Pedobiologia 21: 117-124.
- RAIGNIER, A. 1948: L'économie thermique d'une colonie polycaulyque de la fourmi des bois *Formica rufa polyctena* FOERST. – La Cellule 51: 279-368.
- RISCH, A.C., JURGENSEN, M.F., SCHUTZ, M. & PAGE-DUMROESE, D.S. 2005: The contribution of red wood ants to soil C and N pools and CO<sub>2</sub> emissions in subalpine forests. – Ecology 86: 419-430.
- ROGERS, L.E. 1972: The ecological effect of the western harvester ant (*Pogonomyrmex occidentalis*) in the shortgrass plains ecosystem. – USA/IBP Grassland Biome, Technical report, 206 pp.
- ROSENGREN, R., FORTELIUS, W., LINDSTROM, K. & LUTHER, A. 1987: Phenology and causation of nest heating and thermoregulation in red wood ants of the *Formica rufa* group studied in coniferous forest habitats in southern Finland. – Annales Zoologici Fennici 24: 147-155.
- ROSENGREN, R. & SUNDSTROM, L. 1991: The interaction between wood ants, *Cinara* aphids and pines. A ghost of mutualism past? In: HUXLEY, C.R. & CUTLER, D.F. (Eds.): Ant plant interactions. – Oxford University Press, Oxford, New York & Tokyo, pp. 81-91.
- ROTHANZL, J., KOTOUČOVÁ, M., HRABINOVÁ, I., PLAČKOVÁ, I. & HERBEN, T. 2007: Genetic differentiation of *Agrostis capillaris* in a grassland system with stable heterogeneity due to terricolous ants. – Journal of Ecology 95: 197-207.
- SNYDER, S.R., CRIST, T.O. & FRIESE, C.F. 2002: Variability in soil chemistry and arbuscular mycorrhizal fungi in harvester ant nests: the influence of topography, grazing and region. – Biology and Fertility of Soils 35: 406-413.
- STADLER, B., SCHRAMM, A. & KALBITZ, K. 2006: Ant-mediated effects on spruce litter decomposition, solution chemistry, and microbial activity. – Soil Biology & Biochemistry 38: 561-572.



- STEINER, A. 1924: Über den sozialen Wärmehaushalt der Waldameise (*Formica rufa*). – Zeitschrift für Vergleichende Physiologie 2: 23-56.
- STERNBERG, L.D., PINYON, M.C., MORERA, M.Z., MOUTINHO, P., ROJAS, E.I. & HERRE, E.A. 2007: Plants use macronutrients accumulated in leaf-cutting ant nests. – Proceedings of the Royal Society B-Biological Sciences 274: 315-321.
- STRADLING, D.J. 1978: Food and feeding habits of ants. In: BRIAN, M.V. (Ed.): Production ecology of ants and termites. – Cambridge University Press, Cambridge, MA, pp. 81-106.
- TALBOT, M. 1953: Ant of old field community on the Edwin S. Geirge Reserve, Livingston country, Michigan. – Contributions from the Laboratory of Vertebrate Biology of the University of Michigan 69: 1-9.
- TSCHINKEL, W.R. 2004: The nest architecture of the Florida harvester ant, *Pogonomyrmex badius*. – Journal of Insect Science 4: article number 21.
- VINSON, S.B. 1991: Effect of the red imported fire ant (Hymenoptera, Formicidae) on a small plant-decomposing arthropod community. – Environmental Entomology 20: 98-103.
- WAGNER, D. 1997: The influence of ant nests on *Acacia* seed production, herbivory and soil nutrients. – Journal of Ecology 85: 83-93.
- WAGNER, D. & JONES, J.B. 2004: The contribution of harvester ant nests, *Pogonomyrmex rugosus* (Hymenoptera, Formicidae), to soil nutrient stocks and microbial biomass in the Mojave Desert. – Environmental Entomology 33: 599-607.
- WAGNER, D. & JONES, J.B. 2006: The impact of harvester ants on decomposition, N mineralization, litter quality, and the availability of N to plants in the Mojave Desert. – Soil Biology & Biochemistry 38: 2593-2601.
- WAGNER, D. & JONES, J.B. & GORDON, D.M. 2004: Development of harvester ant colonies alters soil chemistry. – Soil Biology & Biochemistry 36: 797-804.
- WANG, D., MCSWEENEY, K., LOWERY, B. & NORMAN, J.M. 1995: Nest structure of ant *Lasius neoniger* EMERY and its implications to soil modification. – Geoderma 66: 259-272.
- WAY, M.J. & KHOO, K.C. 1992: Role of ants in pest management. – Annual Review of Entomology 37: 479-503.
- WELLENSTEIN, G. 1952: Zur Ernährungsbiologie der Roten Waldameise. – Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 59: 430-451.
- ZACHAROV, A.A., IVANICKAJA, E.F. & MAKSIMOVA, A.E. 1981: Nakoplenie elementov v gnezdach ryžich lesnych muravev. – Pedobiologia 21: 36-45.
- ZARAGOZA, S.R., WHITFORD, W.G. & STEINBERGER, Y. 2007: Effects of temporally persistent ant nests on soil protozoan communities and the abundance of morphological types of amoeba. – Applied Soil Ecology 37: 81-87.
- ZHOU, H., CHEN, J. & CHEN, F. 2007: Ant-mediated seed dispersal contributes to the local spatial pattern and genetic structure of *Globba lancangensis* (Zingiberaceae). – Journal of Heredity 98: 317-324.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Myrmecological News = Myrmecologische Nachrichten](#)

Jahr/Year: 2008

Band/Volume: [011](#)

Autor(en)/Author(s): Frouz J., Jilkova V.

Artikel/Article: [The effect of ants on soil properties and processes \(Hymenoptera: Formicidae\). 191-199](#)