

Aloa WALKER 1855 and Amsacta WALKER 1855, two distinct genera of Arctiidae (Lepidoptera)

by

Werner THOMAS⁽¹⁾ and David GOODGER

Abstract: *Aloa* and *Amsacta* are redefined as distinct genera and a catalogue of species is provided. The genus *Aloa* is reinstated for the type species *lactinea* CRAMER 1777 and its relatives, i. e. the Australasian species formerly included under *Amsacta*. *Amsacta* becomes restricted to the Afrotropical region only.

Aloa WALKER 1855 und *Amsacta* WALKER 1855, zwei getrennte Arctiiden-Gattungen (Lepidoptera)

Zusammenfassung: Die Gattungen *Aloa* und *Amsacta* werden neu definiert, und eine Artenliste (Katalog) wird gegeben. Die Gattung *Aloa* wird für die Typusart *lactinea* CRAMER 1777 und ihre Verwandten wiedereingerichtet, das bedeutet für die australasiischen Arten (Australien, Neuguinea und umliegende Inseln), die früher unter *Amsacta* geführt wurden. *Amsacta* wird hierdurch auf die afrotropische Faunenregion begrenzt.

Introduction

WALKER (1855 b) described *Amsacta* as a monotypic genus within the Liparidae, now Lymantriidae, with *marginalis* WALKER 1855 as the only included species. HAMPSON (1901) used *Amsacta* within the Arctiidae, based on venation characters, for a group of 17 species including *Phalaena lactinea* CRAMER 1777, subsequently cited as type species for *Aloa* WALKER 1855 by MOORE ([1883]) (WATSON et al. 1980). In his supplement, HAMPSON (1920) added another 8 species to *Amsacta*. Today, after further descriptions and transfers from other genera, *Amsacta* currently contains 31 species.

(1) = Contributions to the systematics of the Arctiidae, no. 5. – Beiträge zur Arctiiden-Systematik, Nr. 5.

The disappearance of *Aloa* into synonymy is also due to HAMPSON (1901). HAMPSON (1901) treated *Aloa* as a synonym of *Diacrisia* HÜBNER [1819] 1816 based on his invalid designation of *Spilosoma lineata* WALKER 1855 as type species (WATSON et al. 1980). MOORE ([1883]) had already designated *Phalaena lactinea* CRAMER 1777 as type species of *Aloa*.

Subsequent authors all followed HAMPSON in treating *Phalaena lactinea* CRAMER and its relatives as congeneric with *Amsacta marginalis* WALKER.

The two type species *Amsacta marginalis* and *Aloa lactinea* are generically distinct. The genus *Aloa* is reinstated for the species *lactinea* and its relatives. All Australasian species are extracted from *Amsacta* which becomes restricted to the Afrotropical region only. In addition, many other African species must be placed outside *Amsacta* into other genera. As these species must be subject to a thorough revision to find their correct generic placement they are listed as "AMSACTA SENSU LATO".

Catalogue entries

Each species entry includes the name, authorship, date of publication, pagination and the genus with which the name was originally combined. This is followed by a statement about the type material: Holotype or syntype(s) (the "s" in parentheses indicating that the author of the taxon neither designated a holotype nor stated how many specimens were before him when he described the taxon). Entries inset and in small types placed below a valid name are junior synonyms, homonyms, incorrect spellings or infrasubspecific names.

Country of origin is cited from the original description. Square brackets are used to indicate that we have had to rely on evidence external to the original description in determining the country of origin. Following the country is a quotation from the original description concerning the place of capture within the country.

Catalogue

***Aloa* WALKER, 1855 a: 699.**

Type species *Phalaena lactinea* CRAMER, 1777: 58.

Forewing of type species white, with costa edged in red. Hind wing white, with dark circular markings (Figs. 1, 3 & 4). Abdomen white ven-

trally, yellow with wide dark bands dorsally. Antenna laminate. Other species are similar except for *senegalensis*, *rubricostata*, *nalonyi*, *albistriga* which have the veins on the forewing picked out in brown scales. *A. costalis* has dark fascia on the forewing and *negrita* has black forewings with white streaks.

Uncus broad; weakly bifurcate distally. Juxta triangular. Wide tegumen. Vesica scobinate with cornuti. Valves narrow; pointed apically. Saccus broad and rounded. Eighth abdominal tergite bifurcate posteriorly; eighth sternum with lateral sclerites. Coremata absent.

albistriga WALKER, 1865: 303 (*Aloa*). Comb. rev.

Holotype "South Hindostan".

ab. *confluens* ROTHSCILD, 1914: 251 (*Amsacta*). Comb. n.
Syntypes India.

ab. *sarala* STRAND, 1919: 244 (*Amsacta*). Comb. n.
Holotype [India] "Trichinopoly".

arabicum HAMPSON, 1896: 258 (*Spilosoma*). Comb. n.

Syntypes Aden.

gracilis STAUDINGER, 1899: 354 (*Spilosoma*). Comb. n.
Holotype Palestine.

ab. *pedunculata* STRAND, 1919: 244 (*Creatonotos*). Comb. n.
Holotype Aden.

cardinalis BUTLER, 1875: 22 (*Areas*). Comb. n.

c. *cardinalis* BUTLER, 1875: 22.

Syntype(s) [Philippines] "Philippine Islands".

c. *celebensis* ROTHSCILD, 1910: 158 (*Amsacta*). Comb. n.
Holotype [Sulawesi] "Celebes".

ab. *celebesica* STRAND, 1919: 238 (*Amsacta*). Comb. n.
Holotype [Sulawesi] "Celebes".

c. *luteomarginata* ROTHSCILD, 1910: 158 (*Amsacta*). Comb. n.
Syntypes Timor.

c. *reducta* ROTHSCILD, 1910: 158 (*Amsacta*). Comb. n.
Syntypes [Indonesia] "Toekan Besi Islands".

flavimargo HAMPSON, 1894: 27 (*Creatonotos*). Comb. n.
Syntype(s) [Myanmar] Burma "Bhama".

collaris HAMPSON, 1891: 54 (*Aloa*). Comb. rev.
Syntypes India "Nilgiris".

costalis WALKER, 1865: 301 (*Aloa*). Comb. rev.
Syntypes Northern Australia.

corsima SWINHOE, 1892: 171 (*Aloa*). Comb. rev.
Holotype Australia "Queensland".

lactinea CRAMER, 1777: 58 (*Phalaena*).
Syntype(s) [Indonesia] "Batavia".

sanguinolenta FABRICIUS, 1793: 473 (*Bombyx*).
Syntype(s) "India Orientali"

frederici KIRBY, 1892: 223 (*Aloa*). Comb. rev.
Holotype Nepal.

Replacement name for *marginata* MOORE, 1883: 15.

f. *nigrita* HAMPSON, 1894: 28 (*Creatonotos*). Comb. n., stat. n.
Syntypes India "Cheri-Punji".

marginata MOORE, 1883: 15 (*Aloa*).
Junior homonym of *marginata* DONOVAN, 1805.

marginata DONOVAN, 1805: t. 34 fig. 2 (*Bombyx*). Comb. n.
Australia.

roseicostis BUTLER, 1875: 23 (*Areas*). Comb. n.
Syntype(s) Cape York.


punctipennis BUTLER, 1876: 126 (*Areas*). Comb. n.
Syntypes Australia.

gangara SWINHOE, 1892: 171 (*Aloa*). Comb. rev., syn. n.
Syntypes Australia.

ab. *slaveola* ROTHSCHILD, 1914: 251 (*Amsacta*). Comb. n.
Syntypes numerous localities.

ab. *simplicans* STRAND, 1919: 243 (*Amsacta*). Comb. n.
Syntypes Western Australia

ab. *gangarides* STRAND, 1919: 240 (*Amsacta*). Comb. n.
Syntypes Western Australia.



Figs. 1, 3, 4: *Phalaena lactinea* CRAMER 1777. Type species of *Aloa* WALKER 1855. Male specimen, male genitalia and aedeagus.

Figs. 2, 5, 6: *Amsacta marginalis* WALKER 1855. Type species of *Amsacta* WALKER 1855. Male specimen, male genitalia and aedeagus.

moloneyi DRUCE, 1887: 670 (*Areas*). Comb. n.

Syntype(s) Gambia.

hampsoni ROTHSCILD, 1910: 157.

Holotype Ghana "Gambaga".

senegalensis ROTHSCILD, 1933: 180 (*Amsacta*). Comb. n.

Syntypes Senegal "Sedhiou".

moorei BUTLER, 1875: 23 (*Areas*). Comb. n.

Syntype(s) "Almarah".

sara SWINHOE, 1889: 404 (*Aloa*). Comb. rev.

Syntypes [Pakistan] "Karachi".

***Amsacta* WALKER, 1855 b: 804.**

Type species: *Amsacta marginalis* WALKER, 1855 b: 805.

Unmarked brown wings bordered with yellow. Male antenna bipectinate. Foretibia with well developed terminal spines (Figs. 2, 5 & 6).

Uncus broad; weakly trifurcate posteriorly; dorsal surface spinose posteriorly. Valve digitate. Juxta tubular; emarginate posteriorly; fused with arms of transtilla. Vesica scobinate, but without cornuti. Eighth abdominal sternum with lateral sclerites. Coremata present between seventh and eighth sterna.

fuscosa BARTEL, 1903: 194 (*Acantharctia*). Stat. rev.

Lectotype Togo "Bismarckburg".

latimarginalis ROTHSCILD, 1933: 180 (*Amsacta*). Comb. n.

[Sudan] "Tambura, Southern Bahr-el-Ghazal".

marginalis WALKER, 1855 b: 805 (*Acantharctia*). Comb. n.

Syntypes Sierra Leone.

AMSACTA SENSU LATO

aliena KIRIAKOFF, 1954: 184 (*Amsacta*).

Holotype Zaïre "Kabongo (Lualaba)".

aureolimbata ROTHSCILD, 1910: 157 (*Amsacta*).

Syntypes Angola "Bihe".

bicoloria GÄDE, 1916: 125 (*Acantharctia*).
Holotype Cameroun, north "Garua-Reibuba".

stygioides ROTHSCHILD 1916: 271 (*Estigmene*).
Syntypes Nigeria "Abinsi, River Benue".

duberneti TOULGOET, 1968: 76 (*Amsacta*).
Holotype Madagascar "Nord d'Ankazobe".

emittens WALKER, 1855: 638 (*Creatonotos*).
Syntypes Sri Lanka.

flora SWINHOE, 1885: 296 (*Aloa*).
Syntype(s) India "Bombay".

grammiphlebia HAMPSON, 1901: 323 (*Amsacta*).
Syntypes [Zimbabwe] "Mashonaland, Salisbury".

lineola FABRICIUS, 1793: 465 (*Bombyx*).
India.

punctistriga WALKER, 1855 a: 676 (*Spilosoma*).
Holotype Northern India.

candidula WALKER, 1855 a: 704 (*Aloa*).
India "Kimorak".

diminuta WALKER, 1855 a: 705 (*Aloa*).
Holotype India.

strigata WALKER, 1869: 10 (*Spilosoma*).
"?Punjab".

rubicosta MOORE, 1872: 573 (*Creatonotos*).
Syntypes India "Manpuri & Bombay".

insolata SWINHOE, 1889: 404 (*Aloa*).
Syntypes India.

felderii ROTHSCHILD, 1910: 124 (*Diacrisia*).
India "Kotegurh".

octomaculata ROTHSCHILD, 1933: 185 (*Estigmene*).
Syntypes "Solane".

ab. *assamica* STRAND, 1919: 242 (*Amsacta*).
Syntypes India "Assam".

ab. *kaschmirica* STRAND, 1919: 242 (*Amsacta*).
Syntypes "Kashmir".

ab. *isolatana* STRAND, 1919: 240 (*Amsacta*).
Syntypes Burma "Myingyan".

melanogastra HOLLAND, 1897: 410 (*Cycnia*).

Syntype(s) [Somalia or Ethiopia] "Gof".

epicaste FAWCETT, 1915: 93 (*Diacrisia*).

Syntype(s) [Kenya] "Masongaleni"

occidentalis BARTEL 1903: 191 (*Spilosoma*).

Syntype(s) [Namibia] "Outjo".

nivea HAMPSON, 1916: 237 (*Amsacta*).

Holotype [South Africa] "E. Transvaal, Karino".

paolii BERIO, 1936: 27 (*Amsacta*).

Holotype Somalia "Villaggio Duca degli Abruzzi".

References

- BARTEL, M. (1903): Neue aethiopische Arctiidae der Sammlung des Kgl. Museums für Naturkunde in Berlin. — Deutsche Entomologische Zeitschrift Iris, Dresden, **16**: 170–220.
- BERIO, E. (1936): Specie nuove di Lepidotteri della Somalia Italiana. — Bollettino della Società Entomologica Italiana **68**: 27.
- CRAMER, P. (1777): De Uitlandsche Kapellen, vol. 2. — 152 pp., Amsterdam, Utrecht.
- DONOVAN, E. (1805): Insects of New Holland. — 41 pls., 51 pp., London.
- DRUCE, H. (1887): Descriptions of some new species of Heterocera. — Proceedings of the Zoological Society of London **1887**: 777–782.
- FABRICIUS, J. C. (1793): Entomologiae systematicae et auctae, vol. 3. — [vi] + 488 pp., Copenhagen.
- FAWCETT, J. M. (1915): Notes on a collection of Heterocera made by Mr W. FEATHER in British East Africa, 1911–12. — Transactions of the Zoological Society of London **17**: 165–184.
- GAEDE, M. (1916): Neue afrikanische Lepidoptera des Berliner Zoologischen Museums. — Internationale Entomologische Zeitschrift **9**: 125–126.
- HAMPSON, G. F. (1894): The Fauna of British India including Ceylon and Burma, Moths, vol. 2. — 609 pp., London.
- (1896): On moths collected at Aden and in Somaliland. — Proceedings of the Zoological Society of London **1896**: 257–283.
- (1901): Catalogue of the Lepidoptera Phalaenae in the British Museum, vol. 3. — xix + 690 pp., London.
- (1916): Descriptions of new species of the family Arctiidae in the British Museum. — Novitates Zoologicae **23**: 230–240.
- (1920): Catalogue of the Lepidoptera Phalaenae in the British Museum, vol. 2 (supplement). — xxiii + 619 pp., London.
- KIRIAKOFF, S. G. (1954): Hétérocères nouveaux ou peu connues du Katanga. — Revue de Zoologie et de Botanique Africaines **50**: 169–188.
- KIRBY, W. F. (1892): A synonymic catalogue of Lepidoptera Heterocera. — xi + 951 pp., London.

Sonderheft/Special Issue in memoriam Werner Thomas

- MOORE, F. D. (1872): Descriptions on new Indian Lepidoptera. – Proceedings of the Zoological Society of London **1872**: 555–583.
- (1883) The Lepidoptera of Ceylon, vol. 2. – viii + 162 pp., London.
- ROTHSCHILD, L. W. (1910): Catalogue of the Arctiinae in the Tring Museum with notes and descriptions of new species. – Novitates Zoologicae **17**: 113–188.
- (1914): Arctiidae, in: SEITZ, A. (ed.), Die Gross-Schmetterlinge der Erde **10**: 238–263. – Stuttgart.
- (1933): New species and sub-species of Arctiinae. – Annals and Magazine of Natural History (10) **11**: 167–194.
- STAUDINGER, O. (1899): Ueber Lepidopteren aus dem östlichsten Thian Schan-Gebiet. – Deutsche Entomologische Zeitschrift Iris, Dresden, **12**: 329–351.
- STRAND, E. (1919): Arctiidae; subfam. Arctiinae, in: STRAND, E. (ed.), Lepidopterorum Catalogus, vol. 22. – 416 pp., Berlin.
- SWINHOE, C. (1885): On the Lepidoptera of Bombay and the Deccan. – Proceedings of the Zoological Society of London **1885**: 287–307.
- (1892): Catalogue of Eastern and Australian Lepidoptera Heterocera in the collection of the Oxford University Museum, vol. 1. – vii + 324 pp., Oxford.
- WALKER, F. (1855 a): List of the Specimens of Lepidopterous Insects in the Collection of the British Museum, vol. 3. – pp. 583–775, London.
- (1855 b): List of the Specimens of Lepidopterous Insects in the Collection of the British Museum, vol. 4. – pp. 776–976, London.
- (1865): List of the Specimens of Lepidopterous Insects in the Collection of the British Musem, vol. 31. – pp. 1–321, London.
- (1869): Characters of undescribed Lepidoptera Heterocera. – 112 pp., London.
- WATSON, A., FLETCHER, D. S., & NYE, I. W. B. (1980): Noctuoidea (part): Arctiidae, Lymantriidae, Cocytiidae, Notodontidae, Ctenuchidae, Strepsimanidae, Dilobidae, Thaumetopoeidae, Diptidae, Thyretidae, in: NYE, I. W. B. (ed.), The generic names of moths of the world, vol. 2. – xiv + 228 pp., London.

Address of the author:

(Dr. Werner THOMAS died on February 28th, 1991)

David GOODGER, Department of Entomology, The Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom

Reprint requests to D. GOODGER.

ENTOMOLOGISCHE NOTIZ

**"Phragmatobia" (besser *Spilosoma*?) caesarea im Raum Schlüchtern,
Main-Kinzig-Kreis (Lepidoptera, Arctiidae)**

Zuerst einige systematische Vorbemerkungen: Das kleine Bärenspinnerchen wird in Bestimmungsbüchern meist als "Arctinia" caesarea geführt, jedoch ist

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nachrichten des Entomologischen Vereins Apollo](#)

Jahr/Year: 1992

Band/Volume: [13](#)

Autor(en)/Author(s): Thomas Werner, Goodger David

Artikel/Article: [Aloa Walker 1855 and Amsacta Walker 1855, two distinct genera of Arctiidae 297-305](#)