

Orthonama obstipata FABRICIUS, 1794 (Geometridae: Larentiinae) — new for the fauna of Borneo

Christian H. SCHULZE and Konrad FIEDLER

Dr. Christian H. SCHULZE and Prof. Dr. Konrad FIEDLER¹, Department of Animal Ecology I, University of Bayreuth, D-95440 Bayreuth, Germany; email: christian.schulze@uni-bayreuth.de

Abstract: The first records of *Orthonama obstipata* FABRICIUS, 1794 (Geometridae: Larentiinae, Xanthorhoini) — a geometrid moth with almost worldwide distribution — for Borneo are presented. One specimen each was collected at the south-eastern border of Mount Kinabalu Park (Sabah, East Malaysia) in the vicinity of Poring Hot Springs in March and September 1997, respectively.

***Orthonama obstipata* FABRICIUS, 1794 (Geometridae: Larentiinae) — neu für die Fauna von Borneo**

Zusammenfassung: Die ersten Nachweise der fast weltweit verbreiteten Wanderfalterart *Orthonama obstipata* FABRICIUS, 1794 (Geometridae: Larentiinae, Xanthorhoini) für die Insel Borneo werden berichtet. Je ein Falter wurde im März und September 1997 an der Südostgrenze des Kinabalu-Nationalparks (Sabah, Ostmalaysia) in der Umgebung von Poring Hot Springs gefunden.



Fig. 1: *Orthonama obstipata* (FABRICIUS, 1794) — ♀, East Malaysia, Sabah, Kinabalu Park, Poring Hot Springs, 14. III. 1997.

Introduction

In the years 1997–1999 we intensively surveyed the moth fauna of Mount Kinabalu Park. The major aim of our studies was to assess how moth communities respond to anthropogenic changes in land-use (e.g. SCHULZE 2000, BECK et al. 2002, FIEDLER & SCHULZE in press). For these purposes, we did light trapping at a variety of elevational zones and in habitats ranging from cultivated areas to primary rainforest. One focal point for our studies was the vicinity of Poring Hot Springs, situated at the south-eastern border of Mount Kinabalu National Park at an altitude of 450 to 500 m a.s.l. (Sabah, East Malaysia) (for details about the area see WONG & PHILLIPPS 1996 and HÄUSER et al. 1997).

Results

Our samples include two ♀ specimens of *Orthonama obstipata* FABRICIUS, 1794 (Geometridae: Larentiinae: Xanthorhoini). Both vouchers are preserved in coll. C. H. SCHULZE, University of Bayreuth. To our knowledge, these are the first specimens of this widespread geometrid moth recorded from Borneo. The species is not mentioned by HOLLOWAY (1997). One specimen (Fig. 1) was attracted by light between 20:00 h and 20:30 h on 14. III. 1997, the other between 20:30 h and 21:00 h on 8. IX. 1997. Both individuals were collected in disturbed habitats. One light trapping site was situated within a secondary forest, the second one in an open cultivated area adjacent to a bamboo thicket and secondary vegetation growing at the edge of a small river.

Because of its characteristic appearance, it is surprising that *O. obstipata* has not been reported earlier from Borneo. The species did not evolve any pronounced geographical variation in wing colouration or any other morphological features within its vast distributional range, and just one subspecies (*contrariata* WALKER, 1862, described from Isle Tristan da Cunha in the southern Atlantic ocean) is accepted by SCOBLE (1999). However, there is a conspicuous sexual dimorphism (see colour plates in COVELL 1984 and SKOU 1984). In ♂♂ the wing colouration is yellowish brown with a darker area between the wing base and the median line. A small black discoidal spot is located in the centre of the forewing. The lines on both wings are generally obscure. The wing colouration in ♀♀, in contrast, is much darker and the black discoidal spot on the forewing is characteristically encircled with white (Fig. 1). Published data on wingspan range between 1.5 and 2.2 cm (e.g. COVELL 1984). The two Bornean specimens with a wingspan of ca. 2.0 cm fit very well into this range.

O. obstipata is distributed almost worldwide including North and South America, Africa, Europe, and throughout Southern and Eastern Asia from India to China, the Philippines, Taiwan and Japan (e.g. COVELL 1984, HEPPNER & INOUE 1992, KARSHOLT & RAZOWSKI 1996). However, records of *O. obstipata* from the Malay Archipelago as well as from Australia and New Zealand are apparently absent (for Australia: NIELSEN et al. 1996). COVELL (1984) mentioned that *O. obstipata* dies out each year in the colder areas of North America, but repopulates rapidly the following spring. Migratory behaviour which may

¹ New address after September 2004: Division of Population Ecology, Institute of Ecology and Conservation Biology, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria; email: konrad.fiedler@univie.ac.at

result in annual colonization of areas, but without establishing permanent populations is also documented for many parts of central and northern Europe (e.g. VOJNITS 1966, SKOU 1984, KAABER 1997, PALMQVIST 1999, EBERT 2001). So far as we know, migration of *O. obstipata* has not been documented from South East Asia. However, the apparent rarity of this species on Borneo and the fact that it was recorded as late as 1997 for the first time, may indicate that this Geometridae species has reached northern Borneo just recently (for example via the Philippines?). It should be noted that our samples were taken before the severe El Niño drought which facilitated the large-scale forest fires in Borneo in the year 1998 (SIEGERT et al. 2001).

With regard to the range of larval foodplants *O. obstipata* must be classified as a generalist feeding on a wide variety of herbaceous as well as woody plant taxa (summarised in SCOBLE 1999). These generalized feeding habits for certain facilitate the wide geographical distribution of the moth. Remarkably, many of its host plants known to be utilized in Europe and North America are weeds growing in agricultural areas, such as *Senecio*, *Eupatorium* (Asteraceae), *Convolvulus* (Convolvulaceae), or *Polygonum* (Polygonaceae). Thus, *O. obstipata* should find appropriate host plants also at the localities around Poring. Since our two records were separated by almost half a year, it seems safe to assume that the species has established at Poring for at least a couple of generations (generation time of the moth under warm climatic conditions less than 2 months).

Acknowledgements

We thank Mrs. Munirah Abd. MANAN (Economic Planning Unit, Kula Lumpur) for granting permission to conduct our research. Special thanks are due to Prof. Dr. Mohamed MARYATI for helpful comments and advice during our stay in Sabah and to Datuk Lamri ALI and Frances LIEW for granting permission to work at Mount Kinabalu National Park. Additionally, we thank the staff of Sabah Parks for their kind support of our studies and the supply of accommodation during the stay. The light trapping in 1997 would have not been possible without the help of Jan BECK, Sabine BRAMMERTZ, André HERZIG, Jutta KLEIN, Annette MALSCH, Evi and Roswitha MÜHLENBERG, Dr. Wolfgang A. NÄSSIG and Edda RODDEWIG. Our research was supported by the Deutsche Forschungsgemeinschaft (Fi 547/4-1 and 547/4-2).

References

- BECK, J., SCHULZE, C. H., LINSENMAIR, K. E., & FIEDLER, K. (2002): From forest to farmland: diversity of geometrid moths along two habitat gradients on Borneo. — *Journal of Tropical Ecology* **18**: 33–51.
- COVELL, C. V. jr. (1984): *A field guide to the moths of eastern North America*. — Boston (Houghton Mifflin), xv + 496 pp.
- EBERT, G. (ed.) (2001): *Die Schmetterlinge Baden-Württembergs, Band 8: Nachtfalter VI*. — Stuttgart (Eugen Ulmer), 541 pp.
- FIEDLER, K., & SCHULZE, C. H. (in press): Forest modification affects diversity, but not dynamics of speciose tropical pyraloid moth communities. — *Biotropica* (in press).
- HÄUSER, C. L., SCHULZE, C. H., & FIEDLER, K. (1997): The butterfly species (Insecta: Lepidoptera: Rhopalocera) of Kinabalu Park, Sabah. — *The Raffles Bulletin of Zoology* **45**: 281–304.
- HEPPNER, J. B., & INOUE, H. (1992): *Lepidoptera of Taiwan, Vol. 1, Part 2: Checklist*. — Gainesville (Association of Tropical Lepidoptera & Scientific Publishers), xlix + 276 pp.
- HOLLOWAY, J. D. (1997): The moths of Borneo: family Geometridae, subfamilies Sterrhinae and Larentiinae. — *Malayan Nature Journal* **51**: 1–242.
- KAABER, S. (1997): Observations on three migrations of Lepidoptera to the Faroe Islands in 1996. — *Entomologiske Meddelelser* **65** (3): 109–118.
- KARSHOLT, O., & RAZOWSKI, J. (eds.) (1996): *The Lepidoptera of Europe. A distributional checklist*. — Stenstrup (Apollo Books), 380 pp.
- NIELSEN, E. S., EDWARDS, E. D., & RANGSI, T. V. (eds.) (1996): *Checklist of the Lepidoptera of Australia*. — Monographs on Australian Lepidoptera, vol. 4. Collingwood (CSIRO Publ.), xiv + 529 pp.
- PALMQVIST, G. (1999): Remarkable records of Macrolepidoptera in Sweden 1998. — *Entomologisk Tidskrift* **120** (1/2): 59–74.
- SCHULZE, C. H. (2000): *Auswirkungen anthropogener Störungen auf die Diversität von Herbivoren – Analyse von Nachfalterzönosen entlang von Habitatgradienten in Ost-Malaysia*. — Ph.D. thesis, University of Bayreuth, 349 pp.
- SCOBLE, M. J. (1999): *Geometrid moths of the world*. — Collingwood (CSIRO Publ.), xxv + 1016 + 129 pp.
- SIEGERT, F., RUECKER, G., HINRICH, A., & HOFFMANN, A. A. (2001): Increased damage from fires in logged forests during droughts caused by El Niño. — *Nature* **414**: 437–440.
- SKOU, P. (1984): *Nordens Målere. Danmarks Dyreliv Bind 2*. — Copenhagen, Svendborg (Apollo Bøger), 332 pp.
- VOJNITS, A. (1966): *Wanderfalter in Ungarn 1964*. — *Atalanta* **1**: 7.
- WONG, K. M., & PHILLIPPS, A. (eds.) (1996): *Kinabalu: summit of Borneo. A revised and expanded edition*. — Kota Kinabalu (The Sabah Society & Sabah Parks), xxviii + 544 pp.

Received: 5. III. 2004

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nachrichten des Entomologischen Vereins Apollo](#)

Jahr/Year: 2004

Band/Volume: [25](#)

Autor(en)/Author(s): Schulze Christian H., Fiedler Konrad

Artikel/Article: [Orthonama obstipata Fabricius, 1794 \(Geometridae: Larentiinae\) - new for the fauna of Borneo 153-154](#)