

- und bemerkenswerte Funde aus neuen Aufsammlungen. — Atalanta, Würzburg, **18**: 121–167.
- KARSHOLT, O., & RAZOWSKI, J. (Hrsg.). (1996): The Lepidoptera of Europe. A distributional checklist. — Stenstrup (Apollo Books), 380 S. + CD.
- KOBES, L. W. R. (2007): Erster Nachweis des Eulenfalters *Chortodes extrema* (HÜBNER, 1809) in Niedersachsen. — Beiträge zur Naturkunde Niedersachsens, Hannover, **60**: 29–31.
- (2008): Nachweis von *Cryphia algae* (FABRICIUS, 1775), einer in Südniedersachsen seltenen Flechtereule. — Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. **28** (3/4): 11–13.
- (2010): *Helicoverpa armigera* (HÜBNER, [1803–1808]), ein Zufallsbefund in Mitteldeutschland oder Speerspitze einer Arealerweiterung? (Lepidoptera, Noctuidae). — Atalanta, Marktleuthen, **41** (1/2): 193–196, 289 [Farbtaf. 1, Abb. 2–5].
- KOCH, M. (1991): Wir bestimmen Schmetterlinge, 3. Auflage (bearb. von W. HEINICKE). — Radebeul (Neumann), 792 S.
- MEINEKE, T. (1984): Untersuchungen zur Struktur, Dynamik und Phänologie der Großschmetterlinge (Insecta, Lepidoptera) im südlichen Niedersachsen. — S. 267–268 in: Biologische Schutzgemeinschaft (Hrsg.), Mitteilungen zur Fauna und Flora Süd-Niedersachsens, Göttingen, Band **6**. — 453 S.
- POOLE, R. W. (1989): Noctuidae. — In: HEPPNER, J. B. (Hrsg.), Lepidopterorum Catalogus (New Series), Fascicle **118** (3 Teile). — Leiden, New York (Brill/Flora & Fauna Publ.), xii + 1314 S.
- SPEYER, Ad., & SPEYER, Aug. (1862): Die geographische Verbreitung der Schmetterlinge Deutschlands und der Schweiz. Zweiter Theil. Die Noctuinen im weiten Sinne. — Leipzig (W. Engelmann), viii + 320 S.
- STAUDINGER, O., & REBEL, H. (1901): Catalog der Lepidopteren des palaearctischen Faunengebietes. I. Theil: Famil. Papilionidae–Hepialidae. — Berlin (R. Friedländer), xxxii + 411 S.

Eingang: 30. VII. 2010

## Entomologische Notiz

### Supplementary information on the *luzonica*-group of the genus *Cricula* on the Philippines — further mislabelled specimens (Lepidoptera, Saturniidae)

Dr. Wolfgang A. NÄSSIG F.R.E.S., Entomologie II, Forschungsinstitut Senckenberg, Senckenbergenanlage 25, D-60325 Frankfurt am Main, Germany; wolfgang.naessig@senckenberg.de

Colin G. TREADAWAY F.R.E.S., Entomologie II, Forschungsinstitut Senckenberg, Senckenbergenanlage 25, D-60325 Frankfurt am Main, Germany; colin.treadaway@web.de

NÄSSIG & TREADAWAY (2011: 191) reported on two specimens of the *luzonica*-group of the genus *Cricula* WALKER, 1855 in coll. S. NAUMANN, Berlin (discovered just shortly before finishing the manuscript), with label data: “Negros, Mt. Canlaon, x. 2007, leg. local collector, via A. SALDAITIS”. These specimens were clearly members of the *luzonica*-group (s. str.), and we expected that they possibly might represent an autochthonous population from Negros island.

However, the results of the mtDNA (COI gene, see NÄSSIG & TREADAWAY 2011; BC nos. 2147 + 2148 SNB) barcode analysis received in April 2011 showed that these two specimens fully fall within the variability range of *Cricula kareli* NÄSSIG & TREADAWAY, 1997 (no significant BC difference to other specimens; the BC sequences of both specimens are identical to that of BC no. 1061 SNB, see NJ trees in figs. 1a + 1b of NÄSSIG & TREADAWAY 2011: 190). Also the ♂ genitalia of these “Negros” specimens (GP nos. 2219/10 + 2220/10 SNB), differing slightly from those of the taxon *Cricula leyteana bayani* NÄSSIG & TREADAWAY, 2011 from Panay island, do not show significant differences to *C. kareli*.

We, therefore, have to state that these two “Negros” specimens, just like the singleton from “Mindoro” listed by NÄSSIG & TREADAWAY (2011: 189, note below type material of *C. halconensis*), are most likely mislabelled trader’s specimens, in both cases of *C. kareli* from Mindanao island, and do not represent autochthonous populations of the *luzonica*-group from other islands.

The tentative records for *C. leyteana bayani* for Negros island on pages 191, 192 (map), 193 (checklist) and 194/195 (legends) of NÄSSIG & TREADAWAY (2011) have to be deleted for that taxon. *C. leyteana bayani* is restricted (according to present knowledge) to Panay island; we do not know any reliable record of a member of the *luzonica*-group from Negros so far.

### Reference

- NÄSSIG, W. A., & TREADAWAY, C. G. (2011): Revisional notes on the *luzonica*-group of the genus *Cricula*, with the description of new taxa from Panay and Mindoro (Philippines; Lepidoptera: Saturniidae). — Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. **31** (4): 187–196

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nachrichten des Entomologischen Vereins Apollo](#)

Jahr/Year: 2011

Band/Volume: [32](#)

Autor(en)/Author(s): Redaktion

Artikel/Article: [Entomologische Notiz 59](#)