An illustrated and annotated checklist of *Arhopala* BOISDUVAL, 1832, taxa occurring in North Maluku and Maluku, Indonesia (Lepidoptera: Lycaenidae) – Part 4: The *fulla* species-group

Andrew Rawlins, Alan Cassidy and Stefan Schröder

Andrew Rawlins, 392 Maidstone Road, Rainham, Kent, ME8 0JA, UK; acrawlins@hotmail.com (corresponding author) Alan Cassidy, 18 Woodhurst Road, Maidenhead, Berkshire, SL6 8TF, UK; accassidy@aol.com Stefan Schröder, Auf dem Rosenhügel 15, D-50997 Köln (Cologne), Germany; ste.schroeder@gmx.net

Abstract: This paper is the 4th in the series covering the species and subspecies of the lycaenid genus *Arhopala* BOISDUVAL, 1832 that occur in the Indonesian provinces of North Maluku and Maluku and deals with the *fulla* species-group, *sensu* EVANS (1957). Nine described taxa (5 species) are recognised as occurring there. Two new synonyms are proposed. We briefly discuss the taxonomy and distribution of the taxa *admete* HEWITSON, 1863, *eucolpis* KIRSCH, 1877 and *sudesta* EVANS, 1957. The taxon *sosias* FRUHSTORFER, 1914 from Obi, placed by EVANS (1957) as a synonym of *A. fulla canulia* HEWITSON, 1869 is restored to a distinct subspecies (stat. rev.). A map shows the islands discussed in the text and all taxa are illustrated in colour.

Keywords: Lepidoptera, Lycaenidae, Theclinae, Arhopala, fulla species-group, new synonyms, Indonesia, North Maluku, Maluku.

Illustriertes und kommentiertes Verzeichnis der Arhopala-Arten (Lepidoptera: Lycaenidae, Theclinae), die in den Nordmolukken und Molukken (Indonesien) vorkommen – Teil 4: Die *fulla*-Artengruppe

Zusammenfassung: Dies ist die vierte Publikation einer Serie über die Arten und Unterarten der Lycaenidengattung Arhopala BOISDUVAL, 1832 aus den indonesischen Provinzen Nordmaluku und Maluku. Sie befaßt sich mit der Artengruppe von Arhopala fulla sensu Evans (1957). 9 beschriebene Taxa (mit 5 anerkannten Arten) sind von dort bekannt. Zwei neue Synonyme werden vorgeschlagen. Taxonomie und Verbreitung der Taxa admete HEWITSON, 1863, eucolpis KIRSCH, 1877 und sudesta Evans, 1957 werden kurz diskutiert. Das Taxon sosias FRUHSTORFER, 1914 von Obi, von Evans (1957) als Synonym of A. fulla canulia HEWITSON, 1869 behandelt, wird als separate Unterart wieder aufgewertet (stat. rev.). Eine Karte zeigt die besprochenen Inseln, und alle Taxa werden farbig illustriert.

Introduction

Arhopala BOISDUVAL, 1832 (Lycaenidae, Theclinae, Arhopalini) is the fifth genus to be published in NEVA in the series on the lycaenid genera of the Indonesian provinces of North Maluku (Maluku Utara) and Maluku. As Arhopala is a large group we have split the genus into sections for publication. Previous parts (RAWLINS et al. 2018a, 2018b, 2018c) have covered an introduction to the genus and the anthelus, theba, democritus, eumolphus and centaurus species-groups.

This is the 4th part and covers the *fulla* species-group, sensu Evans (1957). We recognise nine taxa, comprising five species, as occurring in the Maluku area. Two new synonyms are introduced. The taxon sosias FRUHSTOR-FER, 1914 from Obi, placed by Evans (1957) as a synonym of A. fulla canulia Hewitson, 1869 is restored back to a distinct subspecies. We briefly discuss the taxonomy and distribution of the taxa *admete* HEWITSON, 1863, *eucolpis* KIRSCH, 1877 and *sudesta* EVANS, 1957.

For the biogeography of the region see RAWLINS et al. (2014: 5-8). For the purposes of this paper we make the following key points:

- We use the term Maluku to include both the Indonesian political Provinces of North Maluku (= Maluku Utara) and Maluku.
- The province North Maluku comprises: the Sula islands, the islands we term "northern Maluku" (see below), Obi and Gebe.
- The province Maluku comprises: the islands we term "central Maluku" (see below), the Gorong, Watubela and Tayandu Island groups, the Banda Islands, the Kei Islands, the islands of Southwest Maluku (including Wetar), the Tanimbar Islands and the Aru Islands.
- We use the biogeographical term "northern Maluku" to mean the islands of Morotai, Halmahera, Ternate, Bacan, Kasiruta and Mandioli and some associated smaller islands.
- We use the biogeographical term "central Maluku" to mean the islands of Buru, Ambelau, Manipa, Kelang, Buano, Seram, Ambon, Haruku, Saparua, Nusa Laut, Geser and Seram Laut.

The map shows these islands of Maluku and North Maluku. Here we note that the Indonesian western half of the Island of New Guinea along with its associated offshore islands (previously variously known as Irian, Irian Jaya, West Irian, Irian Barat) now consists of two political provinces: West Papua and Papua. We use the term "New Guinea" in its geographical sense to mean the whole island including these two Indonesian provinces along with the mainland part of the country of Papua New Guinea.

Where available, both surfaces of both sexes of each taxon are illustrated. To reduce the number of colour plates needed, the specimens are illustrated "halved", showing the upperside on the left and the underside on the right. In most cases we have depicted the left half of the butterfly, but where the right side is in significantly better condition we have shown this and flipped the image to allow easier comparison of similar taxa.

We have examined the collections of the Natural History Museum, London (NHMUK), and examined specimens and photographs from some private collections.



Map: Provinces of North Maluku and Maluku showing the island names used in the text.

Abbreviations used

AME	Allyn Museum of Entomology, Gainesville, Florida,
	USA.

- CARR Coll. Andrew RawLINS, Rainham, Kent, UK.
- coll. Collection.
- CSSK Coll. Stefan Schröder, Köln (Cologne), Germany.
- HT Holotype.
- LT Lectotype.
- NHMUK The Natural History Museum, London, UK.
- PT Paratype.
- SMT Senckenberg Museum für Tierkunde, Dresden, Germany.
- stat. rev. Status revivisco (status revised).
- syn. n. Synonymus novus (new synonymy).
- TL Type locality.

Arhopala BOISDUVAL, 1832

Type species: *phryxus* BOISDUVAL, 1832, designated by SCUD-DER (1875: 120). Plate 1, Figs. 1-28: subspecies of Arhopala admete; A. eucolpis, A. sudesta, A. acron. - Figs. 1-16: subspecies of Arhopala admete. Figs. 1-12: A. admete admete: 1: 3, ups./uns., Halmahera (Baru, Ibu, viii. 2002, CARR). 2: ♀, ups./uns., Morotai (Daeo, vi.-xi. 1994, CARR). 3: ♂, type, ups./uns., Seram (Ceram, HEWITSON Coll., NHMUK). 4: Q, ups./uns., Seram (Kanike, North Ceram, 600 m, 1912, E. STRESEMANN, NHMUK). 5: d, ups./uns., Bacan (Makian, xı. 2002, CARR). 6: ♀, ups./uns., Kasiruta (v. 2005, CARR). 7: ♂, ups./uns., Kelang (Tihu, xı. 2003, CARR). 8: Q, ups./uns., Kelang (Tihu, xı. 2003, CARR). 9: 3, ups./uns., Obi (I. 2013, CARR). 10: 9, ups./uns., Obi ("type" admetina = admete, H. FRUHSTORFER, NHMUK). 11: 3, ups./uns., Obi ("type" admetina = admete, H. FRUHSTORFER, NHMUK). 12: Q, ups./ uns., Obi ("type" pallata = admete, H. FRUHSTORFER, NHMUK). - Figs. 13–14: A. admete ?cora: 13: ♂, ups./uns., Waigeo (2016, CSSK). 14: ♀, ups./ uns., Waigeo (2016, CSSK). Figs. 15-16: A. admete cora: 15: 3, ups./uns., New Guinea (Sorong, 2005, CSSK). 16: ♀, ups./uns., New Guinea (Sorong, 2005, CSSK). - Figs. 17-22: Arhopala eucolpis: 17: 3, ups./uns., Waigeo (type waigeoensis, Waig. PLAT[EN], NHMUK). 18: ♀, ups./uns., Waigeo (type waigeoensis, Waig. PLAT[EN], NHMUK). 19: ♂, ups./uns., New Guinea (Sorong, 2005, CSSK). 20: ♀, ups./uns., New Guinea (Sorong, 2005, CSSK). 21: З, ups./uns., Tagula (Mt. Riu, 2000 ft., Sudest Isl., III. 1916, ЕІСННОRN Bros., NHMUK). 22: Q, ups./uns., Tagula (Sudest Isl., II. 1916, EICHHORN Bros., NHMUK). - Figs. 23-24: Arhopala sudesta: 23: 3, HT, ups./uns., Tagula (Sud-Est I., IV. [18]98, Meek, NHMUK). 24: ♀, ups./uns., Tagula (Sud-Est I., IV. [18]98, МЕЕК, NHMUK). Figs. 25-28: Arhopala acron: 25: д, ups./ uns., Bacan (VIII. 2010, CARR). 26: 9, ups./uns., Morotai (Daeo, III. 2004, CARR). 27: d type, ups./uns., Bacan (Batchian, HEWITSON Coll., NHMUK). 28: Q, type, ups./uns., Bacan (Batchian, HEWITSON Coll., NHMUK). - All NHMUK specimen photographs on both plates are © Trustees of the Natural History Museum London, reproduced with permission.



Annotated checklist of the *Arhopala "fulla* speciesgroup" taxa of North Maluku and Maluku

Introduction to the fulla species-group

EVANS (1957: 124–126) included four species in his *fulla* group - acron HEWITSON, 1862, admete HEWITSON, 1863, disparilis FELDER, 1860 and *fulla* HEWITSON, 1862. We note here that EVANS placed this species-group under the genus Narathura MOORE, 1879, which is now considered a synonym of Arhopala, as discussed in Arhopala part 1 (RAWLINS et al. 2018a).

Under the heading "The *fulla* species group of Evans (1957)", PARSONS (1998: 393) stated: "A group of 4 distinctive tailless species were treated by Evans (1957) as the *fulla* (HEWITSON, 1863) species group." PARSONS noted that only *Arhopala disparilis* was absent from mainland New Guinea. However he then only listed one species - fulla - in that section. *A. azenia* and *A. admete* were the last two species included at the end of the previous section on "The *centaurus* species group of Evans (1957)". It seems this was just a layout error.

PARSONS (1998: 393) raised *azenia* HEWITSON, 1863 from a subspecies of *acron* to a full species and we concur.

Both Evans (1957: 125) and Parsons (1998: 393) treated the taxa *eucolpis* KIRSCH, 1877 and *sudesta* Evans, 1957 as subspecies of *A. admete*.

SCHRÖDER (2015: 13) raised *eucolpis* to a full species referring to sympatry with *admete* in Maluku and New Guinea. We have seen no specimens or records of *eucolpis* from Maluku and SCHRÖDER (pers. comm.) now agrees, so we exclude *eucolpis* from the Maluku checklist. SCHRÖDER treated *sudesta* as a subspecies of *eucolpis*.

RAWLINS et al. (2018d) reviewed the status, relationships and distribution of the taxa *eucolpis*, *sudesta* and *admete*. In brief, they demonstrated sympatry of *eucolpis* and *admete* in Sorong, West Papua and Waigeo and of *eucolpis* and *sudesta* in Tagula Island, concluding that all three taxa represented distinct species.

We agree and consider the *fulla* group to comprise seven species – *acron*, *azenia*, *admete*, *eucolpis*, *sudesta*, *disparilis* and *fulla*.

Neither *A. eucolpis* nor *A. sudesta* is recorded from Maluku but we note here their distribution:

Arhopala eucolpis (Figs. 17–22) – from Misool, Waigeo, Yapen, New Guinea, Goodenough, Rossel and Tagula (= Sudest) (Evans 1957; NHMUK).

Arhopala sudesta (Figs. 23-24) – from Tagula (= Sudest) and Rossel (1 \eth) (NHMUK).

Five species of the *fulla* species-group occur in Maluku.

Checklist

Arhopala admete (Hewitson, 1863)

Amblypodia admete: HEWITSON (1863: 7, pl. 3, figs. 18, 19); TL: Seram – see note 1. Range: Maluku – Halmahera, Bacan, Obi, Seram, Ambon (NHMUK), Morotai, Kasiruta, Kelang (TENNENT & RAWLINS 2010), Waigeo, New Guinea (CSSK) – see note 1.

Note 1: EVANS (1957: 125) listed 3 subspecies for *admete* – the nominotypical, *eucolpis* KIRSCH, 1877 and *sudesta* EVANS, 1957. He considered that *admete* was restricted to Maluku and *eucolpis* to New Guinea and offshore islands. PARSONS (1998: 393) followed EVANS in treating *eucolpis* and *sudesta* as subspecies of *Arhopala admete* but noted that *sudesta* might be a distinct species. However, as noted above, RAWLINS et al. (2018d) considered *eucolpis*, *sudesta* and *admete* represented 3 distinct species. They described a new subspecies *A. admete cora* from Sorong in West Papua. They tentatively included the *admete* population on Waigeo with *cora* but noted that this may represent a further undescribed race.

Note 2: Underside hindwing blue tornal spots are present (on occasions vestigial) in *Arhopala admete admete* from Maluku (Figs. 1-12) but absent in *A. admete cora* known from Sorong (Figs. 15-16) and Waigeo (Figs. 13-14) and *A. eucolpis* (Figs. 17-22).

Note 3: Surprisingly we have seen no records of either *admete* or *eucolpis* from Aru, but consider it possible that one or other (more likely *eucolpis*), or both, occur there.

Arhopala admete admete (HEWITSON, 1863)

(Fig. 1: \eth , Halmahera; Fig. 2: \bigcirc , Morotai; Fig. 3: \eth type, Seram; Fig. 4: \bigcirc , Seram; Fig. 5: \eth Bacan; Fig. 6: \bigcirc , Kasiruta; Fig 7: \eth , Kelang; Fig. 8: \bigcirc , Kelang; Fig. 9: \eth , Obi; Fig. 10: \bigcirc type *admetina = admete*, Obi; Fig. 11: \eth type *admetina = admete*, Obi; Fig. 12: \bigcirc , type *pallata = admete*, Obi.)

- Amblypodia admete: HEWITSON (1863: 7, pl. 3, figs. 18, 19); TL: Seram – see note 1.
- = Arhopala admete admetina: FRUHSTORFER syn. n. see note 2.
- = Arhopala admete admetina f. hellava (= pallata?): FRUHS-TORFER - syn. n. - see note 2.

Range: Maluku – Halmahera, Bacan, Obi, Seram, Ambon (NHMUK), Morotai, Kasiruta, Kelang (TENNENT & RAWLINS 2010).

Note 1: HEWITSON described only the \mathcal{S} and illustrated both upperside and underside, noting that the specimen(s) was in the collection of A. R. WALLACE and came from Seram. EVANS (1957: 125) noted that the \mathcal{S} "type" was in the BMNH (Fig. 3).

Note 2: In his publication on FRUHSTORFER types, TALBOT (1923: 84) listed "Arhopala admete admetina" ($\mathcal{J} \otimes \mathcal{Q}$) and "Arhopala admete admetina f. hellava" (\mathcal{Q}), both from Obi. We have found these 3 specimens in the NHMUK main collection amongst the series of Obi admete specimens. The \mathcal{J} (Fig. 11) bears four labels:

- 1. A handwritten "admete admetina FRUHST.".
- 2. A red ringed circular type label.
- 3. "Fruhstorfer Coll. NHMUK 1933-131"
- 4. "Obi, H. Frustorfer".

One Q bears labels 2-4 as above. This *admetina* pair are indistinguishable from the other Obi *admete* specimens. The 2nd Q (Fig. 12) is very small with very pale underside markings and bears labels 2-4 as above and a further handwritten label stating: "*pallata* FRUHST.". This surely corresponds to the f. *hellava* recorded by TALBOT. We consider this specimen to be a minor aberration of *admete*, rather than a stable form. We have been unable to find any published description of these FRUHSTORFER "types" or any further reference to *admetina*. Including these 3 "types", the NHMUK contains 11 $\partial \partial$ and 6 QQ of *admete* from Obi. EVANS also noted 11 $\partial \partial$ and 6 QQ so he must have seen these "type" specimens. He clearly included them with *admete*, but without creating synonymy, so it seems likely that he considered they were unpublished names. Thus, it seems *admetina* and *pallata/hellava* are probably *nomina nuda*, but we cannot be certain and therefore we

synonymise *admetina* FRUHSTORFER and *pallata/hellava* FRUHSTORFER with *admete* HEWITSON, 1863.

Note 3: As Evans (1957: 125), we include Obi specimens with *admete*, but note they are generally smaller than *admete* from other Maluku localities. The uppersides of both sexes exhibit the narrower black borders typical of specimens from central Maluku – see note 4.

Note 4: Northern Maluku *admete* generally differ from those from central Maluku. We have examined 20 $\eth \eth \ref{s}$ & 16 $\circlearrowright \ref{s}$ from northern Maluku (Morotai, Halmahera, Bacan) and 5 $\eth \ref{s}$ & 2 $\circlearrowright \ref{s}$ from central Maluku. In both sexes the upperside dark borders are usually broader in northern Maluku specimens (except 1 of the 6 $\circlearrowright \ref{s}$ and 2 of the 7 $\eth \ref{s}$ from Morotai). In addition northern Maluku $\circlearrowright \ref{s}$ exhibit a darker and more matt blue (except 1 from Halmahera and 1 from Morotai – not the same specimen that has narrower borders). Although in general the northern Maluku and central Maluku specimens can be easily phenotypically separated, the few exceptions incline us not to describe the northern Maluku population as a new race.

Arhopala acron (Hewitson, 1862)

(Fig. 25: ♂, Bacan; Fig. 26: ♀, Morotai; Fig. 27: ♂ type, Bacan; Fig. 28: ♀ type, Bacan.)

Amblypodia acron: HEWITSON (1862: 9, pl. 5, figs. 53, 54); TL: Bacan – see note 1.

Range: endemic to northern Maluku – Halmahera, Bacan (NHMUK), Morotai (TENNENT & RAWLINS 2010).

Note 1: HEWITSON (1862) described both sexes and illustrated the \eth underside and the \wp upperside, noting that the specimens were in the NHMUK. EVANS (1957: 125) stated that the \eth "type" was in the NHMUK (Fig. 27).

Note 2: Evans (1957: 125) placed *azenia* HEWITSON, 1863 as a subspecies of *acron* but PARSONS (1998: 393) recognised that the 2 taxa were distinct species without specifying how they differed. TENNENT & RAWLINS (2010: 15) described the differences in detail and separated the Obi population as a new subspecies of *azenia* – see below.

Note 3: The extent of white markings in the underside hindwing postdiscal area varies between *acron* specimens, with occasional specimens lacking it completely – see Figs. 25–28.

Arhopala azenia (Hewitson, 1863)

Amblypodia azenia: HEWITSON (1863: pt. 1: 7, illustrations pt. 2: pl. 3, figs. 22, 23); TL: Waigeo.

Range: Maluku, New Guinea and various off shore islands.

Note: There are 2 subspecies – both occur in Maluku; one is endemic to Obi.

Arhopala azenia azenia (Hewitson, 1863)

(Fig. 29: ♂, Aru; Fig. 30: ♀, Aru; Fig. 31: ♂ HT, Waigeo; Fig. 32: ♀ PT, Waigeo.)

Amblypodia azenia: HEWITSON (1863: pt. 1: 7, illustrations pt. 2: pl. 3, figs. 22, 23); TL: Waigeo – see note 1.

Range: Aru, Waigeo, Misool, Yapen, New Guinea (NHM UK), Normanby Island in the D'Entrecasteaux group and Sariba Island in the Louisiades (PARSONS 1998). ?Seram – see note 2.

Note 1: HEWITSON described both sexes and illustrated both surfaces of the male of *azenia*, noting that the specimens were in the collections of A. R. WALLACE and W. C. HEWITSON and came from Waigeo. The HT σ and AT (i.e. PT) Q are in the NHMUK (Figs. 31 & 32).

Note 2: Evans (1957: 125) recorded 1 \circ and 1 \circ from Seram in the NHMUK. We have examined these specimens and confirm they are typical *azenia*. The \circ bears 3 labels:

A. Handwritten "Ceram"

B. "EX. MUSAEO D^{ris} BOISDUVAL"

C. "Ex Oberthür Coll. Brit. Mus. 1927-3"

The Q beside it has only labels B and C and no location label. We know of no further specimens from Seram or anywhere in central Maluku so consider this Seram record as uncorroborated.

Arhopala azenia patsyae Tennent & RAWLINS, 2010

(Fig. 33: ♂ PT, Obi; Fig. 34: ♀ PT, Obi; Fig. 35: ♂ HT Obi; Fig. 36: ♀ PT, Obi.)

Arhopala azenia patsyae: TENNENT & RAWLINS (2010: 15, figs. 38-41); TL: Obi – see note.

Range: endemic to Obi.

Note: TENNENT & RAWLINS (2010) described and illustrated both sexes of *patsyae*. The HT ♂ and a series of PTs of both sexes are in the NHMUK (Figs. 33–36).

Arhopala disparilis (C. Felder, 1860)

(Fig. 37: ♂, Morotai; Fig. 38: ♀, Ambon; Fig. 39: ♂ type, Ambon; Fig. 40: ♀, Haruku.)

- Amblypodia disparilis: C. FELDER (1860: 453); TL: Ambon see note 1.
- = Amblypodia courvoisieri: RIBBE (1901: 335, pl. 6, fig. 2); TL: Seram – see note 3.

Range: Endemic to Maluku – Ambon (NHMUK), Seram, Haruku, Morotai (CARR; TENNENT & RAWLINS 2010.

Note 1: C. FELDER briefly described both sexes of *disparilis* in Latin. He noted that the specimens from Ambon were in the FELDER collection. EVANS (1957: 125) noted that the \eth "type" was in the NHMUK (Fig. 39).

Note 2: C. & R. FELDER (1865: 230, pl. 28, figs. 4 & 5) then also described and illustrated both surfaces of the Q.

Note 3: In his paper on new Lepidoptera from Seram, RIBBE (1901) described only the *courvoisieri* \eth , illustrating both surfaces. Evans (1957: 125) listed *courvoisieri* as a synonym of *disparilis* and we agree. TAKANAMI (1989: 51) designated a \eth LT of *courvoisieri* in the SMT.

Arhopala fulla (Hewitson, 1862)

Amblypodia fulla: Неwitson (1862: 10, pl. 6, figs. 67, 68); TL: Buru.

Range: Andaman Islands, Burma, Thailand, Peninsular Malaysia, Borneo, Indonesia including Maluku, Philippines, New Guinea and some offshore islands.

Note: EVANS (1957: 125-126) listed 7 subspecies. He considered *sosias* FRUHSTORFER from Obi as a synonym of *canulia* HEWITSON. We consider *sosias* is a distinct subspecies – see below. Four subspecies are found in Maluku.

Arhopala fulla fulla (Hewitson, 1862)

(Fig. 41: ♂ type *prasiae* = *fulla*, Ambon; Fig. 42: ♀ type *prasiae* = *fulla*, Ambon; Fig. 43: ♂ HT, Buru; Fig. 44: ♀, Buru.)

Amblypodia fulla: Hewitson (1862: 10, pl. 6, figs. 67, 68); TL: Buru – see note 1.

= Arhopala fulla prasiae [emendation for original spelling prasiä]: FRUHSTORFER (1914: 157); TL: Ambon – see note 2.

Range: endemic to central Maluku – Buru, Ambon (NHMUK) – see note 3.

Note 1: HEWITSON (1862) described only the \eth of *fulla* and illustrated both surfaces and noted "In the collection of Dr. BOISDUVAL from Boirou". The HT \eth is now in the NHMUK (Fig. 43).

Note 2: FRUHSTORFER (1914) described both \mathcal{J} and \mathcal{Q} prasia [sic] from Ambon and noted the specimens were in his collection. He

differentiated it from *fulla* and noted that the upperside was a richer blue-violet than in *fulla* from Buru. There are \eth and \heartsuit types in the NHMUK (Figs. 41 & 42). EVANS (1957: 126) listed "*prasia*" [sɪc] as a synonym of *fulla*. We have compared the \eth types (Figs. 41 & 43) and concur.

Note 3: This taxon is very likely also to occur on Seram.

Arhopala fulla babsi JOICEY & TALBOT, 1917

(Fig. 45: ♂, Aru; Fig. 46: ♀, Misool; Fig. 47: ♂ HT, Waigeo; Fig. 48: ♀ PT, Waigeo.)

Arhopala fulla babsi: JOICEY & TALBOT (1917: 219); TL: Waigeo – see note 1.

Range: Waigeo, Misool, New Guinea, Tagula (labelled Sudest Island) (NHMUK), Aru – Wokam Island (CARR) Aru – Wamar Island (AME) – see note 2.

Note 1: JOICEY & TALBOT (1917) first described the male of *babsi* then stated: "Q. This sex of *fulla* does not appear to be known." But he then appeared to describe the female and noted: "*Hab.* Waigeu, 6 $\partial \partial$, 2 QQ". The HT ∂ and a PT Q (labelled as AT) are in the NHMUK (Figs. 47 & 48).

Note 2: PARSONS (1998: 393) recorded specimens of *fulla* from Aru in the Allyn Museum (AME), without specifying the subspecies. We have examined photographs of a \eth in the AME, courtesy of Andy WARREN. The specimen is typical *babsi* and was taken by R. G. & C. M. WIND in Dobo (Wamar Island) 30. IV. 1939. We confirm the presence of *babsi* in Aru (1 \eth , Wokam Island, III. 2003, CARR – Fig. 45).

Arhopala fulla canulia (Hewitson, 1869)

(Fig. 49: ♂, Halmahera; Fig. 50: ♀, Halmahera; Fig. 51: ♂ type, "Phillipine [*sic*] Isles" *recte* Halmahera; Fig. 52: ♀, Halmahera.)

Amblypodia canulia: HEWITSON (1869: 14f, pl. 3c, fig. 54); TL: Halmahera? – see notes 1 & 3.

Range: northern Maluku – Halmahera, Ternate (NHMUK), Bacan (Bethune-Baker 1903, Fruhstorfer 1914 – see note 3), Morotai (Tennent & Rawlins 2010).

Note 1: HEWITSON (1869) described the 3 and illustrated only the underside. He noted: "In the collection of W. C. HEWITSON, from the Philippines." EVANS (1957: 126) stated "Philippines', recte Halmaheira" without giving any explanation and noted that the ♂ "type" was in the NHMUK. This specimen (Fig. 51) bears only a type label, a HEWITSON coll. label and a handwritten label stating "Phillipine [sic] Isles". On the same page, Evans described a new subspecies - santa - from Luzon in the Philippines, noting 3 $\partial \partial$ and 1 Q in the NHMUK. This clearly distinct subspecies has a brown underside rather than the grey of canulia. The canulia type clearly matches specimens in the NHMUK from Halmahera and Ternate but Evans gave no reason why he chose Halmahera as the TL. We suspect he chose Halmahera on the balance of probability, as he noted that the NHMUK held 18 canulia specimens from Halmahera, 2 from Ternate and 3 from Obi (treated by us as sosias). See also note 3 re TL.

Note 2: EVANS (1957: 126) listed *sosias* (TL Obi) as a synonym of *canulia*, but as discussed below, we consider the Obi population – *sosias* – represents a distinct subspecies.

Note 3: In his revision of the "Amblypodia group", BETHUNE-BAKER (1903: 149, pl. 3, fig. 26) listed Arhopala canulia and gave the habitat as "Philippines; Batchian", but he did not record where the specimens were held. He illustrated a \eth typical of northern Maluku canulia without stating its collection locality. This is the first record we have seen recording the taxon from Bacan. In describing sosias, FRUHSTORFER (1914: 157) compared it to "A. canulia Hew. von Batjan" – indicating either that he considered HEWITSON's canulia type was from Bacan, or that he was familiar with canulia from Bacan. Arhopala fulla canulia is recorded from the other northern Maluku islands of Morotai, Ternate and Halmahera, so is very likely to be present on Bacan as well.

Note 4: TREADAWAY (1995: 77) records only the subspecies A. *fulla santa* EVANS, 1957, in his checklist of the butterflies of the Philippines.

Arhopala fulla sosias FRUHSTORFER, 1914 stat. rev.

(Fig. 53: 3 type, Obi; Fig. 54: 3, Obi; Fig. 55: 3, Obi.)

Arhopala canulia sosias: FRUHSTORFER (1914: 157); TL: Obi – see note 1.

Range: endemic to Obi.

Note 1: FRUHSTORFER (1914) described sosias from 3 \overrightarrow{O} from Obi in his collection. EVANS (1957: 126) listed sosias as a synonym of *canulia* and noted that the \overrightarrow{O} "type" was in the NHMUK (Fig. 53). We have compared the *canulia and sosias* \overrightarrow{O} types in the NHMUK (Figs. 51 & 53) and small series of *A. fulla* from Obi and Halmahera and we note that the undersides of Obi and Halmahera specimens clearly and consistently differ. Obi specimen undersides are a much lighter grey colour and the pale postdiscal and submarginal streaky bands are whiter. Both subspecies have a submarginal black spot, edged inwardly with blue, in spaces 1a and 2 on the hindwing underside and in some specimens a vestigial spot in space 3. In Obi specimens this space 3 spot is usually more developed. Therefore, we consider that *sosias* is not a synonym of *canulia* and re-establish the Obi population as a distinct subspecies *Arhopala fulla sosias*.

Note 2: We have not seen any QQ of this subspecies.

Acknowledgements

Thanks to Hiromi DETANI for information and advice; also to Chris MÜLLER for the map. Thanks also to NHMUK staff, particularly Dr. Blanca HUERTAS for access to the collections and the NHMUK for permission to take photographs of specimens. All NHMUK specimen photographs are © Trustees of the Natural History Museum London, reproduced with permission.

Plate 2, Figs. 29-55: subspecies of Arhopala azenia; A. disparilis, subspecies of A. fulla. - Figs. 29-36: subspecies of Arhopala azenia. Figs. 29–32: A. azenia azenia: 29: ♂, ups./uns., Aru (IV. 2007, CARR). 30: ♀, ups./uns., Aru (Wokam, x. 2006, CARR). 31: 3, HT, ups./uns., Waigeo (Waigiou, HEWITSON coll., NHMUK). 32: ♀, PT, ups./uns., Waigeo (Waigiou, HEWITSON coll., NHMUK). - Figs. 33-36: A. azenia patsyae: 33: J, PT, ups./uns., Obi (vii. 1992, CARR). 34: Q, PT, ups./uns., Obi, (H. FRUHSTORFER, NHMUK). 35: 3, HT, ups./uns., Obi, (H. FRUHSTORFER, NHMUK). 36: ♀, PT, ups./uns., Obi, (H. FRUHSTORFER, NHMUK). - Figs. 37-40: Arhopala disparilis: 37: 3, ups./uns., Morotai (Daeo, III. 1994, CARR). 38: ♀, ups./uns., Ambon (Laha, 31. viii. 1993, CARR). 39: ♂, type, ups./uns., Ambon (Amboina, DOLESCHALL type, FELDER Colln., NHMUK). 40: Q, ups./uns., Haruku (x. 2006, CARR). - Figs. 41-55: subspecies of Arhopala fulla. Figs. 41-44: A. fulla fulla: 41: 3, ups./uns., Ambon (type prasiä, Amboina, H. FRUHSTORFER, NHMUK). 42: Q, ups./ uns., Ambon (type prasiä, Amboina, H. FRUHSTORFER, NHMUK). 43: 3, HT, ups./uns., Buru (Bourou, Ex Musaeo Dris BOISDUVAL, NHMUK). 44: Q, ups./uns., Buru (Bouru, JOICEY Bequest, NHMUK). - Figs. 45-48: A. fulla babsi: 45: ♂, ups./uns., Aru (Wokam, III. 2003, CARR). 46: ♀, ups./ uns., Misool (Misol, 29. 1. [18]99, Н. КÜHN, NHMUK). 47: д, HT, ups./ uns., Waigeo (Waigeu, II.–III. 1915, A. C. & F. PRATT, NHMUK). 48: Q, PT, ups./uns., Waigeo (Waigeu, II.-III. 1915, A. C. & F. PRATT, NHMUK). -Figs. 49–52: A. fulla canulia: 49: 3, ups./uns., Halmahera (Halmaheira, WATERSTRADT, NHMUK). 50: Q, ups./uns., Halmahera (Gunung Rotang, Oba, I. 2016, CARR). 51: 3, HT, ups./uns., Halmahera ("Phillipine Isles" recte Halmahera, Неwitson Coll., NHMUK). **52:** ^о, ups./uns., Halmahera (Gunung Rotang, Oba, I. 2016, CARR). - Figs. 53-55: A. fulla sosias: 53: ੋ, type, ups./uns., Obi (H. FRUHSTORFER, NHMUK). 54: ਨੋ, ups./uns., Obi (H. FRUHSTORFER, NHMUK). 55: 3, HT, ups./uns., Obi (H. FRUHSTORFER, NHMUK).



References

- BOISDUVAL, J. B. A. D. (1832): Faune entomologique de l'Océan Pacifique avec l'illustration des insects nouveaux recueillis pendant le voyage. Première Partie: Lépidoptères. – In: DUMONT D'URVILLE, J., Voyage de découvertes de l'Astrolabe, Exécuté par ordre du Roi, Pendant les années 1826-1827-1828-1829, sous le Commandement de M. J. DUMONT D'UR-VILLE. – Paris (J. Tastu), 267 pp., Atlas: 5 pls. [biodiversitylibrary. org/ page/ 2071230 = text, biodiversitylibrary.org/ page/ 51854032 = plates.]
- BETHUNE-BAKER, G. T. (1903): A revision of the Amblypodia group of butterflies of the family Lycaenidae. – Transactions of the Zoological Society of London, London, 17 (1): 1–164, pls. 1–5. [biodiversitylibrary.org/page/33729745 = text, biodiversitylibrary. org/page/33729855 = plates.]
- EVANS, W. H. (1957): A revision of the Arhopala group of Oriental Lycaenidae (Lepidoptera: Rhopalocera). – Bulletin of the British Museum (Natural History), Entomology, London, 5 (3): 85-141. [biodiversitylibrary.org/page/2361223.]
- FELDER, C. (1860): Lepidopterorum Amboinensium species novae diagnosibus collustratae a Dr. C. FELDER. (1) Rhopalocera.
 Pp. 448-462 *in*: FRAUENFELD, G., Diagnosen einiger neuer Insecten und Untersuchung mehrerer Sandproben verschiedener Küstenpunkte, gesammelt während der Reise Sr. Maj. Fregatte Novara. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften zu Wien, 40 (11): 447-468. [biodiversitylibrary.org/page/6442184.]
- —, & FELDER, R. (1865-1875): Reise der Österreichischen Fregatte Novara um die Erde in der Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von WÜLLERSTORF-URBAIR. Zoologischer Theil. 2. Band: Zweite Abtheilung: Lepidoptera. – Kaiserlich-Königliche Hof- und Staatsdruckerei, in Commission bei Carl Gerold's Sohn, Vienna, vi + 549 pp., Atlas: 140 pls. [biodiversitylibrary.org/page/2520784 = text, biodiversitylibrary.org/page/1221095 = plates.]
- FRUHSTORFER, H. (1914): Neue Arhopala-Rassen. Deutsche Entomologische Zeitschrift Iris, Dresden, 27 (4): 155–166. [biodiversitylibrary.org/page/12722708].
- HEWITSON, W. C. (1862): Specimen of a catalogue of Lycaenidae in the British Museum. — London (British Museum), 15 pp, 8 pls. [biodiversitylibrary.org/page/23824029 = text, biodiversitylibrary. org/page/23823995 = plates.]
- (1863-1878): Illustrations of diurnal Lepidoptera. Lycaenidae. – London (John van Voorst), x + 228 + 47 (Supplement + Index) pp., 107 pls. [biodiversitylibrary. org/ page/ 30120889 = text. biodiversitylibrary.org/page/30336394 = plates.]
- KIRSCH, T. (1877): Beitrag zur Kenntnis der Lepidopteren-Fauna von Neu Guinea. – Mittheilungen aus dem K. Zoologischen Museum zu Dresden, Dresden, Heft 2: 101-134, pls. 5-7. [biodiversitylibrary.org/page/34978667.]
- JOICEY, J. J., & TALBOT, G. (1917): New Lepidoptera from Waigeo, Dutch New Guinea, and Biak. – The Annals and Magazine of Natural History, London, (8) **20** (117): 216–229. [biodiversitylibrary.org/page/18593550.]
- PARSONS, M. (1998): The butterflies of Papua New Guinea: their systematics and biology. San Diego, London, etc. (Academic Press), xvi + 928 pp.
- RAWLINS, A., CASSIDY, A., MÜLLER, C., SCHRÖDER, S., & TENNENT, W. J. (2014): An illustrated and annotated checklist of *Jamides* HÜBNER, 1819, taxa occurring in the Indonesian provinces of North Maluku and Maluku (Lepidoptera: Lycaenidae). – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. **35** (1/2): 5–39.

- —, —, & SCHRÖDER, S. (2018a): An illustrated and annotated checklist of *Arhopala* BOISDUVAL, 1832 taxa occurring in North Maluku and Maluku, Indonesia (Lepidoptera: Lycaenidae) – Part 1: Introduction, *anthelus* and *theba* speciesgroups. – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. 38 (4): 191–201.
- , —, & (2018b): An illustrated and annotated checklist of Arhopala BOISDUVAL, 1832, taxa occurring in North Maluku and Maluku, Indonesia (Lepidoptera: Lycaenidae)
 Part 2: The democritus and eumolphus species-groups. – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. 39 (1): 21–37.
- —, —, & (2018c): An illustrated and annotated checklist of Arhopala BOISDUVAL, 1832 taxa occurring in North Maluku and Maluku, Indonesia (Lepidoptera: Lycaenidae) – Part 3: The centaurus species-group. – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. 39 (2): 85–102.
- —, SCHRÖDER, S., & CASSIDY, A. (2018d): A new subspecies of Arhopala admete (HEWITSON, 1863) from Sorong, West Papua, Indonesia and a brief review of the taxonomy of the Arhopala fulla species-group (sensu Evans, 1957) (Lepidoptera: Lycaenidae). – Butterflies, Tokyo, 77: 39–42.
- RIBBE, C. (1901): Neue Lepidopteren von Ceram, Niederländisch-Ostindien. – Deutsche Entomologische Zeitschrift Iris, Dresden, 13: 334-337, pl. 6. [biodiversitylibrary. org/ page/ 12770904 = text, biodiversitylibrary.org/page/12770953 = plate.]
- SCUDDER, S. H. (1875): Historical sketch of the generic names proposed for butterflies: A contribution to systematic nomenclature. – Proceedings of the American Academy of Arts and Sciences, Boston, **10**: 91–293. [biodiversitylibrary. org/ page/10425091.]
- SCHRÖDER, S. (2015): The Arhopala centaurus-group of Evans (1957) in Papua, Indonesia (Lepidoptera: Lycaenidae). – SUGAPA (Suara Serangga Papua), Jayapura, 9 (1): 1-22. [www. researchgate. net/ publication/ 274718793_ The_ Arhopala_ centaurus-group_ of_ Evans_ 1957_ in_ Papua_ Indonesia_ Lepidoptera_Lycaenidae.]
- TAKANAMI, Y. (1989): On some type specimens of Lycaenidae from South East Asia. – Tyô to Ga, Osaka, **40** (1): 23-80. [www. jstage.jst.go.jp/article/lepid/40/1/40_KJ00006597983/_pdf.]
- TALBOT, G. (1923): The FRUHSTORFER collection of butterflies. Catalogue of types with general account and list of the more interesting forms. – Nice (J. Gastaud), 8 + 135 + ix pp. (N.B. Sometimes cited as "MARTIN, 1922" – see TENNENT 2008.) [biodiversitylibrary.org/page/10082505.]
- TENNENT, W. J. (2008): Citation of the contents of "The FRUHSTOR-FER collection of butterflies: Catalogue of types with general account and list of the more interesting forms" published in the 1920s. – Entomologist's Gazette, Iver, Bucks, **59**: 41–53.
- —, & RAWLINS, A. (2010): Notes on Arhopala BOISDUVAL, 1832 from Sulawesi and Maluku, including new subspecies of A. argentea STAUDINGER, 1888, A. chamaeleona BETHUNE-BA-KER, 1903, and A. azenia (HEWITSON, [1863]) (Lepidoptera, Lycaenidae). – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N.F. 31 (1/2): 9-16.
- TREADAWAY, C. G. (1995): Checklist of the butterflies of the Philippine Islands (Lepidoptera: Rhopalocera). – Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, Suppl. 14: 7-118, 14 pls.

Received: 30. viii. 2018

¹²⁸

[©] Entomologischer Verein Apollo e. V., Frankfurt am Main, April 2019

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Nachrichten des Entomologischen Vereins Apollo

Jahr/Year: 2018

Band/Volume: 39

Autor(en)/Author(s): Rawlins Andrew, Cassidy Alan, Schröder Stefan

Artikel/Article: An illustrated and annotated checklist of Arhopala Boisduval, 1832, taxa occurring in North Maluku and Maluku, Indonesia (Lepidoptera: Lycaenidae) — Part 4: The fulla species-group 121-128