

## Diagnosen neuer Meladomus.

Von

Dr. W. Kobelt.

### *Meladomus (Lanistes) fultoni* n.

T. umbilicata, globosa, subtiliter striatula, nitida, viridi-fuscescens, basin versus lutescens, fasciis angustis in anfr. ultimo 4 ornata; spira breviter conica, gradata, apice valde erosa; sutura distincta, linearis. Anfr. persist. 3—4 supra breviter contabulati, dein distincte angulato-carinati, rapide crescentes, ult. permagnus, initio medio quoque angulatus, angulo et carina supera aperturam versus evanescentibus, circa umbilicum profundum acute carinatus, carina usque ad peristoma persistente. Apert. mediocriter obliqua, irregulariter ovata, supra leviter lunata, intus fuscescens fasciis vix translucentibus, albido labiata; peristoma acutum, basi subeffusum, marginibus haud junctis, columellari supra tantum leviter dilatato et reflexo. Diam 45, alt. obl. 47, alt. apert. obl. 30, diam 15 mm.

Hab. Victoria Nyanza (cfr. M. Ch. II t. 27 fig. 1, 2).

### *Meladomus (Lanistes) senegalensis* n.

T. sinistrorsa, late et pervie umbilicata, depresso subglobosa, solida, vix nitens, oblique et sat ruditer striata, sculptura spirali vix conspicua, viridi-fusca, fasciis saturatioribus latiusculis obsoletis ornata, et in anfractu ultimo usque ad aperturam erosa; spira gradata, depresso conica, undique erosa. Anfr. superst. 4, supremi convexi, penultimus infra suturam contabulatus, angulatus, ultimus multo major, modice dilatatus, supra planatus, obsolete tantum angulatus, dein convexus, circa umbilicum iterum carinatus. Apert. late ovato-rotundata, modice obliqua: columella stricta, faucibus fuscis late albido-coeruleo limbatis, fasciis externis translucentibus; peristoma tenuer, acutum, mar-

ginibus callo tenui vix junctis, externo primum horizontali, dein regulariter areuato, intus distinete labiato, columellari supra albo, parum dilatato et reflexo. — Diam. maj. 45, alt. 35, dimens. apert. 27:20 mm.

Meladomus (Lanistes) senegalensis (? carinatus var.) Kobelt, in Mart. Chemnitz II p. 31 t. 28 fig. 13, 14.

Hab. Senegal.

*Meladomus libycus chaperi* n.

Difffert a typo t. minore, plus minusve spiraliter lirata, globosa vel ovato-globosa, spira gradata, acute angulata, saepe profunde erosa, anfr. ultimo circa umbilicum latum pervium acute carinato. — Diam. maj. 25—27, alt. 26—27 mm.

Meladomus libycus chaperi Kobelt in: Mart. Chemnitz II p. 30 t. 27 f. 3—6.

Hab. Dahomey Africæ occidentalis.

*Meladomus libycus assiniensis* n.

T. anguste et obtecte<sup>9</sup> umbilicata, elongato-ovata, obsolete striata, sculptura spirali fere nulla, unicolor nigro fusca, solida, crassa; spira distinete gradata; sutura linearis. Anfr. superst. 4 supra plani, dein distinctissime angulato-carinati, infra carinam vix convexiusculi; ultimus major, elongatus, basi carina mediocri umbilicum cingente munitus. Apertura angusta et subirregulariter ovata, supra acute angulata, infra compressa, intus nigrofusca; peristoma acutum, margine columellari vix dilatato sed intus albo-caloso, in umbilicum impresso eumque fere obtegente. — Alt. 29, diam. maj. 24, dim. apert. 20:10 mm.

Meladomus libycus assiniensis Kobelt, in: Mart. Chemnitz II p. 30 t. 26 f. 6, 7.

Hab. Assinie Africæ occidentalis.

*Meladomus boettgeri* n.

T. ovato-globosa, aperte umbilicata, laeviuscula, vix striatula, sculptura spirali nulla, nitidula, viridi-grisea, fas-

ciis latis fuscis pulcherrime ornata, epidermide cutacea viridi lutescente induta; spira depressa, subgradata, valde erosa. Anfr. superst.  $4\frac{1}{2}$  convexi, supra planati sed vix angulati, ult. multo major, rotundatus, circa umbilicum vix levissime compressus. Apert. oblonga, supra et infra angustata, haud dilatata, faucibus fuscescentibus pulcherrime fasciatis; peristoma tenue, acutum, haud incrassatum, marginibus callo tenuissimo junctis. — Operculum normale, annulis prominentibus 3—4 insigne. — Diam. 35, alt. 35, alt. apert. obl. 25, lat. 16 mm.

Meladomus boettgeri Kobelt, in: Mart. Chemn. ed. II t. 30 f. 6, 7.

Hab. Gabun.

---

### Diagnosen neuer Cyclotiden.

Von

Dr. W. Kobelt.

---

#### 1. *Cyclotus (Opisthoporus) bialatus* (Moellendorff mss.).

T. parva, late umbilicata, depressa, solidula, subtiliter striatula, epidermide tenui, lutescente, adhaerente induta, strigis fulminulatis supra castaneis, infra peripheriam pallidioribus dein evanescentibus ornata. Spira breviter conica, prominula, summo unicolor castaneo, apice acutissimo, sutura impressa. Anfr.  $4\frac{1}{2}$  regulariter accrescentes, inferi ad suturam planati, dein convexi, ultimus circa umbilicum compressus, antice valde descendens, 3 mm pone aperturam tubulo erecto retrorsum spectante insignis. Apertura peroliqua, circularis, plano irregulari; peristoma fusco-albidum, subduplex: internum continuum, haud porrectum, externum tenuissimum, extus late dilatatum et reflexum, supra et ad basin alato-productum, ad columellam subite evanescens. — Operculum normale.

# ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft](#)

Jahr/Year: 1912

Band/Volume: [44](#)

Autor(en)/Author(s): Kobelt Wilhelm

Artikel/Article: [Diagnosen und Meladomus 6-8](#)