

D/58

Bibliothek
Museum WIT

NEUE ENTOMOLOGISCHE NACHRICHTEN

aus dem Entomologischen Museum
Dr. Ulf Eitschberger

Beiträge zur Ökologie, Faunistik
und Systematik von Lepidopteren

37. Band

ISSN 0722-3773

März 1996

VADIM V. DUBATOLOV

Three contributions to the
knowledge of palearctic Arctiinae

(Lepidoptera, Arctiidae)

Verlag: Dr. Ulf Eitschberger, Humboldtstr. 13a, D-95168 Marktleuthen

Einzelpreis: DM 98,-

NEUE ENTOMOLOGISCHE NACHRICHTEN

aus dem Entomologischen Museum
Dr. Ulf Eitschberger

Beiträge zur Ökologie, Faunistik
und Systematik von Lepidopteren

Herausgeber und Schriftleitung:

Dr. ULF EITSCHBERGER,

Humboldtstr. 13a, D-95168 Marktleuthen

Die Zeitschrift und alle in ihr enthaltenen einzelnen Beiträge und Abbildungen sind urheberrechtlich geschützt. Jede Verwertung außerhalb der engen Grenzen des Urheberrechts gesetzes ist ohne Zustimmung des Verlages unzulässig und strafbar. Das gilt insbesondere für Vervielfältigungen auf foto mechanischem Wege (Fotokopie, Mikrokopie), Übersetzungen, Mikroverfilmungen und die Einspeicherung und Verarbeitung in elektronischen Systemen.

ISSN 0722-3773

D.E.

NEUE ENTOMOLOGISCHE NACHRICHTEN

aus dem Entomologischen Museum
Dr. Ulf Eitschberger

Beiträge zur Ökologie, Faunistik
und Systematik von Lepidopteren

37. Band

ISSN 0722-3773

März 1996

VADIM V. DUBATOLOV

Three contributions to the
knowledge of palearctic Arctiinae

(Lepidoptera, Arctiidae)

Verlag: Dr. Ulf Eitschberger, Humboldtstr. 13a, D-95168 Marktleuthen

Three contributions to the knowledge of palearctic Arctiinae

(Lepidoptera: Arctiidae)

by

VADIM V. DUBATOLOV

1. <i>Sinowatsonia</i> , a new genus of Arctiidae from Tibet (China) (on the systematics of the genus <i>Micractia</i> SEITZ s.l., Part 2) .	5
2. A review of the genus <i>Palearctia</i> FERGUSON (on the systematics of the genus <i>Micractia</i> SEITZ s.l., Part 3) .	9
3. A list of the Arctiinae of the territory of the former U.S.S.R.	39

**1. *Sinowatsonia*, a new genus of Arctiidae from Tibet (China)
(on the systematics of the genus *Micractia* SEITZ s. l., Part 2)**
(Lepidoptera: Arctiidae)

by

V. V. DUBATOLOV

The genus *Micractia* SEITZ, 1910 s.l. was divided by FERGUSON (1984) and DUBATOLOV (1987) into several genera: *Micractia* SEITZ, 1910 (*M. trigona* (LEECH, 1899)), *Sinoarctia* DUBATOLOV, 1987 (*S. kasnakovi* DUBATOLOV, 1987, *S. sieversi* (GRUM-GRSHIMAILO, 1891), *S. forsteri* (DANIEL, 1943) and (a personal communication of the late W. THOMAS) *S. mussoti* (OBERTHÜR, 1911)), *Ocnogynodes* DUBATOLOV, 1987¹ (*O. y-albula* (OBERTHÜR, 1886)), *Sibirarctia* DUBATOLOV, 1987 (*S. kindermannii* (STAUDINGER, 1867), *S. bureatica* (O. BANG-HAAS, 1927)), and *Palearctia* FERGUSON, 1984 (a series of Central-Asian species). However, there are two species described by DANIEL (1943) from East Tibet, which do not belong to any of these genera: *Micractia hoenei* DANIEL, 1943, and *Micractia batangi* DANIEL, 1943. They turned out to belong to a separate genus within the tribe Spilosomini. This new genus is described in this paper.

The author expresses his sincere gratitude to Dr. A. WATSON at The Natural History Museum, London, for offering the male and female of *M. hoenei* (DANIEL)² and the photographs of *M. batangi* (DANIEL) for this study. I wish to express my thanks to Dr. O. E. KOSTERIN (Novosibirsk) for his help in translating the manuscript. This work was partly supported by the International Science Foundation and the Russian Academy of Natural Sciences.

***Sinowatsonia* gen. nov.**

Type-species: *Micractia batangi* DANIEL, 1943.

The wings are well developed in both sexes. The forewing is rather narrow, its apex slightly prominent (colour plate fig. 1a, b). The pattern consists of fused spots forming an Y-like pattern, with one branch oriented along the cubitus to the wing apex and the other from the cubitus to the tip of vein 1 (A). Vein 11 (R₁) starts from the apex of the cell, vein 10 (R₂) from the 7+8 (R₄₊₅) vein. The hindwing colouration varies from white to carmine-red.

In the males the head is densely covered with long erect hairs, in females they are substantially shorter. The palpae are moderately long, covered with thin hairs, in the males their underside is covered with long erect hairs. The eyes are naked, in the males relatively large, hemispheric, in the females oval, slightly convex. The antennae are bipectinate in males and serrate in females. The proboscis is reduced. The body is rather narrow, covered with sleeky hairs in males and sleeky scales in females. The fore tibia is simple, a little longer than half of the femora, the epiphysis is shorter than the tibia. The middle tibia is armed with a single pair, the hind tibia with two pairs of almost wedge-shaped spurs, which are slightly shorter than the tibia diameter.

Male genitalia (figs 1, 2). The anterior part of the tegumen is wide, forming a "collar", the valva is narrow, with a wide rounded tooth on the internal edge.

The female genitalia have not been studied.

1 The genus *Ocnogynodes* DUBATOLOV, 1987 was united with *Lithosarctia* DANIEL, 1954 by DE FREINA & WITT (1994). Now this genus includes three species: *L. hoenei* DANIEL, 1954, *L. y-albula* (OBERTHÜR, 1886) and *L. thomasi* DE FREINA & WITT, 1994.

2 It should be noted that the female of *M. hoenei* (DANIEL) in the collection of the The Natural History Museum (London) was determined as "*Micractia sieversi* GR.-GR."




Fig. 1: Male genitalia of *Sinowatsonia batangi* (DANIEL, 1943). Holotype. a – general view, b – the aedeagus with the vesica everted, c – VIII. sternite. Photo by courtesy of Dr. A. WATSON.

Notes. Concerning the structure of the male genitalia, the two species of the new genus, *S. hoenei* (DANIEL) and *S. batangi* (DANIEL), are very similar to the species of the genus *Ocnogyna* LEDERER, 1853, especially to *O. parasita* (HÜBNER, [1790]). However, they have two pairs of spurs on the hind tibia, the fore tibia is simple, and the females have normally developed wings (while all *Ocnogyna* species have a single pair of spurs on the hind tibia, a short fore tibia much widening to the tip, where it almost lacks scales and bears a large tooth; the females are wingless or brachypterous). These characters are shared with the genus *Maurica* DE FREINA & WITT, 1984. However, our species differ from the representatives of the latter genus by more prolonged forewings with the apices drawn off, by the presence of a tooth on the valva, and the presence of a “collar” on the tegumen (the *Maurica* species have rather short and broad wings, stick-shaped valvae, and the tegumen without a “collar”). Whereas the species of the genus *Lithosactria* DE FREINA & WITT (= *Ocnogynodes* DUBATOLOV) resemble *S. hoenei* (DANIEL) and *S. batangi* (DANIEL) by wing colouration, they lack the “collar” on the tegumen (the presence of which is a characteristic of several genera of the tribe Spilosomini) and have the valva of quite a different shape. The wing colouration of the two species of *Sinowatsonia* gen. nov. resembles to some extent that of the representatives of the genera *Eospilarctia* KÔDA, 1988, and *Paraspilarctia* KÔDA, 1988, which also have the “collar” on the tegumen, but greatly differ from them by the shape of the valva, aedeagus, its vesica and by some other characters.




Fig. 2: Male genitalia of *Sinowatsonia hoenei* (DANIEL, 1943). Paratype. "Li-kiang ca. 2000 m Prov. Nord-Yuennan 1.6.1934 H. HÖNE" a – general view, b – the aedeagus with the vesica everted.

Key to the species of *Sinowatsonia*

- 1 (2) The hindwing is evenly pinkish-red in males and carmine-red in females. The distal part of the valva is three times narrower than the narrowest point of its proximal part.
S. hoenei (DANIEL, 1943) **comb. nov.** (China: Sichuan, Yunan, Tibet (FANG, 1982))
- 2 (1) The hindwing is white with pink bases in males, females are unknown. The distal part of the valva is two times narrower than the narrowest point of its proximal part.
S. batangi (DANIEL, 1943) **comb. nov.** (China: Sichuan (FANG, 1982))

References

- DANIEL, F. (1943): Beiträge zur Kenntnis der Arctiidae Ostasiens unter besonderer Berücksichtigung der Ausbeuten H. Hönes aus diesem Gebiet (Lep., Het.). II. Teil. Hypsinae, Micrarctiinae, Spilosominae, Arctiinae. – Mitt. Münch. Ent. Ges. **33** (2/3): 673–759, Taf. 14–22.
- DUBATOLOV, V. V. (1987): K sistematike roda *Micrarctia* SEITZ, s.l. (Lepidoptera, Arctiidae) [On the systematics of the genus *Micrarctia* SEITZ, s.l. (Lepidoptera, Arctiidae)]. In: Nasekomye, klestshi i gelminki [Insects, mites and helminths]. – Novosibirsk: "Nauka", Sib. Dept., p. 30–47 (in russian).
- FANG CHENG-LAI (1982): Arctiidae. – Iconographia heterocerorum sinicorum **2**: 190–277 (in chinese).
- FERGUSON, D. C. (1984): Two new generic names for groups of holarctic and palaearctic Arctiini (Lepidoptera, Arctiidae). – Proc. entomol. Soc. Wash. **86** (2): 452–459.
- FREINA, J. J. DE & T. J. WITT (1984): Taxonomische Veränderungen bei den Bombyces und Sphinges Europas und Nordwestafrikas. Zwei neue Arctiidaegattungen für den westpaläarktischen Faunenbereich: *Watsonarctia* gen. n. und *Maurica* gen. n. (Lepidoptera, Arctiidae). – Entomofauna **5** (28): 323–334.

- FREINA, J. J. DE & T. J. WITT (1994): Zur Kenntnis der Gattung *Lithosarcia* DANIEL, 1954 mit Beschreibung zweier neuer Taxa. – *Atalanta* **25** (3/4): 535–542, Farbtafel XVIIIa.
- KÔDA, N. (1987/1988): A Generic Classification of the Subfamily Arctiinae of the Palaearctic and Oriental Regions based on the Male and Female Genitalia (Lepidoptera, Arctiidae). Part I. – *Tyô to ga* **38** (1987): 153–237; Part II. *Tyô to Ga* **39** (1988): 1–79.

Explanation of the colour plate (p. 89):

1a	1b	1b

Fig. 1a: *Sinowatsonia batangi* (DANIEL, 1943). Holotype ♀. "Batang (Tibet) Im Tal des Yangtze (ca. 2800 m) 26.7.1936 H. HÖNE". Photo by courtesy of Dr. A. WATSON.

Fig 1b: *Sinowatsonia hoenei* (DANIEL, 1943). The female from the collection of The Natural History Museum (London) and its labels. Photo by courtesy of Dr. A. WATSON.

**2. A review of the genus *Palearctia* FERGUSON
(on the systematics of the genus *Micractia* SEITZ s.l., Part 3)**
(Lepidoptera: Arctiidae)

by

V. V. DUBATOLOV

The genus *Palearctia* FERGUSON, 1984 was described by D. C. FERGUSON while dividing the genus *Micractia* SEITZ, 1910, s.l. At first it included seven species: *P. glaphyra* (EVERSMANN, 1843), *P. buraetica* (O. BANG-HAAS, 1927), *P. kindermannii* (STAUDINGER, 1867), *P. rupicola* (GRUM-GRSHIMAILO, 1890), *P. postflavida* (HAMPSON, 1894), *P. erschoffii* (ALPHERAKY, 1882), and *P. ladakensis* (O. BANG-HAAS, 1927). However, the genus in this sense still remained heterogeneous. Later (DUBATOLOV, 1987) two species, *P. kindermannii* (STGR.) and *P. buraetica* (B.-H.), were separated into a new genus *Sibiractia* DUBATOLOV, 1987, while four more species from the former genus *Micractia* SEITZ were added to the genus *Palearctia* FERG., namely *Micractia wagneri* (PÜNGELE, 1918), *M. gratiosa* (GRUM-GRSHIMAILO, 1890), *M. variabilis* DANIEL, 1966, and *M. mongolica* (ALPHERAKY, 1888). In the same paper the genus *Palearctia* FERG. was erroneously united with *Tancrea* PÜNGELE, 1898. Indeed, these two genera are closely related, as can be inferred from the male genitalia structure and the wing pattern. However, *T. pardalina* PÜNGELE, 1898 has the hind tibiae with only one pair of spurs, whereas all the species of *Palearctia* FERG. have two pairs of spurs on the hind tibiae; this was noted later in the description of *P. mira* DUBATOLOV & TSHISTJAKOV, 1989. Besides, these genera also differ by the presence or absence of the tympanal apparatus. The representatives of *Palearctia* FERG. lack a tympanal apparatus, while *T. pardalina* PÜNGELE shows an apparatus consisting of two large chitinized bubbles on either side of the abdomen base. Thus, I include into the genus *Palearctia* FERG. the majority of species formerly considered within *Micractia* SEITZ, s.l., except for: *Micractia* (s. str.) *trigona* (LEECH, 1899), the species of the genera *Sinoactia* DUBATOLOV, 1987, *Ocnogynodes* DUBATOLOV, 1987¹, *Sibiractia* DUBATOLOV, 1987 and *Sinowatsonia* DUBATOLOV, 1996. The systematic position of *Oroncus ladakensis* O. BANG-HAAS, 1927 deserves special discussion. This species was referred to the genus *Micractia* SEITZ s.l. by DRAUDT (1931) and, later, to *Palearctia* FERG. by FERGUSON (1984). Thanks to Prof. H. J. HANNEMANN I have received the type specimen, a female, for investigation, and found out that this species should be placed into the genus *Arctia* SCHRANK, 1802—*Arctia ladakensis* (O. BANG-HAAS, 1927) **comb. nov.**—because it is closely related to *A. intercalaris* (EVERSMANN, 1843). These two species can be easily distinguished by the female antennae, which are simple in *A. ladakensis* (B.-H.) and serrate in *A. intercalaris* (Ev.).

In the course of this revision of the genus *Palearctia* I have studied the collections of the Zoological Museum of the Institute of Animal Systematics and Ecology (former: Biological Institute) of the Siberian Branch of the Russian Academy of Sciences in Novosibirsk, Russia, the Zoological Institute of the Russian Academy of Sciences in St.-Petersburg, Russia, the Zoological Museums of Moscow State University, Russia, and Kiev State University, Ukraine, the Schmalhausen Institute of Zoology of the Ukrainian Academy of Sciences in Kiev, Ukraine, and the Institute of Biology and Pedology of the Far East Branch of the Russian Academy of Sciences in Vladivostok. I am very thankful to Drs. A. L. LVOVSKY, A. V. SVIRIDOV, V. RAEVSKII and Yu. A. TSHISTJAKOV for their help during my work with these collections. Some type specimens were kindly sent to me by Prof. H. J. HANNEMANN of the Zoological Museum of the Humboldt University, Berlin, Germany, and the colour slides of some taxa by Dr. A. WATSON of The Natural History Museum (London), England. Besides, I am greatly indebted for receiving materials to P. V. BOGDANOV, V. A. GANSON, L. V. KAABAK, V. G. MAKHAT, V. S. MURZIN,

1 The genus *Ocnogynodes* DUBATOLOV, 1987 was united with *Lithosactia* DANIEL, 1954 by DE FREINA & WITT (1994). Now this genus includes three species: *L. hoenei* DANIEL, 1954, *L. y-albula* (OBERTHÜR, 1886) and *L. thomasi* DE FREINA & WITT, 1994.

A. V. NEKRASOV, S. K. SAZONOV, G. A. SHAPIRO, E. A. TARASOV (Moscow, Russia), A. I. IVANOV, L. V. PETRIKEVICH, V. N. PRASOLOV (St.-Petersburg, Russia), V. N. OLSCHWANG (Ekaterinburg, Russia), Yu. E. PERUNOV (Barnaul, Russia), I. G. PLYUSTSH (Kiev, Ukraine), K. ČERNÝ (Innsbruck, Austria). I also wish to express my gratitude to Dr. O. E. KOSTERIN (Novosibirsk) for correcting the original manuscript.

This work was partly supported by the International Science Foundation and the Russian Academy of Natural Sciences.

Genus *Palearctia* FERGUSON, 1984

Proc. Entomol. Soc. Wash. **86**(2): 454–456.

Type-species: *Arctia glaphyra mannii* ALPHERAKY, 1881.

The genus was carefully described by FERGUSON, so there is no reason to make a new description.

Key to Subgenera

- 1(2) The body is covered with light, whitish or yellowish hairs and scales. The tegumen is very long, dorsally with a knob-like projection. The uncus is strongly broadened proximally. The moths inhabit semideserts and deserts of Central (Inner) Asia. *Centractia* DUBATOLOV

2(1) The body is covered with dark hairs and scales. The tegumen is short, without a projection. The uncus is not broadened proximally. The moths inhabit high mountains of Middle, Central (Inner) and South Asia. *Palearctia* FERGUSON

Palearctia FERGUSON, 1984

Subgenus *Centrarctia* DUBATOLOV, 1990

Chlenistonogie i gel'minty. Novosibirsk: 157.

Type-species: *Arctia mongolica* ALPHERAKY, 1888.

***Palearctia mongolica* (ALPHERAKY, 1888)**

Stett. Ent. Ztg. 49: 67 (*Arctia mongolica*).

= *Arctia serarum* GRUM-GRSHIMAILO, 1899, Ann. Mus. Zool. Imp. Acad. Sci. St.-Petersb. 4: 463–464,
SYN. DOV.

= *Arctia serum* GRUM-GRSHIMAILO, 1902, Ann. Mus. Zool. Imp. Acad. Sci. St.-Petersb. 7:204. *syn. nov.*

Type locality: "Dschungor; 16. August 1884. (POTANIN)" (ALPHERAKY, 1888). The type specimen has labels in Russian: "16.VIII.1884 / Kil'ya / Dzhongor", "Mongolica / ♂. ALPH. / "Original", "kol. Vel. Kn. / NIKOLAYA / MIKHAILOVICH" [coll. Grand Duke NIKOLAI MIKHAILOVICH]. The species was described on the materials of G. N. POTANIN's expedition in 1883–1884. Judging from POTANIN's book (1950), in these days (11.–17. August 1884) he was in the headquarters of DZHUNGAR (Dzhongor), the prince of the Ordos nomads, nearly 180 km SE of Baotou (Nei Mongol, China), 55 "versts" (i. e. nearly 59 km) from the Huang He River.

Distribution. Tuva: the southern part of the republic within the Ubsu Nur intermontane hollow. Mongolia: Uvs, Chovd, Gov'-Altaj (Gobi Altai), Ömnögov' (South Gobi), and Dornogov' (East Gobi) aimaks; Töv (Central) aimak (DANIEL, 1968). Dornod (East, or objalsan) aimak (DANIEL, 1967), Bajanchongor and Öbörchhangai aimaks (DANIEL, 1965). China: Nei Mongol (Ordos and Yin Shan).

Description. In addition to the characters mentioned in the key, it should be noted that the forewings are white with a light yellowish-rose tint. The hindwings are rose-red with bright contrasting dark-brown spots (figs 7a–c). The forewing pattern varies considerably: the subbasal costal spot (M_1 band²) may be reduced to two strokes, the discoidal spot ($D+M_2$) to 2–3 separate spots, the submarginal spot may also be reduced (figs 7b, c). The hindwings always have the discoidal spot fused with the costal one (M_2), a row of submarginal spots (E_3) and, sometimes, the costal spot of the band M_1 . Genitalia. See fig. 1.




Fig. 1: Genitalia of *P. mongolica* ALPH. a — male (Lake Ubsu-Nur, Tuva), b — female (S. Mongolia).

Biology. The moths fly from the end of June until the middle of August. The caterpillars have been found on *Artemisia sieversiana* by A. L. Lvovsky in the South Gobi aimak of Mongolia; in captivity they feed on various dicotyledonous plants (DUBATOLOV, 1985).

Notes. A moth with a split dark pattern of the forewing was described as *Arctia serarum* GRUM-GRSHIMAILO, 1899. Later, this author (GRUM-GRSHIMAILO, 1902) wrote that this name resulted from a misprint, the correct name would be *Arctia serum* Gr.-Gr. (from Latin "Sera"—China). The type locality was reported as "... montibus Mongoliae meridionalis, 1871, ..., N. M. PRZEWALSKI ..." I have studied the type specimen, a male (fig. 7b) which is supplied with the following labels (in Russian): "17930", "Dolina Khuan-khe / i In'-shan' / PRZHEVALSK. 71" [the Huang He valley / and the Yin Shan / PRZHEVALSKY]. In genitalia structure it appears to be identical with the type specimen of *Arctia mongolica* ALPH. The type-localities of both taxa are situated not far from each other and from the Huang He river in Nei Mongol (China). The light form (f. *serarum* Gr.-Gr.) occurs both in Mongolia and North China (Nei Mongol). It was described by DANIEL (1965) in detail. For a long time *P. mongolica* (ALPH.) was confused with *Sibirarctia kindermanni* (STGR.), for example, by FANG CHENG-LAI (1982). The

2 I use the terminology of SOKOLOV (1936).

figure of *P. mongolica* (ALPH.) (FANG CHENG-LAI, 1982: t. 68, fig. 1564) is given with the erroneous identification "Micractia kindermanni (STAUDINGER)". In the description of the species' range, FANG CHENG-LAI (1982:211) listed Heilongjiang, Liaonin, Nei Mongol, Hebei, Ningxia-Huizu, Xinjiang, Mongolia, and the USSR, evidently confusing the data of the two species, since *P. mongolica* (ALPH.) does not seem to occur in Heilongjiang and Liaonin. A note by KURENTZOV (1965) that "Micractia mongolica ALPH." occurs in Transbaikalia is also misleading. The examination of that particular specimen (♂, [the Chita region], Kruchina, forest-steppe, 20.VIII.1955, A. I. KURENTZOV), now kept in the Institute of Biology and Pedology in Vladivostok, has shown that it is in fact *Sibirarctia kindermanni* (STGR.).

Material. Tuva: 1 ♂, Ovyurskii distr., Sachan, 6.VII.1971 (KOROTYAEV); 2 ♂♂, Khol'-Oozhu, 7.VII.1968 (KOSTYUK), 22.VII.1971 (KOROLEVSKAYA); 1 ♂, Lake Ubsa (Uvs or Ubsu) [northern side], 6.VII.1914 (TOMASHINSKII). Mongolia: Uvs aimak: 2 ♂♂, 1 ♀, 50 km E of Ulaangom, 5., 6.VIII.1970 (KERZHNER, CHOGSOMDZHAV, EMELYANOV); 2 ♂♂, 40 km S of Naryn-Bulak, 6.VII.1969 (KOSTYUK); Chovd aimak: 1 ♂, 17.VII.1969 (KOSTYUK); Gov'-Altaj (Gobi Altai) aimak: 3 ♂♂, Ushiin-Bulak spring, 30 km NW of Beger, 13.VII.1970 (EMELYANOV); Ömnögov' (South Gobi) aimak: 1 ♂, Daurgan-mod terrain, 30 km SE of Mt. Khan-Bogdo, 24.VI.1971 (EMELYANOV); 1 ♀, Bulgan, 24.VI.1972 (MEDVEDEV); 1 ♀, ex larva, 30 km S of Shine-Dzhinst, mid VII.1981, 1 pupa 16.–18.VIII., 1 imago 1.IX.1981 (LVOVSKY); Dornogov' (East Gobi) aimak: 1 ♂, Aragut, 27.VI.1971 (KERZHNER); 1 ♂, 30 km SSE of Lake Shokhoi-Nur, Namt-Ula, 4.VIII.1971 (KERZHNER); 1 ♂, 20 km W of Altai, 14.VII.1980 (KERZHNER). China, Nei Mongol: 1 ♂ (the type specimen of *Arctia mongolica* ALPH.), Kil'ya Dzhongor [the headquarters of prince DZHUNGAR], 16.VIII.1884 (POTANIN); 1 ♂ (the type specimen of *Arctia serarum* GR.-GR.), N 17930, Huang He valley and Yin Shan (PRZHEVALSKY).

Subgenus *Palearctia* FERGUSON, 1984

Proc. Entomol. Soc. Wash. **86** (2): 454–456.

Type species: *Arctia glaphyra mannii* ALPHERAKY, 1881.

Key to the species

- 1(2) On the forewing upperside the dark pattern consists of longitudinal strokes divided by light veins. Vein 2 (Cu_2) is light. The species inhabits the Altai.
P. mira DUBATOLOV & TSHISTJAKOV
- 2(1) On the forewing upperside the pattern consists of angular spots or transversal bands. If it consists of broad longitudinal strokes, then vein 2 (Cu_2) is dark.
- 3(8) The forewing upperside has a V-shaped stroke distally of the cell, near the beginning of veins 2 and 3 (Cu_1 and Cu_2). If this stroke is divided into two branches, then there is a short longitudinal stroke distally of the beginning of vein 2 (Cu_2), which is parallel to the veins 1 and 2 (A and Cu_2).
- 4(5) On the forewing upperside the dark pattern is of even intensity. The hindwing is not white. The valvae are rhombus-shaped. The aedeagus bears spurs distally.
P. glaphyra (EVERSMANN)
- 5(4) On the forewing upperside the medial and basal spots are darker than the marginal ones. The valvae are not rhombus-shaped, otherwise the hindwing is white.
- 6(7) The valvae are rhombus-shaped. The hindwing is white. The aedeagus bears no spurs distally. The species inhabits the Kirghiz Ala-Tau (Alexander Mts.). *P. golbecki* spec. nov.

- 7(6) Valvae with a costal projection. The hindwing may be of various colours. The aedeagus bears spurs distally. *P. gratiosa* (GR.-GR.)
- 8(3) The forewing upperside has neither V-shaped nor a longitudinal stroke distally of the cell (near the beginning of the veins 2 and 3 (Cu_1 and Cu_2)).
- 9(10) The basal third of the hindwing upperside, up to the tornal angle, is contrasting dark. The species inhabits the Himalayas. *P. marxi* (B.-H.)
- 10(9) The basal part of the hindwing upperside is light or has dark veins, otherwise the dark basal spot is not contrasting and does not reach the tornal angle.
- 11(12) On the forewing upperside the vein 2+3 (Cu_{1+2}) is very dark at the point of dichotomy. The median part of the forewing has two dark transversal bands. The species inhabits Central Tien Shan, the Sarydzhaz region. *P. wagneri* (PÜNGELER)
- 12(11) On the forewing upperside the vein 2+3 (Cu_{1+2}) is not dark or has only a few dark scales at the point of dichotomy.
- 13(14) The hindwing underside has a dark transversal stroke near the costa base. *P. gracilis* spec. nov.
- 14(13) The hindwing underside has no dark transversal stroke near the costa base.
- 15(16) On the forewing the dark medial transversal bands usually are discontinuous between veins 1 and 2 (A and Cu_2). The hind angle of the fore spot of the first median band (M_1) is not sharp. The aedeagus is without spurs distally. *P. erschoffii* (ALPH.)
- 18(17) On the forewing the dark medial bands are not interrupted, otherwise the hind angle of the fore spot of the first median band (M_1) is sharp. The aedeagus bears spurs distally. *P. ferghana* (STGR.)

***Palearctia mira* DUBATOLOV & TSHISTJAKOV, 1989**

Zool. Zhurnal 68(11):141–143.

Type locality: Altai, Kurai Range, the upper course of the Yarly-Yary river near the town Aktash.

Distribution. Known only from the Kurai Range in the Altai.

Biology. The moths were collected on alpine meadows from the 10th of July until the beginning of August, at 2700 m above sea level.

Description (fig.2). In the males the wing pattern is similar to that of *Grammia quenseli* (PAYKULL, 1793) (DUBATOLOV & TSHISTJAKOV, 1989: 142, fig. 1), but the strokes in the cell and in front of vein 1 (A) on the forewing are darker. In general, the pattern is not contrasting, but is rather diffuse. In the females the pattern is reduced, consisting of strokes on the basal half of the cell, in front of vein 1 (A), two stroke-like spots between veins 2 and 3 (Cu_1 and Cu_2), one spot between veins 5 and 6 (M_1 and M_2), and two apical spots. The hindwing is dark-grey in males and yellowish-grey in females, with weakly expressed diffuse spots.

Genitalia. See DUBATOLOV & TSHISTJAKOV, 1989: 143, f. 2a, b.



Fig. 2: *P. mira*, holotype ♂, Altai

Palearctia glaphyra (EVERSMANN, 1843)

Bull. Soc. Imp. Nat. Mosc. **16** (3): 544, t. 10, f. 4 (*Euprepia glaphyra*).

Type locality: “ promontoriis Altaicus australibus” [in front of the mountains of the South Altai]. Two specimens were identified by Prof. E. EVERSMANN in his collection as *Euprepia glaphyra* Ev. (BREMER, 1870). The second one, a male from Kyakhta (Transbaikalia), which is not the type, in fact turned out to be *Sibirarctia buraetica* (B.-H.). The first one, the type specimen, has two labels: “Noor Saisan / gegend” and “coll. EVERSMANN”

Many authors doubted that the materials of EVERSMANN, including the types of *P. glaphyra* (Ev.), *Arctia intercalaris* (Ev.), *Parnassius actius* Ev. and *P. delphinus* (Ev.), and some others, originated from the Altai, because all these species do not inhabit these mountains. That was admitted long ago by LEDERER (1853) and BANG-HAAS (1927). The former author (LEDERER, 1853: 353), according to a report given by A. KINDERMANN, mentioned that these materials were collected by “ Herrn SCHRENCK, einem Botaniker aus Petersburg in der Tarbagataischen und Allakanischen [possibly, Alakol'? – V. D.] Gebirgen ” In my opinion, the environs of Lake Zaisan should be assumed to be the type locality, as it is indicated on the type's label. Most likely, this specimen was collected in the mountains south of this lake. However the data available indicates the Dzhungarian Alatau as the north-eastern most point of the range of *P. glaphyra* (Ev.).

Distribution. Kazakhstan: mountains near Lake Zaisan (?); Dzhungarian Alatau, including the Tyshkantau³ and the Trans-Ili Alatau. Kyrgyzstan: Kirghiz Ala-Tau (Alexander Mts.), Kungei Ala-Tau, Terskei Ala-Tau; Central and Inner Tien Shan west to Naryn and the Dolon pass. China, Xinjiang: East Tien Shan, Kuruktag, Hotan (?). The nominative subspecies occurs in the mountains of East Kazakhstan from the Dzhungarian Alatau to Lake Zaisan (?) and, probably, in neighbouring regions of Xinjiang (China).

Biology. The moths fly from the middle of June until 20th of August at 2200–3000 m above sea level (according to my material), or at 6,000–10,000 feet (ALPHERAKY, 1891). Caterpillars, according to observations by S. K. SAZONOV (pers. comm.), are polyphagous on various dicotyledonous plants.

Description. The main characters of the species are indicated in the key. The groundcolour of the forewing may vary from white to yellow, the dark pattern is variable, the spots may be fused along the veins to form dark strokes, or across the veins to form two median bands, often being narrowly split between veins 1 and 2 (A and Cu₂). Sometimes the vein tips have light scales on the dark marginal band (E₁₊₂). The hindwing is often bright red but may vary from yellowish-orange to black. The usual arrangement of the hindwing spots is as follows: a dark discoidal spot always present, three submarginal spots (E₃), a narrow dark margin (E₁) from the costa to vein 3 (Cu₁), sometimes to the tornal angle. The pattern of the underside is poorly developed, reduced or, on the hindwing, sometimes absent.

In females the wing pattern is more reduced than in males, the marginal band (E₁₊₂) is strongly reduced and sometimes absent, as well as the spots along the costa, the hindwing as in males. Several aberrations have been described: ab. *obscurata* BÖTTCHER (= *tristis* SEITZ, = *fumosa* NIEPELT), characterized by a dark forewing with spots fused along the margin; ab. *amabilis* BÖTTCHER, characterized by the forewing with a large white space between bands M₁ and M₂; ab. *rosearia* SEITZ, characterized by a rose forewing; ab. *flava* O. BANG-HAAS, characterized by a yellow forewing; ab. *illustrata* O. BANG-HAAS, characterized by a white hindwing; ab. *nigroradiata* O. BANG-HAAS, characterized by the hindwing veins suffused by black scales; ab. *puengeleri* BÖTTCHER, characterized by a black hindwing; ab. *gratiosata* BÖTTCHER, characterized by the absence of the discoidal spot on the hindwing.

The nominative subspecies is characterized by its small size (forewing length 13 mm) and by more marked orange hindwings (fig. 8a).

3 The author saw a male specimen of *P. glaphyra* (Ev.) at the Tyshkantau Mts., 46 km N of Dzharkent, the headwaters of Chiliksa-1 brook (the 2nd left tributary to the Sargin river, the left tributary to the Bolshoi [Great] Usek river), at 3000 m, on a subalpine meadow with juniper bushes, 31.VII.1994.

Genitalia. See figs 3–4; the nominative subspecies: 3a, 4a.

While studying the taxa *manni* STGR. and *naryna* B.-H., FERGUSON (1984) found out that their vesicae of the aedeagus are different. For this reason he separated *P. naryna* (B.-H.) as a good species. As it appeared from my materials collected from different regions of the Tien Shan, there are some differences in the structure of the vesica between Transilian, Central Tian Shanian and East Tian Shanian specimens. Based on a strong similarity of the wing pattern and other features of genitalia, and taking into account the allopatry of distribution, I consider the moths inhabiting these mountain systems to represent a series of subspecies.

Material. Kazakhstan: 1 ♂ (the holotype of *Euprepia glaphyra* Ev.), Noor Saisan [Lake Zaisan] gegend (coll. EVERSMANN); Dzhungarian Alatau: 3 ♂♂, 40 km ENE of Tekeli, the upper reaches of the Kora river, NW slope of the Semenov-Tian-Shanskij Pass, 2950–2970 m, an alpine meadow with small juniper bushes, 6.VIII.1994 (DUBATOLOV, KOSTERIN).

***Palearctia glaphyra dublitzkyi* (O. BANG-HAAS, 1927)**

Horae Macrolepid. 1: 63, t. 8, f. 29, 30 (*Micrarctia glaphyra dublitzkyi*).

Type locality: "Berg Khoum Bel beim Ort Alma Ata" [Kumbel' Mt. near Alma-Ata].

Distribution. Kazakhstan: Trans-Ili Alatau; Kungei? Ala-Tau in Kazakhstan and Kyrgyzstan.

Description (fig. 8b). On the forewing the groundcolour is usually golden-yellow, the spots are smaller than in the nominotypical subspecies. The dark spots on the hindwing underside are well developed. The forewing length is 15–17 mm.

Material. Kazakhstan: Trans-Ili Alatau: 1 ♂ (TSVETAEV); 5 ♂♂, a ridge top, 3 km SW of Medeo, 24.VII.1980 (DUBATOLOV); 1 ♂, near the Tuyuksu glacier, 2300 m, 1.VII.1975 (coll. PLYUSTSH); 1 ♂, Srednii [Middle] Talgar river, 26.VII.1978 (coll. PERUNOV); 1 ♂, same locality, 2600 m, 10.VII.1985 (KOSTYUK); Issyk [town]: 2 ♂♂, 15.–27.VII.1936, 2 ♂♂ of ab. *puengeleri* B.-H., 24.VII.1937, 3 ♂♂, 10.VII.1940 (coll. SHELJUZHKO), 1 ♂, 25.VIII.1954 (coll. TSVETAEV); 9 ♂♂, 2 ♀♀, same locality, Kaskasu terrain (IONOV); 1 ♂, Lake Mus-Kol', 3000 m, 23.VII.1936 (coll. TSVETAEV); 6 ♂♂, Kumbel' terrain, 24.VII.–12.VIII.1924, 25.VII.1925 (KRASNITSKII).

***Palearctia glaphyra aksuensis* (O. BANG-HAAS, 1927)**

Horae Macrolepid. 1: 63, t. 8, f. 26, 27 (*Micrarctia glaphyra aksuensis*).

= *Micrarctia glaphyra naryna* O. BANG-HAAS, 1927, Horae Macrolepid. 1: 63, t. 8, f. 28, **syn. nov.**

Type locality: "Gebirge nördlich von Aksu" [the mountains north of Aksu] (China, Xinjiang).


Distribution. Kyrgyzstan and China, Xinjiang: Inner and Central Tien Shan north to the Terskei Alatau; the southern part of the East Tien Shan; Kuruktag, ?Hotan.

Description (figs 8c, 8d, 10f). The dark pattern on the forewing is well developed, the spots being sometimes fused. The dark spots on the hindwing are usually not large, often reduced on the underside. The forewing length is 13–16 mm.


Note. I consider two subspecies described by BANG-HAAS (1927), *naryna* B.-H. and *aksuensis* B.-H., as conspecific, because:

- 1) in some parts of the species' range, for example at the Dolon pass or in the environments of Naryn (see: BÖTTCHER, 1905: f. 4–14), specimens occur either with the pattern typical for *aksuensis* B.-H. or with fused spots (*naryna* sensu FERGUSON, 1984, which differs by the wing pattern from *naryna* B.-H. and is identical to ab. *tristis* SEITZ or *obscurata* BÖTTCHER);
- 2) the vesica of the aedeagus of all studied specimens from Inner and Central Tien Shan and Kuruktag are almost identical and correspond to fig. 13a in FERGUSON (1984:455), which shows *P. naryna* sensu FERGUSON. It should be noted that the published figure of the *P. naryna* imago (FERGUSON, 1984: 453, f. 6) represents not a typical specimen, but an extreme variant of ab. *obscurata* BÖTTCHER;


3a




3b



3c



3d






Fig. 3: Genitalia of *Palearctia* species. a — *P. g. glaphyra* Ev., holotype; b, e — *P. g. manni* STGR. (East Tien Shan); c, d — *P. gratiosa flava* sub-spec. nov.; f — *P. wagneri* PÜGL. (Kaingdy-Katta mountain range, Central Tien Shan); g — *P. golbecki* spec. nov., holotype. b, d — females, others: males.

3) since *aksuensis* B.-H. and *naryna* B.-H. were described in the same work on the same page, I assume as a senior name *aksuensis* B.-H., as it was described before *naryna* B.-H.

Material. Kyrgyzstan: Terskei Ala-Tau: 1 ♂, VII.1974 (BAKUROV); 5 ♂♂, Pokrovka, 27., 30.VII.1955, 13.VII., 2.VIII.1967 (TSVETAEV); 1 ♂, Borskaun river, 9.VII.1978 (PETRIKEVICH); 1 ♂, 10 km SW of Przhevalsk, 13.VII.1980 (DUBATOLOV); 2 ♂♂, 20 km of the Karakol gorge, 28.VII.1985; 2 ♂♂, Ken-suу (PLYUSTSH); 8 ♂♂, Naryn, 11., 15.VII.1906 (AKULIN), 10.VI.1910 (KSENZOPOLSKY), 9.VII.1904, 20., 27.VI.1915 (NEZHIVOV), 26.VII.1967 (TSVETAEV); 6 ♂♂, Dolon pass, 31.VII.1967, 2.–5.VIII.1969 (TSVETAEV), 9.VII.1979 (STSHERBINA), 20.VII.1989; 28 ♂♂, 5 ♀♀, Kaingdy-Katta mountain range, Tashkoro, 19.–28.VII.1982, 2.–9.VII.1989, 6., 19.VII.1986 (PLYUSTSH, MAKHAT, SAZONOV), 4.–6.VII.1988 (BOGDANOV); 1 ♂, 2 ♀♀, Inyl'chek-Tau mountain range, Tashkoro, 19.VII.1982 (PLYUSTSH). China, Xinjiang: 1 ♂, Aksu, 1914 (G. RÜCKBEIL); 1 ♂, 1 ♀, "Kuruk-tag, pascus Ak-Tash, init. VII 1914" (RÜCKBEIL); 2 ♀♀, Hotan (BABKIN).

***Palearctia glaphyra manni* (STAUDINGER, 1881)**

Stett. Ent. Ztg. **17**: 402 (*Arctia glaphyra manni*).

= *Arctia glaphyra manni* ALPHERAKY, 1882, Horae Soc. Entomol. Ross. **17**: 29, t. 1, f. 32.

Type locality: Kungess (Kunes He) river upper course, Arshan (Arxan) terrain in China, Xinjiang.

Distribution. North-East Tien Shan at the Ili river headwaters.

Description (figs 8e, f). A rather large moth, the forewing length being 16–17 mm. The wing pattern is well developed and consists of separate spots on the forewing, and (the main character!) large spots on the hindwings.

Notes. The type series of the subspecies was collected by S. ALPHERAKY in the upper course of the Kungess river (Kunes He) in 1874. Several specimens, which were supplied with the name *mannii*, were sent to O. STAUDINGER. The latter author described them in his work, but with another spelling — *manni*. As the paper of S. ALPHERAKY was published one year later, the subspecies should be named according to the spelling of STAUDINGER and with his authorship.

Material. China: Xinjiang: 1 ♂, the lectotype of *A. g. manni* ALPH., Arshan, 1.VII.1874 (ALPH[ERAKY]); 1 ♂, "Thian-chan", N 568 (STAUD[INGER]); 1 ♂, "Alatau", N 835, 1882 (STAUD[INGER]); 1 ♂, "Thianchan", N 520, 16.VII.1912 (coll. A. MEINHARD); 4 ♂♂, Jeldus (E. LE MOULT); 1 ♂, Kuldja [Gulja] (coll. Zoological Museum of Helsinki University, Finland); 1 ♀, same locality.

***Palearctia wagneri* (PÜNGELER, 1918)**

Z. öster. Ent. Ver. **3**: 46, textfig. (*Arctia wagneri*).

Type locality: "Sary-Djas" [Sarydzhaz] (Central Tien Shan) in Kyrgyzstan.

Distribution. Since the type had been collected all the known specimens of this species originated from the Kaingdy-Katta Range in Central Tien Shan in Kyrgyzstan.

Biology. The moths were collected at 3200 m above sea level in the first half of June.

Description (fig. 8g). This species is mostly related to *P. glaphyra* (Ev.), but differs by the presence of the fused narrow transversal bands M₁ and M₂ on the forewing upperside and also by the presence of black scales on the cubital veins.

Genitalia. See figs 3f, 4d.

Notes. According to the male genitalia structure, including the vesica, *P. wagneri* (PÜNL.) and *P. glaphyra* (Ev.) are very similar. Nevertheless, the differences in wing pattern, with respect to a full sympatry of both taxa at the Kaingdy-Katta Range, proved them to be different species.

Material. Kyrgyzstan: 3 ♂♂, 1 ♀, Kaingdy-Katta mountain range, Tashkoro, 5.–6.VII.1984, 11., 18.VII. 1986 (SAZONOV, MAKHAT, GANSON, PLYUSTSH).

***Palearctia golbecki* spec. nov. (figs. 10a, b)**

Male. The forewing length is 9.5 mm in the holotype and 11.0 mm in the paratype, the wing expanse being 20 and 23.5 mm, respectively. The head is covered with bristly hairs, which are dark brown and light grey. The antennae are bipectinate, with equal branches 4 times longer than the diameter of an antenna unit. The body is short, 7.5–9.0 mm, stumpy, covered with thick brownish-black hairs. The margins of the patagia, tegulae, two longitudinal bands on the thorax, the lateral sides, end and underside of the abdomen are covered with light-grey hairs. The femorae are covered with long dark-brown and light-grey hairs, the tibiae and tarsi with light-grey scales. The fore tibia is short, 1.5 mm in length, with a large epiphysis. The single pair of spurs on the middle tibia and the two pairs of spurs on the hind tibia are slightly shorter than the tibia diameter. The wings of the holotype are narrow, those of the paratype are slightly wider, with sharp apices. The wing pattern is as in *P. gratiosa* (GR.-GR.), but the basal, medial and discoidal spots are very dark, while the margin is much lighter. The hindwing is white with dark veins, a discoidal spot and a wide margin. The pattern on the underside is similar, but the bands M₁ and M₂ on the forewing are reduced; the hindwing has a diffuse brown spot at the costa base.

Genitalia. See fig. 3g. The general structure as in *P. glaphyra*, but the aedeagus has dorsally no spurs at the distal part, and the vesica structure is different.

Female. The forewing length is 9.5 mm, the wing expanse being 21 mm. The head is covered with brownish-grey sleeky hairs. The palpi are stretched forward. The antennae are biserrate. The body is thick, cylindrical, 11.5 mm in length, covered with dense brownish-black hairs. The margins of the tegulae, patagia, two longitudinal bands on the thorax and abdomen underside, as well as the evenly coloured legs, are covered with brownish-grey scales. The spurs are short, not longer than half of the tibia diameter. The wings are very narrow, with contrasting black strokes: one in the middle of the cell, another behind the base of vein 2 (Cu₂); a discoidal spot, a transversal spot between the bases of veins 2 and 3 (Cu₁ and Cu₂), a spot between veins 5 and 6 (M₁ and M₂), another one at the distal part of vein 2 (Cu₂), and three spots along vein 1 (A) and in front of it. The marginal part of the wing is slightly darkened. The hindwing is almost entirely dark-brown with a clear lightened space between veins 3 and 4 (M and Cu₁) and a weakly lightened space behind vein 2 (Cu₂). The pattern of the wing underside is similar, but strongly lightened.

Material. Holotype ♂, "Prov. Syr-Darja, Jug. Alexandrinum, Loc. Tshai-Sandyk, 910.VI.25, A. GOLBECK" (TSCHETVERIKOV collection in the Zoological Institute, St.-Petersburg). Paratypes: 1 ♀, same locality and date; 1 ♂, Kirgizskaya SSR, Kirgiz Range (Alexander Mts.), Tyuz-Ashu pass, 3400 m, 17 VII.1977 (V. MURZIN).

***Palearctia gratiosa* (GRUM-GRSHIMAILO, 1890)**

Mém. lépidop. Ed. N. M. ROMANOFF 4: 533, T. 14, f. 5 (*Arctia glaphyra gratiosa*).


Type locality: "Artcha-Basch", 9,500 p. (Alai Range). The type specimen could not be found in the Zoological Institute, St.-Petersburg.

Distribution. Uzbekistan: West Tien Shan (Chatkal mountain range), Alai-Pamir (Turkestan and Zeravshan mountain ranges); Kyrgyzstan: Chatkal, Kirghiz Alatau (Alexander Mts.), Terskei Alatau, and Alai Ranges; Tajikistan: Turkestan, Hissar, Peter-the-Great and Trans-Alai Ranges, Pamirs); Afghanistan (Karakoram); India (Kashmir). West China, including Sichuan (DANIEL, 1961). It should be noted that the moth depicted on fig. 1565 (t. 68) in FANG CHENG-LAI (1982), in my opinion, is not "*Micrarctia glaphyra* (EVERSMANN)", as it is mentioned, but instead *P. gratiosa* (GR.-GR.). The nominotypical subspecies occurs in the Alai, Peter-the-Great Ranges and the West Pamir.


Biology. The moths were found at 2200–4500 m above sea level, the caterpillars are polyphagous on dicotyledonous plants.

Description. The main characters are noted in the key. The groundcolour varies from white to yellowish, pinkish or greyish. The forewing pattern is very variable (figs 10c–d). The hindwings may be of


4a




4b




4c




4d




4f



4e



4h



4g




Fig. 4: Aedeagus of *P. glaphyra*, *P. wagneri* and *P. gratiosa*. a — *P. g. glaphyra* Ev., holotype; b — *P. g. dublitzkyi* O. BANG-HAAS (Trans-Ili Alatau); c — *P. g. manni* STGR. (East Tien Shan); d — *P. wagneri* PÜNGL. (Kaingdy-Katta mountain range, Central Tien Shan); e — *P. g. aksuensis* O. BANG-HAAS (Kaingdy-Katta mountain range); f — *P. gr. caroli* subspc. nov., holotype; g — *P. gr. sergei* subspc. nov., holotype; h — *P. gr. gratiosa* GR.-GR. (Alai Range); i — *P. gr. rupicola* GR.-GR. (Pamirs); j — *P. gr. flavala* subspc. nov., holotype.








Fig. 5: Variation of male valvae of *P. glaphyra* (a-k) and *P. gratiosa* (l-q). a-b — *P. g. dublitzkyi* O. BANG-HAAS (Trans-Ili Alatau); c-g, i-j — *P. g. aksuensis* O. BANG-HAAS (c — "Alatau", d — Przhevalsk, e-g — Terskei Ala-Tau, i — "Thian-Shan", j — Kaingdy-Katta mountain range); h — *P. g. "ab. illustrata"* O. BANG-HAAS, Tien Shan; k — *P. g. manni* STGR. (East Tien Shan); l — *P. gr. caroli* subspc. nov. (Chatkal mountain range); m — *P. gr. sergei* subspc. nov. (Terskei Ala-Tau); n — *P. gr. gratiosa* GR.-GR. (Alai Range); o — *P. gratiosa* GR.-GR. ?ssp. (Turkestan mountain range); p — *P. gr. flavala* subspc. nov. (Hissar mountain range); q — *P. gr. rupicola* GR.-GR. (Trans-Alai Range).




6a




6b



6d



6c



6e




Fig. 6: Genitalia of *Palearctia*. a, c — *P. erschoffii* ALPH. (East Thian-Shan); b — lectotype of "Arctia erschoffi issyka STGR."; d, e, i — *P. ferghana* STGR. (Alai Range); f — *P. f. schottlaenderi* STRAND (Tien Shan); g — *P. f. susamyrta* subspec. nov., holotype; h — *P. f. turkestana* subspec. nov., holotype; j — *P. gracilis* spec. nov., paratype (Chatkal mountain range); k — *P. marxi* O. BANG-HAAS, holotype. c, i: females, e — h: male aedeagus.

different colouration, normally red, sometimes orange, yellow, greyish-white or of different colours: pinkish at the base, yellowish at the margin. The discoidal spot may be well developed or fully absent. The marginal spots also vary. The veins are usually invisible, but often set with black scales at the wing base. Several aberrations have been described: ab. *flava* B.-H., with yellow hindwings, ab. *nigromarginalis* B.-H., with a continuous black margin at the hindwing. The nominotypical subspecies is characterized by a bright red hindwing without a discoidal spot. The V-shaped stroke is well developed, sometimes narrowly interrupted.

Genitalia. See figs 3–5.

Material. Kyrgyzstan: 1 ♂, "mts. Alaiensis centr., cl. merid., fl. Koksu, pr. Kosch-Dube, 3900 m, 8.VIII. 1964" (BUNDEL); Alai Range: 1 ♂⁴, Iordan, the Ulyumur terrain, 16.VII.1935 (TSVETAEV); 2 ♂♂, Gao-mysh pass, above Khaidarken, 3500–3600 m, 16.–17.VII.1983 (SAZONOV); 1 ♂, Kuruksai Mountain Range, Alaudin river, 3600 m, 27.VII.1987 (TARASOV). Tajikistan: 1 ♂, "Buchara or., Jg. Petri Magni, val. fl. Kara-schura, Ic. Gursy-Tasch", 11.VII.1911 (coll. TSCHETVERIKOV); Peter-the-Great Range: 1 ♀, Tupehek, 13000 feet, 14.VII.1913 (KOCHUBEI); 1 ♂, Ganoshou near Tadzhikabad, 3500 m, 16.VII. 1971 (TSVETAEV); 1 ♂, terrain Sus (Darai-Nazarak), 3000 m, 8.VII.1984 (PETRIKEVICH); Pamirs: 1 ♂, Vanch Range, terrain Lyangar, Kashtiga river, 3500 m, 31.VII.1973; 1 ♂, Shugnan Range, terrain Sangou-Dara, 3500 m, 18.VII.1965 (TSVETAEV); 1 ♂ of ab. *flava* B.-H., "Pamir S.W., Jugum Want-schesis, fl. Gudjiwas, 3000 m, 4.VIII.1955" (BUNDEL); 1 ♂, Russian Geographical Society glacier, 29.VII.1983 (KAABAK).

***Palearctia gratiosa caroli* subspec. nov.**

Distribution. Uzbekistan: Chatkal Mountain Range; Kyrgyzstan: Chatkal Mountain Range, Kirghiz Alatau (Alexander Mts.). Most probably this subspecies will be found on other ranges of the West Tien Shan.

Description (figs 9a–c). The hindwing usually has an uneven groundcolour: darker reddish at the base and the center and lighter yellowish at the external and costal margins. This yellowish zone may be very narrow or extremely wide, nearly reaching the wing base. The discoidal spot is narrow or absent. The veins bear few black scales at the base. On the forewing upperside the V-shaped stroke is always interrupted behind the cell and the origin of vein 2 (Cu_2), forming two separate medial bands (M_1 and M_2).

Material. Holotype ♂, ex larva, "UdSSR – Kirghisia, Thian-Shan Mts., Alaarcha valley, 2200–2700 m, 6.83, ČERNÝ". Paratypes: 1 ♀ ex ovo, "USSR – Kirghisia, Frunzenskaya region, Kirghizskij Ala-Tau, Alaarcha valley, 2600–3000 m, ČERNÝ"; the Kirghiz Ala-Tau: 1 ♀, Uzun-Gyr, 31.VII.1981 (PLYUSTSH); 1 ♂, Ala-Archa, 3500 m, 31.VII.1986 (NEKRASOV); 1 ♂, Chatkal mountain range, Aksu river, 3300 m, 3.VIII.1950 (BUNDEL). The name is given in honour of KAREL ČERNÝ, Innsbruck, Austria.

***Palearctia gratiosa sergei* subspec. nov.**

Distribution. The southern slopes of the Terskei Ala-Tau in Inner Thian-Shan, Kyrgyzstan.

Description (fig. 8h). The largest subspecies, the forewing being 15 mm in length, the wing expanse is 31.5 mm. The hindwing is uniformly orange with a large contrasting discoidal spot, submarginal spots and dark veins. The spots of the forewing are contrasting, the V-shaped spot being well developed.

Material. Holotype ♂, SE of the Terskei Ala-Tau, 6 km upstream of the Molo gorge, 12 km of the Kuilyu river (Sary-Dzhaz tributary), 19.VII.1984 (SAZONOV). Paratypes: 3 ♂♂, 2 ♀♀, same locality (SAZONOV). The name is given in honour of SERGEI SAZONOV (Moscow).

4 This specimen may have originated from Uzbekistan, because the Iordan town belongs to this country, but the neighbouring mountains to Kyrgyzstan.

***Palearctia gratiosa flava* subspec. nov.**

Distribution. Uzbekistan: Zeravshan and Turkestan mountain ranges; Tajikistan: Zeravshan, Turkestan, and Hissar mountain ranges.

Description (figs 9e–g). The forewing as in *P. g. gratiosa*, but the hindwing is always yellow. The discoidal spot is weakly expressed or fully absent.

Material. Holotype ♂, "Samarkand", [Magiyan river, northern slope of the Hissar Range, 30 km SE of Pendzhikent], No 34, 1886 (HABERHAUER). Paratype ♀, same locality.

Additional material: 1 ♂, Samark[and] region, [Hissar mountain range], Laudan pass, 17.VII.1912 (KIRICHENKO); 1 ♀, Zeravshan mountain range, Sary-ob (pr. Vory), 26.VI.1925; 3 ♂♂, Turkestan mountain range, Kumbel pass, 3000–3200 m, 7.VII.1976, 7.VII.1982 (GANSON, TSVETAEV, SAZONOV); 2 ♂♂, Hissar mountain range, Anzob pass, 3300 m, 5.VIII.1962 (TSVETAEV), 12.VII.1979 (PRASOLOV); 1 ♂, "Fanske Gory" Mts., Zigrat pass, 6.VII.1981 (FILEVA).

***Palearctia gratiosa rupicola* (GRUM-GRSHIMAILO, 1890)**

Mém. lépidop. Ed. N. M. ROMANOFF 4: 535, t. 19, f. 6 (*Arctia rupicola*).

= *Arctia glauca* STAUDINGER, 1892, Dtsch. Entomol. Zeit. IRIS 4:251, t. 3, f. 5.

Type locality: Aram [Aram-Kungei], Trans-Alai Range.

Distribution. Kyrgyzstan: eastern part of the Trans-Alai Range. Tadzhikistan: eastern part of the Trans-Alai Range, East and South-West Pamirs, westwards to the eastern part of the Shugnan Range.

Description (figs 9d, 9i, 10e). The hindwings are white, greyish-white, or black, the wing pattern is rather contrasting, as in the nominate subspecies.

Material. Trans-Alai Range: 1 ♀ (the holotype of *Arctia rupicola* GR.-GR.), "Arom-Kungei", 17.VII.1886 (kol. b. Vel. KN. NIKOLAYA MIKHAILOVICH) [coll. former Grand Duke NIKOLAI MIKHAILOVICH] (labels in russian); 1 ♀, same locality, 3.VII.1908 (AVINOV); 1 ♂, Kzhyr-Art pass, 20.VII.1982 (TARASOV). Pamirs: 1 ♂, Koitezek pass, 24.VII.1982 (SAZONOV); 1 ♂, Shugnan Range, Dzhilandy, 4500 m, 11.VIII.1989 (BOGDANOV).

***Palearctia gratiosa kashmirica* FERGUSON, 1985 comb. nov.**

Entomography, Ann. Rev. Biosyst., Sacramento 3:222, f. 44, 101–102 (*Palearctia kashmirica*).

Type locality: Baltistan, Kashmir, India.

Description (colour plate fig. 2). This race is very similar to *P. g. rupicola* (GR.-GR.) but the wing pattern is very diffuse.

Genitalia. Do not differ from those of *P. gratiosa* (GR.-GR.) (FERGUSON, 1985: f. 101, 102).

***Palearctia gratiosa postflavida* (HAMPSON, 1894)**

Moths Ind. 2: 18, f. 5; Cat. Arct. Agar. coll. Br. Mus.: 238, 132 (*Ocnogyna postflavida*).

Type locality: Kashmir, Skoro-la, 15,000 f.

Description. The hindwing is orange-yellow, the forewing pattern is diffuse.

Genitalia. Do not differ from those of *P. gratiosa* (GR.-GR.) (cf. FERGUSON, 1984: 456, f. 114, 14a).

Notes. DANIEL (1961) reported on a male of "Micrarctia postflavida Hmps." from Sichuan (China), Omeishan, leg. STÖTZNER. This specimen was not figured and described, so it belonging to subspecies *P. g. postflavida* (Hmps.) has to be confirmed, but I have no doubt that it belongs to the species *P. gratiosa* (GR.-GR.).

***Palearctia gratiosa lochmatti* (REICH, 1933) comb. nov.**Int. Ent. Zeit. 27: 312 (*Micrarctia lochmatti*).

Type locality: "Drittes Seitental des Shyok-Tals, 4800 m, ..., 3.VIII.1929 von Fr. LOCHMATTER"

Description. I haven't examined this taxon. The original description as follows: "Steht in Größe, Gestalt und Aussehen etwa in der Mitte zwischen *Micr. glaphyra gratiosa* ab. *flava* O. BANG-HAAS und *Phragmatobia (Oroncus SEITZ) wagneri* PÜNGELER. Flügelspannung: 31 mm. Beide Flügel stark schwärzlich überstäubt, so daß die hellen Zeichnungen sehr zurücktreten. Vorderflügel schwarzbraun. Schwarze rundliche Flecken in der Mitte und am Ende der Zelle, ferner in der Mitte des Innenrandes und am Außenwinkel und unterhalb der Spitze. Die Mittelbinde ist verschmälert und unvollständig, nur bis zu der weißen Submediana reichend. Die äußere Binde sehr verschmälert, scharf nach außen gewinkelt. Hinterflügel blaß gelblich, mit einem deutlichen Discalfleck, der bei *wagneri* ganz fehlt, und einer 2 mm breiten ganz zusammenhängenden Randbinde. Unterseite der Vorderflügel mit einer breiten dunklen Binde schräg von der Costa zur Querrippe und einer dunklen Randbinde. Unterseite der Hinterflügel wie oben, aber verwaschener und nicht so stark schwarz gesäumt. Körper oben schwarz, seitlich und unten gelb. Palpen schwarz mit orangegelber Spitze." According to this description I consider this taxon most probably belonging to *P. gratiosa* (GR.-GR.), but as a distinct subspecies.

***Palearetia erschoffii* (ALPHERAKY, 1882)**Horae Soc. Entomol. Ross. 17: 29, t. 1, f. 33 (*Arctia erschoffii*).= *Arctia erschoffi issyka* STAUDINGER, 1887, Stett. Ent. Ztg. 48: 82, **syn. nov.**= *Micrarctia kindermannii korlana* BANG-HAAS, 1927, Horae Macrolepid. 1: 67, t. 8, f. 47.

Type locality: "Julduz" [a river, Xinjiang, China].

Distribution. Uzbekistan?: Chatkal mountain range. Kyrgyzstan: Chatkal and Sussamyr mountain ranges, Kirghiz Alatau, Kungei Alatau, Terskei Alatau, Inner Tien Shan (Naryn), Central Tien Shan (Sarydzhab, Kaingdy-Katta mountain range). Kazakhstan: Trans-Ili Alatau, ?Kungei Alatau. China in Xinjiang: East Tien Shan: Julduz, Kuruktag Range. The nominotypical subspecies occurs in almost all parts of the species' range excluding the Naryn region.

Biology. The moths of the nominotypical subspecies have been collected at altitudes of 3000–3400 m above sea level, and, according to ALPHERAKY (1891), from 8000 to 9000 feet. The imagines fly from July to September, seldom already in June.

Description The forewing pattern is rather peculiar (figs 7f, g; colour plate fig. 5). The groundcolour varies from white to yellowish. The dark spots may vary slightly in size and form, rarely the costal spots of bands M_1 and M_2 are fused with the corresponding hind spots of these bands. Sometimes the costal and hind spots are stretched along the wing margins. The groundcolour of the hindwing is red, sometimes yellowish (ab. *flava* B.-H.). The discoidal spot is absent or, rarely, present only as few black scales. The submarginal spots (E_3) are narrow. The pattern of the hindwing underside is the same as on the upperside, but at the middle of the costa there may be a black stroke, sometimes fused with the discoidal spot. The nominotypical subspecies is characterized by its larger size (forewing length 12–15 mm, wing expanse 23–29 mm). The forewing groundcolour varies from white to slightly yellowish with black, less frequently brownish spots, but always more pointed than in the next subspecies. The hind spots on the forewing are usually fused along the margin.

Genitalia. See figs 6a–c.

Fig. 7: Imagines of *Palearctia*. a — *P. mongolica* ALPH., ♂, S. Tuva; b — holotype ♂ of "*Arctia serarum* GR.-GR.>"; c — *P. mongolica* ALPH., ♀, S. Mongolia; d — *P. ferghana ferghana* STGR., lectotype ♀; e — *P. erschoffii selmonsi* BÖTTCH., ♂, At-Bashi; f — *P. erschoffii erschoffii* ALPH., lectotype ♂; g — *P. erschoffii erschoffii* ALPH., ♀, East Tien Shan; h — *P. ferghana turkestana* subsp. nov., holotype ♂.




7a



7e




7b



7f



7c



7g



7d



7h

Notes. I have studied the type of *Arctia erschoffii issyka* STGR., described from the mountains southwest of Lake Issyk-Kul. This specimen, a male, has the following labels: "Issyk-Kul / mer.-occ. / 88 RCKBL.", "erschoffii v. issyka STGR.", "Zool. Mus. / Berlin". It does not differ from *P. erschoffii erschoffii* (ALPH.) and should be synonymized with it.

Material. Kazakhstan: Trans-Ili Alatau: 1 ♂, Oj-dzhejlau, 13.VIII.1935 (IONOV); 1 ♂, Turgen' river, 3000 m, 22.VIII.1983 (GAMAYUNOVA). Kazakhstan/Kyrgyzstan: Kungei Ala-Tau: 1 ♂, Kshi-Yuryukty river, 3200 m, 21.VII.1974 (coll. SHAPIRO). Kyrgyzstan: 1 ♂, Chatkal mountain range, Lake Sary-Chelek, 3000 m, 4.VIII.1983 (MURZIN); Kirghiz Ala-Tau (Alexander Mts.): 1 ♂ (TRUSSEVICH); 1 ♂, Pishpek [Bishkek], 25.V.1907; 1 ♂, Kegety-pass, 17.VII.1927 (JANKOWSKY); Terskei Ala-Tau: 1 ♂, Pokrovka, 3000 m, 2.VIII.1935 (TSVETAEV); 2 ♂♂, Molo gorge, 12 km upstream the Kuilyu river, 3300 m, 19.VII.1984 (SAZONOV); Issyk-kul: 3 ♂♂, (TANCRE); Central Tien Shan: 34 ♂♂, Kaingdy-Katta Range near Tashkoro, 10.VII.1982 (TARASOV), 21.VII.1985, 5.–22.VII.1986 (PLYUSTSH, MURZIN). China, Xinjiang, East Tien Shan: 1 ♂, the lectotype of *Arctia erschoffii* ALPH., Juldus, ".7.79" (ALPH[ERAKY]); 2 ♂♂, 1 ♀, same locality; 1 ♂, Tsamma, 9.VIII.1879 (ALPHERAKY).

Palearctia erschoffii selmonsi (BÖTTCHER, 1905)

Ent. Zeit. 19: 62, f. 18–21 (*Arctia erschoffii selmonsi*).

Type locality: Naryn, Inner Tien Shan, Kyrgyzstan.

Distribution. Inner Tien Shan in the upper courses of the Naryn River and its tributaries. A specimen of this subspecies, which was reported erroneously from Lake Issyk-kul, Rybach'e (DUBATOLOV, 1985), originated in fact from the At-Bashi region not far from Naryn.

Biology. This subspecies occurs at lower altitudes than the nominate subspecies, at about 2400 m above sea level.

Description (fig. 7e). A small subspecies, the forewing length being 9.5–10 mm, the wing expanse is 19.5–21 mm. The groundcolour of the forewing is always yellowish, the spots are brownish, slightly rounded. The spots on the underside of the forewing are not fused along the margin.

Material. Kyrgyzstan: Inner Tien Shan: 2 ♂♂, Naryn, 16.VII.1914 (NEZHIVOV), 1 ♂, same locality, 2400 m, 15.VII.1976; 1 ♂, At-Bashi, Char river, IX.1969 (material of D. I. BERMAN's expedition).

Palearctia ferghana (STAUDINGER, 1887) comb. nov.

Stett. Ent. Ztg. 48: 82 (*Arctia erschoffii ferghana*).

Type locality: "Margelan, Alai mont." [Kyrgyzstan / Uzbekistan?].

Distribution. Kyrgyzstan: Alai and Sussamyr Ranges. Tajikistan: Turkestan mountain range, Pamirs? Tien Shan (an unknown locality). Afghanistan (Anjuman pass), ?Nepal. The nominotypical subspecies occurs in the Alai Range.

Biology. According to V. S. MURZIN (pers. comm.) the moths occur in July at altitudes higher than 3000 m above sea level.

Description (figs 7d, 10i; colour plate fig. 6). In wing pattern this species is very similar to the previous one; the wing pattern can vary to a great extent (DANIEL, 1966: Tab. 3). The moths of the nominotypical


Fig. 8: Imagines of *P. glaphyra* Ev. (a–f), *P. wagneri* PÜNGL. (g) and *P. gratiosa* (Gr.-Gr.) (h). a — *P. g. glaphyra* Ev., holotype ♂; b — *P. g. dublitzkyi* O. BANG-HAAS, ♂, Trans-Ili Alatau; c — *P. g. aksuensis* O. BANG-HAAS ab. *obscurata* BÖTTCH., ♂, Naryn; d — *P. g. aksuensis* O. BANG-HAAS ab. *puengeleri* BÖTTCH., ♂, Terskei Ala-Tau; e — *P. g. manni* STGR., lectotype ♂ of "*Arctia glaphyra manni* ALPH.>"; f — *P. g. manni* STGR., ♀, East Tien Shan; g — *P. wagneri*, ♂; h — *P. gratiosa sergei* subsp. nov., holotype ♂.




8a




8b



8c



8d




8e



8f



8g



8h

subspecies are almost identical with *P. erschoffii* (ALPH.), but the forewing spots are more angular. The hindwings are usually orange-red with small spots.

Genitalia. See figs 6d–h. They are very characteristic, therefore by their structure the species is easily distinguishable.

Material. 1 ♀⁵, the lectotype of *Arctia erschoffii ferghana* STGR., "48", "origin", "Horae Macrolep. Vol. 1 abgebildet t. 8, f. 50 beschrieb. p. 67" Kyrgyzstan: 1 ♂, Alai Range, Dugoba river, 3000 m, 17 VII. 1984, 1 ♀, ex larva, same locality, 5.VIII.1985 (MURZIN); 1 ♂, "mts. Alaiensis cent., cl. merid., fl. Koksu, pr. fl. Kere-Kasyk, 4.VIII.1954" (BUNDEL).

***Palearctia ferghana schottlaenderi* (STRAND, 1912)**

Int. Ent. Zeit. (Guben) 6:2 (*Arctia schottlaenderi*).

Type locality: "Thianschangebirge", without exact locality.

Distribution. Not known by certainty, because all the examined specimens of this subspecies are old and labelled "Tien Shan". I haven't found any specimens from our days.

Description (fig. 10j; colour plate figs 3, 4). The groundcolour of the forewing is rose, the spots are diffuse. The hindwing is carmine-red, the submarginal spots (E₃) are very small.

Notes. Dr. A. WATSON kindly sent me a colour slide of the type specimen of *Arctia schottlaenderi* ab. *infrapicta* STRAND (colour plate fig. 3), which depicts a female, not a male, as mentioned in the original description (STRAND, 1912), and a colour slide of a water-colour figure of the *Arctia schottlaenderi* STRAND type specimen (colour plate fig. 4). The two specimens studied from the collection of the Zoological Institute (St.-Petersburg) do not differ from the specimens on these slides.

Material. 2 ♂♂, Tien Shan, NN 516, 517 (coll. MEINHARD).

***Palearctia ferghana susamyra* subsp. nov.**

Description (fig. 10k). This subspecies is similar to *P. f. schottlaenderi* STRAND, but the groundcolour of the forewing is pure white, and the transverse bands are fused. The black submarginal spots (E₃) and the discoidal spot on the hindwing are large; on the costal margin and at the bases of the veins there are black scales; the hindwing above vein 7 (Rs) is yellowish.

Material. Holotype ♂, "Kirghizskaya SSR, Susamyr Range, the Ala-bel' pass, 3300 m", 14.VII.1977 (MURZIN).

***Palearctia ferghana turkestanica* subsp. nov.**

Description (fig. 7h). The groundcolour of the forewing is white, while the spots are increased so as to fuse into transverse bands. On the hindwing there are a narrow discoidal spot and large spots along the margin, the anal vein bases are suffused with black scales.

Material. Holotype ♂, Tadzhikistan, Turkestan mountain range, the Shakhristan pass, 16.VII.1980 (PRASOLOV).

***Palearctia ferghana variabilis* (DANIEL, 1966), comb. nov.**

Mitt. Münch. Ent. Ges. 56 (7): 161–163, t. 3, f. 4–16 (*Micractia variabilis*).

Type locality: Afghanistan, Hindu Kush, Anjuman pass.

5 In the original description it was erroneously considered as a male.

Distribution. Afghanistan: Hindu Kush and South Badakhshan.

Description. According to the figures in the original description (DANIEL, 1966: f. 4–16), the taxon is extraordinarily variable in forewing pattern. The hindwing has a distinct and contiguous black margin or is fully black. Regarding the male genitalia structure (DANIEL, 1966: 162, Abb. 2) this taxon is conspecific with *P. ferghana* (STGR.).

Biology. The moths were collected from 12.VII. to 12.VIII. 1963 at 2900 m above sea level in Badakhshan and at 3900–4300 m in Hindukush.

***Palearctia ?ferghana mustangbhoti* (DANIEL, 1961)**

Veröff. Zool. Staatssaml. München 6 (4): 155, t. 11, f. 8 (*Micrarcacia mustangbhoti*).

Type locality: Nepal, Mustangbhot, Tange-Khola, 5000 m, 31.VIII.55.

Description. I haven't studied the type specimen of this taxon. The original description as follows: "Wenig größer als das Bild von *postflavida* bei SEITZ, plumper. Auf der schwarzbraunen Vorderflügel-Grundfarbe sind die Querbinden im Diskus und Außenrand der Vorderflügel wesentlich reicher, während der Wurzelstrahl fehlt. Die Hinterflügel-Grundfarbe ist rot, an der Basis und am Innenrand breit, am Außenrand schmal schwarz tingiert, mit kräftigem Zellschlüßfleck. Am Außenrand ist zwischen Mediana 2 und 3 ein roter Zahn, der den Saum erreicht. Die Fransen beider Flügel sind dunkel mit schwacher Teilungslinie (*postflavida* hat leuchtend gelbe Hinterflügel-Fransen). Die Vorderflügel-Unterseite ist heller, alle Zeichnungen breiter und verschwommener. Die Hinterflügel sind hellrot mit schwächerer Zeichnung als oberseits. Die Fühler sind schwach gezähnt wie bei *maculosa* GERN. Der Kopf, Thorax und das Abdomen sind kräftig dunkel behaart, ohne laterale Aufhellungen, Der Aderverlauf entspricht völlig dem der Urbeschreibung von *postflavida* beigegebenen Bild." According to this description and to the photograph (DANIEL, 1961: t. 11, f. 8), it is possible to conclude that this taxon is related to *P. ferghana* (STGR.) and probably conspecific with it. In this case it differs from the other subspecies of *P. ferghana* (STGR.) by the expanded white groundcolour and a contiguous band M₂ on the forewings, by large discoidal and marginal (E₃) spots on the hindwing and, mostly, by a strong black suffusion at the base and anal spaces of the hindwing.

***Palearctia gracilis* spec. nov. (fig. 10g–h)**

Male. The forewing length is 12–13.5 mm, the wing expanse being 23–26 mm. The head is covered with long dark brown and yellow-grey hairs, bristly on the vertex and semisleeky on the frons. The palpi are stretched forward, slightly longer than the frons hairs. The antennae are bipectinate, the short branches being three times and the long ones four times longer than an antenna unit diameter. The body is narrow, 8.5–11 mm long. The thorax is covered with long bristly black-brown and yellowish hairs, the latter concentrating along the margins of the patagiae and tegulae. The abdomen is covered with scales of the same colour, its tip being covered with hairs. On the dorsal side the light scales concentrate along the hind margins of the segments. The ventral part of the abdomen has two rows of black spots on the sternite sides. The legs are covered with yellow scales, the femora with bristly hairs. Sometimes the femorae, and, rarely, the tibiae are black laterally. The fore tibia is 1.5 mm long, with a big epiphysis. One pair of spurs is present on the middle tibia and two pairs of spurs on the hind tibia, nearly equal to the tibia diameter or slightly shorter. The wings are wide, the forewing is rounded apically. The groundcolour of the forewing is yellowish. There are two basal spots, often fused; the costal spot of the band M₁ is triangular or quadrangular in shape, sometimes narrow, often with rough margins, extended behind the cubitus. The costal spot of the band M₂ is fused with the discoidal one forming a rounded quadrangular or reniform spot with the hind inner angle drawn off and crossing vein 2 (Cu₂). The hind spots of the median bands are fused into a trapezoid or parallelogram-formed spot with one or two incisions on its fore margin. There are three spots of the submarginal band (E₃): the first one on the costa is quadrangular or composed of two fused

triangles, it reaches vein 4 (M_3); the second one at the middle is triangular; the third one, lying on the hind margin, is also triangular; sometimes all these spots are fused into a single band. Some specimens have the spots fused along the hind margin. The external margin (E_{1+2}) has widenings at the apex and between veins 4–5 ($M_2–M_3$) and 1–2 ($A–Cu_2$). The hindwings of the holotype are rose, those of the paratypes vary from yellow (ab. *flava* SHELUZHKO in litt.) to bright red. The hindwing base has black veins or is entirely black up to 2/3 of the anal margin length. Sometimes the band M_1 is developed. The discoidal spot is dark, rounded or reniform. The submarginal spots (E_3) are large, the costal one being reniform, the middle (between veins 2–3 or $Cu_1–Cu_2$) and tornal ones are rounded. The margin is narrow and black. The wing underside is yellow, on the forewing the pattern is weakly developed and consists of two basal spots, a rounded spot at the middle of the cell, and a large discoidal spot. The costal spot of the E_3 band is of the same shape as on the upperside, the medial one is reduced, the margin (E_{1+2}) is developed up to vein 2 (Cu_2), it is crossed by light veins. The hindwing underside has a costal stroke at the base or, instead of the band M_1 , a discoidal bracket-shaped stroke and three submarginal spots of band E_3 .

Genitalia. See fig. 6j. The valva have elongate tips, they are prominent basally, the aedeagus has no teeth.

Material. Holotype ♂, "holotypus ♂ *gracilis* SHELU.", "Fergana S., Jug. Tshatkal, fontes flum. Ning-Dzilki", 19.VII.1929 (JANKOWSKI, Mus. Zool. Univers. Kiev). Paratypes: 5 ♂♂, same locality, 20.VII.1929 (JANKOWSKI); 1 ♂, "Fergana S., Jug. Tshatkal, the Tshanatsh-pass, 7.VII.1932" (MOTOV); 1 ♂, Fergana, [Sary-Chelek], Khadzhi-Ata river, 8,000 [feet], 29.VIII.1904 (GRUM-GRSHIMAILO); 4 ♂♂, Kyrgyzstan, Chatkal Range, Sary-Chelek, 2700–3000 m, 26.VII.1983, 5.VIII.1983 (MURZIN).

Note. The new species differs strongly from the others by wing pattern and male genitalia structure. For the first time, the species was found in the collection of the Zoological Institute (St.-Petersburg), some specimens were kindly given for investigation by V. S. MURZIN. Later the species was found in the collection of the Zoological Museum of Kiev State University as an unpublished subspecies "*Micrarctia erschoffi gracilis*". So, I preserve SHELUZHKO's name for this species.

Palearctia marxi (O. BANG-HAAS, 1927) comb. nov.

Horae Macrolep. 1: 68, t. 8, f. 51 (*Micrarctia kindermannii marxi*).

Type locality: "Bashahr Staten, Poo nordöstlich Simla" [India, Himachal-Pradesh]. The type specimen, a male, was kindly loaned for investigations by Prof. H. J. HANNEMANN. It bears the labels: "49", "Bashahr / India sept.", "Type / O. BANG-HAAS", "*Micrarctia kindermannii / marxi* O. B.-HAAS", "Horae Macrolep. / Vol. I. abgebildet / t. 8 f. 51 / beschrieb. p. 68", "Zool. Mus. / Berlin"

Description (colour plate fig. 7). A good colour figure of this species was published in the original description. The forewing have contiguous narrow transversal bands; the hindwing is red with a contrasting black base and large discoidal and submarginal spots (E_3).

Genitalia. See fig. 6k. Very similar to *P. gracilis* spec. nov., but the tips of the valva and uncus are rounded.


Fig. 9: Imagines of *P. gratiosa* Gr.-Gr. *P. gr. caroli* subspec. nov. (a–c): a — ♂, Chatkal mountain range, b — holotype ♂, c — paratype ♀, Kirghiz Ala-Tau; d — *P. gr. rupicola* Gr.-Gr., ♂; e–g — *P. gr. flava* subsp. nov. (e — holotype ♂, f — paratype ♀, g — ♂, Anzob pass, Hissar mountain range); i — *P. gr. rupicola* Gr.-Gr., holotype ♂.



9a




9e



9b



9f




9c



9g



9d




9h




10a




10b



10c



10d




10e



10f



10g



10h




Fig. 10: Imagines of *Palearctia*. a–b — *P. golbecki* spec. nov. (a — holotype ♂, b — paratype ♀); c–d — *P. gratiosa gratiosa* Gr.-Gr. (c — ♂, Alai Range, d — ♂, Pamirs, glacier of the Russian Geographical Society); e — *P. gratiosa rupicola* Gr.-Gr., ♂, Pamirs, Shugnan Range; f — *P. glaphyra aksuensis* O. BANG-HAAS, ♂, Kaingdy-Katta mountain range, Central Tien Shan; g–h — *P. gracilis* spec. nov. (g — holotype ♂, h — ♂, Khadzhi-Ata river Bary-Chelek); i — *P. ferghana ferghana* STGR., ♂, Alai Range; j — *P. ferghana schottlaenderi* STRAND, ♂, Tien Shan; k — *P. ferghana susamyra* subspec. nov., holotype ♂.

References

- ALPHÉRAKY, S. (1882): Lépidoptères du district de Kouldjà et des montagnes environnantes. — Horae Soc. Entomol. Ross. **17**: 15–103, t. 1–3.
- ALPHERAKY, S. (1888): Neue Lepidopteren. — Stett. Ent. Ztg. **49**: 66–69.
- ALPHERAKY, S. (1891): Kuldzha i Tyan-Shan'. Putevye zametki [Gulja and Thian Shan. Travel notes]. — St.-Petersburg, 182 pp. (in russian).
- BANG-HAAS, O. (1927): Horae Macrolepidopterologicae Regionis Palaearcticae. I. — Dresden, 128 pp.
- BÖTTCHER, E. (1905): Neue und wenig bekannte Arctiiden aus Turkestan. — Entomol. Zeit. **19**: 53–55, 61–63.
- BREMER, O. (1870): Katalog kollektssi cheshuekrylykh professora EVERSMANNA, prinadlezhastshei Russkomu Entomologicheskому Obstshestvu [A Catalogue of the Lepidoptera collection of Professor EVERSMANN, belonging to the Russian Entomological Society]. — Trudy Russ. Entomol. Obstsh. **4** (1–2), separate pagination, p. 1–23 (in russian).
- DANIEL, F. (1961): Zygaenidae-Cossidae. In: Lepidoptera der Deutschen Nepal-Expedition 1955. — Veröff. zool. Staatssaml. München **6** (4): 151–162, Pl. 15.

- DANIEL, F. (1965): 53. Bombyces et Sphinges. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (Lepidoptera). – Reichenbachia, Staatl. Museum Tierkunde Dresden **7**(10): 93–102.
- DANIEL, F. (1966): Zwei neue Arctiidae aus Afghanistan (Lep.). – Mitt. Münch. entomol. Ges. **56** (7): 161–164, T. 3.
- DANIEL, F. (1967): 117. Bombyces et Sphinges II. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (Lepidoptera). – Reichenbachia, Staatl. Museum Tierkunde Dresden **9** (23): 201–208.
- DANIEL, F. (1968): 165. Bombyces et Sphinges III. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (Lepidoptera). – Reichenbachia, Staatl. Museum Tierkunde Dresden **11** (25): 265–277.
- DE FREINA, J. J. & T. J. WITT (1994): Zur Kenntnis der Gattung *Lithosarcia* DANIEL, 1954 mit Beschreibung zweier neuer Taxa. – *Atalanta* **25** (3/4): 535–542.
- DRAUDT, M. (1931): 4 Familie: Arctiidae, Bärenspinner. In: SEITZ, A., Die Gross-Schmetterlinge der Erde. – Stuttgart: Alfred Kernen, Bd. **6**, Suppl. zu Bd. 2, S. 61–94, T. 5–7.
- DUBATOLOV, V. V. (1985): Vysshye medveditsy (Lepidoptera, Arctiidae: Arctiinae) gor Yuzhnoi Sibiri. Soobstshenie 1 [Arctiinae (Lepidoptera, Arctiidae) of the South Siberian mountains. Part 1]. In: Chlenistonogie Sibiri i Dalnego Vostoka [Arthropoda of Siberia and the Far East]. – Novosibirsk: "Nauka", Sib. Dept., p. 134–159 (in russian).
- DUBATOLOV, V. V. (1987): K sistematike roda *Micractria* SEITZ, s.l. (Lepidoptera, Arctiidae) [On the systematics of the genus *Micractria* SEITZ, s.l. (Lepidoptera, Arctiidae)]. In: Nasekomye, klestshi i gelminty [Insects, mites and helminths]. – Novosibirsk: "Nauka". Sib. Dept., p. 30–47 (in russian).
- DUBATOLOV, V. V. (1990): Vysshye medveditsy (Lepidoptera, Arctiidae: Arctiinae) gor Yuzhnoi Sibiri. Soobstshenie 2 [Arctiinae (Lepidoptera, Arctiidae) of the South Siberian mountains. Part 2]. In: Chlenistonogie i gelminty [Arthropoda and helminths]. – Novosibirsk: "Nauka", Sib. Dept., p. 139–169 (in russian).
- DUBATOLOV, V. V. & J. A. TSHISTJAKOV (1989): Novyi vid roda *Palearctia* (Lepidoptera, Arctiidae) interesnyi primer konvergentnogo skhodstva u medvedits [A new species of the genus *Palearctia* (Lepidoptera, Arctiidae) – an interesting example of convergent similarity in tiger moths]. – *Zool. Zhurnal* **68** (11): 141–143 (in russian).
- EVERSMANN, E. (1843): Quaedam lepidopterorum species novae, in montibus Uralensibus et Altaicus habitantes, nunc descriptae et depictae. – *Bull. Soc. Imp. Nat. Mosc.* **16** (3): 535–553.
- FANG CHENG-LAI (1982): Arctiidae. – *Iconographia heterocerorum sinicorum*. Beijing: Institute of Zoology, Acad. Sinica, Science Press, p. 190–277, t. 63–73.
- FERGUSON, D. C. (1984): Two generic names for groups of holarctic and palaearctic Arctiini (Lepidoptera, Arctiidae). – *Proc. entomol. Soc. Wash.* **86** (2): 452–459.
- FERGUSON, D. C. (1985): Contribution toward reclassification of the world genera of the tribe Arctiini, part 1. – Introduction and a revision of the *Neoarctia*-*Grammia* group (Lepidoptera: Arctiidae, Arctiinae). – *Entomography, Ann. Rev. Biosystematics*, Sacramento **3**: 181–275.
- GROUM-GRISHIMAILO, G. [GRUM-GRZHIMAILO, G. E.] (1890): Le Pamir et sa faune lépidoptérologique. – *Mém. lepidop. Ed. N. M. ROMANOFF*. St. Petersb.: M. H. STASSULÉWITSCH, 576 p., 21 pl.
- GRUM-GRSHIMAILO, G. (1890): Lepidoptera nova vei parum cognita regionis palaearcticae. 1. – *Ann. Mus. Zool. Acad. Imp. Sciences St.-Petersburg* **4**: 455–472.
- GRUM-GRSHIMAILO, G. (1902): Adnotatio. – *Ann. Mus. Zool. Acad. Imp. Sciences St.-Petersburg* **7**: 204.
- HAMPSON, G. (1894): Moths. – Fauna of British India including Ceylon and Burma. London, t. **2**, XXII + 609 pp.
- HAMPSON, G. (1901): Catalogue of the Arctiidae (Arctianae) and Agaristidae in the collection of the British Museum (Nat. Hist.). – London, vol. **3**, Xix + 690 pp.
- KURENTZOV, A. I. (1965): Zoogeografiya Priamurya [Zoogeography of the Amur Region]. – Moscow-Leningrad: Nauka, 155 pp. (in russian).
- LEDERER, J. (1853): Lepidopterologisches aus Sibirien. – *Verh. Zool.-Bot. Ges.* **3**: 351–386.
- NIEPELT, W. (1911): Neue paläarktische Macrolepidopteren. – *Int. Ent. Zeit.* **58**: 274–275.

- POTANIN, G. N. (1950): Tangutsko-tibetskaya okraina Kitaya i Tsentral'naya Mongolia [The Tangut-Tibet outlying district of China and Central Mongolia]. – 2nd edition. Moscow: Gos. izd-vo geograf. lit., 652 pp. (in Russian).
- PÜNGELER, R. (1918): Zeit. öster. Ent.-Ver. Wien **3**: 44–47.
- REICH, P. (1933): Eine neue Arctiidae (Micrarctiinae) der holländischen Karakorum Expedition 1929–1930. – Int. ent. Zeit. **27**: 312.
- SEITZ, A. (1910): 4. Familie: Arctiidae, Bärenspinner. In: SEITZ, A., Die Gross-Schmetterlinge der Erde. Stuttgart: Alfred Kernen, Abt. 1, Bd. **2**, S. 43–103, T. 10–18, 56.
- SOKOLOV, G. N. (1936): Die Evolution der Zeichnung der Arctiidae. – Zool. Jahrb. Anat. **61** (2): 139–194.
- STAUDINGER, O. (1881): Beitrag zur Lepidopterenfauna Central-Asiens. – Stett. Ent. Ztg. **42**: 253–300, 393–424.
- STAUDINGER, O. (1887): Centralasiatische Lepidopteren. – Stettiner Ent. Ztg. **68**: 49–102.
- STAUDINGER, O. (1892): Neue Arten und Varietäten von Lepidopteren des paläarktischen Faunengebiets. – Dtsch. ent. Zeit. IRIS **4**: 224–339, T. 3.
- STRAND, E. (1912): Ein neuer Bär von Tianschan. – Int. Ent. Zeit. Guben **6**: 2.

Explanation of the colour plate (p. 89):

2	5	
3	6	
4	7	

Fig. 2: *Palearctia kashmirica* FERGUSON, 1985. Holotype. Now — *P. gratiosa kashmirica* FERG. Photo by courtesy of Dr. A. WATSON.

Fig. 3: *Arctia schottlaenderi* ab. *infrapicta* STRAND, 1912. Holotype. Now — *P. ferghana schottlaenderi* (STRAND). Photo by courtesy of Dr. A. WATSON.

Fig. 4: *Arctia schottlaenderi* STRAND, 1912. A water-colour figure of the holotype. Now — *P. ferghana schottlaenderi* (STRAND). Photo by courtesy of Dr. A. WATSON.

Fig. 5: *Arctia erschoffii issyka* STAUDINGER, 1887. Type. Now a synonym of *P. erschoffii erschoffii* (ALPH.).

Fig. 6: *Arctia erschoffii ferghana* STAUDINGER, 1887. Lectotype. Now *P. ferghana* (STGR.).

Fig. 7: *Micrarcia kindermannii marxi* O. BANG-HAAS, 1927. Holotype. Now *P. marxi* (O. B.-H.).

3. A list of the Arctiinae of the territory of the former U.S.S.R.

(Lepidoptera, Arctiidae)

by

V. V. DUBATOLOV

Summary: This is the first survey of the Arctiinae fauna of the former USSR territory, which occupied the major part of the Palearctic. It contains 98 species of these moths. This paper is the third of a series of checklists of Heterocera from the former USSR territory (as it was before the year 1990) and concerns the tiger-moths, Arctiinae. The moth fauna of the Palearctic is still insufficiently known. The western part (Western Europe and North-West Africa) is better studied because of DE FREINA & WITT's nice monograph¹ (1987) dealing with Bombycoidea, Sphingidae and some Noctuoidea, while the Asiatic fauna is rather poorly studied yet.

The first part includes a list of species and subspecies with their synonymy and geographic distribution, based on the collections of the Zoological Institute (St.-Petersburg, Russia), the Zoological Museum of Moscow State University (Moscow, Russia), the Zoological Museum of the Institute of Animal Systematics and Ecology (former: Biological Institute) (Novosibirsk, Russia), the Institute of Biology and Pedology (Vladivostok, Russia), the Yakutian Institute of Biology (Yakutsk, Sakha-Yakutia: Russia), the Zoological Museum of Kiev State University (Kiev, Ukraine), the Zoological Museum of the Zoological Institute (Kiev, Ukraine), and, partly, from literature. The main faunistic reviews are: ALPERAKY (1876 and 1908) on the Taganrog region (near the Azov Sea), BRYK (1942) on the Kuril Is., BUDASHKIN (1987) on the Crimea, CHRISTOPH (1877, 1884, 1887, 1889) on South Turkmenistan, DERZHAVETS et al. (1986) on the St.-Petersburg region, DIDMANIDZE (1978) on Georgia in Transcaucasia, DARITSHEVA & DUBATOLOV (1989) on Turkmenistan, DUBATOLOV (1985b, 1990d) on the mountains of South Siberia, DUBATOLOV (1991a) on southern Sakhalin and the Kunashir Is. (Kuril Is.), DUBATOLOV (1994) giving a distribution list of Arctiinae throughout the regions and countries of the former USSR, DUBATOLOV & ZOLOTARENKO (1991) on the West Siberian plain, EFETOV & BUDASHKIN (1987 and 1990) on the Crimea, FALKOVITSH (1969) on Central Kazakhstan, GROSSER (1983) on Bashkiria, INOUE (1961) on Sakhalin, KAISILA (1947) on Karelia, KONOVALOVA (1968) on the Kuril Is., KOSHANTSCHIKOV (1923–1925) on the Minusinsk region in South Siberia, KUMAKOV & KORSHUNOV (1979) on the Saratov region (Lower Volga), KOSTJUK & GOLOVUSHKIN (1994) on East Transbaikalia (Chita region), KUZNETSOV (1960) on the Kopetdagh Mts. in Turkmenistan, KUZNETSOV & MARTYNOVA (1954) on the Ural river in Western Kazakhstan, LVOVSKY (1971) on the Astrakhan region (Lower Volga), MERZHEEVSKAYA et al. (1976) on Belarus, LAVROV (1927) on the Omsk region in West Siberia, MIRZOYAN & ARUTYUNYAN (1980) on Armenia in Transcaucasia, MOLTRECHT (1929) on southern Far East, REMM & VIIDALEPP (1986) on Estonia, SACHKOV (1992) on the Samara region, SEDYKH (1974) on the Komi republic in North-Eastern Europe, SEDYKH (1979) on Kamchatka, SHAPOSHNIKOV (1904) on the North-Western Caucasus, SHELUZHKO (1941) on the Ukraine, STSHTETKIN (1960) on South-Western Tajikistan, SUVORTSEV (1894) on the Semipalatinsk region in North-Eastern Kazakhstan, TCHARUSHINA & SHERNIN (1974) on the Vyatka region in North-Eastern Europe, TSCHETVERIKOV (1993) on the Gorkii

1 Meanwhile two new generic names have been described in the Arctiidae. The first one is *Canararcinia* DUBATOLOV, 1990, Redkie gelminki, klestshi i nasekomye. Novosibirsk: 85, with the type species *Liparis rufescens* BRULLE, 1836 (= *Rhypania rufescens* (BRULLE)) from Canary Is. This genus was attributed to the tribe Spilosomini as relating to the genus group *Ocnogyna* LED. The second one is *Atlantarchia* DUBATOLOV, 1990, Novosti faunistiki i sistematiki. Kiev: 90 (Arctiini), from the Western Mediterranean, with *Euprepia dido* WAGNER, 1941 from E. Algeria and Tunisia and the type species, *Arctia oberthueri* (OBERTHÜR, 1890) from Algeria and Tunisia, *A. ungemachi* LE CERF, 1924 from Morocco, and *A. fasciata* (ESPER, 1785) from S. France, N. Italy and the Iberian Peninsula.

(Nizhnii Novgorod) region (Middle Volga), TSHISTJAKOV (1992) on the Khingan Nature Reserve in the Amur Region, SHLYKOV (1988) on the Penza region, SULCS & VIIDALEPP (1967) on the Baltic Provinces, VIIDALEPP (1979) on Tuva in Siberia, and VIIDALEPP & REMM (1982) on Sakhalin.

The distribution of the Arctiinae species throughout the regions and countries of the former USSR is summarized in DUBATOLOV (1994). Information on the Arctiinae distribution abroad was taken mainly from the works of DE FREINA & WITT (1987) for Europe, FANG CHENG-LAI (1982) for China, and INOUE (1982) for Japan.

The second part of this paper includes descriptions of new species and subspecies.

This work was partly supported by the International Science (Soros) Foundation and the Russian Academy of Natural Sciences.

The author would like to express his gratitude to Dr. O. KOSTERIN for preparing photographs and for correcting English in this paper.

Part I: List of Arctiinae

Callimorphini

***Callimorpha* LATREILLE, 1809**

Genera Crust. Insect. 4: 220.

(= *Panaxia* TAMS, 1939, *Entomologist* 72: 73).

1. *Callimorpha dominula* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia, north to Southern Karelia (KAISILA, 1947), the headwaters of the river Volga, Nizhnii Novgorod, and Bashkiria; Caucasus; Transcaucasia; South Turkmenistan (Kushka). West Europe, north to Southern England, South Scandinavia (except for the southern part of the Iberian peninsula); Turkey; Iraq; Iran.

a. *Callimorpha dominula dominula* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1: 509 (*Phalaena*).

Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia. Central Europe.

b. *Callimorpha dominula rossica* KOLENATI, 1846

Mel. Ent. fasc. 5: 95 (*Callimorpha*).

(= *Callimorpha dominula lutea* STAUDINGER, 1861, *Cat. Ed.* 1: 214; = *C. d. teberdina* SHELUZHKO, 1934, *Ent. Zeit.* 48: 74; = *C. d. swanetica* REICH, 1935, *Ent. Rundschau* 53: 9).

Caucasus; Transcaucasia, except for the Talysh Mts.

c. *Callimorpha dominula philippii* BARTEL, 1906

Soc. Ent. 21 (6): 41 (*Callimorpha philippii*).

(? = *Callimorpha dominula persica* LE CERF, 1913, *Ann. Hist. Nat. Paris Ent.* 2: 82, t. 1, f. 10).

Azerbaijan (Talysh Mts.); South Turkmenistan (Kushka).

Euplagia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 180.

2. *Euplagia quadripunctaria* (PODA, 1761)

Baltic Provinces (Lithuania, Latvia); Belarus; Ukraine, Crimea; Moldova; European Russia, north to the St.-Petersburg region, Middle Volga, and Southern Urals; Caucasus; Transcaucasia; South Turkmenistan (Kopetdagh Mts.). South and Middle Europe, north to Southern England (except for the North Sea and Baltic coasts); Turkey; Syria; North Iran.

a. *Euplagia quadripunctaria quadripunctaria* (PODA, 1761)

Insecta Mus. Graec.: 89, No. 20 (*Noctua*).

(= *Phalaena hera* LINNAEUS, 1767, Syst. Nat. (Edn. 12) 1 (2): 834).

Baltic Provinces; Belarus; Ukraine; Moldova; European Russia; Caucasus. West Europe.

b. *Euplagia quadripunctaria fulgida* (OBERTHÜR, 1896)

Et. Ent. 20: 56, t. 10, f. 174 (*Callimorpha hera*).

Transcaucasia; South Turkmenistan (Kopetdagh Mts.). S. Greece; Turkey; Syria; North Iran.

3. *Euplagia splendidior* (TAMS, 1922)

Entomologist 55: 196 (*Callimorpha quadripunctaria*).

(= *Callimorpha quadripunctaria tkatshukovi* SHELJUZHKO, 1935, Zeitschr. Öster. Ent. Ver. Wien 20: 21).

South Transcaucasia (Armenia, Azerbaijan: Nakhichevan). East Turkey; North Iraq; West Iran (Zagros Mts.).

Eucallimorpha DUBATOLOV, 1990

Taksonomiya nasekomykh i gelmintov. Novosibirsk: 99–100.

4. *Eucallimorpha principalis* (KOLLAR, 1844)

In: HÜGEL, Kaschmir 4 (2): 465, t. 20, f. 2 (*Euprepia*).

Tajikistan (Pamirs). Afghanistan; Pakistan; India (Kashmir, Himalaya); Nepal; China (Zhejiang, Jiangxi, Sichuan, Yunan, Tibet).

a. *Eucallimorpha principalis fedtschenkoi* (GRUM-GRSHIMAILO, 1902)

Ezheg. zool. Muz. 7: 197 (*Callimorpha principalis*).

Tajikistan (Pamirs).

Cymbalophora RAMBUR, 1866

Cat. syst. Lepid. Andalousie (2): 231.

5. *Cymbalophora rivularis* (MÉNÉTRIËS, 1832)

Cat. raison.: 263 (*Chelonia*).

Transcaucasia (Armenia, West Azerbaijan); East Caucasus (Daghestan). Central Italy; the Balkans; Turkey; ? West Iran ("Turkey", Balaban, 70 km East of Van, 8.X.1986, leg. MOBERG & HILLMAN, in the collection of the Természettudományi Múzeum, Állattára, Budapest, Hungary. I have found this locality on the map of Turkey to be in fact on the Iran territory, 70 km east of Lake Van).

Carcinopyga FELDER, 1874

in FELDER & ROGENHOFER, Reise öst. Fregatte Novara (Zool.) 2 (Abt. 2): t. 101, f. 3.
 (= *Euarctia* STAUDINGER, 1887, Stett. Ent. Ztg. 48: 79).

6. *Carcinopyga lichenigera* FELDER, 1874

In FELDER & ROGENHOFER, Reise öst. Pregatte Novara (Zool.) 2 (Abt. 2): t. 101, f. 3.
Tajikistan: Pamirs (DUBATOLOV, 1990). East Afghanistan; North Pakistan; India (Kashmir, Ladakh).

7 *Carcinopyga proserpina* (STAUDINGER, 1887)

Stett. Ent. Ztg. 48: 79 (*Euarctia*).
Tajikistan (Zeravshan, Hissar, and Darvaz Mts., Pamirs). East Afghanistan. It can also be found in Uzbekistan, for example, in the western parts of the Zeravshan and Hissar Mts.

8. *Carcinopyga lindti* ČERNÝ, 1986

Neue Entomol. Nachrichten 19 (1/2): 31.

Uzbekistan (West Tien Shan: Chatkal, Karzhantau, and Pskem Mts.); Kyrgyzstan (Chatkal, Kirghiz Alatau Mts., Issyk-Kul: Kungei Alatau and Terskei Alatau Mts.). Possibly it can be found also in South Kazakhstan, for example, in the Karzhantau Mts.

Taxonomical note: I do not agree with THOMAS (1989) and consider *C. lindti* ČERNÝ a good species.

Axiopoena MÉNÉTRIÈS, 1842

Bull. Acad. Imp. Sci. St.-Petersb. 9: 42.

9. *Axiopoena maura* (EICHWALD, 1830)

Zool. spec. 2: 196 (*Bombyx*).

(= *Axiopoena fluviatilis* SWINHOE, 1885, Trans. Ent. Soc. Lond.: 351, t. 9, f. 7).

Transcaucasia: Azerbaijan (Nakhichevan: Ordubad); Turkmenistan (Great Balkhan and Kopet-dagh Mts., Badhyz: Gezgyadyk Mts.). Iran; Afghanistan; North Pakistan.

10. *Axiopoena karelini* MÉNÉTRIÈS, 1863

Cat. lep. Petersb. 3: 160, t. 17, f. 5 (*Axiopoena*).

(= *Axiopoena maura transcaucasica* SHELJUZHKO, 1926, Dtsch. Ent. Zeit. IRIS 40: 58).

Western Caucasus (Sochi in Russia; Abkhasia); Transcaucasia (Georgia, Armenia, Azerbaijan; Nakhichevan). East Turkey; North Iraq.

Note: In my revision of the genus *Axiopoena* MÉN. (DUBATOLOV, 1989) there is a mistake in fig 1. It must be read as follows: a – *A. karelini*; b – *A. maura*.

Tyria HÜBNER, [1819] 1816

Verz. bekannter Schmett.: 166.

(= *Hipocrita* HÜBNER, [1806], Tentamen determinationis digestionis : [1] (rejected and invalid work); = *Euchelia* BOISDUVAL, 1828, Eur. Lepid. Index meth.: 39).

11. *Tyria jacobaeae* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1: 511 (*Phalaena*).

Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia, north to Vologda and Vyatka; Caucasus; Transcaucasia (Georgia, Armenia, West Azerbaijan); West Siberia, north to Tyumen and Tomsk, east to Krasnoyarsk and Minusinsk; North Kazakhstan and the mountains of East Kazakhstan; the mountains of eastern Middle Asia (in South Kazakhstan, Uzbekistan, Kyrgyzstan and North Tajikistan). West Europe, north to Ireland, southern parts of Scotland and Scandinavia; Asia Minor; China (Xinjiang); North America: from the end of '50s it was introduced into Canada (British Columbia) and the USA (Oregon, California) (NAGEL & ISAACSON, 1974).

Dodia DYAR, 1901

Jl. N. Y. ent. Soc. 9: 85.

(= *Hyalocoa* HAMPSION, 1901, Cat. Lep. Phal. Br. Mus. 3:xi, 5, 202).

12. *Dodia diaphana* (EVERSMANN, 1848)

The mountains of South Siberia from the Altai to Transbaikalia and the Stanovoi Mts.; Amur region; Sikhote-Alin Mts.; Central and East Yakutia; Magadan region (Upper Kolyma and Omolon rivers). North Mongolia (Chovsgol region).

a. *Dodia diaphana diaphana* (EVERSMANN, 1848)

Bull. Soc. Imp. Nat. Moscou 21 (3): 212 (*Lithosia*).

The mountains of South Siberia; Amur region; Sikhote-Alin Mts.; Central Yakutia. North Mongolia.

b. *Dodia diaphana arctica* TSHISTJAKOV, 1988

Entomol. Obozr. 67: 641, f. 7

East Yakutia; southern part of the Magadan region.

13. *Dodia albertae* DYAR, 1901

Jl. N. Y. ent. Soc. 9: 85.

The mountains of South Siberia from the East Sayan Mts. to Transbaikalia and the Stanovoi Mts.; Polar Siberia (North Urals, southern part of the Taimir peninsula); Yakutia; Magadan region (Upper Kolyma). North Mongolia; Alaska; Canada, east to Quebec.

a. *Dodia albertae atra* (A. BANG-HAAS, 1912)

Dtsch. Ent. Zeit. IRIS 26: 109 (*Hypocrita*).

(= *Hyperborea kozhantshikovi* SHELJUZHKO, 1918, Neue Beitr. syst. Insektenk. 1 (13): 104).

Mountains of South Siberia; Yakutia; southern part of the Magadan region. North Mongolia.

b. *Dodia albertae eudiopota* TSHISTJAKOV, 1988

Entomol. Obozr. 67: 638, f. 5.

Polar Urals; Taimir; North-West Yakutia.

14. *Dodia sazonovi* DUBATOLOV, 1990

Chlenistonogie i gelmnty. Novosibirsk: 148–149, f. 2 1.

Altai (Aktash).

15. *Dodia kononenkoi* TSHISTJAKOV & LAFONTAINE, 1984

Transbaikalia (Khamar-Daban Mts., Chita region: Sokhondo Mt., Yablonovy, and Udokan Mts.); Sikhote-Alin Mts.; Magadan region (Upper Kolyma). Canada (Yukon territory).

a. *Dodia kononenkoi kononenkoi* TSHISTJAKOV & LAFONTAINE, 1984

Canad. Entomol. **116**: 1553, f. 5, 6, 9.

Magadan region (Upper Kolyma). Canada (Yukon territory).

b. *Dodia kononenkoi transbaikalensis* TSHISTJAKOV, 1988

Entomol. Obozr. **67**: 634, f. 2.

Transbaikalia.

Note: The preservation place of the subspecies holotype was given incorrectly in the description: it must be read as the Zoological Institute (St.-Petersburg).

c. *Dodia kononenkoi sikhotensis* TSHISTJAKOV, 1988

Entomol. Obozr. **67**: 636, f. 3.

Primorye (Sikhote-Alin Mts.).

Lacydes WALKER, 1855

List Specm. lepid. Ins. Colln. Br. Mus. **3**: 685.

(= *Acymba* RAMBUR, 1866, Cat. syst. Lepid. Andalousie (2): 235; = *Palparctia* SPULER, 1906, Schmett. Eur. **2**: 133; = *Volgarctia* ALPHERAKY, 1908, Horae Soc. Ent. Ross. **38**: 606).

16. *Lacydes spectabilis* (TAUSCHER, 1806)

South-East Ukraine (PLJUSTSH, KOSTJUK, 1988); eastern part of European Russia, north to Penza and Vyatka; southern part of West Siberia (steppe and forest steppe belts: Kurgan region, Karasuk in the Novosibirsk region, and the Kulunda steppe in the Altaiskii Krai); Kazakhstan; Middle Asia; Southern Transcaucasia (Armenia). East Turkey; North Iraq; Iran; Afghanistan; China (Xinjiang); South Mongolia (a male in the collection of the Természettudományi Múzeum, Állattára, Budapest, Hungary).

a. *Lacydes spectabilis spectabilis* (TAUSCHER, 1806)

Mem. Soc. Nat. Univ. Imp. Mosc. **1**: 212, f.

(= *Eyprepia intercissa* FREYER, 1842, Neuere Beitr. Schmett. **4**: 118, pl. 356; = *Eyprepia incissa* FREYER, 1842, Neuere Beitr. Schmett. **4**: pl. 356).

South-East Ukraine; eastern part of European Russia; West Siberia; Kazakhstan; Middle Asia, including the plane part of Turkmenistan. Afghanistan; China (Xinjiang); South Mongolia.

b. *Lacydes spectabilis annelata* (CHRISTOPH, 1887)

Mem. lepidop. Ed. N. M. ROMANOFF **3**: 55, t. 3, f. 2 (*Arctia spectabilis*).

(= *Volgarctia kendevani* SCHWINGENSCUSS, 1937, Zeit. öst. Ent. Ver. **22**: 60, t. 3, f.)

Turkmenistan (Kopetdagh and Great Balkhan Mts.). Iran (Elburs).

c. *Lacydes spectabilis sheljuzhkoi* DUBATOLOV, 1996 subspec. nov. (see p. 70)

Southern Transcaucasia (Armenia). East Turkey.

Spiris HÜBNER, [1819] 1816

Verz. bekannter Schmett.: 169.

(= *Emydia* BOISDUVAL, 1828, Eur. Lepid. Index meth.: 39).

17. *Spiris striata* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) **1**: 502 (*Phalaena*).

(= *Phalaena grammica* LINNAEUS, 1758, Syst. Nat. (Edn. 10) 1: 822; = *Euprepia funerea* EVERSMANN, 1847, Bull. Soc. Imp. Nat. Moscou 20 (3): 77, t. 5, f. 5; = *Coscinia striata strandi* OBRAZTSOV, 1936, Festschr. E. STRAND 2:239; = *Coscinia striata wisniewskii* WOJTUSIAK & NIESOLOWSKI, 1946, Acta Mus. Hist. nat. Acad. polon. 6: 62).

Baltic Provinces; Belarus; Ukraine, Crimea; European Russia, north to the St.-Petersburg region (Luga) and the Upper Volga; Caucasus; Transcaucasia (Georgia, Azerbaijan); North and East Kazakhstan; South Siberia, north to Kurgan, Tomsk, Krasnoyarsk, Irkutsk, and Central Yakutia; it is extreme rare in Transbaikalia (1 ♂, Petrovskii Zavod in the Chita region, in the collection of the Kiev State University). South and Central Europe (except for the North Sea coast and the western and southern parts of the Iberian peninsula); Asia Minor; Syria; China (Xinjiang); West Mongolia; it is rare in Central Mongolia, too.

18. *Spiris bipunctata* (STAUDINGER, 1892)

Dtsch. Ent. Zeit. IRIS 5:345 (*Emydia striata*).

(= *Spiris bipunctata* ab. *nigrina* DUBATOLOV, 1985, Chlenistonogie Sibiri i Dalnego Vostoka. Novosibirsk: 137, an aberration with uniformly black wings).

Siberia: southern slopes of the West Sayan Mts. (Sayano-Shushenskii Nature Reserve) in the southern part of Krasnoyarskii Krai; Tuva; Transbaikalia; Middle Amur. Central and Eastern Mongolia; China (Xinjiang, Qinghai, Shanxi, Heilongjiang).

Coscinia HÜBNER, [1819] 1816

Verz. bekannter Schmett.: 169.

19. *Coscinia cibraria* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:507 (*Phalaena*).

(= *Phalaena cibrarium* LINNAEUS, 1761, Fauna Suecica (Edn. 2):302; = *Emydia cibrum sibirica* STAUDINGER, 1892, Dtsch. Ent. Zeit. IRIS 5:346; = *Coscinia cibraria fumidaria* O. BANG-HAAS, 1927, Horae Macrolepid. 1:58, t. 8, f. 3; = *Coscinia cibraria nikitini* O. BANG-HAAS, 1938, Entomol. Zeit. 52 (22): 179).

Baltic Provinces; Belarus; North Ukraine; European Russia, north to Karelia and Vyatka, south to Samara; North Kazakhstan; South Siberia, north to Tyumen, Tomsk, the Ob-Enisei canal (Aleksandrovskii Shlyuz, 59°N), the middle part of the Angara river, and Central Yakutia; it is rare in the Chita region and Middle Amur. West Europe, north to South-Eastern England and South Scandinavia (except for the south-western part of the Iberian peninsula); North Africa; North Mongolia; China (North Xinjiang, Heilongjiang).

Epimydia STAUDINGER, 1892

Dtsch. Ent. Zeit. IRIS 5:346.

20. *Epimydia dialampra* STAUDINGER, 1892

Dtsch. Ent. Zeit. IRIS 5:346, t. 3, f. 2.

The mountains of South Siberia from the south-eastern Altai and the Kuznetskii Alatau to the Chita region; Yakutsk; mountains of East Yakutia. North Mongolia. ?Tien Shan (see: DUBATOLOV, 1985b).

Uteheisa HÜBNER, [1819] 1816

Verz. bekannter Schmett.: 168.

(= *Deiopeia* CURTIS, 1827, Br. Ent. 4:169).

21. *Utetheisa pulchella* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:534.

Transcaucasia; Middle Asia. Migrates north to Latvia, Belarus, Karelia, the lower part of the Kama river in Europe and Semipalatinsk in Kazakhstan. The Mediterranean, migrates north to South England and South Scandinavia; Africa; West and South Asia; not recorded in China (FANG, 1982), Korea (WITT, 1980, 1985) and Japan (INOUE, 1982).

Arctiini

Parasemia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 181.

(= *Nemeophila* STEPHENS, 1828, Illust. Br. Ent. (Haustellata) 2: 55, 72).

22. *Parasemia plantaginis* (LINNAEUS, 1758)

(= *Nemeophila macromera* BUTLER, 1881, Trans. ent. Soc. Lond.: 5; = *Nemeophila macromera leucomera* BUTLER, 1881, Trans. ent. Soc. Lond.: 5; = *Nemeophila melanomera* BUTLER, 1881, Trans. ent. Soc. Lond.: 5; = *Parasemia plantaginis japonica* INOUE & KOBAYASHI, 1956, Tinea 3: 138, t. 17, f. 4–6; = *Parasemia plantaginis jezoensis* INOUE, 1976, Bull. Fac. domest. Sci. Otsuma Wom. Univ. 12: 170, pl. 2, f. 47).

Baltic Provinces; Belarus; North Ukraine, mostly in the Carpathians; European Russia, north to Murmansk and Ukhta (Komi republic), south to Krasnodar and Saratov; Caucasus; Transcaucasia; North-East Kazakhstan (Altai and Saur Mts.); Siberia, north to Beryozov (lower course of the Ob river), the Nizhnyaja Tunguska, Vilui, and Aldan rivers; the South of the Magadan region; Kamchatka, north to Verkhoturov Is. (about 60°N); Amur basin; Primorye; Sakhalin; Northern and Southern Kuril Is. West Europe (except for South Italy and South Greece); Asia Minor; North Iran (Elburz Mts.); Mongolia; China (Xinjiang, Qinghai, Sichuan, Shanxi, Nei Mongol, Dunbei) Korea; Japan (Hokkaido, Honshu); ?Alaska; North and West Canada; Western USA.

a. *Parasemia plantaginis plantaginis* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:501 (*Phalaena*).

(= *Phalaena hospita* [DENIS & SCHIFFERMÜLLER], 1775, Ankündung syst. Werkes Schmett. Wien-ergegend: 316; = *Parasemia plantaginis uralensis* KRULIKOVSKY, 1904, Rus. Entomol. Obozr. 4(1): 30; = *P. p. carpathica* DANIEL, 1939, Mitt. Münch. ent. Ges. 29: 358).

Baltic Provinces; Belarus; North Ukraine; European Russia; South Siberia, except for high mountains. West Europe, excluding the Balkans; Mongolia, except for high mountains.

b. *Parasemia plantaginis caucasica* (MÉNÉTRIÈS, 1832)

Cat. raison.: 262 (*Chelonia*).

(= *Parasemia plantaginis passanauriensis* ALBERTI, 1973, Atalanta 4(6): 389).

Caucasus; Transcaucasia. Asia Minor.

c. *Plantaginis plantaginis sifanica* GRUM-GRSHIMAILO, 1891

Horae Soc. Ent. Ross. 25: 462.

(= *Parasemia plantaginis altaica* SEITZ, 1910, Gross-Schmett. Erde 2: 81, f. 16f).

High mountains of the Altai, Sayan and Tannu-Ola. Mongolia (high mountain areas); China (Qinghai).

d. *Parasemia plantaginis nycticans* (MÉNÉTRIÈS, 1859)

Bull. Acad. Imp. Sci. St. Petersb. 17: 217 (*Lithosia*).

(= *P. p. nycticans* auct., nec MÉNÉTRIÈS, 1859; = *Nemeophila plantaginis* ab. *melas* CHRISTOPH,

1893, Dtsch. Ent. Zeit. IRIS 6:88; = [*P.p.*] *trybomi* BRYK, 1942, Dtsch. Ent. Zeit. IRIS 56:31, nomen nudum).

The northern part of West Siberia; Evenkia; Yakutia; Magadan region.

e. *Parasemia plantaginis floccosa* (GRAESER, 1888)

Berl. Entomol. Ztg. 32: 115 (*Nemeophila plantaginis*).

Middle Amur; Primorye. North-East China; Korea.

f. *Parasemia plantaginis sachalinensis* MATSUMURA, 1927

J. Coll. Agr. Hokk. Imp. Univ. 19:59.

Sakhalin.

g. *Parasemia plantaginis araitensis* MATSUMURA, 1929

Ins. mats. 3: 168 (*Parasemia plantaginis* f. *araitensis*).

(= *Parasemia plantaginis paramushira* BRYK, 1942, Dtsch. ent. Zeit. IRIS 56:30; = *Parasemia plantaginis kamtschadalus* BRYK, 1942, Dtsch. ent. Zeit. IRIS 56:29, nomen nudum).

Kamchatka; Northern Kuril Is. (Alaid (= Araito or Atlasov) and Paramushir Is.).

h. *Parasemia plantaginis kunashirica* BRYK, 1942

Dtsch. Ent. Zeit. IRIS 56:29, t. 1, f. 11.

Southern Kuril Is. (from Urup to Kunashir).

Hyphoraia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 182.

23. *Hyphoraia aulica* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine; Moldova; European Russia, north to South Karelia and Vyatka; Caucasus; Transcaucasia (Georgia); North Kazakhstan; South Siberia, north to Tyumen, Omsk, Novosibirsk, Minussinsk, the middle part of the Angara river; Transbaikalia; Amur basin; Primorye. Central Europe, west to South Belge and East France, south to Austria, Pannonia and the North Balkans; South Scandinavia; northern part of Asia Minor; ?Mongolia, China (Xinjiang, Heilongjiang, Liaonin); Korea; Japan (Hokkaido).

a. *Hyphoraia aulica aulica* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:505.

(= *Hyphoraia aulica rishiriensis* MATSUMURA, 1927, J. Coll. Agr. Hokk. Imp. Univ. 19:59, t. 4, f. 3).

Baltic Provinces; Belarus; Ukraine; Moldova; European Russia; Siberia; the south of the Far East. West Europe; ?Mongolia, China; Korea; Japan.

b. *Hyphoraia aulica testudinariooides* (SOVINSKY, 1905)

Rus. Entomol. Obozr. 5(3–4): 109–110 (*Arctia aulica*).

Caucasus; Transcaucasia (Georgia). ?Turkey.

Pararctia SOTAVALTA, 1965

Suom. hyont. Aikak. [Acta Ent. Fenn.] 31:173.

24. *Pararctia lapponica* (THUNBERG, 1791)

European Russia, polar regions only: Kola peninsula, Mesen, Kanin peninsula, Pechora river (Ust-Tsilia), Polar Urals; Yamal peninsula; Taimir, south to Igarka; East Sayan Mts.; mountains of

Transbaikalia, Stanavoi Mts., Yakutia; Magadan region; Chukotka; Wrangel Is.; Kamchatka. Polar Scandinavia; North America (Alaska; North-West Canada, east to Quebec).

a. *Pararctia lapponica lapponica* (THUNBERG, 1791)

Diss. ent. sistens Insecta Suecica 2: 40, f. 7

(= *Bombyx festiva* BORKHAUSEN, 1790, Eur. Schmett. 3: 191; = *Bombyx avia* HÜBNER, 1804, Samml. europ. Schmett. 2: 230, 247; = *Hyphoraia festiva rosea* SHELJUZHKO, 1929, Ent. Anz. 9: 424).

Entire polar Eurasia; Yakutia south to the Verkhoyanskii Mts.; Kamchatka. ?Mountains of East Sayan and Transbaikalia.

b. *Pararctia lapponica lemniscata* (STICHEL, 1911)

Berl. Ent. Ztg. 56: 99 (*Arctia festiva*).

The systematic position of the East Sayan specimens needs revision.

Mountains of East Yakutia (Suntar-Khayata and Cherskii Mts.), Vilui river.

25. *Pararctia tundrana* TSHISTJAKOV, 1990

Novosti sistematiki nasekomykh Dalnego Vostoka. Vladivostok: 97–98, f. 1–3.

Polar Ural; Yamal; Gydan; Taimir; Polar Yakutia; ?mountains of South Yakutia (Stanovoi Mts.: Tokko); Chukotka; northern part of Koryakia.

Note: The species was formerly considered as *Pararctia subnebulosa* (DYAR, 1899), Ent. News Philad. 10: 130 (*Hyphoraia*), which inhabits Polar America: Alaska and the Yukon territory. These species differ in male genitalia structures (TSHISTJAKOV, 1990).

Borearctia DUBATOLOV, 1984

Entomol. Obozr. 63: 337–339.

26. *Boreoarctia menetriesii* (EVERSMANN, 1846)

Bull. Soc. Imp. Nat. Moscou 19(3): 84 (*Euprepia*).

Karelia; lower course of the Ob river (Oktyabrskoe); North-East Kazakhstan ("Songoria"); Altai and Sayan Mts.; Evenkia; Yakutia; Middle Amur; Primorye (Sikhote-Alin Mts.); Central Sakhalin: Poronai river (HORI, 1926). Finland. ?China (I think that "*Callimorpha principalis*" of FANG CHENG LAI (1984) from Heilongjiang may belong to this species).

Acerbia SOTAVALTA, 1963

Suom. hyont. Aikak. 29: 263.

27. *Acerbia alpina* (QUENSEL, 1802)

In: ACERBI, Travels through Sweden, Finland and Lapland to the North Cape, 1798 & 1799, 2: 253. (= *Arctia thulea* DALMAN, 1823, Anlecta Entom. Holmiae: 92; = *Hyphoraia alpina sibirica* O. BANG-HAAS, 1927, Horae Macrolepid. 1: 70, t. 9, f. 2).

Russia: Polar Urals; southern part of Yamal peninsula; Taimir peninsula; Wrangel Is.; Altai, Sayan, and Stanovoi Mts.; mountains of East Yakutia. North-East Kazakhstan: Altai Mts. Mountains of polar Scandinavia (Torne Lappmark in Sweden and Lapponia enontekiensis in Finland); North Mongolia (Chovsgol region); Alaska; North-West Canada.

28. *Acerbia seitzi* (A. BANG-HAAS, 1910)

Dtsch. Ent. Zeit. IRIS **24**: 30, t. 3, f. 17 (*Arctia*).

(= *Arctia strandi* NIEPELT, 1911, Int. Ent. Ztg. **5**: 274; = *Phragmatobia niepeltiana* STRAND, 1919, Lep. Cat. **22**: 416; = *Hyporaia seitzi khumbeli* O. BANG-HAAS, 1927, Horae Macrolep. **1**: 70, t. 9, f. 4).

Tien Shan: Zailiiskii (Trans-Ili) Alatau Mts. (Kazakhstan); Kungei Alatau Mts., Naryn (Kyrgyzstan); Chimgan Mt. (prope Tashkent in Uzbekistan).

Platarctia PACKARD, 1864

Proc. ent. Soc. Philad. **3**: 109.

29. *Platarctia atropurpurea* (O. BANG-HAAS, 1927)

The middle course of the Yenisei river; Altai and Sayan Mts.; Tuva; Transbaikalia; Yakutia (except for the polar part); the southern part of the Magadan region. Mongolia.

a. *Platarctia atropurpurea atropurpurea* (O. BANG-HAAS, 1927)

Horae Macrolepид. **1**: 117 (*Hyporaia ornata*).

(= *Arctia ornata* STAUDINGER, 1896, Dtsch. Ent. Zeit. IRIS **9**: 188, t. 4, f. 1, 2).

The middle part of the Yenisei river; mountains of South Siberia; South Yakutia. Mongolia.

b. *Platarctia atropurpurea sotavaltae* DUBATOLOV, 1996 subspec. nov. (see p. 70)

Mountains of East Yakutia; the southern part of the Magadan region.

Stauropolia SKALSKI, 1988

Vestnik zoologii (4): 22.

† *S. nekrutenkoi* SKALSKI, 1988

Vestnik zoologii (4): 22.

Russia: Stavropol region (Miocene).

Oroncus SEITZ, 1910

Gross-Schmett. Erde **2**: 82.

30. *Oroncus tancrei* (STAUDINGER, 1887)

Stett. Ent. Ztg. **48**: 81 (*Arctia*).

(= *Phragmatobia urania* PÜNGELER, 1904, Soc. Ent. **19**: 121).

Kyrgyzstan: Inner and Central Tien Shan (Naryn, Sarydzhab). China: Kinjiang (East Tien Shan: a mountain area north of Korla).

31. *Oroncus fasciata* O. BANG-HAAS, 1927

Horae Macrolep. **1**: 61, t. 8, f. 17 (*Oroncus tancrei*).

East Kazakhstan (Dzhungarian Alatau Mts.).

32. *Oroncus alaica* O. BANG-HAAS, 1927

Horae Macrolep. **1**: 61, t. 8, f. 19, 20 (*Oroncus tancrei*).

The Turkestan mountain range (the border between Uzbekistan and Tajikistan); Alai (Kyrgyzstan); Trans-Alai (Zaalaiskii Mts.), Pamirs (Tajikistan).

Gonnerda MOORE, 1879

Proc. zool. Soc. Lond.: 395.

Gonnerda perornata MOORE, 1879

Proc. zool. Soc. Lond.: 395, t. 32, f. 2.

India (Kashmir). In the collection of the Zoological Institute (St.-Petersburg) is a male of this species labelled "Przhevalsk" (now Karakol in Kyrgyzstan); I consider this label to be a mistake.

Arctia SCHRANK, 1802

Fauna Boica 2(2): 152.

(= *Hypercompe* HÜBNER, [1806], Tentamen determinationis digestionis [1] (rejected and invalid work); = *Eyprepia* OCHSENHEIMER, 1810, Schmett. Eur. 3: 299; = *Euprepia* HÜBNER, [1819] 1816, Verz. bekannter Schmett.: 181; = *Chelonia* GODART, [1823] 1822, Hist. nat. Lepid. Papillons Fr. 4: 299; = *Arctinia* EICHWALD, 1830, Zool. spec. 2: 195; = *Callarctia* PACKARD, 1864, Proc. ent. Soc. Philad. 3: 114).

33. *Arctia caja* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia, north to Arkhangelsk and Ukhta (the latter in the Komi republic); Caucasus; Transcaucasia; North Kazakhstan; mountains of eastern Middle Asia and East Kazakhstan; South Siberia, north to Khanty-Mansiisk, Ket river in the Tomsk region, Angara river, Central Yakutia and the Magadan region (upper course of the Kolyma river); Kamchatka; Amur basin; Primorye; Sakhalin; Southern Kuril Is. (from Iturup to Kunashir). West Europe (except for the polar regions, the southern part of the Iberian peninsula and Greece); Asia Minor; north-western Iran; mountains of Afghanistan and Pakistan; Mongolia; China (Xinjiang, Henan, Nei Mongol; Hebei; Dunbei); Korea; Japan (Hokkaido, Honshu); North America (Canada; USA, including Alaska).

a. *Arctia caja caja* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1: 500 (*Phalaena*).

(= *Arctia caja confluens* REBEL, 1910, in: BERGE, Schmett. Buch (ed. 9): 430; = *Arctia caja rebeli* VNUKOVSKY, 1929, Zool. Anzeiger 83(9/10): 223).

Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia; Caucasus; North and East Kazakhstan; South Siberia, east to the Angara river. West Europe; North-West China (Xinjiang).

b. *Arctia caja ossetica* DUBATOLOV, 1996 subspec. nov. (see p. 71)

Western Caucasus.

c. *Arctia caja wiskotti* STAUDINGER, 1879

Horae Soc. Ent. Ross. 14: 333.

Transcaucasia. Asia Minor.

d. *Arctia caja tshimgana* SHELUZHKO, 1935

Mitt. Münch. ent. Ges. 25: 31, t. 3, f. 7.

Uzbekistan: West Tien Shan. Possibly also in Kazakhstan and Kyrgyzstan.

e. *Arctia caja pamiroalaica* STSHESTKIN, 1982

Izv. Akad. Nauk Tadz. SSR, Otd. biol. nauk (1): 39–43.

Alai-Pamirs in Tajikistan and, possible, in South Kyrgyzstan. Afghanistan.

f. *Arctia caja sajana* O. BANG-HAAS, 1927

Horae Macrolepid. 1: 72, t. 9, f. 6.

East Sayan Mts.; Baikal; Transbaikalia; Yakutia (excluding the polar regions); Magadan region. Mongolia.

g. *Arctia caja kamtschadalis* DRAUDT, 1931

In: SEITZ, Macrolepid. World, Suppl. 2: 87, f. 7f.

Kamchatka.

h. *Arctia caja phaeosoma* (BUTLER, 1877)

Ann. Mag. Nat. Hist. (4)20: 395 (*Euprepia*).

Amur basin; Primorye; Sakhalin; Southern Kuril Is. (from Iturup to Kunashir). East China; Korea; Japan.

34. *Arctia olschwangi* DUBATOLOV, 1990

Taksonomiya nasekomykh i gelmintov. Novosibirsk: 89–93, f. 1a, 2 l.

Polar Ural; Yamal peninsula; Yakutia: Lena river delta.

35. *Arctia flavia* (FUESSLY, 1779)

Mag. f. Liebhaber d. Ent. 2: 70, t. 1, f. 11 (*Phalaena (Bombyx)*).

(= *Arctia flavia campestris* GRAESER, 1892, Berl. Ent. Ztg. 37: 212; = *A. f. uralensis* HEYNE, 1899, Soc. Ent. 14: 98; = *A. f. sibirica* HEYNE, 1899, Soc. Ent. 14: 98; = *A. f. lederi* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 75, t. 9, f. 12; = *A. f. baicalensis* O. BANG-HAAS, 1927, Horae Macrolepid. 1: 74, t. 9, f. 11).

North Ukraine², including the Carpathians; European Russia east of Tver, north to Syktyvkar (Komi republic), south to the Saratov region and Orenburg; North Kazakhstan; South Siberia, north to Khanty-Mansiisk, Tomsk, and the Podkamennaya Tunguska river; Central Yakutia; Magadan region (Upper Kolyma); Amur basin; Shantar Is.; Primorye. West Europe (Alps, Balkan peninsula: Rila Mts.); Mongolia; China (Xinjiang, Nei Mongol, Hebei).

36. *Arctia rueckbeili* PÜNGELER, 1901

Dtsch. Ent. Zeit. IRIS 14: 190, t. 3, f. 11.

Central Tien Shan in Kyrgyzstan (Sarydzhaz river); Turkestan Mts. at the border between Uzbekistan and Tajikistan; Alai in Uzbekistan and Kyrgyzstan; Alai valley in Kyrgyzstan (DUBATOLOV, 1987b). China (Xinjiang).

37. *Arctia intercalaris* (EVERSMANN, 1843)

East Kazakhstan (Dzhungarian Alatau Mts.; Trans-Ili (Zailiiskii Alatau) Mts.); Tien Shan and Alai-Pamirs in South Kazakhstan, Uzbekistan, Kyrgyzstan, and Tajikistan (excluding the south-western region). Mountains of Afghanistan and North-West Pakistan; India (from Kashmir to Kulu).

2 Material: 1 ♂, Prov. Kijev, Teterjev, 1914, A. HUTSHEK leg. (SHELJUZHKO coll.); 1 ♀, the Carpathians, Krasnogorsk, 14.VII.1984, TIKHOMIROV leg. (KAABAK coll.). Both localities need confirmation.

a. *Arctia intercalaris intercalaris* (Eversmann, 1843)

Bull. Soc. Imp. Nat. Moscou **16** (3): 544, t. 10, f. 1a, b (*Euprepia*).

(= *Arctia intercalaris boettcheri* O. Bang-Haas, 1927, Horae Macrolepid. **1**: 72).

Mountains of East Kazakhstan (south from Lake Zaisan); Tien Shan in South Kazakhstan, Kyrgyzstan, and North-East Uzbekistan.

b. *Arctia intercalaris alpherakyi* STAUDINGER, 1886

Ent. Month. Mag. **22**: 258.

(= *Arctia intercalaris badakhshana* WILTSHERE, 1961, Beitr. naturk. Forsch. SW Deutschl. **19** (3): 340).

Alai-Pamirs Mts. in South-Eastern Uzbekistan, South Kyrgyzstan, and Tajikistan (excluding the south-west region). Afghanistan (Badakhsban).

Arctia ladakensis (O. BANG-HAAS, 1927)

Horae Macrolep. **1**: 116 (*Oroncus*).

I have seen three female specimens of this species from the Aksu river in Central Tien Shan (China: Xinjiang) (VI.1912, RÜCKBEIL leg.), in the collection of the Zoological Institute (St.-Petersburg). This locality is situated near the border to Kyrgyzstan. They are all similar to the type specimen of this species which is kept in the collection of the Zoological Museum of the Humboldt University (Berlin). These specimens have simple, not dentate antennae as in *Arctia intercalaris* (Ev.).

Epicallia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 182.

38. *Epicallia villica* (LINNAEUS, 1758)

South Baltic Provinces (Lithuania and Latvia); Belarus; Ukraine, Crimea; Moldova; European Russia north to St.-Petersburg and Vyatka; Caucasus; Transcaucasia; West Siberia (Kurgan). Central and South Europe, north to South England; North-West Africa; Asia Minor; the Near East; North-West and North Iran, east of the Shahkuh Mts..

a. *Epicallia villica villica* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1: 501 (*Phalaena*).

South Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia; West Siberia; Caucasus; Transcaucasia (excluding the Talysh Mts.). Central Europe.

b. *Epicallia villica confluens* (ROMANOFF, 1884)

Mem. lepidop. Ed. N. N. ROMANOFF 1: 87, t. 4, f. 9 (*Arctia villica*).

South-East Transcaucasia (Azerbaijan: Talysh Mts.). North Iran.

Eucharia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 181.

(= *Ammobiota* WALLENGREN, 1885, Skand. Heterocer-Fjarilar **2**: 304).

39. *Eucharia festiva* (HUFNAGEL, 1766)

Baltic Provinces (South Lithuania, Latvia); Belarus; Ukraine, Crimea; Moldova; European Russia, north to Kaluga, Sormovo 30 km N of Nizhnii Novgorod (TSCHETVERIKOV, 1993), and the lower

course of the Kama river; Caucasus; Transcaucasia; Kazakhstan; the eastern part of Middle Asia (Uzbekistan: Tashkent, Alai); West Siberia, north to Kurgan, Omsk, Novosibirsk; West Altai; Tuva; South Transbaikalia. Central and South Europe (except for the southern part of the Iberian peninsula, West France, the North Sea and Baltic coasts, the Alps, the north-western and southern parts of the Balkans); Asia Minor; Mongolia; China (Xinjiang, Nei Mongol; Hebei).

a. *Eucharia festiva festiva* (HUFNAGEL, 1766)

Berl. Mag. 2: 416, 437 (*Phalaena*).

(= *Phalaena hebe* LINNAEUS, 1767, Syst. Nat. (Edn. 12) 1 (2): 820; = *Chelonia interrogationis* MÉNÉTRIÉS, 1863, Cat. lep. Pet. 3: 147, t. 15, f. 1; = *Arctia hebe sartha* STAUDINGER, 1886, Stett. Ent. Ztg. 47: 401; = *Arctia festiva nivea* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 75, t. 9, f. 14, 15).

South Baltic Provinces; Belarus; Ukraine, Crimea; Moldova; European Russia; Caucasus; Transcaucasia; North Kazakhstan; West Siberia. Central and South Europe; Asia Minor; West China.

Note: The type locality of *Chelonia interrogationis* MÉN. is Kolyvan in Altaiskii Krai (West Altai). I have seen the type specimen in the collection of the Zoological Institute (St.-Petersburg) as well as other specimens from this region. They are all similar to *E. festiva festiva* (HFN.) but differ from Transbaikalian, Tuvanian, Mongolian and North Chinese specimens which I consider as *E. festiva collaris* (GR.-GR.).

b. *Eucharia festiva iliensis* (WAGNER, 1913)

Int. Ent. Zeit. 7(1): 4 (*Arctia hebe*).

(= *Arctia festiva philippi* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 76, t. 9, f. 20; = *A. f. interposita* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 76, t. 9, f. 21, 22).

South Kazakhstan; Uzbekistan (Tashkent and the Alai Mts.).

c. *Eucharia festiva collaris* (GRUM-GRSHIMAILO, 1899)

Ezegod. Zool. Muz. Imp. Akad. Nauk 4: 462 (*Arctia hebe*).

(= *Chelonia culoti* OBERTHÜR, 1912, Et. lep. comp. 6: 322–323, 116, f. 10–28).

Tuva; South Transbaikalia. Mongolia; China (Nei Mongol, Hebei).

Pericallia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 182.

(= *Pleretes* LEDERER, 1853, Verh. zool.-bot. Ver. Wien 2 (Abh.): 77).

40. *Pericallia matronula* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1: 509 (*Phalaena*).

(= *Phalaena matrona* HÜBNER, [1803]; Eur. Schmett. 2: f. 238, 239; = *Pericallia matronula central-asiae* O. SCHULTZ, 1905, Ent. Zeit. 18(31); = *Pleretes matronula agassizi* SPULER, 1906, Schmett. Eur. 2: 140; = *Pericallia matronula amurensis* SHELJUZHKO, 1926, Dtsch. Ent. Zeit. IRIS 40: 56; = *P. m. sachalinensis* DRAUDT, 1931, in: SEITZ, Macrolep. World, Suppl. 2: 84, f. 7).

Baltic Provinces; Belarus; North and Central Ukraine (including the Carpathians); Moldova; European Russia, north to St.-Petersburg, the Upper Volga, and the lower Kama rivers, south to Saratov; North Kazakhstan; South Siberia, north to Tobolsk (SITNIKOV, 1992), the Chulym river (Tomsk region), the middle course of the Angara river, Irkutsk, and Transbaikalia; Middle Amur; Primorye; Southern Sakhalin; Southern Kuril Is. (Kunashir). Central Europe (East France, South and Central Germany, the Alps, East Europe from Central Poland south to Hungary and Romania; in the Carpathians); North Mongolia; China (Dunbei, Hebei), Korea, Japan (Hokkaido, Honshu).

Micractiini

Divarctia DUBATOLOV, 1990

Regkie gelminty, klestshi i nasekomye. Novosibirsk: 84.

41. *Divarctia diva* (STAUDINGER, 1887)

Stett. Ent. Ztg. **48**: 84 (*Ocnogyna*).

(= *Arctia haberhaueri* ALPHERAKY, 1888, Stett. Ent. Ztg. **49**: 67).

Mountains of Middle Asia: South-West Kyrgyzstan (Ferghanian mountain range); Uzbekistan (the West Tien Shan: Chatkal, Pskem, and Karzhantau Mts.; Alai-Pamirs: Turkestan, Zeravshan, and Hissar Mts.); Tajikistan (Turkestan, ?Hissar, Vakhsh, Peter-the-Great, and Khozratishokh mountain ranges).

Tancrea PÜNGELER, 1898

Soc. Ent. **13**: 57.

42. *Tancrea pardalina* PÜNGELER, 1898

Soc. Ent. **13**: 57.

Kazakhstan: deserts of the Ili river basin; Turkmenistan: Kara-kum desert (DUBATOLOV, 1991b). China (Xinjiang: Ili and Kungesse rivers).

Holoarctia FERGUSON, 1984

Proc. Entomol. Soc. Wash. **86** (2): 452–454.

43. *Holoarctia cervini* (FALLOU, 1864)

Ann. Soc. Ent. France **4**(4): 23, t. 1, f. 2 (*Nemeophila*).

Khibiny Mts. (Kola peninsula); South Urals, Altai, East Sayan, Khamar-Daban, and Stanovoi Mts.; mountains of East Yakutia (Verchoyanskii mountain range); Kamchatka (Ichinskaya Sopka Mt.); Chukotka. West Europe: the Alps; mountains of polar Scandinavia (Torne Lappmark in Sweden); North-West Mongolia; North Korea (KÔDA, 1988: as “*Hyperborea czekanowskii*”); Alaska. Canada: Alberta (ssp. *sordida* McDUNNOUGH, 1921, Canad. entomol. **53**: 167).

a. *Holoarctia cervini fridolini* (TORSTENIUS, 1971)

Entomol. Ts. Arg. **92** (3–4): 173–177, f. 1, 2 (*Orodemnias cervini*).

Khibiny Mts.; South Urals; Verkhoyanskii Mts.; Chukotka; ?Kamchatka. Polar Scandinavia; Alaska.

b. *Holoarctia cervini perunovi* DUBATOLOV, 1990

Chlenistonogie i gelminty. Novosibirsk: 152.

Altai. North-West Mongolia.

c. *Holoarctia cervini puengeleri* (O. BANG-HAAS, 1927)

Horae Macrolepidop. **1**: 60, t. 8, f. 14 (*Orodemnias*).

East Sayan, Khamar-Daban, Stanovoi Mts. (there is a female in the V. A. GANSON (Moscow) collection, which is labelled “Okhotskoe Sea”).

44. *Holoarctia marinae* DUBATOLOV, 1985 (January)

Chlenistonogie Sibiri i Dalnego Vostoka. Novosibirsk: 152, fig. 2v, 3b; 1985 (December), Ann. Ent. Fenn. 51 (2): 57 (the holotype below, nec above!).
Altai (Terektskii and Kuraiskii Ranges).

Palearctia FERGUSON, 1984

Proc. Entomol. Soc. Wash. 86 (2): 454–456.

45. *Palearctia mira* DUBATOLOV & TSHISTJAKOV, 1989

Zool. Zhurnal 68 (11): 141, f. 1a, b, 2.
Russia: the south-eastern part of the Altai (Kuraiskii Range).

46. *Palearctia glaphyra* (EVERSMANN, 1843)

Mountains of East Kazakhstan: "Lake Zaisan", Dzhungarian Alatau, Trans-Ili (Zailiiskii) Alatau mountain ranges; Kyrgyzstan: Kirghiz Alatau, Kungei Alatau, and Terskei Alatau Mts.; Central Tien Shan, west to Naryn and the Dolon pass. China (Xinjiang: East Tien Shan, Kuruktag Mts., ?Khotan).

a. *Palearctia glaphyra glaphyra* (EVERSMANN, 1843)

Bull. Soc. Imp. Nat. Moscou 16 (3): 544, t. 10, f. 4 (*Euprepia*).

East Kazakhstan, from Lake Zaisan to the Dzhungarian Alatau, including the Tyshkantau.

b. *Palearctia glaphyra dublitzkyi* (O. BANG-HAAS, 1927)

Horae Macrolepidop. 1: 63, t. 8, f. 29, 30 (*Micrarcacia glaphyra*).

Kazakhstan: Trans-Ili (Zailiiskii) Alatau Mts.; Kyrgyzstan: Kungei Alatau, ?Kirghizskii Alatau mountain ranges.

c. *Palearctia glaphyra aksuensis* (O. BANG-HAAS, 1927)

Horae Macrolepidop. 1: 63, t. 8, f. 26, 27 (*Micrarcacia glaphyra*).

(= *Micrarcacia glaphyra naryna* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 63, t. 8, f. 28).

Kyrgyzstan: Central Tien Shan, south from the Terskei Alatau mountain range and east from Naryn. China (Xinjiang: Aksu region).

47. *Palearctia wagneri* (PÜNGELER, 1918)

Z. Öster. Ent. Ver. 3: 46, textfig. (*Arctia*).

Kyrgyzstan: Central Tien Shan (Sarydzhaz region).

48. *Palearctia golbecki* DUBATOLOV, 1996

Neue Ent. Nachr. 37: 19.

Kyrgyzstan: Kirghiz Alatau mountain range.

49. *Palearctia gratiosa* (GRUM-GRSHIMAILO, 1890)

Uzbekistan: West Tien Shan (Chatkal mountain range), Alai-Pamirs (Turkestan and Zeravshan mountain ranges); Kyrgyzstan: Chatkal, Kirghiz and Terskei Alatau, Alai Mts.; Tajikistan: Turkestan, Hissar, Peter-the-Great, and Trans-Alai mountain ranges, Pamirs. Afghanistan (Karakorum); India (Kashmir). West China (fig. 1565 with "*Micrarcacia glaphyra* (EVERSMANN)" of FANG, 1982).

a. *Palearctia gratiosa caroli* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 24.

Uzbekistan: Chatkal Mts.; Kyrgyzstan: Chatkal and Kirghizskii Alatau Mts.

b. *Palearctia gratiosa sergei* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 24.

Kyrgyzstan: south-eastern part of the Terskei Alatau mountain range.

c. *Palearctia gratiosa gratiosa* (GRUM-GRSHIMAILO, 1890)

Mem. lepidop. Ed. N. N. ROMANOFF **4**: 533, t. 19, f. 5 (*Arctia glaphyra*).

Kyrgyzstan: Alai; Tajikistan: Alai and Peter-the-Great mountain ranges, North-West Pamirs.

d. *Palearctia gratiosa flava* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 25.

Uzbekistan: Zeravshan and Turkestan mountain ranges; Tajikistan: Zeravshan, Turkestan, and Hissar mountain ranges.

e. *Palearctia gratiosa rupicola* (GRUM-GRSHIMAILO, 1890)

Mem. lepidop. Ed. N. M. ROMANOFF **4**: 535, t. 19, f. 6 (*Arctia*).

(= *Arctia glauca* STAUDINGER, 1892, Dtsch. Ent. Zeit. IRIS **4**: 251, t. 3, f. 6).

Kyrgyzstan: the eastern part of the Trans-Alai; Tajikistan: the eastern part of the Transalai mountain range, East and South-West Pamirs.

50. *Palearctia erschoffii* (ALPHERAKY, 1882)

Uzbekistan?: Chatkal Mts.; Kyrgyzstan: Chatkal, Sussamyr, Kirghizskii Alatau, Kungei Alatau, and Terskei Alatau mountain ranges, Inner Tien Shan (Naryn), Central Tien Shan (Sarydzhab); Kazakhstan: Trans-Ili (Zailijskii) Alatau, ?Kungei Alatau mountain ranges. China (Xinjiang): East Tien Shan (Julduz).

a. *Palearctia erschoffii erschoffii* (ALPHERAKY, 1882)

Horae Soc. Ent. Ross. **17**: 29, t. 1, f. 33 (*Arctia*).

(= *Arctia erschoffi issyka* STAUDINGER, 1887, Stett. Ent. Ztg. **48**: 82).

Entire Tien Shan within Uzbekistan, Kyrgyzstan, Kazakhstan, and China, except for the Naryn region in Kyrgyzstan.

b. *Palearctia erschoffii selmonsi* (BÖTTCHER, 1905)

Ent. Zeit. **19**: 62, f. 18–21 (*Arctia erschoffi*).

Kyrgyzstan: Inner Tien Shan (Naryn region).

51. *Palearctia ferghana* (STAUDINGER, 1887)

Kyrgyzstan: Alai, Sussamyr Mts.; Tajikistan: Turkestan mountain range, ?Pamirs. Tien Shan (an unknown locality). Afghanistan (Anjuman pass), ?Nepal.

a. *Palearctia ferghana schottlaenderi* (STRAND, 1912)

Intern. Ent. Zeit. (Guben) **6**: 2 (*Arctia*).

Tien Shan (unknown locality).

b. *Palearctia ferghana sussamyla* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 30.

Kyrgyzstan: Sussamyr Mts.

c. *Palearctia ferghana ferghana* (STAUDINGER, 1887)

Stett. Ent. Ztg. **48**: 82 (*Arctia erschoffi*).

Kyrgyzstan: Alai Mts.

d. *Palearctia ferghana turkestanica* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 30.

Tajikistan: Turkestan mountain range.

? *Palearctia ferghana variabilis* (DANIEL, 1966)

Mitt. Münch. Ent. Ges. **56** (7): 161, t. 3, f. 4–16 (*Micrarctia*).

It can be found in the Pamirs within Tajikistan.

52. *Palearctia gracilis* DUBATOLOV, 1996

Neue Ent. Nachr. **37**: 31.

Uzbekistan and Kyrgyzstan: West Tien Shan (Chatkal Mts.).

Centrarctia DUBATOLOV, 1990

Chlenistonogie i gelminty. Novosibirsk: 157

53. *Centrarctia mongolica* (ALPHERAKY, 1888)

Stett. Ent. Ztg. **49**: 67 (*Arctia*).

Southern Tuva (Ubsu-nur hollow). Mongolia; China (Nei Mongol).

Sibirarctia DUBATOLOV, 1987

Nasekomye, klestshi i gelminty. Novosibirsk: 36–40.

54. *Sibirarctia kindermannii* (STAUDINGER, 1867)

South Urals; Omsk; Novosibirsk; Krasnoyarsk; Altai; Khakasia; the southern banks of Lake Baikal; Transbaikalia; Middle Amur; South Primorye. Mongolia; China (Qinghai, Shanxi, Heilongjiang; Liaonin).

a. *Sibirarctia kindermannii kindermannii* (STAUDINGER, 1867)

Stett. Ent. Ztg. **28**: 102–103 (*Arctia*).

South Urals; South-West Siberia east to the Yenisei river.

b. *Sibirarctia kindermannii pomona* (STAUDINGER, 1897)

Dtsch. Ent. Zeit. IRIS **10**: 326, t. 9, f. 23 (*Arctia kindermannii*).

(= *Micrarctia kindermannii roseni* O. BANG-HAAS, 1927, Horae Macrolepidop. **1**: 65, t. 8, f. 40).

The southern banks of Lake Baikal; Transbaikalia; Upper Amur. Mongolia.

c. *Sibirarctia kindermannii pretiosa* (STAUDINGER, 1887)

Mem. lepidop. Ed. N. M. ROMANOFF **1**: 65, t. 8, f. 40 (*Arctia kindermannii*).

(= *Micrarctia kindermannii ussuriensis* O. BANG-HAAS, 1927, Horae Macrolepidop. **1**: 65, t. 8, f. 42).

The Middle Amur; Primorye. China (Heilongjiang).

55. *Sibirarctia buraeatica* (O. BANG-HAAS, 1927)

The south-eastern part of the Altai; Tuva; East Sayan Mts.; Baikal; Transbaikalia; Central Yakutia; mountains of East Yakutia. North and Central Mongolia.

a. *Sibirarctia buraeatica buraeatica* (O. BANG-HAAS, 1927)

Horae Macrolepidop. 1: 64, t. 8, f. 35, 36 (*Micrarczia*).

(= *Micrarczia buraeatica elwesi* O. BANG-HAAS, 1927, Horae Macrolepidop. 1: 64, t. 8, f. 38).

The south-eastern part of the Altai; Tuva; Baikal; Transbaikalia; Central Yakutia. Mongolia.

b. *Sibirarctia buraeatica validus* (O. BANG-HAAS, 1927)

Horae Macrolepidop. 1: 64, t. 8, f. 37 (*Micrarczia buraeatica*).

South-western Transbaikalia (Kyakhta district in Buryatia).

c. *Sibirarctia buraeatica chajataensis* DUBATOLOV, 1996 subspec. nov. (see p. 71)

Mountains of East Yakutia (Suntar-Khayata mountain range, Oimyakon region).

Chelis RAMBUR, 1866

Cat. syst. Lepid. Andalousie (2): 256.

(= *Cletis* KIRBY, 1892, Synonymic Cat. Lepid. Heteroc. 1: 263).

56. *Chelis maculosa* (GERNING, 1780)

Frankf. Beitrag. Wiss. 2: 862, t. 2, f. 1–3 (*Phalaena*).

Ukraine; Crimea; European Russia: Don and Volga rivers, north to the lower course of the Kama river; South Urals; North Kazakhstan; South-West Siberia (Karasuk). West Europe: Central Spain, Central and Southern France, Central and Southern Germany, Austria, Hungary; South Poland; the northern and south-eastern parts of the Balkans (Macedonia, Bulgaria, West Turkey); China (Xinjiang).

a. *Chelis maculosa mannerheimi* (DUPONCHEL, 1836)

Hist. nat. Lep. France, Suppl. 3: 49, t. 4, f. 2 (*Chelonia*).

Ukraine; Crimea; the southern part of European Russia; North Kazakhstan; South-West Siberia. China (Xinjiang).

57. *Chelis caecilia* (KINDERMANN in LEDERER, 1853)

Verh. zool.-bot. Ges. Wien 3: 364 (*Arctia maculosa*).

(= *Cletis maculosa insularia* W. KOSHANTSCHIKOV, 1924, Ezhegod. Gos. Muz. im. Martjanova 2(1): 69, syn. nov., according to a good redescription of "Arctia maculosa" var. *caecilia* KINDERM. in litt." in: STAUDINGER, 1867, Stett. Ent. Ztg. 28: 105–106).

North-East Kazakhstan: Altai Mts.; Russia: Altai Mts.; Khakasia, West Buryatia. Mongolia.

58. *Chelis reticulata* (CHRISTOPH, 1887)

Caucasus (Teberda; South Daghestan); Transcaucasia; Turkmenistan (Kopetdagh Mts.). Asia Minor; Syria; North Lebanon; North Iran.

a. *Chelis reticulata reticulata* (CHRISTOPH, 1887)

Stett. Ent. Ztg. 48: 163 (*Arctia maculosa*).

Turkmenistan (Kopetdagh Mts.).

b. *Chelis reticulata transcaucasica* DUBATOLOV, 1988

Taksonomiya zhivotnykh Sibiri. Novosibirsk: 92, f. 2d.

Caucasus; Transcaucasia.

59. *Chelis ferghana* DUBATOLOV, 1988

Taksonomiya zhivotnykh Sibiri. Novosibirsk: 92, f. 1n, 2e.

West Tien Shan: Uzbekistan (Chatkal, Mts., Chimgan Mt.); Kazakhstan (Talas Alatau Mts. in the Aksu-Dzhebagly Nature Reserve); Kyrgyzstan: (Chatkal Mts., east of Toktogul).

60. *Chelis thianshana* DUBATOLOV, 1988

Taksonomiya zhivotnykh Sibiri. Novosibirsk: 93, f. 1k, 1l, 2g.

Kyrgyzstan: Kirghiz Alatau, Sussamyr, and Terskei Alatau mountain ranges, Inner Tien Shan; Kazakhstan: Trans-Ili (Zailijskii) Alatau, Dzhungarian Alatau, and Tarbagatai mountain ranges. China (Xinjiang: East Tien Shan).

61. *Chelis dahurica* (BOISDUVAL, 1832)

Icones hist. Lep. 2: 126, t. 60, f. 1 (*Chelonia*).

(= *Arctia gruneri* STAUDINGER, 1867, Stett. Ent. Ztg. 28: 104; = *Arctia maculosa sojota* TSCHETVERIKOV, 1904, Rus. Entomol. Obozr. 4 (2–3): 79).

South Urals; South-West Siberia (Kurgan and Omsk regions; Novosibirsk region: Lake Chany; Barnaul); North-East Kazakhstan (Altai and Saur Mts.); mountains of South Siberia: Altai, Khakasia, Tuva, Sayan Mts., west and south of the Irkutsk region, and Transbaikalia; ?the south-western part of the Primorye territory (Pogranichnyi (Grodekovo)). North and Central Mongolia.

Note: *Phragmatobia maculosa* ab. 5. *strigulosa* (STGR. ined.), HAMPSHON, 1901, Cat. Lep. Phal. Br. Mus. 3: 236 was established with a poor diagnosis: "Fore wing with the black spots larger, the veins more clearly streaked" and without the designation of the type locality. So, I consider it impossible to use this name (DUBATOLOV, 1988).

Grammia RAMBUR, 1866

Cat. syst. Lepid. Andalousie (2): 261.

(= *Orodemnias* WALLENGREN, 1885, Skand. Heterocer-Fjarilar 2 (3): 315).

62. *Grammia quenseli* (PAYKULL, 1793)

Kola peninsula; Polar Urals; North-East Kazakhstan (Saur Mts.); Altai; Tuva; East Sayan Mts.; Transbaikalia; the middle part of the Amur basin; the northern part of the Khabarovsk region, Yakutia; Magadan region; Wrangel Is.; Kamchatka. The Alps; the Carpathians (Transylvanian Alps); Polar Scandinavia (Tromso, Finmark, Lule Lapmark, Torne Lapmark, Lapponia kemensis, Lapponia enontekiensis); North Mongolia (Chovsgol aimak); Japan (Hokkaido: Daisetsu Mt.); polar regions and mountains of North America: from Labrador throughout the North-West and Yukon territories to Alaska, south to Manitoba and Quebec.

a. *Grammia quenseli quenseli* (PAYKULL, 1793)

Skr. Naturh.-Selsk. Kiobenhavn 2 (2): 99, f. (Bombyx).

Kola peninsula; Polar Urals; Wrangel Is.; Saur Mts. (Kazakhstan). Polar Scandinavia.

b. *Grammia quenseli liturata* (MÉNÉTRIÈS, 1859)

Bull. Acad. ph.-mat. St. Pet. 17: 500 (*Chelonia quenseli*).

(= *Orodemnias quenseli daisetsuzana* MATSUMURA, 1927, Ins. matsumur. 1: 110, 113).

East Altai; Tuva; East Sayan Mts.; Transbaikalia; Middle Amur; the northern part of the Khabarovsk region (Khaikan river, the left tributary of the upper Uchur is the type locality of this subspecies); Yakutia; Magadan region; Kamchatka. North Mongolia; Japan (Hokkaido).

63. *Grammia philippiana* FERGUSON, 1985

Entomography. Ann. Rev. Biosystematics. Sacramento, **3**:229–230, fig. 53–54.
Wrangel Is.; Alaska.

a. *Grammia philippiana olga* DUBATOLOV, 1990, **comb. nov.**

Redkie gelminty, klestshi i nasekomye. Novosibirsk: 79, f. 1a, 2a (*G. olga*).
Wrangel Is.

64. *Grammia turbans* (CHRISTOPH, 1892)

Horae Soc. Ent. Ross. **26**:460 (*Arctia*).

Khakasia; East Sayan Mts.; the southern part of the Baikal region; Transbaikalia; the middle part of the Amur basin; Central Yakutia. Mongolia; Canada (Alberta, Saskatchewan, Manitoba, North-West territories).

Hyperborea GRUM-GRSHIMAILO, [1900] 1899

Ezhegod. zool. Muz. Imp. Acad. Nauk **4**:464.

65. *Hyperborea czekanowskii* GRUM-GRSHIMAILO, [1900] 1899

Ezhegod. zool. Muz. Imp. Acad. Nauk **4**:464.

Evenkia (Nizhnyaya Tunguska river); North Yakutia (Olenek and Yana rivers, Suntar-Khayata mountain range, Oimyakon region); Kolyma river; Koryakia; Chukotka; Wrangel Is.; Kamchatka; Stanovoi mountain range; the northern part of the Chita region (Udokan mountain range). Alaska.

Diacrisia HÜBNER, [1819] 1816

Verz. bekannter Schmett.: 169.

66. *Diacrisia sannio* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Karelia, Arkhangelsk region, and Ukhta (Komi republic); Caucasus; Transcaucasia (east to West Azerbaijan); North Kazakhstan; mountains of East Kazakhstan, North Uzbekistan and Kyrgyzstan (south to the Alai valley); South Siberia, north to Oktyabr'skoe (the lower course of the Ob river), Ket river (Tomsk region), and Angara river; Transbaikalia; Central Yakutia. West Europe, north to Middle Scandinavia (except for the south-western part of the Iberian peninsula, Greece, and West Turkey); North-East Turkey; West China; Mongolia.

a. *Diacrisia sannio sannio* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) **1**:506 (*Phalaena*).

(= *Phalaena russula* LINNAEUS, 1758, Syst. Nat. (Edn. 10) **1**:510; = *Phalaena vulpinaria* LINNAEUS, 1758, Syst. Nat. (Edn. 10) **1**:520; = *Nemeophila russula pallida* STAUDINGER, 1892, Dtsch. Ent. Zeit. IRIS **5**:347).

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia; Kazakhstan; Siberia. Western Europe; Mongolia.

b. *Diacrisia sannio caucasica* SCHAPOSNIKOFF, 1904

Ezhegod. Zool. Mus. Imp. Akad. Nauk **9**: 253.

(= *Diacrisia sannio caucasiana* STRAND, 1919, Lep. Catal. **22**: 416).

Caucasus; Transcaucasia. North-East Turkey.

c. *Diacrisia sannio mortua* (STAUDINGER, 1887)

Stett. Ent. Ztg. **48**: 78 (*Nemeophila russula*).

(= *Diacrisia sannio uniformis* A. BANG-HAAS, 1907, Dtsch. Ent. Zeit. IRIS **20**: 69; = *D. s. syrdarja* STRAND, 1919, Lep. Catal. **22**: 416).

Mountains of Middle Asia (within North Uzbekistan, Kyrgyzstan, and South Kazakhstan).

67 *Diacrisia irene* BUTLER, 1881

Trans. Entomol. Soc. Lond.: 6.

(= *Diacrisia russula amuri* STAUDINGER, 1892, Mem. lepidop. Ed. N. M. ROMANOFF **6**: 277; = *Diacrisia sannio rubrocentralis* BRYK, 1948, Ark. Zool. **41A**(1): 44; = *Diacrisia sannio rishiriensis* MATSUMURA, 1930, Ins. matsumur. **5**: 35, t. 1, f. 2).

Middle part of the Amur basin; Primorye, Sakhalin; Southern Kuril Is. (Kunashir). East China; Korea; Japan (Rishiri, Hokkaido, Honshu).

Rhypariooides BUTLER, 1877

Ann. Nag. nat. Hist. (4)**20**: 395.

68. *Rhypariooides nebulosa* BUTLER, 1877

Ann. Mag. nat. Hist. (4)**20**: 396.

(= *Rhypariooides simplicior* BUTLER, 1881, Trans. ent. Soc. Lond.: 6).

South Primorye; Southern Kuril Is. (Kunashir). China (Dunbei, Nei Mongol); ?Korea; Japan (Hokkaido, Honshu).

69. *Rhypariooides amurensis* (BREMER, 1861)

Bull. Acad. Imp. Sci. St. Petersb. **3**: 477 (*Chelonia rubescens*).

Middle part of the Amur basin; Primorye; Southern Kuril Is. (Kunashir). China (Dunbei, Hebei, Shanxi, Shaanxi, Zhejiang, Fujian, Jiangxi, Jiangsu, Hubei, Hunan, Guangxi, Sichuan, Yunan); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu).

70. *Rhypariooides metelkana* (LEDERER, 1861)

Wien. entomol. Monatschr. **5**(5): 162, t. 3, f. 12 (*Nemeophila*).

(= *Chelonia flava* BREMER, 1861, Bull. Acad. Imp. Sci. St. Petersb. **3**: 477).

South Ukraine (Kherson region); Russia: Azov sea (Taganrog); Daghestan (Derbent); Volga delta near Astrakhan (KÖNIG, 1985); South-West Siberia: Karasuk (DUBATOLOV, 1985c); the middle part of the Amur basin; Primorye; ?Shantar Is. Local in West Europe (South-West and North France, South Belgium, Germany: Berlin region, South Slovakia, Hungary, West Romania and the Danube delta); China (Heilongjiang, Jiangxi, Jiangsu, Nei Mongol); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Yaku, Okinawa).

Rhyparia HÜBNER, [1820] 1816

Verz. bekannter Schmett.: 183.

71. *Rhyparia purpurata* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Karelia, the Upper Volga, and Vyatka; Caucasus; Transcaucasia; North Kazakhstan; mountains of East Kazakhstan; North Kyrgyzstan, including the territory around Lake Issyk-kul; South Siberia, north to Khanty-Mansiisk, Tomsk, Krasnoyarsk, and the Angara river; Baikal area; Transbaikalia; Amur basin; Primorye. West Europe, north to Southern Scandinavia (except for the North Sea coast, North France, central and southern parts of the Iberian peninsula and South Greece); Asia Minor; Syria; Mongolia; China (Xinjiang, Jilin, Jiangxi, Heilongjiang); Korea; Japan (Honshu).

a. *Rhyparia purpurata purpurata* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:505 (*Phalaena*).

(= *Phalaena purpurea* LINNAEUS, 1767, Syst. Nat. (Edn. 12) 1 (2): 828; = *Rhyparia purpurata uralensis* SPULER, 1906, Schmett. Europ. 2: 131; = *Rhyparia purpurata barteli* KRULIKOWSKII, 1909, Materialy k poznaniyu fauny i flory Ross. Imp. Otd. Zool. 9: 172).

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia; Kazakhstan; North Kyrgyzstan; Siberia. West Europe; Asia Minor; Mongolia.

b. *Rhyparia purpurata caucasica* (ALPHERAKY, 1867)

Horae soc. Ent. Ross. 10: 14 (*Arctia*).

Caucasus; Transcaucasia. ?Turkey.

c. *Rhyparia purpurata gerda* WARNECKE, 1918

Int. Entomol. Zeit. 12: 81.

Middle part of the Amur basin; Primorye. East Mongolia; North-East China; Korea; Japan.

Amurrhyparia DUBATOLOV, 1985

Sistematika i biologija tshlenistonogikh i gelmintov. Novosibirsk: 66–68.

72. *Amurrhyparia leopardinula* (STRAND, 1919)

Lep. Catal. 22: 185 (*Diacrisia*).

(= *Chelonia leopardina* MÉNÉTRIÈS, 1859, Bull. Acad. Sci. St. Petersb. 17: 218; = *Diacrisia leopardinula mandschurica* O. BANG-HAAS, 1936, Entomol. Zeit. & Int. Ent. Zeit. 50 (30): 348).

South-East Transbaikalia; the middle part of the Amur basin; Primorye. China (Heilongjiang, Shanxi, Gansu, Qinghai, Tibet, Sichuan).

Spilosomini

Ocnogyna LEDERER, 1853

Verh. zool.-bot. Ver. Wien 2 (Abh.): 78.

(= *Somatrichia* KIRBY, 1892, Synonomic Cat. Lepid. Heterocera 1: 274; = *Trichosoma* RAMBUR, 1832, Annls ent. Soc. Fr. (1)1: 272).

73. *Ocnogyna parasita* (HÜBNER, [1790])

Moldova; Crimea; Lower Volga; ?Caucasus (?Abkhasia), ?Armenia. Local in West Europe: South-East France, the Alps, Central Europe from South Slovakia to the North Balkans; Zakynthos Is. (Greece); Asia Minor; ?Iraq.

- a. *Ocnogyna parasita parasita* (HÜBNER, [1790])
Beitr. Gesch. Schmett. 2 (2): 42, t. 2, f. I (*Phalaena*).
Moldova, Crimea. The Alps and Central Europe.
- b. *Ocnogyna parasita rothschildi* A. BANG-HAAS, 1912
Dtsch. Ent. Zeit. IRIS 26: 108, t. 6, f. 3 (*Ocnogyna*).
Lower Volga (Samara region).

74. *Ocnogyna armena* STAUDINGER, 1871

Daghestan; East Transcaucasia (Armenia; Azerbaijan); Turkmenistan (Kopetdagh Mts., Badhyz); South Uzbekistan: Kashkadarinskaya region (STSHETKIN, 1975; TILAVOV, 1978); South-West Tajikistan (STSHETKIN, 1975). ?East Turkey; ?Iraq; Iran; North Afghanistan.

a. *Ocnogyna armena armena* STAUDINGER, 1871

Cat. Lepidop. Europ. Faunengeb. (Edn. 2): 59 (*Ocnogyna loewii armena*).
East Transcaucasia (Armenia, Azerbaijan). ?East Turkey.

b. *Ocnogyna armena daghestana* DUBATOLOV, 1996 subspec. nov. (see p. 71)
East Caucasus (Daghestan).

b. *Ocnogyna armena pallidior* CHRISTOPH, 1884

Mem. Lepidop. Ed. N. M. ROMANOFF 1: 109, t. 7, f. 1a, 1b (*Ocnogyna loewii pallidior*).
Turkmenistan (Kopetdagh Mts., Badhyz); South Uzbekistan; South-West Tajikistan. North Iran;
Afghanistan.

Note: According to the male genitalia structure (fig. 1), *Ocnogyna loewii* (ZELLER, 1846), Entomol. Zeit. 7:9–11 (*Trichosoma*) (= *Trichosoma clathrata* LEDERER, 1855, Verh. zool.-bot. Ges. Wien 5: 202, t. 2, f. 7) from Asia Minor (Taurus Mts.) and the Near East differs well from *O. armena* STGR., including *O. a. pallidior* CHR. The uncus of the former species is gradually tapering to the tip, the apex is not wider than 0.1 mm; the latter species has the uncus forming a short and wide processus at the tip, 0.2 mm wide or wider, its tip is blunt. *O. cypriaca* O. BANG-HAAS, 1934, Ent. Zeit. 48: 48 (*Ocnogyna loewii cypriaca*) from Cyprus Is. is a good species, differing better from the both mentioned species by the presence of many cornuti in the aedeagus vesica and a different structure of the valvae.

Tajigyna DUBATOLOV, 1990

Redkie gelminty, klestshi i nasekomye. Novosibirsk: 82–83.

75. *Tajigyna gansoni* DUBATOLOV, 1990

Redkie gelminty, klestshi i nasekomye. Novosibirsk: 83–84, f. 1b, 2g.
Tajikistan (Peter-the-Great mountain range near Komsomolabad).

Watsonarctia DE FREINA & WITT, 1984

Entomofauna 5 (28): 324.
(= *Eurachia* DUBATOLOV, 1985, Chlenistonogie Sibiri i Dalnego Vostoka. Novosibirsk: 147; = *Eucastana* LERAUT, 1985, Linneana belg. 10: 43).

76. *Watsonarctia deserta* (BARTEL, 1902)

North and Central Ukraine; European Russia: Penza, Simbirsk (= Ul'yanovsk), and Samara regions; Bashkirie; Caucasus; Transcaucasia; North Kazakhstan; mountains of East Kazakhstan; South Siberia: Barnaul region, Altai, Khakasia, Minusinsk; the middle part of the Angara river; Irkutsk. Central and South Europe (except for the western and central parts of the Iberian peninsula, the Alps; the Adriatic coast and Greece); Asia Minor; North Mongolia; China (Xinjiang).

a. *Watsonarctia deserta deserta* (BARTEL, 1902)

Dtsch. Ent. Zeit. IRIS 15: 226–227 (*Arctia casta*).

(= *Bombyx casta* ESPER, 1784, Schmett. in Abbildungen nach der Natur 3: 177, t. 33, f. 2; = *Phragmatobia esperi* KOÇAK, 1980, Comm. Fac. Sci. Univ. Ankara, C, 24(2):10; = *Eucharia casta sibirica* W. KOSHANTSCHIKOV, 1924, Ezhegod. Gos. Muz. im. Nartjanova 2(1):68).

Ukraine, the south part of European Russia; Caucasus; North-West Kazakhstan; South Siberia. West Europe; North Mongolia.

b. *Watsonarctia deserta karduchchena* (DE FREINA, 1983)

Mitt. Münch. ent. Ges. 72: 105, Abb. 20 (*Eucharia deserta*).

Transcaucasia. East Turkey.

c. *Watsonarctia deserta centralasiae* (O. BANG-HAAS, 1927)

Horae Macrolepidop. 1: 59, t. 8, f. 9 (*Eucharia casta*).

Mountains of East Kazakhstan. China (Xinjiang: East Tien Shan).

Chionarctia KÔDA, 1988

Tyô to Ga 39(1):54–58.

(= *Gigantospilosoma* DUBATOLOV, 1990, Taksonomia nasekomykh i gelmintov. Novosibirsk: 95–96).

77. *Chionarctia niveum* (MÉNÉTRIËS, 1859)

Bull. phys.-math. Acad. Sci. St. Petersb. 17: 218 (*Dionychopus nivea*).

?Irkutsk; Middle Amur; Primorye; Southern Sakhalin; Southern Kuril Is. (Kunashir). China (Dunbei, Hebei, Nei Mongol, Shaanxi, Henan, Shandong, Zhejiang, Fujian, Hubei, Hunan, Guangxi, Sichuan); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima).

Alphaea WALKER, 1855

List Spec. lepid. Insects Colln. Br. Mus. 3: 683.

78. *Alphaea melanostigma* (ERSCHOFF, 1872)

Horae Soc. Ross. 8(4): 316 (*Spilosoma*).

(= *Spilarctia karakorumica* DANIEL, 1961, Mitt. Münch. ent. Ges. 51: 159).

Uzbekistan: Zeravshan, Alai mountain ranges; Kyrgyzstan: Alai; Tajikistan: Hissar mountain range, Pamirs. Afghanistan; Pakistan; North India (Himalaya, south to Sikkim and Assam). ?China (Qinghai).

Andala WALKER, 1855

List Spec. lepid. Insects Colln. Br. Mus. 3: 774.

79. *Andala guttata* (ERSCHOFF, 1874)

In: FEDTSCHENKO, Puteshestvje v Turkestan **2**:32, t. 2, f. 28 (*Arctia*).

Uzbekistan: West Tien Shan (Chirman Mt.), ?Zeravshan mountain range; Tajikistan: Hissar mountain range. East Afghanistan.

80. *Andala transversa* (MOORE, 1879)

Proc. Zool. Soc. Lond.:398 (*Cycnia*).

(= *Spilosoma puella* STAUDINGER, 1887, Stett. Ent. Ztg. **48**:85).

Tajikistan: Peter-the-Great and Darwaz mountain ranges, West Pamirs. East Afghanistan; West Pakistan.

Hyphantria HARRIS, 1841

Rep. Insects Mass. injurious to Vegr.:255.

81. *Hyphantria cunea* (DRURY, 1773)

Illust. nat. Hist. exot. Insects **2**:index to vol. 1; 1770, ibidem **1**:36, t. 18, f. 4 (*Phalaena*).

(= *Hyphantria textor* HARRIS, 1828, Rep. Insects Mass. injurious to Vegr.:255).

Moldova (since 1970); Ukraine (since 1952); Transcarpathians, Odessa, Kherson, Nikolaev, and Zaporozhje regions, Crimea; the south part of European Russia (since 1975): Krasnodar and Stavropol regions, North-West Caucasus, east to the west coast of the Caspian Sea (SHAROV & IZHEVSKIY, 1987) and the southern parts of the Samara region (since the '90s) (SACHKOV, 1992); Baltic Provinces (Lithuania and Estonia, since the '80s). South-West France; South Italy; Switzerland, Central Europe, north to South Germany, East Austria and South Slovakia; North Turkey; South Mongolia; North-East China: Liaonin (FANG, 1981), Japan (Honshu, Shikoku, Kyushu); North America (South Canada and USA).

Diaphora STEPHENS, 1827

in: Anonymous, Retrospective Review (2)**1**:244.

82. *Diaphora mendica* (CLERCK, 1759)

Icon. Insect. rariorum **1**: t. 3, f. 5 (*Phalaena*).

(= *Phalaena rustica* HÜBNER, 1791, Beiträge zur Geschichte der Schmett. **2**:64, pl. 3, 2, pl. 3, 2, H;
= *Diaphora mendica malatiana* BYTINSKI-SALTZ, 1936, Ent. Rec. **48**:2 (sep.).

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Karelia and the southern part of Komi republic; Caucasus; Transcaucasia; North Kazakhstan; South Siberia north to Tobolsk, Tomsk, the middle part of the Angara river, and Irkutsk; south of the Baikal region. West Europe, north to South Scandinavia (except for North Ireland, North Scotland, the southern regions of the Iberian peninsula and Greece); North Turkey; Syria; Lebanon.

Eudiaphora DUBATOLOV, 1990

Taksonomiya nasekomykh i gelmintov. Novosibirsk: 93–95.

83. *Eudiaphora turensis* (ERSCHOFF, 1874)

In: FEDTSHENKO, Puteschestvje v Turkestan **3**:33, t. 2, f. 29 (*Spilosoma*).

(= *Diaphora turensis maracandica* SEITZ, 1910, Gross-Schmett. Erde **2**: 92, f. 17d; = *Diaphora afghanistanensis* DANIEL, 1966, Mitt. Münch. ent. Ges. **56**(7):163, t. 3, f. 1).

Turkmenistan: Kopetdagh and Kuhitang mountain ranges, Amu-Darja river (Chardzhou), Kazakhstan: Syr-Darja, Chu and Ili river valleys; Ketmen mountain range; Kyrgyzstan: Ferghanian mountain range; Uzbekistan: the Karzhantau and Zeravshan mountain ranges, Tajikistan: Hissar and Turkestan mountain range, West Pamirs. Afghanistan; China (Xinjiang).

Spilosoma CURTIS, 1825

Br. Ent. 2: f. 92.

84. *Spilosoma lubricipedum* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:505 (*Phalaena lubricipeda*).

(= *Bombyx menthastris* [DENIS & SCHIFFERMÜLLER], 1775, Ankündung syst. Werkes Schmett. Wiener-gegend: 54; = *Diacrisia menthastris chishimana* MATSUMURA, 1929, Ins. matsumur. 3: 167; = *Diacrisia masuensis* MATSUMURA, 1930, Ins. matsum. 5: 34, t. 1, f. 1; = *S. menthastris elegans* BRYK, 1948, Ark. Zool. 41A(1): 43).

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Karelia, Arkhangelsk region and Komi republic; Caucasus; Transcaucasia (except for East Azerbaijan); North Kazakhstan; South Siberia, north to Tobolsk, Ket river (Tomsk region), and Angara river; Transbaikalia; Amur basin; Primorye; Sakhalin; Southern Kuril Is. (from Urum to Kunashir). West Europe, north to Middle Scandinavia (except for the central and south-western parts of the Iberian peninsula, South Italy, South Greece); northern part of Asia Minor; ?Mongolia; China (Dunbei, Hebei, Nei Mongol, Shaanxi, Jiangsu, Zhejiang, Anhui, Jianxi, Fujian, Hubei, Sichuan, Guizhou, Yunan); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Yaku).

85. *Spilosoma punctarium* (STOLL in CRAMER, [1782])

Uitlandsche Kapellen 4: 233, t. 398, f. D (*Bombyx*).

(= *Arctia punctigera* MOTSCHULSKY, [1861], Et. Ent. 9: 31, t. 1, f. 7; = *Spilosoma roseiventer* SNELLEN, 1863, Tijds. 6: 143; = *Spilosoma dornesii* OBERTHÜR, 1879, Diagnoses Espèces nouv. Lepid. Ille d'Ascoli: 6; = *Spilosoma doerriesi* OBERTHÜR, [1881] 1880, Et. Ent. 5: 31, t. 1, f. 7; = *Spilosoma punctarium miserata* BRYK, 1942, Dtsch. Ent. Zeit. IRIS 56: 33).

Middle Amur; Primorye; Southern Kuril Is. (from Urum to Kunashir). China, west to Tibet; Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yaku).

86. *Spilosoma urticae* (ESPER, 1789)

South Lithuania; Ukraine; Crimea; Moldova; European Russia, north to Bryansk, Moscow, and the Middle Volga, there is some old material from St.-Petersburg, too; Caucasus; Transcaucasia; Kazakhstan; Kyrgyzstan (Bishkek, Osh); Uzbekistan: West Tien Shan and Ferghana valley; South Siberia, north to Kurgan, Chulym river (Tomsk region), and the middle part of the Angara river; Transbaikalia (Ulan-Ude); Middle Amur (Blagovestshensk); South Primorye; ?Southern Kuril Is. (?Kunashir). West Europe, north to South England and South Scandinavia, south to North Italy and the Northern Balkans; ?Turkey; North Iran (Chelus); China (Xinjiang, Sichuan, Jiangsu).

a. *Spilosoma urticae urticae* (ESPER, 1789)

Schmett. 3: 20, t. 83, f. 2 (*Phalaena*).

South Lithuania; Ukraine, Crimea; Moldova; European Russia; Caucasus; Transcaucasia; Kazakhstan; Kyrgyzstan; North Uzbekistan; South Siberia; Middle Amur. West Europe.

b. *Spilosoma urticae mandli* SCHÄWERDA, 1922

Zeit. Öst. Ent. Ver. 7: 11 (*Spilosoma mandli*).

(= ?*Spilosoma sangaica* WALKER, [1865], List. Spec. Lepid. Insects Colln. Br. Mus. 31: 294³).

South Primorye; ?Kunashir.

- 87 *Spilosoma streltzovi* DUBATOLOV, 1996 spec. nov. (see p. 76)
Middle Amur, Primorye.

Spilarctia BUTLER, 1875

Cistula ent. 2: 39.

Note. "Spilarctia tschitaensis DANIEL", 1953, Mitt. Münch. ent. Ges. 43: 252, t. 7, f. 11, which was described from Chita (Transbaikalia) is a synonym of *Isochlora grumi* ALPHERAKY, 1892, Horae Soc. Ent. Ross. 26:448 (Noctuidae).

88. *Spilarctia luteum* (HUFNAGEL, 1766)

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Karelia and Kotlas; West Caucasus; West Transcaucasia (West Georgia); North Kazakhstan; South Siberia, north to Tobolsk, Chulym river (Tomsk region), Krasnoyarsk, and Irkutsk; Middle Amur; Primorye; Sakhalin; Southern Kuril Is. (from Urup to Kunashir); possibly absent from Transbaikalia. West Europe, north to South Scandinavia (except for the central, southern and eastern parts of the Iberian peninsula); the northern part of Asia Minor; China (Dunbei, Hebei, Shaanxi); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu).

a. *Spilarctia luteum luteum* (HUFNAGEL, 1766)

Berlin. Mag. 2:412 (*Phalaena*).

(= *Phalaena zatima* STOLL, 1781, Pap. Exot. 4:182, t. 381, f. F).

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia; West Caucasus; West Georgia; North Kazakhstan; South Siberia. West Europe; the northern part of Asia Minor.

b. *Spilarctia luteum japonicum* (ROTHSCHILD, 1910)

Novit. Zool. 17: 131 (*Diacrisia lutea japonica*).

(= *Spilarctia obliqua ursulina* BRYK, 1942, Dtsch. Ent. Zeit. IRIS 56:31–32; = *S. obliqua bergmani* BRYK, 1942, Dtsch. Ent. Zeit. IRIS 56:31; = *Diacrisia lutea rhododactyla* BRYK, 1948, Ark. Zool. 41A(1):41).

Middle Amur; Primorye; Southern Sakhalin; Southern Kuril Is. (from Urup to Kunashir). East China; Korea; Japan.

89. *Spilarctia seriatopunctatum* (MOTSCHEULSKY, [1861])

Et. Ent. 9:31 (*Spilosoma seriatopunctata*).

(= *Spilarctia ione* BUTLER, 1875, Cistula ent. 2:41; = *Spilarctia mollicula* BUTLER, 1877, Ann. Mag. nat. Hist. (4)20: 395; = *Spilosoma striatopunctata* OBERTHÜR, 1879, Diagnoses especes nouv. Lepid. Ile d'Askold: 6; = *Spilarctia casigneta flavómaculosa* KARDAKOFF, 1928, Ent. Mitt. 17:415; = *S. casigneta sjoquisti* BRYK, 1942, Dtsch. Ent. Zeit. IRIS 56: 32; = *Diacrisia seriatopunctata varians* BRYK, 1948, Ark. Zool. 41A(1):41).

Middle Amur; Primorye; Southern Sakhalin; Southern Kuril Is. (from Urup to Kunashir). China (Heilongjiang, Jilin, Jiangxi, Fujiang, Sichuan); Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Yaku, Amami, Okinawa).

3 According to the type-specimen (?) colour figure (BUTLER, 1879b: pl. XLII, fig. 5) it resembles *Spilosoma urticae* (ESPER) very much by the wings (elongate fore wing with well marked black dots and almost white hind wing). *S. urticae mandli* SCHAW. has the same wings. However, the introduction of the new synonymy demands the investigation of the specimen in the collection of The Natural History Museum (London).

90. *Spilarctia obliquizonatum* (MIYAKE, 1910)

Jl. Coll. Agric. Imp. Univ. Tokyo **2**: 208 (*Diacrisia obliquizonata*).

Southern Sakhalin; Southern Kuril Is. (Kunashir). Japan (Hokkaido, Honshu, Shikoku, Kyushu).

91. *Spilarctia subcarneum* (WALKER, 1855)

List Spec. lepid. Insects Colln. Br. Mus. **3**: 675–676 (*Spilosoma subcarnea*).

(= *Aloa bifrons* WALKER, 1855, List Spec. lepid. Insects Colln. Br. Mus. **3**: 705; = *Spilosoma rybakovi* ALPHERAKY, 1897, Mem. lepidop. Ed. N. M. ROMANOFF **9**: 171, t. 10, f. 9).

South Primorye. China, from Dunbei south to Taiwan, Guangdong and North Yunan; Korea; Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Yaku, Okinawa).

a. *Spilarctia subcarneum charbini* (DANIEL, 1943)

Mitt. Münch. ent. Ges. **33**(3): 696, t. 19, f. 7, 8 (*Spilarctia subcarnea*).

South Primorye. North-East China (Dunbei).

Notes. 1) This species was reported from the Kamchatka peninsula by KURENTSOV (1966) as a pest of *Betula ermanii*. I haven't found any specimen from Kamchatka in KURENTZOV's collection in the Institute of Biology and Pedology (Vladivostok). It is quite possible that he determined the species by the larvae only, and that this determination might be wrong. 2). There are two male specimens of this species originating from South Primorye, Kaimanovka, 24.VIII. 1981, PETRIKEVITSH leg., with a yellow, not red abdomen. These specimens have been collected together with the typical form. I haven't found any significant differences between them in the male genitalia. I propose a new name for specimens with yellow abdomen: *Spilarctia subcarneum* f. *lydia* DUBATOLOV, **forma nov.**

Lemyra WALKER, 1856

List Spec. lepid. Insects Colln. Br. Mus. **7**: 1690.

(= *Thanatarctia* BUTLER, 1877, Ann. Mag. nat. Hist. (4)**20**: 395).

93. *Lemyra inaequalis* (BUTLER, 1879)

Ann. Mag. nat. Hist. (5) **4**: 351 (*Spilarctia*).

(= *Diacrisia rhodophila japonensis* ROTHSCHILD, 1914, in: SEITZ, Gross-Schmett. Erde **10**: 239, f. 21e).

L. inaequalis (BTL.) is distributed in Japan only (Honshu, Shikoku, Kyushu, Tsushima, Yaku). In the Lower Amur, the mountain taiga of Primorye and Korea occurs a related species. It will be described by Yu. A. TSHISTJAKOV and Y. KISHIDA. Such a specimen was figured by FANG CHENG-LAI (1982: t. 69, fig. 1587) as "*Spilarctia japonensis* ROTH." The mentioned author gave its distribution in China as follows: Heilongjiang, Liaonin, Jilin, Zhejiang,

94. *Lemyra jankowskii* (OBERTHÜR, [1881] 1880)

Et. Ent. **5**: 31–32, t. 8, f. 3 (*Spilosoma*).

(= *Diacrisia jankowskii korearctica* BRYK, 1948, Ark. Zool. **41A**(1): 42).

Middle Amur; Primorye. China (Heilongjiang, Liaonin, Hebei, Shandong, Shaanxi, Jiangsu, Zhejiang, Shanxi, Sichuan, Tibet, Yunan).

Note: This species was reported to occur also in the Kuril Is.: Kunashir (KONOVALOVA, 1968). I consider this a mistake, because this species does not inhabit Japan.

Phragmatobia STEPHENS, 1828

Illust. Br. Ent. (Haustellata) **2**: 55, 73.

95. *Phragmatobia fuliginosa* (LINNAEUS, 1758)

Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; European Russia, north to Murmansk, Arkhangelsk region, and Komi republic; Caucasus; Transcaucasia; Kazakhstan; all the countries of Middle Asia, except for deserts; Siberia, north to Beryozov (the lower course of the Ob river), Podkamennaya Tunguska river, Central Yakutia, and the upper course of the Kolyma river; Kamchatka; Bering Is. (DYAR, 1899). The south-easternmost locality in Russia is the Nerchinsk surroundings in Transbaikalia. West Europe; North-West Africa; Asia Minor; the Near East; Iraq; North Iran; Afghanistan; China (Xinjiang, Qinghai, Nei Mongol).

a. *Phragmatobia fuliginosa borealis* (STAUDINGER, 1871)

Cat. Lep. Europ. Faunengeb. (Edn. 2): 59 (*Spilosoma fuliginosa*).

The northern part of Eurasia (the northern regions of European Russia; Siberia, except for the southern part; Scandinavia).

b. *Phragmatobia fuliginosa fuliginosa* (LINNAEUS, 1758)

Syst. Nat. (Edn. 10) 1:509 (*Phalaena*).

(= *Spilosoma fuliginosa fervida* STAUDINGER, 1871, Cat. Lep. Europ. Faunengeb. (Edn. 2): 59; = *Phragmatobia fuliginosa lurida* ROTHSCILD, 1910, Novit. Zool. 17: 115).

?Baltic Provinces; Belarus; Ukraine; Crimea; Moldova; central and southern regions of European Russia; North Kazakhstan; South Siberia. Central and South Europe.

c. *Phragmatobia fuliginosa taurica* DANIEL, 1970

Zeitschr. der Arbeitsgemeinschaft Österr. Entomologen 22:9–10.

Caucasus; Transcaucasia (excluding the main part of Azerbaijan and Armenia). East Turkey; the Near East; mountains of northern and north-western Iran.

d. *Phragmatobia fuliginosa paghmani* LENEK, 1966

Zeit. Wiener Ent. Ges. 51: 105.

South-East Transcaucasia (Azerbaijan and Armenia); all the countries of Middle Asia; South Kazakhstan. Iran; Afghanistan. China (West Xinjiang).

Note: *P. f. pulverulenta* ALPHERAKY, 1889, Mem. Lepidop. Ed. N. M. ROMANOFF 5:84 (*Spilosoma fuliginosa*) = *P. f. pallida* ROTHSCILD, 1910, Novit. Zool. 17: 116 = *P. f. thibetica* STRAND, 1919, Lep. Cat. 22:416, occurs only in China: East Xinjiang, Qinghai, Nei Mongol; and South Mongolia (DARITSHEVA, DUBATOLOV, 1989).

96. *Phragmatobia amurensis* SEITZ, 1910

Middle Amur; Primorye; Southern Sakhalin; Southern Kuril Is. (from Iturup to Kunashir). China (Dunbei, Hebei); Korea; Japan (Hokkaido, Honshu).

Note: For a review of distinguishing characters see DUBATOLOV (1990d). The males of *Ph. amurensis* SEITZ have pectinate antennae, those of *Ph. fuliginosa* (L.) simple ones.

a. *Phragmatobia amurensis amurensis* SEITZ, 1910

Gross-Schmett. Erde 2: 79, f. 16b (*Phragmatobia fuliginosa*).

(= *Phragmatobia fuliginosa chosensis* BRYK, 1948, Ark. Zool. 41A(1): 40).

Middle Amur; Primorye. North-East China; Korea.

b. *Phragmatobia amurensis japonica* ROTHSCILD, 1910

Novit. Zool. 17(2): 116 (*Phragmatobia fuliginosa*).

Southern Sakhalin; Southern Kuril Is.; Japan.

97 *Phragmatobia placida* (FRIVALDSZKY, 1835)

Mag. Tuds. Evk. 2:271, t. 7, f. 5 (*Euprepia*).

South Crimea (KOSTJUK & PLJUSTSH, 1987); Transcaucasia (Armenia). The Balkans; Turkey; Syria; North Iraq; North Iran.

Epatolmis BUTLER, 1877

Trans. ent. Soc. Lond.: 348.

98. *Epatolmis caesarea* (GOEZE, 1781)

Ent. Beyträge 3 (3):63 (*Phalaena*).

(= *Phalaena luctifera* ESPER, 1784, Schmett. Abbild. Natur. 3: 222, t. 43, f.1–5; = *Atolmis japonica* WALKER, [1865] 1864, List Specm. lepid. Insects Colln. Br. Mus. 31: 223; = *Estigmene moerens* BUTLER, 1885, Cist. Ent. 3: 114).

Baltic Provinces; Belarus; Ukraine; Moldova; European Russia, north to Luga (St.-Petersburg region), Vasil'sursk (120 km east of Nizhnii Novgorod) (TSCHETVERIKOV, 1993), and Vyatka, south to the Azov Sea and ?Saratov; Transcaucasia (Borzhomi in Georgia and Kirovabad (Gyandzha) in West Azerbaijan); South Siberia: Tobolsk, Barnaul, Khakasia, Minusinsk; southern banks of the Baikal region; Transbaikalia; Middle Amur (Blagowestshensk); South Primorye. Central Europe: Central and South France, South Germany, Austria and Czechoslovakia, south to North Italy and the North-East Balkans; Asia Minor; Mongolia; China (Heilongjiang, Jiangxi, Hebei, Nei Mongol, Shanxi, Shandong, Shaanxi, Jiangsu, Hunan, Sichuan, Yunan); Korea, Japan (Honshu, Shikoku, Kyushu).

Part II: Descriptions of New Species and Subspecies

***Lacydes spectabilis sheljuzhkoi* subspec. nov.**

(Fig. 2a)

The wing expanse of the holotype is 33 mm. The wing pattern is as in *L. spectabilis spectabilis* (TAUSCHER), the spots on the forewing lack distinct borders of black scales, which is typical for *L. spectabilis annelata* (CHRISTOPH). The main character of the new subspecies is a clear yellowish, not whitish groundcolour of the wings, and the spots on the forewing being dark yellow, without a greyish-brown tint, which is typical for both of the other known subspecies.

Material. Holotype ♂, Armenia, "Transcauc. m., pag. Engidzha ad fl. Arax (prope stat. Norashen), 12.IX.1931, M. RJABOV leg." Paratypes: 1 ♂, same data; 4 ♂♂, Norashen, fl. Arax, 12.–13.IX.1931, M. RJABOV leg.; 2 ♂♂, Erivanj, 2.–23.IX.1938. All the specimens belong to L. SHELJUZHKO's collection, kept in the Zoological Museum of Kiev State University.

***Platarctia atropurpurea sotavaltae* subspec. nov.**

(Fig. 2g)

The wing expanse is 33–38 mm in females, and 32 mm in the male. The forewing is greyish-brown with the pattern typical for this species. The main character of the new subspecies is the size of the yellow antemedian and postmedian spots in the space 1 (between veins 1 and 2) on the forewing. These spots are small, either of equal size or the antemedian spot being smaller than the postmedian one. *P. atropurpurea atropurpurea* (BANG-HAAS) has a large antemedian spot in the space 1, elongated between the veins, and usually this spot is twice as long as the postmedian one. In addition, the hindwing in the new subspecies has separate rose spots, being brighter in the females. The

corresponding spots in *P. atropurpurea atropurpurea* are usually fused, being bright yellow in the females and yellow, but rose along the costa in the males.

Material. Holotype ♀, E. Yakutia, 180 km ENE of the settlement Khandyga, the upper course of the East Khandyga River, the stream Kyurbelyakh, 7.VII.1985, V. DUBATOLOV, L. POPOVA leg. Paratypes: 2 ♀♀, same locality, 8.VII.1985, V. DUBATOLOV leg., 4.VII.1986, L. POPOVA leg.; 1 ♀, Yakutia, the southern part of the Verkhoyansky mountain range, the upper part of the Kele river, the stream Charkymbal, 11.VII.1989, L. POPOVA leg.; 1 ♂, Magadan region, Kolyma River, near Mt. Aborigen, a black larva 8.–11.VIII.1986, pupa 26.VII.1986, imago 10.IX.1986, V. DUBATOLOV leg.

This subspecies is named in honour of Prof. Dr. O. SOTAVALTA (Finland), who firstly mentioned this species from North-East Siberia (SOTAVALTA, 1965).

***Arctia caja ossetica* (O. BANG-HAAS et SHELUZHKO, in litt.) subspec. nov.**

(Fig. 2d)

The wing expanse is 60–64 mm in males, 75 mm in the female. This subspecies is transitional between the European *A. caja caja* (LINNAEUS) and the Transcaucasian *A. caja wiskotti* ROMANOFF. The colour of the forewing and the shape of the spots are the same as in the mentioned subspecies, but the colour of the hindwing is peculiar, it is bright yellowish-orange in the new subspecies, bright red in *A. caja caja*, and pale yellow in *A. caja wiskotti*.

Material. Holotype ♂, Caucasus, "Ossetia s.", Buron, 1250 m, 10.VIII.1940 (lumine), L. SHELUZHKO leg." Paratypes: 2 ♂♂, 2 ♀♀, same locality, 10., 13.VIII.1940, L. SHELUZHKO leg.; 1 ♂, Ossetia, Sadovskije Rudniki, 31.VII.1904. All the specimens belong to L. SHELUZHKO's collection kept at the Zoological Museum of Kiev State University.

This subspecies was firstly designated by O. BANG-HAAS and L. SHELUZHKO in his collection. For unknown reasons a description was never published.

***Sibirarctia buraetica chajataensis* subspec. nov.**

(Fig. 2j)

The wing expanse is 26–28 mm. The forewing is very dark, the white bands being reduced to very narrow stripes: the subterminal one is W-shaped, sometimes poorly developed, not wider than 0.25 mm. The postmedial band is better developed but does not exceed a 0.4 mm width. The medial band is reduced to a single small spot inside the cell. Other transversal bands are absent, but there is a good developed stripe between veins 1 and 2, being 0.25 mm wide. The fused part of vein 2+3 and the base of vein 1 are covered with light scales. The hindwing as in *S. b. buraetica* (BANG-HAAS).


Material. Holotype ♂, E. Yakutia, 180 km ENE of the Khandyga settlement, the upper course of the East Khandyga River, near the mouth of the stream Kyurbelyakh, a shingly river bank, 29.VII.1985, V. DUBATOLOV & L. POPOVA leg. Paratypes: 1 ♂, same locality, 17.VII.1985, L. POPOVA leg., 1 ♂ ex larva 29.VI.1989, S. SAZONOV leg.

***Ocnogyna armena daghestana* subspec. nov.**


(Fig. 2i)

The wing expanse is 28 mm. The wing pattern is characteristic for this species, but all the spots are light-brown with darker borders, not evenly dark-brown as in the other *O. armena* STAUDINGER subspecies. The spots on the hindwing are strongly widened, the subterminal one forming a wide band from the costa to vein 2, with an additional spot in the tornal angle. The discoidal spot is also large.


Material. Holotype ♂, East Caucasus, "Dagestan, Derbent, 16.X.1910, Coll. A. XIEZOPOLSKY (*armena*, L. SHELUZHKO det.)" The type belongs to L. SHELUZHKO's collection kept at the Zoological Museum of Kiev State University.




1a




1b



1c



1d



1e






Fig. 1: Male genitalia and VIII. tergites of *Ocnogyna (loewii* (ZELL.)). a – *O. loewii* (ZELL.), Taurus; b – *O. armena armena* STGR., Armenia, Erevan; c – *O. a. daghestana* subsp. nov., Caucasus, Daghestan, Derbent, holotype; d – *O. a. pallidior* CHR., Turkmenistan, West Kopetdagh, Sumbar river; e – *O. cypriaca* O. BANG-HAAS, Cyprus, Limassol, co-type (paratype).







Fig. 3: Male genitalia of *Spilosoma streitzovi* spec. nov. a – general view, b – aedeagus with everted vesica.







2a




2d




2b



2e



2c




2f

Fig. 2: Imagines of tiger moths. a – *Lacydes spectabilis sheljuzhkoi* subspec. nov., holotype; b – *L. s. annelata* (CHR.), Turkmenistan, Kopetdagh Mts., Dushak Mt.; c – *L. s. spectabilis* (TAUSCH.), N.E. Kazakhstan, Semipalatinsk region, Glukhovka; d – *Platarctia atropurpurea sotavaltai* subspec. nov., East Yakutia, upper East Khandyga river, holotype; e – *P. a. atropurpurea* (O. BANG-HAAS), Chita region, Kalar district, Udonkan; f – *Arctia caja ossetica* subspec. nov., Ossetia, Buron, holotype; g – *A. c. wiskotti* STGR., Armenia, Lake Sevan, Semenovka; h – *A. c. caja* (L.), Novosibirsk region, Lake Chany; i – *Sibiractia buraetica chajataensis* subspec. nov., East Yakutia, upper East Khandyga river, holotype; j – *Ocnogyna armena daghestana* subspec. nov., Daghestan, Derbent, holotype; k – *Spilosoma streltzovi* spec. nov., male, Blagovestshensk, paratype; l – *S. streltzovi* spec. nov., female, Blagovestshensk, paratype; m – *Lemyra (inaequalis* (BTL.)), male, Primorye, Ternei; n – *L. (inaequalis* (BTL.)), female, Nikolaevsk-na-Amure.



2g



2h




2i




2j




2k



2l



2m



2n

i j
Lege
verta

Spilosoma streltzovi spec. nov.

(Fig. 2k, l)

The wing expanse is 37–38.5 mm in males, and 43 mm in the female. In males the antennae are uniformly black, long, only slightly shorter than the length of the cell. The fore femur is yellow, the middle and hind ones white. All the tibiae are white, dorsally with a contrasting black stripe, the tarsi are black. The abdominal tergites are yellow with rows of black spots on the dorsal and lateral sides. The sternites are whitish, the traces of two rows of black spots being visible only in the female. In males the wings are opaque, in the female semitransparent, they are all white with small black spots. The forewing has a basal and a subbasal spot on the costa; another one, if visible, is situated behind the median vein. Two or three antemedian spots are present near the costa, sometimes one more spot is added at the middle of vein 1. The female has one more antemedian spot proximally of vein 2. The discoidal vein bears two black spots. The subterminal row is very variable, but always has a stroke-like longitudinal spot between veins 5 and 6. A full subterminal row consists of two spots in the space 1b, and one spot in spaces 4, 5, 6, 7 and 9. There are no terminal spots. The hindwing has a black spot in the fore angle of the cell and usually three subterminal spots in the spaces 1c, 5, and 7; rarely an additional spot is present near vein 1b, and another one in the space 6. The underside is identical to the upperside.

The male genitalia are shown on fig. 3.

Systematical notes. The new species to some extent resembles *S. lubricipedum* (LINNAEUS) by external appearance: a yellow abdomen and white wings with small black spots. It is distinguishable by the black stroke-like longitudinal spot in the middle of the space 5, 1.5 mm off the forewing margin. In *S. lubricipedum* the black spots in the space 5 are always placed along the veins only. There are additional distinctive characters: *S. streltzovi* spec. nov. has longer and black antennae, evenly black tarsi, a black dorsal stripe on the hind tibiae and lacks the terminal spots on the wings. On the contrary, *S. lubricipedum* has shorter antennae, no more than 2/3 of the cell length, covered with black and white scales, white tarsi, sometimes with an incomplete black dorsal stripe, the black terminal dots on the forewings are often visible in the Far Eastern specimens. By these characters, as well as the valva shape, *S. streltzovi* spec. nov. mostly resembles the chinese *S. mienshanica* DANIEL, 1943, Mitt. Münch. Ent. Ges. **33**(3): 732, t. 20, f. 20–21, ab. 27–28 (from Shanxi and Shaanxi), but differs from it by its smaller size, a white thorax and yellow abdomen. *S. mienshanica* is larger, the wing expanse being 41–44 mm in males and 48–49 mm in females, the thorax is cream-white, the abdomen is red. Unfortunately, DANIEL pictured only a male valva, so it is impossible to compare the entire genitalia structure of both species.

Material. Holotype ♂, Amurskaya region, near Blagovestshensk, attracted by light, 6.VI.1990, A. N. STRELTSOV leg. Paratypes: 3 ♂♂, 1 ♀, same locality, 31.V.–8.VI.1990, 15.III.1994, A. N. STRELTSOV leg. I have also seen similar specimens originating from South Primorye.

This species is named in honour of A. N. STRELTSOV from Blagovestshensk.

References

- ALBERTI, B. (1973): Zweite Mitteilung über einige bemerkenswerte Lepidopteren-Formen aus dem Grossen Kaukasus und Transkaukasien (Lepidoptera). – Atalanta **4**(6B): 380–393.
- ALPHERAKY, S. (1867): Cheshuekrylue Severnogo Kavkaza [Lepidoptera of the North Caucasus]. – Trudy Russ. entomol. obstshestva **10**: 1–53 (in russian).
- ALPHERAKY, S. (1876): Cheshuekrylye (Lepidoptera) okrestnostei Taganroga [Lepidoptera of the Taganrog district]. – Trudy Russ. entomol. obstshestva **8**: 150–226 (in russian).
- ALPHÉRAKY, S. (1882): Lépidoptères du district de Kouldjâ et des montagnes environnantes. – Horae Soc. Ent. Ross. **17**: 15–103, pl. 1–3.
- ALPHERAKY, S. (1888): Neue Lepidopteren. – Stett. ent. Ztg. **49**: 66–69.
- ALPHÉRAKY, S. (1897): Lépidoptères de l'Amour et de la Corée. – Mém. l'épidop. Ed. N. M. ROMANOFF **9**: 151–184, T. 10–13.

- ALPHERAKY, S. (1908): Tcheshuekrylye okrestnostei Taganroga. Dobavlenie III. [Lepidoptera of the Taganrog district. Supplement III]. – Horae Soc. ent. Ross. **38**: 558–618 (in russian).
- BANG-HAAS, A. (1907): Neue oder wenig bekannte palaearktische Macrolepidopteren. – Dtsch. Ent. Zeit. IRIS **20**: 69–88, t. 3.
- BANG-HAAS, A. (1910): Neue oder wenig bekannte palaearktische Macrolepidopteren. – Dtsch. Ent. Zeit. IRIS **24**: 27–51, t. 3.
- BANG-HAAS, A. (1912): Neue oder wenig bekannte palaearktische Macrolepidopteren. – Dtsch. Ent. Zeit. IRIS **26**: 103–110, 139–162, 221–230, t. 5–6.
- BANG-HAAS, O. (1927): Horae Macrolepidopterologicae Regionis Palaearticae. Bd. **1**. – Dresden, 128 S., 10 Pl.
- BANG-HAAS, O. (1934): Neubeschreibungen und Berichtigungen der Palaearktischen Macrolepidopterenfauna. – Ent. Zeit. & Int. Ent. Zeit. **48**.
- BANG-HAAS, O. (1936): Neubeschreibungen und Berichtigungen der Palaearktischen Macrolepidoptera. XXV. – Ent. Zeit. & Int. Ent. Zeit. **50** (30): 345–348.
- BANG-HAAS, O. (1938): Neubeschreibungen und Berichtigungen der Palaearktischen Macrolepidoptera. XXXVI. – Ent. Zeit. **52** (22): 177–180.
- BARTEL, M. (1906): *Callimorpha philippii*, eine neue Art aus Zentral-Asien. – Soc. Entom. **21** (6): 41–42.
- BOISDUVAL, J. (1832): Icônes historiques de Lépidoptères d'Europe, nouveaux ou peu connus. Paris, t. **2**, 192 p., 84 pl.
- BÖTTCHER, E. (1905): Neue und wenig bekannte Arctiiden aus Turkestan. – Entomol. Zeitschr. **19**: 53–55, 61–63.
- BREMER, O. (1861): Neue Lepidopteren aus Ost-Sibirien und dem Amur-Lande gesammelt von RADDE und MAACK. – Bull. Acad. Sci. St. Petersb. **3**: 461–495.
- BRYK, F. (1942): Zur Kenntnis der Grossschmetterlinge der Kurilen. – Dtsch. Ent. Zeit. IRIS **56**: 3–90, pl. 1–2.
- BRYK, F. (1848): Zur Kenntnis der Grossschmetterlinge von Korea. Pars II. Macrofrenate II (fins). – Ark. Zool. **41A** (1): 1–225, 7 pls.
- BUDASHKIN, YU. N. (1987): Cheshuekrylye (Soobstshetie 3) [Lepidoptera (Report 3)]. In: Cheshuekrylye Karadagskogo zapovednika [Lepidoptera of the Karadag Nature Reserve]. – Moscow, p. 32–62 (in russian). (Series: "Fauna i flora zapovednikov SSSR" [Fauna and Flora of Nature Reserves of the USSR]).
- BUTLER, A. G. (1877): Descriptions of new species of Heterocera from Japan. – Ann. Mag. Nat. Hist. (4) **20**: 393–404, 473–483.
- BUTLER, A. G. (1879a): Descriptions of new Species of Lepidoptera from Japan. – Ann. Mag. Nat. Hist. (5) **4**: 349–374, 437–457.
- BUTLER, A. G. (1879b): Illustrations of typical specimens of Lepidoptera Heterocera in the collection of the British Museum. Part III. – London, xxiii + 82 p., pl. XLI–LIX.
- BUTLER, A. G. (1881): Descriptions of new Genera and Species of Heterocerous Lepidoptera from Japan. – Trans. Entomol. Soc. Lond., p. 1–23.
- BUTLER, A. G. (1885): Descriptions of moths new to Japan, collected by Messrs LEWIS and PRYER. – Cistula Ent. **3**: 113–139.
- LE CERF, F. (1913): Contribution à la faune lépidoptérologique de la Perse. – Ann. Hist. Nat. Délégation en Perse. T. **2**, entom., fasc. 2, 12 + 85 p., t. 1–2.
- ČERNÝ, K. (1986): *Carcinopyga lindti* spec. nov. (Lepidoptera, Arctiidae) aus Tian Schan. – Neue Entomol. Nachr. **19** (1/2): 31–36.
- CHRISTOPH, H. (1877): Sammelergebnisse aus Nordpersien, Krasnowodsk in Turkmenien und dem Daghestan. – Horae Soc. ent. Ross. **12**: 181–299, T. 5–8.
- CHRISTOPH, H. (1884): Lepidoptera aus dem Achal-Tekke-Gebiete. Erster Theil. – Mém. l'épidop., Ed. N. M. ROMANOFF **1**: 93–138, T. 6–8.
- CHRISTOPH, H. (1887a): Diagnosen neuer Lepidopteren aus Tekke. – Stett. ent. Ztg. **48**: 162–167.
- CHRISTOPH, H. (1887b): Lepidoptera aus dem Achal-Tekke-Gebiete. Dritter Theil. – Mém. l'épidop., Ed. N. M. ROMANOFF **3**: 50–125, T. 3–5.

- CHRISTOPH, H. (1889): Lepidoptera aus dem Achal-Tekke-Gebiete. Vierter Theil. – Mém. lépidop. Ed. N. M. ROMANOFF **5**: 1–58, T. 1–3.
- CHRISTOPH, H. (1892): *Arctia turbans*, eine neue sibirische Art. – Horae Soc. Ent. Rossicae **26**: 460–461.
- CHRISTOPH, H. (1893): Lepidoptera Nova Faunae Palaearcticae. – Dtsch. Ent. Zeit. IRIS **6**: 86–96.
- DANIEL, F. (1939): Gedanken zu einigen Arctiiden-Formen (Lepidoptera). – Mitt. Münch. ent. Ges. **29**: 354–368.
- DANIEL, P. (1943a): Beiträge zur Kenntnis der Arctiidae Ostasiens unter besonderer Berücksichtigung der Ausbeuten H. HÖNE's aus diesem Gebiet (Lep. Het.). I Teil: Callimorphinae und Nycteriminae. – Mitt. Münch. Ent. Ges. **33**: 247–269, T. 13.
- DANIEL, F. (1943b): Beiträge zur Kenntnis der Arctiidae Ostasiens unter besonderer Berücksichtigung der Ausbeuten H. HÖNE's aus diesem Gebiet (Lep., Het.). II Teil. Hypsinae, Micractiinae, Spilosominae, Arctiinae. – Mitt. Münch. Ent. Ges. **33** (2/3): 673–759, T. 14–22.
- DANIEL, F. (1951–1954): Beiträge zur Kenntnis der Arctiidae Ostasiens unter besonderer Berücksichtigung der Ausbeuten von Dr. h. c. H. HÖNE aus diesem Gebiet (Lep.-Het.). – Bonn. zool. Beitr. **2** (3–4)(1951): 291–327, Taf. I; **3** (1–2)(1952): 75–90, Taf. II; **3** (3–4)(1952): 305–324; **4** (1–2)(1954): 89–138, Taf. III.
- DANIEL, F. (1953): Neue Heterocera-Arten und Formen. – Mitt. Münch. Ent. Ges. **43**: 252–261.
- DANIEL, F. (1961): *Spilarctia karakorumica* spec. nov. – Mitt. Münch. Ent. Ges. **51**: 159.
- DANIEL, F. (1966): Zwei neue Arctiidae aus Afghanistan (Lep.). -Mitt. Münch. Ent. Ges. **56** (7): 161–164, T. 3.
- DANIEL, F. (1970): Rassenanalytische Untersuchungen bei *Phragmatobia fuliginosa* L. und *Phragmatobia amurensis* SEITZ (Lep. Arctiidae). – Zeitschr. der Arbeitgemeinschaft Österr. Entomologen **22**: 2–17.
- DARITSHEVA, M. A. & V. V. DUBATOLOV (1989): Fauna i ekologiya cheshuekrylykh semeistva medvedits (Lepidoptera, Arctiidae) Turkmenistana [Fauna and ecology of tiger moths (Lepidoptera, Arctiidae) of Turkmenistan]. – Izvestiya AN Turkm. SSR. Seriya biol. nauk. Ashkhabad **2**: 39–44 (in russian).
- DERZHAVETS, J. A., IVANOV, A. I., MIRONOV, V. G., MISTSHENKO, O. A., PRASOLOV, V. N. & S. J. SINEV (1986): Spisok cheshuekrylykh (Macrolepidoptera) Leningradskoi oblasti [A list of Macrolepidoptera of the Leningrad region]. Fauna cheshuekrylykh (Lepidoptera) SSSR [A fauna of Lepidoptera of the USSR]. – Leningrad: "Nauka". Leningr. Dept., p. 186–270. (Trudy Vsesoyuz. Entomol. Obshch., T. **67**) (in russian).
- DIDMANIDZE, E. A. (1978): Cheshuekrylie aridnykh landshaftov Gruzii [Lepidoptera of the arid landscapes of Georgia]. – Tbilisi: "Me sniereba", 319 pp. (in russian).
- DRAUDT, H. (1931): 4. Family: Arctiidae. In: SEITZ, A. The Palaearctic Bombyces and Sphinges. Div. I. Fauna Palaearctica. Suppl. to vol. 2. Stuttgart: Alfred Kernen, p. 61–94, t. 5–7.
- DUBATOLOV, V. V. (1984): *Borearctia* gen. n. – novyi rod dlya medveditsy *Callimorpha menetriesi* (Ev.) (Lepidoptera, Arctiidae) [*Borearctia* gen. n. – a new genus for the tiger moth *Callimorpha menetriesi* (Ev.) (Lepidoptera, Arctiidae)]. – Entomol. Obozr. **63** (2): 336–339 (in russian).
- DUBATOLOV, V. V. (1985a): Zametki po sistematike medvedits (Lepidoptera, Arctiidae) gruppy roduv *Diacrisia* Hb. – *Rhyparia* Hb. [On the systematics of the tiger moth (Lepidoptera, Arctiidae) genus group *Diacrisia* Hb. – *Rhyparia* Hb.] In: Sistematiika i biologiya chlenistonogikh i gelmintov [Systematics and biology of Arthropoda and helminths]. – Novosibirsk: "Nauka" Sib. Dept., p. 66–71 (in russian).
- DUBATOLOV, V. V. (1985b): Vysshye medveditsy (Lepidoptera, Arctiidae: Arctiinae) gor Yuzhnoi Sibiri. Soobstshenie 1 [Arctiinae (Lepidoptera, Arctiidae) of the South Siberian mountains. Part 1]. In: Chlenistonogie Sibiri i Dalnego Vostoka [Arthropoda of Siberia and the Far East]. – Novosibirsk: "Nauka", Sib. Dept., p. 134–159 (in russian).
- DUBATOLOV, V. V. (1985c): *Rhyparioides metelkana* (LEDERER, 1861) v Sibiri [*Rhyparioides metelkana* (LEDERER, 1861) in Siberia]. – Vestnik zoologii, Kiev **5**: 86 (in russian).
- DUBATOLOV, V. V. (1985d): Notes on the *Holoarctia cervini* complex (Lepidoptera, Arctiidae). 1. *Holoarctia marinae* spec. nov. from Altai. – Ann. Ent. Fenn. **51**: 57–58.

- DUBATOLOV, V. V. (1987a): K sistematike roda *Micractia* SEITZ, s. l. (Lepidoptera, Arctiidae) [On the systematics of the genus *Micractia* SEITZ, s. lat. (Lepidoptera, Arctiidae)]. In: Nasekomye, klestshi i gelminty [Insects, mites and helminths]. – Novosibirsk: "Nauka" Sib. Dept., p. 30–47 (in russian).
- DUBATOLOV, V. V. (1987b): *Arctia rueckbeili* PÜNGELER (Lepidoptera, Arctiidae) v faune SSSR [*Arctia rueckbeili* PÜNGELER (Lepidoptera, Arctiidae) in the fauna of the USSR]. – Vestnik zoologii, Kiev 3: 87 (in russian).
- DUBATOLOV, V. V. (1988): Obzor vidov roda *Chelis* RBR. (Lepidoptera, Arctiidae) fauny SSSR [A review of the species of the genus *Chelis* RBR. (Lepidoptera, Arctiidae)]. In: Taksonomiya zhivotnykh Sibiri [Taxonomy of animals of Siberia]. – Novosibirsk: "Nauka", Sib. Dept., p. 80–98 (in russian).
- DUBATOLOV, V. V. (1989): Reviziya roda *Axiopoena* (Lepidoptera, Arctiidae) [A revision of the genus *Axiopoena* (Lepidoptera, Arctiidae)]. – Vestnik zoologii, Kiev 1:8–13 (in russian).
- DUBATOLOV, V. V. (1990a): Novye taksony vysshikh medvedits (Lepidoptera, Arctiidae: Arctiinae) Palearktiki [New taxa of Arctiinae (Lepidoptera, Arctiidae) from the Palearctic]. In: Redkie gelminty, klestshi i nasekomye [Rare helminths, mites and insects]. – Novosibirsk: "Nauka". Sib. Dept., p. 79–86 (in Russian).
- DUBATOLOV, V. V. (1990b): Novyi dlya fauny SBSR vid medvedits *Carcinopyga lichenigera* C. et R. FELDER, 1874 (Lepidoptera, Arctiidae) [A new species of tiger moth for the USSR fauna: *Carcinopyga lichenigera* C. et R. FELDER, 1874 (Lepidoptera, Arctiidae)]. – Vestnik zoologii, Kiev 2: 87 (in russian).
- DUBATOLOV, V. V. (1990c): Novye taksony vysshikh medvedits Palearktiki. Soobstshenie 2 [New taxa of Arctiinae from the Palearctic. Note 2]. In: Taksonomiya nasekomykh i gelmintov [Taxonomy of insects and helminths]. – Novosibirsk: "Nauka", Sib. Dept., p. 89–101 (in russian).
- DUBATOLOV, V. V. (1990d): Vysshye medveditsy (Lepidoptera, Arctiidae: Arctiinae) gor Yuzhnoi Sibiri. Soobstshenie 2 [Arctiinae (Lepidoptera, Arctiidae) of the South Siberian mountains. Part 2]. In: Chlenistonogie i gelminty [Arthropoda and helminths]. – Novosibirsk: "Nauka", Sib. Dept., p. 139–169 (in Russian).
- DUBATOLOV, V. V. (1990e): Novyi rod semeistva Arctiidae (Lepidoptera) [A new genus of the Family Arctiidae (Lepidoptera)]. In: Novosti faunistiki i sistematiki [News of faunistic and systematic]. – Kiev, Naukova dumka, p. 89–90.
- DUBATOLOV, V. V. (1991a): Moths from Southern Sakhalin and Kunashir, collected in 1989. Part 1. Macroheterocera, excluding Geometridae and Noctuidae – Japan Heteroc. J. 161: 182–187.
- DUBATOLOV, V. V. (1991b): Barsovaya medveditsa, *Tancrea pardalina* PÜNL. (Lepidoptera, Arctiidae: Arctiinae) – novyi vid dlya fauny Turkmenistana [The Pardalina tiger moth, *Tancrea pardalina* PÜNL. (Lepidoptera, Arctiidae: Arctiinae) – a new species for the Turkmenistan fauna]. – Izvestiya AN Turkm. SSR. Seriya biol. nauk. Ashabad, No. 5 (in russian).
- DUBATOLOV, V. V. ([1994] 1993–1994): Tiger-moths (Lepidoptera, Arctiidae: Arctiinae) of the former USSR: the distribution throughout countries and regions – Actias (Moscow) 1(1–2): 11–17.
- DUBATOLOV, V. V. & J. A. TSHISTJAKOV (1989): Novyi vid roda *Palearctia* (Lepidoptera, Arctiidae) – interesnyi primer konvergentnogo skhodstva u medvedits [A new species of the genus *Palearctia* (Lepidoptera, Arctiidae) – an interesting example of convergent similarity in tiger moths]. – Zool. Zhurnal 68 (11): 141–143 (in russian).
- DUBATOLOV, V. V. & G. S. ZOLOTARENKO (1990): K faune medveditseobraznykh cheshuyekrylykh (Lepidoptera: Nolidae, Arctiidae) Zapadno-Sibirskoi ravniny [On the fauna of tiger moths (Lepidoptera: Nolidae, Arctiidae) of the West Siberian plain]. In: Chlenistonogie i gelminty [Arthropoda and helminths]. – Novosibirsk, p. 122–139 (in russian).
- DUPONCHEL, PH. (1836): Supplement à l'histoire naturelle des Lépidoptères d'Europe. Paris, t. 3, 651 p., 50 t.
- DYAR, H. G. (1899): Notes on Alaskan Arctiidae. – Entomol. News 10: 130–131.
- DYAR, H.G. (1901): Diagnosis of a new arctian. – Jl. N. Y. ent. Soc. 9: 85.
- EFEFOTOV, K. A. & J. I. BUDASHKIN (1987): Medveditsy (Lepidoptera, Arctiidae) Kryma [Tiger moths (Lepidoptera, Arctiidae) of the Crimea]. – Vestnik zoologii, Kiev 4: 77–78 (in russian).

- EFETOV, K. A. & J. I. BUDASHKIN (1990): Babotchki Kryma [Lepidoptera of the Crimea]. – Simferopol: "Tavria", 112 pp. (in russian).
- EICHWALD, E. (1830): Zoologia specialis quam expositis animalibus tum vivis, tum fossilibus postissimum Rossiae in Universum, et Poloniae in specie, in usum lectionum publicarum in Universitate Caesarea Vilnensi. Pars alterna. – Vilnae, 323 pp.
- ERSCHOFF, N. (1872): Diagnoses de quelques espèces nouvelles de Lépidoptères appartenant à la faune de la Russiae Asiaticae. – Horae Soc. Entomol. Ross. **8**: 315–318.
- ERSCHOFF, N. (1874): Cheshuekrylye (Lepidoptera). In: FEDTSCHENKO, A., Puteshestvie v Turkestan [Travel to Turkestan]. **2**(5), dept. **3**, 4+6+128 p., 4 pl.
- EVERSMANN, E. (1843): Quaedam lepidopterorum species novae in montibus Uralensibus et Altaicus habitantes nunc descriptae et depictae. – Bull. Soc. Imp. Nat. Mosc. **16**(3): 535–553.
- EVERSMANN, E. (1846): Lepidoptera quaedam nova in Rossia observata. – Bull. Soc. Imp. Nat. Mosc. **19**(3): 83–89.
- EVERSMANN, E. (1847): Lepidoptera quaedam nova Rossiae et Sibiriae indigena descriptis et delineavit. – Bull. Soc. Imp. Nat. Mosc. **20**(3): 66–83, 6 t.
- EVERSMANN, E. (1848): Beschreibung einiger neuer Falter Russlands. – Bull. Soc. Imp. Nat. Mosc. **21**(3): 205–232.
- FALKOVITSH, M. I. (1969): Cheshuekrylye (Lepidoptera) gor Kokshetau i Zharkol-Shoindykolskogo plato [Lepidoptera of the Kokshetau mountains and the Zharkol-Shainsky plateau]. In: Rastitelnye soobstshestva i zhivotnoe naselenie stepei i pustyn Tsentralnogo Kazakhstana [Plant associations and animal populatins of steppes and deserts of Central Kazakhstan]. – Lenigrad: "Nauka". Leningradsk. Dept, p. 444–468 (in russian).
- FALLOU, M. J. (1864): Description d'un nouveau Lépidoptère Hétérocère du genre *Nemeophila* STPH. – Ann. Soc. Ent. France, **4**(4): 23–26, t. 1.
- FANG CHENG-LAI (1981): Distinctions between the fall webworm and *Spilosoma* spp. – Acta entomol. Sinica **24**(3): 340–341 (in chinese).
- FANG CHENG-LAI (1982): Arctiidae. – Iconographia heterocerorum sinicorum **2**: 190–277 (in chinese).
- FERGUSON, D. C. (1984): Two new generic names for groups of holartic and palaearctic Arctiini (Lepidoptera, Arctiidae). – Proc. entomol. Soc. Wash. **86** (2): 452–459.
- FREINA, J. J. DE (1983): 4. Beitrag zur systematischen Erfassung der Bombyces- und Sphinges-Fauna Kleinasiens. Neue Kenntnisse über Artenspektrum, Systematik und Nomenklatur sowie Beschreibungen neuer Taxa (Lepidoptera). – Mitt. Münch. Ent. Ges. **72**: 57–127.
- FREINA, J. J. DE & T. J. WITT (1984): Taxonomische Veränderungen bei den Bombyces und Sphinges Europas und Nordwestafrikas. Zwei neue Arctiidaegattungen für den westpaläarktischen Faunenbereich: *Watsonarctia* gen. n. und *Maurica* gen. n. (Lepidoptera, Artciidae). – Entomofauna **5** (28): 323–334.
- FREINA, J. DE & T. WITT (1987): Die Bombyces und Sphinges der Westpalaearktis. Band 1. – München, p. 333–380.
- FREYER, C. (1842): Neue Beiträge zur Schmetterlingskunde. Bd. **4**. – Augsburg, 167 S., 96 T.
- FRIVALDSZKY, J. (1835): A Balkány vidékén tett természettudományi utazásról. – A Magyar Tudós Társaság Évkönyvei, Buda, t. **2**.
- GRAESER, L. (1888): Beiträge zur Kenntnis der Lepidopteren-Fauna des Amurlandes. – Berl. Entomol. Ztg. **32**: 33–153, 309–414.
- GRAESER, L. (1892): Beiträge zur Kenntnis der Lepidopteren-Fauna des Amurlandes. – Berl. Entomol. Ztg. **37**: 209–234.
- GROSSER, N. (1983): Zur Lepidopteren-Fauna (Macrolepidoptera) Baschkiriens. – Wiss. Z. Univ. Halle **32**(1): 11–21.
- GROUM-GRSHIMAILO, GR. (1890): Le Pamir et la faune lépidoptérologique. – Mém. lépidop. Ed. N. M. ROMANOFF **4**, 576 pp., 21 pls.
- GRUM-GRSHIMAILO, GR. (1891): Lepidoptera nova in Asia centrali novissime lecta et descripta. – Horae Soc. ent. Ross. **25**: 445–465.
- GRUM-GRSHIMAILO, GR. (1899): Lepidoptera nova vel rarum cognita regionis palaearcticae. I. – Ezhegod. Zool Mus. Imp. Akad. Nauk [Ann. Mus. Zool. Acad. Imp. Sc.] **4**: 455–472 (in russian).

- GRUM-GRSHIMAILO, GR. (1902): Lepidoptera nova vel rarum cognita regionis palaearcticae. II. – Ezhegod. Zool. Mus. Imp. Acad. Nauk [Ann Mus. Zool. Acad. Imp. Sc.] 7: 191–204 (in russian).
- HAMPSON, G. (1901): Catalogue of the Arctiidae (Arctiana) and Agaristidae in the collection of the British Museum (Natural History). – Catalogue of the Lepidoptera Phalaenae in the collection of the British Museum (Natural History). London, vol. 3, XII+609 pp., pls. 36–54.
- HEYNE, A. (1899): Neue und wenig bekannte melanistische Formen europäischer Grossschmetterlinge. – Soc. Ent. 14: 97–98, 105–106.
- HORI, H. (1926): *Callimorpha menetriesi* Ev. from Saghalien. – Kontyu 1: 86–87.
- INOUE, H. (1961): Check list of the Lepidoptera of Japan. Part 6: Hyblaeidae-Sphingidae. – Tokyo: Rikusuisha, p. 621–683.
- INOUE, H. (1976): Some new and unrecorded moths belonging to the Bombyces and Sphinges from Japan (Lepidoptera). – Bull. Fac. dormest. Sci. Otsuma Wom. Univ. 12: 153–196.
- INOUE, H. (1982): 68. Arctiidae. In: Moths of Japan, vol. 1: 638–659; vol. 2: 74–79, 136–137, 334–342, pl. 154–162, 346–348.
- KAISILA, J. (1947): Die Makrolepidopterenfauna des Aunus-Gebietes. – Acta ent. Fenn. 1: 1–112.
- KARDAKOFF, N. (1928): Zum Kenntnis der Lepidopteren des Ussuri-Gebietes. – Entomol. Mitt. 17(8): 414–425.
- KOÇAK, A. Ö. (1980): Some notes on the nomenclature of Lepidoptera – Communications Fac. Sci. Univ. Ankara (C.) 24(2): 7–25.
- KÔDA, N. (1987/1988): A Generic Classification of the Subfamily Arctiinae of the Palaearctic and Oriental Regions based on the Male and Female Genitalia (Lepidoptera, Arctiidae). Part I. – Tyô to ga 38(3)(1987): 153–237. Part II. – Tyô to Ga 39(1)(1988): 1–79.
- KOLENATI, F. (1846): Lepidoptera. In: Meletemata entomologica. V. Insecta Caucasi. – Petropoli, 1846, p. 80–112, t. 18.
- KÖNIG, F. (1985): Date noi privind biologia speciei *Diacrisia (Rhyparioides) metelkana* LED. (Lepidoptera, Arctiidae). In: Delta Dunarii. II. – Studii și comunicări de entomologie, 1983, Tulcea, p. 87–90.
- KONOVALOVA, Z. A. (1968): K faune babochek (Heterocera) Kurilskikh ostrovov [On the fauna of Heterocera of the Kuril Islands]. In: Fauna i ekologiya nasekomykh Dalnego Vostoka [Fauna and ecology of insects of the Far East]. – Vladivostok, p. 28–41 (in russian).
- KOSTJUK, I. YU. & M. I. GOLOVUSHKIN (1994): K izucheniyu raznousykh cheshuekrylykh (Lepidoptera, Macroheterocera) Vostochnogo Zabaikal'ya [A contribution to the knowledge of the Macroheterocera (Lepidoptera) of [East] Transbaikalia]. In: Cheshuekrylye Zabaikal'ya [The Lepidoptera of Transbaikalia]. – Kiev, p. 51–57 (in russian).
- KOSTJUK, I. YU. & I. G. PIJUSTSH (1987): Novyi dlya fauny evropeiskoi chasti SSSR vid medvedits – *Phragmatobia placida* FRIV. (Lepidoptera, Arctiidae) [A new species of tiger moths for the European part of the USSR fauna – *Phragmatobia placida* FRIV. (Lepidoptera, Arctiidae)]. – Vestnik zoologii, Kiev, 6: 58.
- KOSHANTSCHIKOV, W. (1923): Materialy k faune cheshuekrylykh Minusinskogo kraja (Sibir, Eniseiskaya guberniya) [Materials to the fauna of Lepidoptera of the Minusinsk region (Siberia, Eniseiskaya guberniya)]. – Ezhegod. Gos. Muz. im. N. A. Mart'yanova [Jahrb. Martjanov. Staatsmus. Minussinsk] 1(1): 2–50 (in russian).
- KOSHANTSCHIKOV, W. (1924): Materialy k faune cheshuekrylykh Minusinskogo kraja, II. [Materials to the fauna of Lepidoptera of the Minusinsk region, II]. – Ezhegod. Gos. Muz. im. N. A. Mart'yanova [Jahrb. Martjanov. Staatsmus. Minussinsk] 2(1): 66–75 (in russian).
- KOSHANTSCHIKOV, W. (1925): Materialy k faune cheshuekrylykh Minusinskogo kraja, III. [Materials to the fauna of Lepidoptera of the Minusinsk region, III]. – Ezhegod. Gos. Muz. im. N. A. Mart'yanova [Jahrb. Martjanov. Staatsmus. Minussinsk] 3(1): 66–83 (in russian).
- KRULIKOVSKY, L. (1904): Notice sur la chasse des Lépidoptères durant l'été 1903 dans le district d Ourjoum, gouv-t de Viatka. – Russ. Entomol. Obozr. 4(1): 27–31 (in russian).
- KRULIKOVSKY, L. (1909): Cheshuekrylye Vyatskoi gubernii [Lepidoptera of Vyatskaya guberniya]. – Materialy k poznaniyu fauny i flory Rossiskoi imperii. Otdel zool. [Materials to the knowledge on fauna and flora of the Russian Empire. Zool. dep.] 9: 48–250 (in russian).

- KUMAKOV, A. P. & J. P. KORSHUNOV (1979): Cheshuekrylye Saratovskoi oblasti [Lepidoptera of the Saratov region]. – Saratov, 240 pp. (in russian).
- KUZNETSOV, V. I. (1960): Materialy po faune i biologii cheshuekrylykh (Lepidoptera) Zapadnogo Kopetdaga [Materials to the fauna and biology of Lepidoptera of West Kopetdagh]. In: Fauna i ekologiya nesekomykh Turkmenskoi SSR [Fauna and ecology of insects of the Turkmenian SSR]. – Trudy Zool. Inst. AS USSR **27**, Moscow-Leningrad: AS USSR, p. 11–93 (in russian).
- KUZNETSOV, V. I. & E. F. MARTYNOVA (1954): Spisok cheshuekrylykh raiona srednego tetseniya r. Ural [A list of Lepidoptera of the middle course of the Ural riv.]. – Trudy Zool. Inst. AN SSSR **16**: 321–350 (in russian).
- LEDERER, J. (1853): Lepidopterologisches aus Sibirien. – Verh. Zool.-Bot. Ges. Wien **3**: 351–386, T. 1–7.
- LEDERER, J. (1855): Beitrag zu einer Schmetterlingsfauna von Cypern, Beirut und einem Theil Kleinasiens. – Verh. Zool.-Bot. Ges. Wien **5**: 177–234, T. 3.
- LEDERER, J. (1861): *Nemeophila metelkana* n. sp. – Wiener Entomol. Monatschr. **5** (5): 162–163, t. 3.
- LÉNEK, O. (1966): Österreichische Iran-Afghanistan-Expeditionen. Beiträge zur Lepidopteren-Fauna, Theil 7. *Phragmatobia fuliginosa* L. ssp. *nova paghmani*. – Zeit. Wien. Entomol. Ges. **51** (8): 105–106.
- LERAUT, P. (1985): *Eucastana* nomen. nov. pour *Eucharia* au sens des auteurs (Lep. Arctiidae). – Linneana Belg. **10**: 43.
- LINNAEUS, C. (1758): [Lepidoptera]. In: *Systema Naturae*, etc. Ed. 10. – Holmiae, p. 458–542.
- LINNAEUS, C. (1761): [Lepidoptera]. In: *Fauna Suecica*, etc., ed. 2. – Holmiae, p. 267–371.
- LINNAEUS, C. (1767): [Lepidoptera]. In: *Systema Naturae*, etc. Ed. 12. – Holmiae, p. 744–900.
- LAVROV, S. D. (1927): Materialy k izucheniyu entomofauny okrestnosti Omska [Materials to a study of the entomofauna of the Omsk district]. – Trudy Sib. Inst. sel. khoz. i lesov. **8** (3): 51–100 (in russian).
- LVOVSKY, A. L. (1971): Materialy po faune cheshuekrylykh (Macrolepidoptera) Astrakhanskoi oblasti [Materials on the fauna of Macrolepidoptera from the Astrakhan Region]. – Entomol. Obozr. **50** (4): 800–810 (in russian).
- MATSUMURA, S. (1927a): New species and subspecies of moths from the Japanese Empire. – J. Coll. Agric. Hokkaido Imp. Univ. **19**: 1–91.
- MATSUMURA, S. (1929): New moths from Kuriles. – Insecta matsum. **3** (4): 165–168.
- MATSUMURA, S. (1930a): New species and forms of Arctiidae from Japan. – Insecta Matsumur. **5**: 31–40, pl. 1.
- MATSUMURA, S. (1930b): A catalogue of the Arctiidae of the Japanese Empire. – Insecta Matsumur. **5**: 58–94, pl. 1.
- McDUNNOUGH, J. (1921): Note on the generic position of two Canadian arctiids with descriptions of new species (Lepid.). – Can. Entomol. **53**: 167–168.
- MÉNÉTRIÈS, E. (1832): Catalogue raisonné des objets de zoologie recueillis dans un voyage au Caucase et jusqu'aux frontières de la Perse. – St.-Pétersbourg, 271 p.
- MÉNÉTRIÈS, E. (1842): Sur un nouveau genre de lépidoptère nocturne de la Russie. – Bull. phys.-mat. Acad. Pétersb. **9** (195): 40–43.
- MÉNÉTRIÈS, E. (1859a): Lépidoptères de la Sibérie orientale et en particulier des rives de l'Amour. – Bull. Acad. Sci. St. Pétersb. **17**: 212–221.
- MÉNÉTRIÈS, E. (1859b): Sur quelques Lépidoptères du gouvernement de Jakoutsk. – Bull. Acad. Sci. St. Pétersb. **17**: 494–500.
- MÉNÉTRIÈS, E. (1863): Descriptions de nouvelles espèces de Lépidoptères de la collection de l'Academie impériale des Sciences. 3ème et dernière partie. In: MÉNÉTRIÈS, E., *Enumeratio corporum animalium Musei Imp. Acad. Sci. etc. Petropolitane. Pars 3*, Lep. Heterocera. – St.-Pétersbourg, p. 145–161, t. 15–18.
- MERZHEEVSKAJA, O. I., LITVINNOVA, A. N. & R. V. MOLCHANNOVA (1976): Cheshuekrylye (Lepidoptera) Belorusii (katalog) [Lepidoptera of Belarus (catalogue)]. – Minsk: "Nauka i tekhnika", 132 pp.
- MIRZOYAN, S. A. & KH. M. ARUTYUNYAN (1980): Medveditsy Armenii (Lep. Arctiidae). – Trudy NII Inst. Zastsh. Rast. Erevan **4**: 96–104 (in russian).

- MOLTRECHT, A. K. (1929): O geograficheskom rasprostranenii cheshuekrylykh Dalnevostochnogo kraya s vydeleniem v osobuyu faunu ussuriiskikh Lepidoptera [Ueber die geographische Verbreitung der Macrolepidopteren des Ussuri- und Amur-Gebietes]. – Zapiski Vadvostokskogo otdela Gosudarstvennogo Russkogo Geograficheskogo obstshestva **3** (XX): 5–70.
- MOORE, F. (1879): Description of new genera and species of Asiatic Lepidoptera Heterocera. – Proc. Zool. Soc. Lond.: 387–417, t. 32.
- MOTSCHULSKY, V. (1860 [1861]): Insects du Japon. – Et. d'Ent. **9**: 4–39.
- NAGEL, W. P. & D. L. ISAACSON (1974): *Tyria jacobaeae* and Tansy Ragwort in Western Oregon. – Journ. Econ. Ent. **67** (4): 494–496.
- NIEPELT, W. (1911): Neue paläarktische Macrolepidopteren. – Int. Ent. Ztg. **5**: 274–275.
- OBERTHÜR, CH. (1879): Diagnoses d'espèces nouvelles de Lépidoptères de l'ile Askold. – Rennes, 16 pp.
- OBERTHÜR, CH. ([1881]): Faune des Lépidoptères de l'ile Askold. Première partie. – Etud. d'Ent. **5**: IX-X, 11–88, pls. 1–9.
- OBERTHÜR, CH. (1896): De la variation chez les Lépidoptères. – Etud. d'Ent. **20**, 20+74 pp., 24 t.
- OBRATZSOV, N. S. (1936): Zur Lepidopterenfauna des Südlichen Transdneprgebietes. – Festschrift für Dr. EMBRIK STRAND **2**: 229–242.
- PAYKULL, G. (1793): Beskrivelse over 5 arter nye nat-sommerfulge. – Skrivi. Naturh. Selsk. Kjøbenhavn **2**: 97–102, t. 2.
- PLJUSTSH, I. G. & I. Y. KOSTJUK (1988): *Acymba spectabilis* TAUSCH. (Lepidoptera, Arctiidae) – novaya medveditsa v faune Ukrayny [*Acymba spectabilis* TAUSCH. (Lepidoptera, Arctiidae) – a new tiger moth for the fauna of the Ukraine]. – Vestnik zoologii, Kiev **4**: 86 (in russian).
- PODA, N. (1761): Insecta musei Graecensis, quae in ordines, genera et species juxta Systema Naturae Caroli Linnaei digessit NICOLAUS PODA, e Societate Jesu, Philosophiae Doctor et Mattheeos Professor. Graecii, Widmanstadii, 18+127 pp., 2 tab.
- PÜNGELE, R. (1898): Diagnosen neuer Lepidopteren aus Centralasien. – Soc. Ent. **13**: 57–58.
- PÜNGELE, R. (1899): Neue Macrolepidopteren aus Central-Asien. – Dtsch. Ent. Zeit. IRIS **12**: 95–106.
- PÜNGELE, R. (1901): Neue Macrolepidopteren aus Centralasien. – Dtsch. Ent. Zeit. IRIS **14**: 171–191, T. 1–3.
- PÜNGELE, R. (1904): Neue Macrolepidopteren aus Centralasien. – Soc. Entomol. **19**: 121–122, 129–131.
- PÜNGELE, R. (1918): [6. *Arctia wagneri* PÜNG. nov. spec.]. – Zeit. Öster. Ent. Ver. **3**: 46.
- QUENSEL, C. (1802): Lepidoptera. In: ACERBI, J., Travels through Sweden, Finland and Lapland, to the North Cape 1798 & 1799. T. 2. – London.
- REBEL, H. (1910): *Arctia caja confluens*. In: BERGE, F., Schmetterlingsbuch, etc. 9. Aufl. – Stuttgart, S. 430.
- REICH, P. (1935): *Callimorpha dominula* L. (Lepidopt. Callimorphinae). – Ent. Rundschau **53** (1): 8–10.
- REMM, H. & J. VIIDALEPP (1986): Catalogus Macrolepidopterorum Estoniae. – Abiks Loodusevaatlejale **84**, Tartu, 1986, 34 pp. (in Estonian).
- ROMANOFF, N. M. (1884): Les Lépidoptères de la Transcaucasie. Première partie. – Mém. lépidop., Ed. N. M. ROMANOFF **1**: 1–92, pl. 1–5.
- ROTHSCHILD, W. (1910): Catalogue of the Arctianae in the Tring museum, with notes and descriptions of new species. – Novitat. Zool. **17**: 1–85, 113–188.
- ROTHSCHILD, W. (1914): Arctiidae, subfamilie Arctiinae. In: SEITZ, A. Die Gross-Schmetterlinge der Erde. Band **10**. – Stuttgart, Alfred Kernen, S. 236–263.
- SACHKOV, S. A. (1992): Fauna vysshikh medvedits (Lepidoptera: Arctiidae, Arctiinae) Samarskoi Luki i sopredelnykh territorii Samarskoi oblasti [The tiger-moth fauna (Lepidoptera: Arctiidae, Arctiinae) of the "Samarskaya Luka" and neighbouring territories of the Samara region]. – Samarskaya Luka. Bulletin **3**: 136–144 (in russian).
- SCHAWERDA, K. (1922): Ussuriensia. – Zeit. österr. Ent. Ver. **7**: 10–11.
- SCHULZ, O. (1905): Ueber die Variabilität von *Pericallia matronula* L. – Ent. Zeit. **18** (31): 125–126 + Beilage.

- SCHWINGENSCHUSS, L. (1937): Weitere Neuheiten aus Nord Persien. – Zeit. Österr. entomol. Ver. **22** (6): 57–61.
- SEDYKH, K. R. (1974): Zhivotnyi mir Komi ASSR. Bespozvonochnye. [Animal world of the Komi ASSR. Invertebrates]. – Syktyvkar, 191 pp.
- SEDYKH, K. R. (1979): Cheshuekrylye (Lep. Macrolepidoptera) fauny Kamchatki i prilegayustshikh oblastei [Lepidoptera (Macrolepidoptera) of Kamchatka and neighbouring regions]. – Entomol. Obozr. **58** (2): 288–298 (in russian).
- SEITZ, A. (1910): 4. Familie: Arctiidae, Bärenspinner. In: SEITZ, A. Die Gross-Schmetterlinge der Erde. I Abt.: Die Gross-Schmetterlinge des Palaearktischen Faunengebietes. Band 2: Die Palaearktischen Spinner & Schwärmer. – Stuttgart: Alfred Kernen, S. 43–103, t. 10–18, 56.
- SHAPOSHNIKOV, CHR. G. (1904): Zametki o Macrolepidoptera tsentralnoi chasti Severo-Zapadnogo Kavkaza [Notes sur les Macrolépidoptères de la partie centrale du Caucase septentrional-occidental]. – Ezhegod. Zool. Mus. Imp. Akad. Nauk [Ann. Mus. Zool. Acad. Imp. Sci. St. Petersb.] **9**: 189–259 (in russian).
- SHAROV, A. A. & S. S. IZHEVSKIY (1987): Kompleks parazitov amerikanskoi beloi babochki *Hyphantria cunea* DRURY (Lepidoptera, Arctiidae) na yuge Evropeiskoi chasti SSSR [The complex of parasites of the fall webworm *Hyphantria cunea* DRURY (Lepidoptera, Arctiidae) in the South of the European part of the USSR]. – Entomol. Obozr. **66** (2): 290–298 (in russian).
- SHELJUZHKO, L. (1918): Diagnoses lepidopterorum novorum Sibiriae. – Neue Beitr. syst. Insektenkunde **1**: 104.
- SHELJUZHKO, L. (1926): Neue palaearctische Heteroceren. – Deutsch. Ent. Zeit. IRIS **40**: 56–65.
- SHELJUZHKO, L. (1934): Eine neue *Callimorpha dominula*-Rasse aus dem Kaukasus. – Ent. Zeit. **48**: 73–75.
- SHELJUZHKO, L. (1935a): Eine neue Rasse von *Callimorpha quadripunctaria* PODA aus Transkaukasien. – Zeitschr. Öster. Ent. Ver. **20**: 21–22, 36, 67.
- SHELJUZHKO, L. (1935b): Einige neue und wenig bekannte Lepidopteren aus dem Westlichen Tianschan. – Mitt. Münch. ent. Ges. **25**: 27–38, t. 3.
- SHELJUZHKO, L. (1941): Materialy do lepidopterofauny Kyivstshiny [Materials to the lepidoptera fauna of the Kiev region]. Bombyces i Sphinges. I. – Trudy Zool. Muz. Kyiv. derzh. un-tu **1**: 1–101 (in ukranian).
- SHLYKOV, O. V. (1988): Spisok cheshuekrylykh (Macrolepidoptera) Penzenskoi obl. [A list of the Macrolepidoptera of the Penza province]. – Entomol. Obozr. **68** (1): 48–61.
- SITNOKOV, P. S. (1992): K sozdaniyu regionalnogo kadastra redkikh nasekomykh Tyumenskoi oblasti [To the creation of the Regional Rare Insects Register of the Tyumen region]. – Ezhegodnik Tyumenskogo oblastnogo kraevedcheskogo muzeya, Tyumen, p. 200–228 (in russian).
- SKALSKI, A. W. (1988): *Stauropolia nekrutenkoi* gen. et spec. nov. (Lepidoptera, Arctiidae) – novaya iskopaemaya medveditsa iz miotsena Kavkaza [*Stauropolia nekrutenkoi* gen. et spec. nov. (Lepidoptera, Arctiidae) – a New Fossil Tiger-Moth from the Miocene of Caucasus]. – Vestnik zoologii, Kiev **4**: 21–25 (in russian).
- SOTAVALTA, O. (1963): The generic position of *Hyphoraia alpina* QUENS. (Lep. Arctiidae). – Suom. hyont. Aikak. [Ann. Ent. Fenn.] **29**: 257–267.
- SOTAVALTA, O. (1965): A revision of the genus *Hyphoraia* HÜBNER, s. lat. (Lepidoptera, Arctiidae). – Suom. hyont. Aikak. [Ann. Ent. Fenn.] **31** (3): 159–197
- SOVINSKY, V. (1905): Tri novykh kavkazskikh raznovidnosti cheshuekrylykh [Trois nouvelles formes de Lepidopteres du Caucase]. – Russ. entomol. Obozr. **5** (3–4): 108–110 (in russian).
- SPULER, A. (1906): XXIII. Fam. Arctiidae STPH. In: SPULER, A., Die Schmetterlinge Europas. Band 2. – Stuttgart, S. 128–152.
- STAUDINGER, O. (1861): [Macrolepidoptera]. In: STAUDINGER, O. & M. WOCKE, Catalog der Lepidopteren Europas und der angrenzenden Länder. – Dresden, XVI+192 S.
- STAUDINGER, O. (1867): Einige neue Lepidopteren (sämtlich aus der Sammlung des verstorbenen O. GRÜNER). – Stett. ent. Ztg. **28**: 100–110.
- STAUDINGER, O. (1871): [Macrolepidoptera]. In: STAUDINGER, O. & M. WOCKE, Catalog der Lepidopteren des europäischen Faunengebietes (Edn. 2). – Dresden, 38+426 S.

- STAUDINGER, O. (1879): Lepidopteren-Fauna Kleinasiens. – Horae Soc. ent. Ross. **14**: 176–482.
- STAUDINGER, O. (1886): Description of a new *Bombyx* allied to *Arctia caja*. – Ent. Month. Mag. **22**: 258.
- STAUDINGER, O. (1887a): Centralasiatische Lepidopteren. – Stett. ent. Ztg. **48**: 49–102.
- STAUDINGER, O. (1887b): Neue Arten und Varietäten von Lepidopteren aus dem Amur-Gebiete. – Mém. lépidop., Ed. N. M. ROMANOFF **3**: 126–232, T. 6–12, 16–17
- STAUDINGER, O. (1891 [1892]): Neue Arten und Varietäten von Lepidopteren des paläarktischen Faunengebiets. – Dtsch. Ent. Zeit. IRIS **4**: 224–339, T. 3.
- STAUDINGER, O. (1892a): Lepidopteren des Kentei-Gebirges. – Dtsch. Ent. Zeit. IRIS **5**: 300–393.
- STAUDINGER, O. (1892b): Die Macrolepidopteren des Amurgebiets. I. Theil. Rhopalocera, Sphinges, Bombyces, Noctuae. – Mém. lépidop., Ed. N. M. ROMANOFF **6**: 83–658, Pl. 4–14.
- STAUDINGER, O. (1896): Drei neue paläarctische Heteroceren. – Dtsch. Ent. Zeit. IRIS **9**: 188–192, t. 4.
- STAUDINGER, O. (1897): Lepidopteren des Apfelgebirges. – Dtsch. Ent. Zeit. IRIS **10**: 320–344, t. 9.
- STICHEL, H. (1911): Zweiter Beitrag zur nordischen Schmetterlingsfauna und anknüpfende Bemerkungen. – Berl. Ent. Zeit. **61**: 33–104, T. 3.
- STRAND, E. (1912): Ein neuen Bär von Tianschan. – Int. Ent. Zeit. (Guben) **6**: 2.
- STRAND, E. (1919): Arctiidae: subfam. Arctiinae. In: Lepidopterorum Catalogus. Pars **22**. – Berlin: W. Junk, 416 S.
- STSHETKIN, J. L. (1960): Vysshye cheshuekrylye Vakhshskoi doliny (Tadzhikistan). Chast' I. Lepidoptera. Rhopalocera i Heterocera [Macrolepidoptera of the Vakhsh valley (Tajikistan). Part I. Lepidoptera. Rhopalocera et Heterocera]. – Stalinabad, 103 pp. (in russian).
- STSHETKIN, J. L. (1975): Pastbistshnaya medveditsa (*Ocnogyna loewii pallidior* Chr.) v Srednei Asii (Lepidoptera: Arctiidae) [Pasturable tiger moth (*Ocnogyna loewii pallidior* Chr.) in Middle Asia (Lepidoptera: Arctiidae)]. – Entomologiya Tadzhikistana [Entomology of Tajikistan]. Dushanbe, p. 134–138 (in russian).
- STSHETKIN, J. L. (1982): Novyi podvid – *Arctia caja pamiroalaica* STSHETKIN, subspec. nov. – iz Srednei Asii (Lepidoptera: Arctiidae) [A new subspecies – *Arctia caja pamiroalaica* STSHETKIN, subspec. nov. – from Middle Asia (Lepidoptera: Arctiidae)]. – Izvestiya Akad. Nauk Tadzh. SSR. Otd. biol. nauk **1**: 39–43 (in russian).
- SULCS, A. & J. VIIDALEPP (1967): Verbreitung der Grossschmetterlinge (Macrolepidoptera) im Balticum. II. Spinnerartige und Schwärmer. – Deutsch. Entomol. Zeit., N.F. **14** (5): 395–431.
- SVORTSEV, M. (1894): Materialy k izucheniyu entomologicheskoi fauny Altaya i Semipalatinskoi oblasti. II [Materials to the study of the entomological fauna of the Altai and the Semipalatinsk region. II]. – Zapiski Zap.-Sib. otd. Imp. Rus. geogr. obstsh. **17** (3): 1–14 (in russian).
- SWINHOE, L.-C. (1885): List of Lepidoptera collected in Southern Afghanistan. – Trans. Entomol. Soc. Lond., part 3, p. 337–355.
- TAMS, W. (1922): A new *Callimorpha* from north-west Persia and Armenia. – Entomologist **55**: 196–197.
- TAUSCHER, A. (1806): Lepidopterorum Rossiae indigenorum observationes sex. – Mem. Soc. Nat. Mosc. **1**: 207–212, t. 1.
- TCHRUSHINA, A. N. & A. I. SHERNIN (1974): Glava 13. Otryad Lepidoptera – Cheshuekrylye [Part 13. Order Lepidoptera]. In: Zhivotnyi mir Kirovskoi oblasti [Animal world of the Kirov region]. Part 2. – Kirov [Vyatka], p. 351–477 (in russian).
- THOMAS, W. (1989): Bemerkungen zur Gattung *Carcinopyga* (Lep., Arctiidae) und zu *Carcinopyga lichenigera* FELDER (Beiträge zur Arctiiden-Systematik 4). – Atalanta **19**: 139–147.
- THOMAS, W. (1990): Die Gattung *Lemyra* (Lepidoptera, Arctiidae). – Nachr. entomol. Ver. Apollo. Suppl. **9**: 1–83.
- TILAVOV, T. (1978): K biologii pastbistshnoi medveditsy (*Ocnogyna loewii pallidior* Chr.) v usloviyakh Kashkadarinskoi oblasti [On the biology of the pasturable tiger moth (*Ocnogyna loewii pallidior*) in the conditions of Kashkadarinskaya oblast]. – Uzbekskii biol. zhurnal [Uzbekian biological journal] **3**: 61–62 (in russian).
- TORSTENIUS, S. (1971): *Orodemnias cervini* FALLOU ssp. *fridolini* n. ssp. (Lepidoptera, Arctiidae). – Entomol. Ts. Arg. **92** (3–4): 173–177

- TSCHETVERIKOV, S. (1904): Lepidoptera palearctica nova. – Russ. Entomol. Obozr. **4** (2–3): 77–79 (in russian).
- TSCHETVERIKOV, S. S. (1993): Babochki Gor'kovskoi oblasti [Lepidoptera of the Gorkii Region]. – Nizhni Novgorod: Nizhnii Novgorod State University Press. 128 pp. (in russian).
- TSHISTJAKOV, J. A. (1988): Medveditsy roda *Dodia* DYAR, 1901 (Lepidoptera, Arctiidae) fauny SSSR [Tiger moths of the genus *Dodia* DYAR, 1901 (Lepidoptera, Arctiidae) of the USSR fauna]. – Entomol. Obozr. **67** (3): 632–643 (in russian).
- TSHISTJAKOV, J. A. (1990): *Pararctia tundrana* sp.n. vikariant *P. subnebulosa* (DYAR, 1899) (Lepidoptera, Arctiidae) v subarkticheskym sektore Azii [Pararetia tundrana sp.n. – a vicariant of *P. subnebulosa* (DYAR, 1899) (Lepidoptera, Arctiidae) in the subarctic sector of Asia]. In: Novosti sistematiki nasekomykh Dalnego Vostoka [News on the systematic of insects of the Far East]. – Vladivostok: DVO AN SSSR, p. 97–99 (in russian).
- TSHISTJAKOV, Y. A. (1992): Sem. Arctiidae [Pam. Arctiidae]. In: Nasekomye Khiganskogo zapovednika. Chast 2 [Insects of the Khingan Nature Reserve. Part 2]. – Vladivostok: Dal'nauka, p. 149–154 (in russian).
- TSHISTJAKOV, Y. A. & J. D. LAFONTAINE (1984): A review of the genus *Dodia* DYAR (Lepidoptera: Arctiidae) with description of a new species from Eastern Siberia and Northern Canada. – Canad. Entomologist **116**: 1549–1556.
- VNUKOVSKY, W. (1929): 3. Einige Nomenklatur-Notizen über die paläarktischen Lepidopteren. – Zool. Anzeiger **83** (9/10): 221–224.
- VIIDALEPP, J. (1979): K faune cheshuekrylykh Tuvinskoi ASSR. II. Raznousye cheshuekrylyye (semeistva Zygaenidae-Cossidae) [On the fauna of Lepidoptera of the Tuva ASSR. II. Sphinges and Bombyces]. – Uchenye zapiski Tartuskogo gos. univ. [Acta et comm. Univ. Tartuensis] **483**: 17–39 (in russian).
- VIIDALEPP, J. & H. REMM (1982): Novye materialy o vysshikh cheshuekrylykh (Macrolepidoptera) Sakhalina [New material of Macrolepidoptera from Sakhalin]. In: Poleznye i vrednye nasekomye Sibiri [Useful and pest insects of Siberia]. – Novosibirsk: "Nauka". Sib. Dept., p. 137–151 (in russian).
- WAGNER, F. (1913): Neue Heteroceræ aus Centralasien. – Int. Ent. Zeit. **7** (1): 2–4.
- WALKER, P. (1855): List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. – London, part 3, p. 583–775.
- WALKER, F. (1856): List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. – London, part 7, p. 1509–1808.
- WALKER, F. (1865): List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. – London, part 31, supl., 321 pp.
- WARNECKE, G. (1918): Eine neue Lokalform von *Rhyptaria purpurata* L. – Intern. Entomol. Zeit. **12** (11): 81.
- WATSON, A., FLETCHER, D. S. & I. W. B. NYE (1980): The Generic Names of Moths of the World. – London: Trustees of the Brit. Mus. (Nat. Hist.), vol. **2**, 228 p.
- WILTSHERE, E. P. (1961): Ergebnisse der Deutschen Afghanistanexpedition 1956 der Landessammlungen für Naturkunde Karlsruhe. Middle East Lepidopteren XV. A second contribution to the Lepidoptera of Afghanistan. – Beitr. naturk. Forsch. SW-Dtschl. **19** (3): 337–371, pl.
- WITT, T. (1980): Bombyces und Sphinges aus Korea. I. (Lepidoptera: Bombycidae, Sphingidae). – Folia ent. Hung. **41** (1): 167–174.
- WITT, T. (1985): Bombyces und Sphinges (Lepidoptera) aus Korea, II. – Folia ent. Hung. **46** (2): 179–194.
- WOJTUSIAK, R. J. & W. NIESIOLOWSKI (1946): Lepidoptera of the Central Caucasus, collected during the Polish Alpine Expedition in 1935, with ecological and zoogeographical remarks. I Part. Macrolepidoptera. – Acta Mus. Hist. nat. Acad. polon., Crakow, **6**: 1–74.
- ZELLER, P. C. (1846): Beschreibung der *Trichosoma lœwii* n. sp. nebst Bemerkungen über CARRENNO's "Insecte, dont l'ordre est incertain". – Entomol. Ztg. **7**: 5–11.

Adress of the author:

V. V. DUBATOLOV

Zoological Museum, Institute of Animal Systematics and Ecology (former: Biological Institute)

Siberian Branch of the Russian Academy of Sciences

Frunze street, 11

Novosibirsk, 91, 630091

Russia.

Colour plate

1a	1b	1b
2		5
3		6
4		7

DUBATOLOV, V. V.: Three contributions to the knowledge of palearctic Arctiinae. 1. *Sinowatsonia*, a new genus of Arctiidae from Tibet (China) (on the systematics of the genus *Micrarczia* SEITZ s. l., Part 2) (Lepidoptera: Arctiidae). – Neue Ent. Nachr. **38**: 5–8.

Fig. 1a: *Sinowatsonia batangi* (DANIEL, 1943). Holotype ♀. "Batang (Tibet) Im Tal des Yangtze (ca. 2800 m) 26.7.1936 H. HÖNE". Photo by courtesy of Dr. A. WATSON.

Fig 1b: *Sinowatsonia hoenei* (DANIEL, 1943). The female from the collection of The Natural History Museum (London) and its labels. Photo by courtesy of Dr. A. WATSON.

DUBATOLOV, V. V.: Three contributions to the knowledge of palearctic Arctiinae. 2. A review of the genus *Palearctia* FERGUSON (on the systematics of the genus *Micrarczia* SEITZ s. l., Part 3) (Lepidoptera: Arctiidae). – Neue Ent. Nachr. **38**: 9–37.

Fig. 2: *Palearctia kashmirica* FERGUSON, 1985. Holotype. Now — *P. gratiosa kashmirica* FERG. Photo by courtesy of Dr. A. WATSON.

Fig. 3: *Arctia schottlaenderi* ab. *infrapicta* STRAND, 1912. Holotype. Now — *P. ferghana schottlaenderi* (STRAND). Photo by courtesy of Dr. A. WATSON.

Fig. 4. *Arctia schottlaenderi* STRAND, 1912. A water-colour figure of the holotype. Now — *P. ferghana schottlaenderi* (STRAND). Photo by courtesy of Dr. A. WATSON.


Fig. 5: *Arctia erschoffi issyka* STAUDINGER, 1887. Type. Now a synonym of *P. erschoffii erschoffii* (ALPH.).

Fig. 6: *Arctia erschoffi ferghana* STAUDINGER, 1887. Lectotype. Now *P. ferghana* (STGR.).

Fig. 7: *Micrarczia kindermannii marxi* O. BANG-HAAS, 1927. Holotype. Now *P. marxi* (O. B.-H.).



micrartia
seivereti, Gr-Gish.
Tunman
1918
G. Forrest
M.J. Mansfield
Coll.
B.M.1950-244.



stretia *schottlaenderi* *SYN.*
Thian-Shan Strand.



Demokratie in Deutschland oder Diktatur der Bürokraten

Jeder Verleger in Bayern ist gezwungen, zwei Freiemplare von jedem Druckerzeugnis kostenfrei an die Bayerische Staatsbibliothek in München und ein weiteres an die Deutsche Bibliothek in Frankfurt am Main zu senden. Dies habe ich als gerade noch tragbar empfunden. Seit der Vereinigung gibt es jetzt zwei nationale Bibliotheken: Eine in Frankfurt/Main und eine in Leipzig. Diese in Leipzig ist nun zusätzlich mit einem Pflichtexemplar zu beschenken. Dies halte ich für unannehmbar, untragbar und für Unrecht. Das gesamte deutsche Schrifttum kann durch eine "Deutsche Nationalbibliographie" schnell und zuverlässig angezeigt werden. Im Zeitalter der Datenverarbeitung und Computer kann auch von beiden Bibliotheken eine gemeinsame Datenbank aufgebaut und genutzt werden. Es existieren ja auch nicht gleichzeitig BND und STASI (oder doch?!) oder zwei Verteidigungsministerien nebeneinander. Statt mit Sparmaßnahmen voranzugehen, fordern Staat und Politiker nur, um sich gleichzeitig selbst maßlos zu bedienen. Gegen das Unrecht vorzugehen ist nun fast aussichtslos, schließlich leben wir in einem Rechtsstaat. Bevor Verfassungsbeschwerde erhoben werden kann, muß der Instanzenweg erschöpft sein, d.h. das Verwaltungsverfahren muß durchlaufen werden. Nach Erfolglosigkeit hat die Verwaltungsgerichtsbarkeit über die Rechtmäßigkeit der Ablieferungspflicht des Pflichtexemplares zu entscheiden. Erst dann ist eine Verfassungsbeschwerde möglich. Bis dahin wären allerdings Prozeßkosten von etwa DM 10 000,-- zu bezahlen. Das jedoch kann ich mir finanziell nicht leisten. Somit wird der Weg, Recht zu bekommen, blockiert, weil wir ja in einem Rechtsstaat leben.

Aus diesem Grund veröffentliche ich obige Zeilen solange, bis man mir verbietet, meine Meinung über diese Demokratie und diesen Rechtsstaat zu äußern, oder bis sich couragierte, vermögende Leute finden, die gegen das Unrechtsgesetz ankämpfen bis es zurückgenommen wird.

Democracy in Germany or dictatorial beaurocracy

Every publisher in Bavaria is made to give two free examples of each publication to the Bavarian State Library (Munich) and a further example to the German Library (Frankfurt/Main). This was just about acceptable for me.

Since the joining of East and West Germany, we now have two national libraries, one in Frankfurt and one in Leipzig. Now the library in Leipzig is to be additionally provided with one. This I find totally unacceptable and unjust. All German literary works could be quickly gathered together in the form of a "German National Bibliography". We live in the age of data processing and computer technology, and it would be possible to build up a data base from both libraries. The BND (Germany's "Secret Service") and the "STASI" (former East Germany's "Secret Service") do not exist side by side, or do they? Two Ministries of Defence also do not exist next to one another, so why two State Libraries? The State and politicians should be setting an example to the people, and yet all they seem to do is serve themselves.

Although we live in a state where the people has the rights, it is pointless to fight such injustice. Through the beaurocracy of our state and in our society, we must go from one stage to the next a bit like an obstacle course. Unless we have a good case it is pointless to spend appr. DM 10,000 required to get to the finish. The chance is there naturally for those with a good case and the finance, but who has this? Those that have the money would rarely attempt this anyway. Our democracy gives us the opportunity, but our beraucracy hinders our attempts. On these grounds I'm publishing the above statement until someone refuses to let me speak my mind or until someone with the finance makes it possible to fight the injustice.

HERBIPOLIANA

Buchreihe zur Lepidopterologie (Herausgeber: Dr. ULF EITSCHBERGER)

Bd. 1: EITSCHBERGER, U.: Systematische Untersuchungen am *Pieris napi-bryoniae*-Komplex (s.l.) (Pieridae). 1984. Textband: 504 S., Tafelband 601 S. mit 110 Farbtafeln. Ungewöhnlich umfangreiche und sehr gut ausgestattete Monographie. Die Farbtafeln zeigen die Tiere in Originalgröße. Format DIN A4. DM 520,— Subskriptionspreis DM 450,— (gilt bei Abnahme aller erscheinenden Teile).

Bd. 2: HACKER, H.: Die Noctuidae Griechenlands. Mit einer Übersicht über die Fauna des Balkanraumes. Die Arbeit behandelt alle 787 am Balkan vorkommenden Noctuidae-Arten. Die griechische Fauna mit 619 Arten wird detailliert dargestellt. Weiterhin erfolgt eine ausführliche Behandlung der Fauna Jugoslawiens (ohne Alpengebiete; 640 Arten), Albaniens (285 Arten), Bulgariens (621 Arten) und Rumäniens (südlicher Teil; 536 Arten) nach modernen taxonomischen und nomenklatorischen Gesichtspunkten. 37 Tafeln, davon 13 in Farbe. Verbreitungskarten für alle in Griechenland vorkommenden Arten. 590 S. Ganzzeilenband. DM 450,—

Bd. 3: LUKHTANOV, V. & A.: Die Tagfalter Nordwestasiens. Butterflies of North-West Asia. Die erste und sehr umfassende Darstellung der in diesem Gebiet Rußlands vorkommenden über 400 Tagfalterarten. Sehr viele neue oder bisher nie publizierte Daten auch zur Biologie. Das Standardwerk auf viele Jahre hinaus. Englische Zusammenfassung bei jedem Taxon. 51 Strichzeichnungen von Genitalpräparaten im Text, 19 Karten zur landschaftlichen Untergliederung des Gebietes, 52 Farbtafeln mit den vergrößert abgebildeten Arten, 4 Farbtafeln mit Biotopaufnahmen, 400 Verbreitungskarten. 440 S. Fadenheftung, fester Einband. DM 248,—

NEUE ENTOMOLOGISCHE NACHRICHTEN

Bd. 22/23: WOLF, W.: Systematische und synonymische Liste der Spanner Deutschlands unter besonderer Berücksichtigung der DENIS & SCHIFFERMÜLLERSchen Taxa (Lepidoptera: Geometridae). 5 Tabellen, 82 S., Kommentare in deutsch und englisch. ARBEITSGEEMEINSCHAFT NÖRDBAYERISCHER ENTOMOLOGEN: Prodromus der Lepidopterenfauna Nordbayerns. 1 Karte, 161 S., umfangreiche Tabellen und Verzeichnis der faunistischen Literatur Nordbayerns. 1988. DM 50,— (Doppelband)

Bd. 25: SCHINTLMEISTER, A.: Zoogeographie der palaearktischen Notodontidae (Lepidoptera). 1989. 20 z. T. ganzseitige Abbildungen, umfangreiche Tabellen, 116 S. DM 42,—

Bd. 26: REISSINGER, E.: Die geographisch-subspezifische Gliederung von *Colias alfacariensis* RIBBE, 1905 unter Berücksichtigung der Migrationsverhältnisse (Lepidoptera: Pieridae). 1989. 82 Tafeln, davon 14 in Farbe, 351 S. DM 145,—

Bd. 27: HACKER, H.: Die Noctuidae Vorderasiens. 1990. 18 Tafeln, davon 6 in Farbe, 740 S. DM 170,—

Bd. 28: RENNER, F.: Neue Untersuchungsergebnisse aus der *Pyrgus alveus* HÜBNER-Gruppe in der Palaearktis unter besonderer Berücksichtigung von Süddeutschland (Lepidoptera: Hesperiidae). 1991. 30 S/W-Tafeln, zahlreiche Textfiguren und Tabellen, 157 S. DM 85,—

Bd. 29: JOHNSON, K.: The Palaearctic "Elfin" Butterflies (Lycaenidae, Theclinae). 1992. 12 S/W-Tafeln, zahlreiche Strichzeichnungen und Verbreitungskarten, 141 S. DM 65,—

Bd. 30: REINHARDT, R. & R. THUST: Zur Entwicklung der Tagfalterfauna 1981–1990 in den ostdeutschen Ländern mit einer Bibliographie der Tagfalterliteratur 1949–1990 (Lepidoptera, Diurna). 1993. Verbreitungskarten für jede Art, 285 S. DM 110,—

Bd. 31: EBERT, K.: Die Großschmetterlinge des Vogtlandes (Insecta, Lepidoptera). 1993. 8 Seiten mit S/W-Abbildungen von Biotopen, 172 S. DM 96,—

Bd. 32: BINDER, A.: Beitrag zur Schmetterlingsfauna Böhmens (Insecta, Lepidoptera). 1994. 137 S. DM 72,—

Bd. 33: FETZ, R.: Larvalmorphologische Beiträge zum phylogenetischen System der ehemaligen Oecophoridae (Lepidoptera, Gelechioidae). 1994. 166 Strichzeichnungen im Text, 270 S. DM 95,—

Bd. 34: HÜBNER, J.: Index Larvarum europaeorum Lepidopterorum earumque nutrimenta quae in Opera Jacobo Hübner effigiaata sunt, adjectis denominationibus auctorum verorum. Verzeichnis europäischer Schmettlinge in ihren verlarvten Ständen. 1995. 20 S. + 24 S. Faksimile, 3 S/W- und 5 Farbtafeln. DM 115,—

Bd. 35: ESCHE, T.: Konkurrieren Nachtschmetterlinge um Blüten? Untersuchungen zu Nischentrennung und Bestäubungseffektivität (Insecta, Lepidoptera). 1996. Zahlreiche S/W-Abbildungen im Text, umfangreicher Tabellenteil, 194 S. DM 120,—

Bd. 36: BECK, H.: Systematische Liste der Noctuidae Europas (Lepidoptera: Noctuidae). 1996. 122 S.

Die Bände 20: HUEMER (Kleinschmetterlinge an Rosaceae unter besonderer Berücksichtigung ihrer Vertikalverbreitung), 21: MÖRTTER (Vergleichende Untersuchungen zur Faunistik und Ökologie der Lepidopteren in unterschiedlich strukturierten Waldflächen im Kottenforst bei Bonn) und 24: SCHURIAN (Revision der *Lysandra*-Gruppe des Genus *Polyommatus* LATR.) sind vergriffen.

Die Subskriptionspreise liegen deutlich unter den angegebenen Einzelheft-Preisen!
Zu beziehen durch: Verlag Dr. ULF EITSCHBERGER, Humboldtstr. 13a, D-95168 Marktleuthen

Ihr Spezialist für Entomologiebedarf

Zu günstigen Preisen erhalten Sie
bei uns:

Für die Zucht

Zuchtkästen, Puppenkästen,
Infrarotstrahler, Zuchtbehälter u. a.

Für den Tag- und Nachtfang

Netze, Gläser, Transportkästen,
Stromaggregate, Lampen, Leuchtröhren
u. a.

Für das Präparieren und Bestimmen

Präparierbestecke, Spannbretter,
Chemikalien, Lupen, Mikroskope und
Binokulare

Für die Sammlung

Insektenkästen in allen Größen mit der
bewährten Moll- oder
Schaumstoffeinlage, Insektenchränke
aller Art u. a.

Literatur

Ein umfangreiches Angebot an neuer und
antiquarischer Literatur ist vorhanden.

Schreiben Sie uns oder rufen Sie uns an.
Wir beraten Sie gern.

- Katalog kostenlos -



**icb —
bioform**
Erzeugnisse

Entomologie • Biologie
Geräte • Lehrmittel
Sammlungen
Institutsmobiliar
Fachbuchhandlung

bioform-Handelsgesell-
schaft Meiser & Co.

**Bittlmairstraße 4
D-85051 Ingolstadt
Telefon 08 41 / 7 55 83**

