## Field notes - Kurze Exkursionsberichte - Excursions en bref

On the presence of *Pyrgus alveus* (HUEBNER, 1803) in the Peloponnesus (Lepidoptera, Hesperiidae)

Enrico Gallo, Mura dello Zerbino 12/16, I-16122 Genoa, Italy.

Recently, while sorting out the *Pyrgus* in my collection, I examined the genitalia of two male specimens taken on a collecting trip to Greece in 1979. They were the only *Pyrgus* caught in the Peloponnesus on that trip. The dissection of the first specimen confirmed the determination based on external characters: *Pyrgus armoricanus persicus* Reverdin taken on Mt. Kyllini. The second specimen, which had proved more difficult to identify, could be determined as *Pyrgus alveus*. This specimen came from the western side of Mt. Chelmos near Kalavryta and was taken on the 23rd July just above the tree-line at about 1800 m, among scattered trees of Greek Fir.

According to DE Jong (1972), the typical subspecies of *P. alveus* is present in the Balkans: he mentions Bulgaria, Yugoslavian Macedonia and Albania, but not Greece. As far as I know, the species has been reported from this country only twice: a single specimen from Drama in Eastern Macedonia (Dacie *et al.*, 1972) and from the Southern Pindus range in Central Greece (Coutsis, 1972). I have found no record from the Peloponnesus. A recent and very thorough work on the butterflies of Mt. Chelmos does not mention it either (Leestmans & Arheilger, 1987).

This new record greatly extends southwards the known Balkanic range of the species. During the last glacial period, the Peloponnesus was probably a refugium for *Pyrgus alveus*, and the colony which has survived there, apparently isolated today, must be considered a relic. We cannot exclude the possibility that it has reached a subspecific level, as in the case of ssp. *centralitaliae* VERITY of Central and Southern Italy, which has a postglacial origin. However, any speculation in this direction is impossible as only one specimen is known, and *P. alveus* is a very variable species. Nevertheless, I think it is useful to give the photographs of my specimen and its genitalia (fig. 1-3). The more evident features are the pointed apex of the forewings and a yellowish flush on the upperside.

In conclusion, we can say that our knowledge on the occurrence of *P. alveus* in Greece is still very patchy; it is certainly one of the rarest *Pyrgus* in this area, and further exploration is needed to define its complete range and variation.

I am grateful to my friend Dr. V. Raineri of the Museo di Storia Naturale of Genoa who photographed the genitalia.



Fig. 1-3. Pyrgus alveus (Hübn.) & Mt. Chelmos, Peloponnesus. 1. Upperside; 2. Underside; 3. Genitalia, left valva removed.

## References

COUTSIS, J. G., 1972. List of Grecian Butterflies: Additional records 1969-1971. Entomologist's Rec. J. Var. 84: 145-151.

DACIE, J. V., DACIE, Margaret K. V. & GRAMMATICOS, P., 1972. Butterflies in Northern and Central Greece, July 1971. *Entomologist's Rec. J. Var.* 84: 257-266.

JONG, R. DE, 1972. Systematics and geographic history of the genus *Pyrgus* in the Palaearctic Region (Lepidoptera, Hesperiidae). *Tijdschr. Ent.* 115: 1-121.

LEESTMANS, R. & ARHEILGER, T., 1987. Les Lépidoptères du massif du Chelmos (Péloponnèse, Grèce): inventaire et considérations zoogéographiques (première partie). Linneana Belgica 11: 150-192.

## ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Nota lepidopterologica

Jahr/Year: 1989

Band/Volume: 12

Autor(en)/Author(s): Gallo Enrico

Artikel/Article: Field notes - Kurze Exkursionsberichte - Excursions en bref

<u>164-165</u>