

# *Trifurcula pallidella* (Duponchel, 1843) (Nepticulidae): distribution, biology and immature stages, particularly in Poland

ERIK J. VAN NIEUKERKEN<sup>1</sup>, ANNA MAZURKIEWICZ<sup>2</sup> & KRZYSZTOF PAŁKA<sup>3</sup>

<sup>1</sup> National Museum of Natural History Naturalis PO Box 9517, NL-2300 RA Leiden, Netherlands;  
e-mail: nieukerken@naturalis.nl (corresponding author)

<sup>2</sup> Warsaw Agricultural University - SGGW, Faculty of Animal Sciences, Department of Animal Biology and Environment, ul. Ciszewskiego 8, 02-786 Warszawa, Poland;  
e-mail: mazurk@alpha.sggw.waw.pl

<sup>3</sup> Lublin University - UMCS, Institute of Biology, ul. Akademicka 19, 20-033 Lublin, Poland;  
e-mail: palka@biotop.umcs.lublin.pl

**Abstract.** *Trifurcula pallidella* (Duponchel, 1843) is recorded for the first time from 30 localities in south-eastern Poland. The biology is described and illustrated in detail for the first time: the larva makes spindle shaped galls in stems of *Chamaecytisus* spp., *Lembotropis nigricans*, and *Cytisus procumbens*. In Poland it is usually found on dry grassland or edges of woodland on calcareous soils. The larva, pupa and galls are described and illustrated. The distribution is revised and mapped on the basis of material and literature: its occurrence in Germany: Bavaria, Italy mainland and Sicily is confirmed, and new records are given for Corsica and Crete. It is widespread from Southeast Poland southwards throughout the Balkans and eastwards to the Volga, in the West to Regensburg in Germany, southern Switzerland, western Italy and Corsica. It is believed to use a different host in Mediterranean localities, possibly *Calicotome*. All known hosts and *Calicotome* belong to the monophyletic *Cytisus* group of recent molecular studies.

**Zusammenfassung.** *Trifurcula pallidella* (Duponchel, 1843) wird erstmals aus 30 Fundorten in Südostpolen gemeldet. Die Biologie wird zum ersten Mal detailliert beschrieben und abgebildet: die Raupe macht spindelförmige Galle im Stengel von *Chamaecytisus* spp., *Lembotropis nigricans*, und *Cytisus procumbens*. In Polen wird die Art meistens in Trockenrasen oder an Waldränder auf Kalkuntergrund gefunden. Die Raupe, Puppe und Galle werden beschrieben und abgebildet. Die Gesamtverbreitung wird auf Grund von Material und Literatur revidiert und auf einer Karte abgebildet: das Vorkommen in Deutschland: Bayern, auf dem Italienischen Festland und Sizilien wird bestätigt, und die Art wird von Korsika und Kreta neu nachgewiesen. *T. pallidella* ist weit verbreitet, von Südostpolen bis über die Balkanhalbinsel im Süden und nach Osten bis zur Wolga, im Westen bis Regensburg in Deutschland, die Südschweiz, Westitalien und Korsika. Es wird angenommen dass *T. pallidella* im Mittelmeerraum andere Futterpflanzen nutzt, wahrscheinlich *Calicotome*. Alle bekannten Futterpflanzen und *Calicotome* gehören zur monophyletische *Cytisus*-Gruppe die auch in rezenten molekularen Studien anerkannt wird.

**Key words.** *Trifurcula*, distribution maps, hostplants, galls, *Cytisus*, *Lembotropis*, *Chamaecytisus*, *Calicotome*, Fabaceae, Genisteae, Braconidae, *Mirax*.

## Introduction

The genus *Trifurcula* Zeller, 1848 includes at least 36 named species in Europe (van Nieukerken 1996), particularly in the Mediterranean region, where also many unnamed species are known. More to the north the number of species is quickly decreasing, still sixteen species are found in the Czech Republic, eleven in Germany, seven in Denmark and five in Sweden. Only six species have been cited from Poland, which probably is an underestimate of the real number. The subgenus *Trifurcula* s. s. comprises a group of rather uniform moths, of which the larvae make stem-mines on Fabaceae (=Leguminosae). Two species groups are recognised: the *T. subnitidella* group which feeds on plants belonging to various Fabaceae tribes (i.e. *Lotus*,



**Figs. 1–4.** Habitats of *Trifurcula pallidella*. 1–2. Poland, Podlasie: Stare Stulno, May 2003, grassland on a sand dune, hostplant *Chamaecytisus ruthenicus*. 3. Poland, Lublin Upland: Brzeźno reserve, May 2003, calciphilous community at the margin of a marsh, hostplant *C. ruthenicus*. 4. Slovakia, Plešivec, Slovenský Kras: Hôrka, October 1992, a dry calcareous hill with grassland, hostplant *C. hirsutus* (not visible on photograph). Photos M. Sielezniew (1–3), E.J. van Nieukerken (4).

*Dorycnium*, *Coronilla*), but not on brooms (tribe Genisteae), and the *T. pallidella* group, which is specialised on brooms.

The type species of *Trifurcula*, *T. pallidella* (Duponchel, 1843) is one of the largest species in the genus and one of the oldest known. Taxonomy and distribution were discussed by van Nieukerken & Johansson (1986), especially in relation to *T. beirnei* Puplesis, 1984, which previously had often been misidentified as *T. pallidella*. At that time the biology was unknown, although it was presumed that *T. pallidella* is associated with *Lembotropis nigricans* (L.) Griseb.

Shortly after that, the larvae were discovered by the senior author during a joint excursion with Aleš and Zdeněk Laštůvka. The larvae make galls in *Lembotropis nigricans* and *Chamaecytisus* spp. and later galls were also found in *Cytisus procumbens* (Waldst. & Kit. ex Willd.) Sprengel (=*Corothamnus procumbens*). The gall is shown and briefly described by A. & Z. Laštůvka (1997). The two Polish authors have found adults and galls of *Trifurcula pallidella* in several places in Poland since 1998. On the basis of these records the species was listed as occurring in Poland by Buszko & Nowacki (2000). We here report the Polish records in particular and provide details on the biology, immature stages and distribution in Poland and in Europe, where many new data have become available since 1986.

## Material and methods

The first two Polish males of *T. pallidella* were found amongst material collected at light in xerothermic habitats in south-east Poland in 1998. In late autumn (20 October) of the same year, several empty galls were found in stems of *Lembotropis nigricans* and *Chamaecytisus ruthenicus* (Fischer ex Wołoszczak) A. Klášková in steppe vegetation in Machnów. The distribution of the host plants (Zajac & Zajac 2001) suggested that the species may have a wider range in Poland. Therefore we started detailed field investigations in 2001-2002 in order to test this hypothesis. A number of localities where the hostplants were expected were selected; these had similar biotic conditions as those mentioned above. Most of these sites are located in existing or planned nature reserves. We visited these sites from mid-September to mid-October: the period when the galls with larvae were expected. Some live material was also sent to the senior author, who reared a couple of specimens as well.

The senior author collected *T. pallidella* in the Czech and Slovak Republics in 1992 in collaboration with A. and Z. Laštůvka on localities where adults had been found previously. Other material studied was found during study of many museum and private collections since 1986, and some data were received from A. and Z. Laštůvka. He also re-examined all known literature citations.

Galls were collected by cutting parts of the stem of the hostplant. The stems were placed in plastic containers lined with tissue. Most larvae easily prepared cocoons on the tissue, after which the stems were removed and dried. Adults and parasites emerged after a winter diapause in an outhouse.


Larvae and pupal skins were examined after macerating in KOH and mounted on slides with euparal, after dehydration with ethanol and staining with chlorazol black and haemaluin. Galls were studied by simply cutting them with razor blades.

Morphological terms of larvae follow Gustafsson (1981), except for the naming of setae on the mesothorax, which follows Hoare (2000), in exchanging resp. the names of D1 and D2 and SV1 and SV2. Homologisation of setae, however, remains problematical. For this reason the ventral group of setae is not labelled in Fig. 21.

Photographs of preparations of immature stages and genitalia were taken with a Zeiss AxioCam digital camera attached to a Zeiss Axioskop H, those of live insects and galls by EvN with the same camera on Zeiss Stemi SV 11, using Carl Zeiss AxioVision 3.0.6 software. Galls were photographed by KP with a Canon D10 digital camera, with a standard lens. The adult moth was photographed with an Olympus DP10 digital camera attached to the stereo-microscope Olympus S260.

The European distribution map was prepared with Dmap 7.0 (Morton 2000). In Tab. 1 and 2 we give summarized locality data, all details of these, dates and UTM coordinates with an accuracy of 10 square kilometres are given in an excel sheet on the website of the senior author (<http://www.naturalis.nl/nieukerken>) and on the website of the journal (<http://www.socleurep.org/council-main.htm>). When coordinates were not available from the source, we obtained most from NIMA (2004).

Depositories of collections are abbreviated according to Evenhuis & Samuelson (2004). For the hostplant names we follow Heywood & Ball (1968). Although since then there



Figs. 5–9. *Trifurcula pallidella*. 5. Male, Poland: Stawska Góra res. 6–7. Live female, Poland, Góra Pieprzowe res., emerged in Leiden. 8. Live pupa, dorsal view, Poland, Stawska Góra res. 9. Cocoon on tissue, Góra Pieprzowe res. Photos: J. Chobotow (5), E.J. van Nieukerken (6–9).

has been much discussion on the generic limits within the *Cytisus* group and the Genistae in general, no consensus has been reached, and there is not another easy single source except the ILDIS database (ILDIS 2003), which uses a somewhat inconsistent nomenclature. The genera *Lembotropis* and *Chamaecytisus* as used here are often (but not always in the same paper) regarded as sections of *Cytisus* (resp. sect. *Lembotropis* and *Tubocytisus*), and the section *Corothamnus* in *Cytisus* (for *C. procumbens*) is in some publications raised to genus. In recent molecular studies, cited below, a large *Cytisus* seems to be paraphyletic, but also the genus *Chamaecytisus* as currently understood appears as polyphyletic (for a revision of this taxon see Cristofolini 1991).

## RESULTS

### *Trifurcula pallidella* (Duponchel)

*Oecophora pallidella* Duponchel, [1843]: 339, pl. 78. Lectotype selected by van Nieukerken & Johansson 1986.

*Trifurcula pallidella*; Zeller 1848: 332.

*Trifurcula incognitella* Toll 1936: 409.

*Trifurcula pallidella*; van Nieukerken & Johansson 1986: 271.

**Diagnosis.** Wingspan 7.5–9 mm. Head: frontal tuft from almost white, yellowish orange to brown ferruginous, collar white to straw, occasionally with a few darker scales. Scape straw to ferruginous orange, flagellum grey-ochreous indistinctly ringed, 42–50 segments. Thorax and forewings uniformly coloured, covered with two types of scales: white and ferruginous-tipped scales, sometimes with golden gloss. The ferruginous-tipped scales usually becoming more abundant towards apex. The general impression of the wing colour varies from white to ochreous, depending on abundance of scale type. Hindwing grey with straw cilia. Abdomen grey dorsally, straw ventrally.

**Similar species.** Of the species with which it can be found together, *Trifurcula pallidella* resembles *T. beirnei* most; this can in many cases be differentiated by the different flying period: *T. pallidella* occurs from May to early (mid) July, *T. beirnei* usually in August to September. However, there are a few earlier records of *T. beirnei*, thus July specimens need to be checked carefully. *T. beirnei* is on average larger (8–11 mm) than *T. pallidella*, and has three types of scales (white, yellow and dark tipped). The characteristic gnathos and large curved valvae of *T. beirnei* can often be seen without dissection.


Most other species likely to confuse with *T. pallidella* belong to the *T. immundella* complex of species, and of these *T. immundella* (Zeller, 1839) and *T. chamaecytisi* Z. & A. Laštůvka, 1994 resemble *T. pallidella* most, but larger and worn specimens of the other species may also be confused. Most of these species, except *T. immundella*, feed on the same hosts as *pallidella*, and can therefore be found in the same localities. These species are on average somewhat smaller (6–8.5 mm, sometimes even smaller), and apart from *T. chamaecytisi*, they are darker when fresh. In collections *T. pallidella* was often confused with the much smaller (5–7 mm) *T. serotinella* Herrich-Schäffer, 1855. The male genitalia offer the best characters, for descriptions and illustrations we refer to van Nieuwerken & Johansson (1986), but we also illustrate here the male genitalia (Fig. 17).

## Life history

**Hostplants.** With certainty galls are now recorded from *Chamaecytisus albus* (Hacq.) Rothm., *C. austriacus* (L.) Link, *C. hirsutus* (L.) Link, *C. ratisbonensis* (Schaeffer) Rothm., *C. ruthenicus* (Fischer ex Wołoszczak) A. Klásková, *Cytisus procumbens* (Waldst. & Kit. ex Willd.) Sprengel (in section *Corothamnus*) and *Lembotropis nigricans* (L.) Griseb.

**Galls.** The galls with full-grown larvae were found from 9 September to early November. Galls occurred at different heights in the stem, both in older shoots and younger shoots of the same year. Occasionally two or three galls were found in the same stem. Old galls remain visible and can still be found in the next spring, at least until June (see Tab. 1 and 2).

The egg is deposited on the stem where later the gall forms, usually difficult to see. The larva first bores into the parenchyma and then feeds in a spiral gallery around the stem, successively moving upwards, also partly boring in the central woody part of the stem (Figs. 10–16). The stem is thickened more or less considerably by the larval




**Figs. 10–16.** *Trifurcula pallidella*, biology, galls and details, 10–14. *Chamaecytisus ruthenicus*, Korphynie (11: Ostowo), 10. Three galls. 11–13. Various sections. 14. Exit hole, indicated by arrow. 15–16. Galls in *Lembotropis nigricans*, Stawska Góra res. Photos E.J. van Nieuwerken (10–14), K. Palka. (15, 16).

activity. The diameter of the galls varies from 3.2–6.0 mm (depending on stem thickness) and the length from 13–26 mm ( $n=20$ ). The frass is deposited in the gallery (Figs. 11–13), in a similar fashion as in stem-mines of related *Trifurcula* species, almost filling the gallery.

The full fed larva quits the gall through a slit in its upper part, usually close to a stem bud (Fig. 14). The pupation takes place in a typical Nepticulidae cocoon (Fig. 9). Cocoon length is 2.7–3.7 mm (mean 3.3 mm,  $n=20$ ), width 1.8–2.5 mm (mean 2.1 mm,  $n=20$ ). Its colour is pale brown to ferruginous brown. In captivity the cocoons were attached to the walls of the rearing tubes, to the surface of a stem or on pieces of paper tissue. After hibernation in outdoor temperature, the adults emerged from the middle of May to early June.

**Voltinism.** The moths fly in central Europe from 9 May to 19 July, most in May and June, in southern Greece and Crete they fly from 9 April to 17 May, but also 10 June in northern Greece. The species is clearly univoltine.

**Parasites.** We reared several parasites of the family Braconidae, all belonging to *Mirax rufilabris* Haliday, 1833 (det. C. van Achterberg, Leiden). They emerged in the



**Fig. 17.** *Trifurcula pallidella*, male genitalia, Germany: Kelheim, slide EJ7919. Scale 100 $\mu$ m.

that he reared it from broom and thus from the only nepticulid known to feed on broom in Ireland: *T. immundella* (see Emmet 1976). According to C. van Achterberg, the ‘real’ *M. rufilabris* was up to now only known from *Trifurcula immundella*, and *T. pallidella* thus comprises a new host record. We may therefore assume that *M. rufilabris* is specialised on *Trifurcula* species that feed in the stem of brooms (see Tab. 1).

The rate of parasitism was high: from the material collected in autumn 2002 we reared 57 specimens of *Mirax*, one of *Chrysocharis* and only thirteen adults of *T. pallidella*.


**Remarks.** The galls were discovered by the senior author during joint field work with Aleš and Zdeněk Laštůvka in Vranovice, Moravia, 5 October 1992. In this locality adults were previously frequently caught around *Chamaecytisus ratisbonensis*. After failing to localise any stem-mines, he opened some spindle-shaped galls, and found to his surprise nepticulid larvae in it. Later several adults were reared from galls of various localities. These galls had also been noted and described before by Baudyš (1925) and Černík (1942), who found them in various localities in Moravia and Slovakia (Tab. 2). The larva was identified as ‘unknown Microlepidopteron’, and also cited in the handbook by Buhr (1964: 422) under *Cytisus* as number 2225. The galls are probably not induced by the egg-laying activity of the adult, but by the feeding larva; since it seems that the larva is not depending on tissue formed in the gall (but this needs to be checked more carefully). This type of thickening should not be called a gall according to some authors.

### Description of immature stages

**Larva** (Figs. 18–21). Mature larvae about 6–8 mm long, extremely slender, yellow when fresh. Body almost completely smooth, small spines (microtrichia) absent. Headcapsule 400–480  $\mu$ m long, 440–515  $\mu$ m wide (n=2), much wider than long. Labrum with 2 pairs of setae; mandibula with 4 strong cusps; labial palpus

laboratory between 9 April and 13 June 2003. We also reared just one specimen of Eulophidae, probably *Chrysocharis* sp. (5 June 2003). All these were reared out of the cocoons and thus both species are koinobionts. *Mirax rufilabris* has been recorded from many Nepticulidae species, including *Trifurcula immundella* (Shaw & Askew 1976), but it is now understood that *Mirax* forms a complex of several species with a narrower host range, of which a revision is in preparation (C. van Achterberg, pers. comm.).

The type locality of *M. rufilabris* is Ireland, probably near Dublin. Haliday renamed it later *M. spartii*, an indication



**Figs. 18–21.** *Trifurcula pallidella*, larva, Stawska Góra res. **18.** Headcapsule, dorsal; scale 100 µm. **19.** Headcapsule and prothorax, ventral; scale 200 µm. **20.** Detail of mouthparts, with labial palps; scale 50 µm. **21.** Seta map, drawn by A. Mazurkiewicz & K. Pałka. Photos E.J. van Nieuwerken.

with 3 segments and long terminal seta, second segment much longer than segment 1 or 3. Prothorax with pair of slender tergites and single indistinct sternite plus two small sternites anterolateral to the medial one; with the full complement of 13 pairs of

setae. Mesothorax with 12 pairs of setae (D1 present, 4 pairs of setae ventral to SV1), metathorax with 10 pairs (D1 and L3 absent). Abdominal segments 1–8 with 6 pairs, A9 with 3 pairs and A10 with 2 pairs. Distribution of setae illustrated in setal map (Fig. 21). Paired ventral ambulatory calli present on T2–3 and A1–7. Anal rods in A10 posteriorly bifid, forming an angle of almost 180°.

The larva differs from other described *Trifurcula* (s. str.) (Gustafsson 1981; Gustafsson & van Nieukerken 1990) by the large headcapsule, which is wider than long, the complete absence of microtrichia, the presence of D1 (named D2 by Gustafsson) on the mesothorax and the labial palpus which has a very long segment 2, but otherwise confirms with the generic description. Only *Trifurcula (Glaucolepis) headleyella* (Stainton, 1854) and *T. (T.) subnitidella* (Duponchel, 1843) have also 12 setal pairs on the mesothorax, the other studied Nepticulidae have only 11 (Gustafsson 1981; Gustafsson & van Nieukerken 1990; Hoare 2000). No setal map of a species of *Trifurcula* (s. str.) had been published previously.

**Pupa** (Figs. 8, 22–24). Pupal exuviae examined. Frons protruding slightly into conical projection, eyecaps large, at eclosion torn from frons. Abdominal tergites 2–8 covered with many spines, in about 3 to 5 rows per segment, but not arranged in distinct rows. Cremaster with two small hooks.


## Distribution

**Localities in Poland** (Fig. 25). As a result of our studies, *T. pallidella* was found in 30 localities in south-east and east Poland. They are listed in Tab. 1 and presented on a map (Fig. 25).

The species reaches in Poland its northern distribution limits. The locality Grzybowce is the northernmost locality of this species in central Europe, but the Russian record in Ul'yanovsk Oblast is just a little further to the North.

**European distribution** (Fig. 26). In Tab. 2 we present all more or less reliable literature records and material seen by EvN or identified by colleagues sorted by country and locality. Since the previous revision of this species (van Nieukerken & Johansson 1986), many more data have become known, from which it appears that the previous suggestion that ‘literature records can not be used at all’ was too rigid. It is clear that the name *T. pallidella* was incorrectly used for *T. beirnei* in Great Britain, Denmark and parts of Germany and Poland. We have now also established that all previous records from France, Portugal and Spain are incorrect (van Nieukerken *et al.* 2004a, 2005): they are misidentifications for a number of pale *Trifurcula* species including *T. immundella*. From the overlapping part of the distributions of *T. pallidella* and *T. beirnei*, the records from Austria: Stelzing (Zeller in Krause 1871), Germany: Altenburg (Krause 1871; Borkowski 1975) and Poland: Sobótka (Zobten) (Wocke 1874; Puplesis 1984) definitely apply to *T. beirnei*. Two of the four localities on the Monte Baldo (Burmann & Huemer 1998), namely those in higher altitudes, refer to an unnamed species that is associated with *Genista radiata* (L.) Scop. (= *Cytisanthus radiatus*) (P. Huemer, *in litt.*) Other sources are reconsidered here in Tab. 2.

Many of the records presented in Tab. 2 are corroborated by either original material



Figs. 22–24. *Trifurcula pallidella*, pupal skin, dorsal view, Korhynie. Scales 500µm, 50µm (24). Photos: E.J. van Nieukerken.

or later findings. Others which have not been corroborated, are quite likely to be correct, but confirmation is still required (Italy: Pisa, Croatia, Macedonia, Rumania): they are all within the area where the known hosts occur and where the occurrence of *T. pallidella* is likely. Most problematic are records at the margin of the distribution area. The occurrence in Germany: Regensburg was previously doubted, and Segerer (1997) wrote: ‘unüberprüfbare Angabe’. However, the occurrence in Germany is here corroborated by three specimens from three localities in Bavaria, including Regensburg (see Fig. 17). The record from Baden-Württemberg (Wörz 1958) has still to be rejected; it is a curious misidentification: upon checking, the specimen (coll. SMNS) appears to be a worn female of *Stigmella floslactella* (Haworth, 1828).



Fig. 25. Distribution of *Trifurcula pallidella* in Poland on 10km squares of the UTM grid.




Fig. 26. Distribution of *Trifurcula pallidella*. Small grey circles: unconfirmed but not rejected literature records, large black circles: records confirmed by material, or reliable literature records. Grey shading gives the combined distribution of the hosts *Lembotropis nigricans* and *Chamaecytisus* spp. of the series *Tubocytisus* s. str. (Cristofolini 1991).

**Tab. 1.** Localities of *Trifurcula pallidella* in Poland and rearing results. UTM coordinates given as 10km squares, all in zone 34U. Hostplants: *Ca* = *Chamaecytisus albus*, *Cra.* = *C. ratisbonensis*, *Cru* = *C. ruthenicus*, *Ln* = *Lembotropis nigricans*. Other abbreviations: Chr. = *Chrysocharis* sp. (parasite), M = *Mirax rufulabris* (parasite), \* = material sent to and reared in Leiden (in coll. RMNH). All material collected by A. Mazurkiewicz and / or K. Pałka.

Region	Locality	UTM	Habitat	Host	Stage	Date	reared
Lublin Upland	Brzeźno village	FB96	xerothermic rail-way embankment	<i>Cru</i>	old galls	8.v.2004	
	Brzeźno res.	FB87	edge of calcareous peat-bog	<i>Cru</i>	larvae	5.x.2002	3♂, 3♀, 5M
					old galls	28.v.2004	
	Czumów	GB03	grassland on loess	<i>Cru.</i>	galls	15.x.2001	
	Gródek	GB03	grassland on loess	<i>Ca</i>	galls	11.xi.2004	
	Podzamcze res.	FB05	grassland on limestone rocks	<i>Cru, LN</i>	galls	7.x.2001	
				<i>Cru</i>	larvae	5.x.2002	2♀, 1M
					1♂, 1♀	24-28.v.2004	
	Rogów res.	FB73	grassland on limestone rocks		1♂, 1♀	25.v.2004	
	Skarpa Dobrska res.	EB68	grassland on loess		1♂	12.vi.1997	
				<i>Cru, LN</i>	galls	7.x.2001	
	Skarpa Dobużańska res.	FB90	grassland on loess	<i>Cru</i>	galls	15.x.2001	
Malopolska Upland	Stawska Góra res.	FB67	grassland on limestone rocks	<i>LN</i>	larvae	6.x.2001	2♂, 2♀
					larvae	3.x.2002	1♂, 21M*
					larvae	10.x.2002	2♂, 5M
	Tarnogóra	FB44	grassland on loess	<i>Cru</i>	galls	8.x.2000	
	Zalesie Kraszeńskie	FB57	xerothermic rail-way emb.	<i>Cru</i>	old galls	13.v.2004	
	Biała Góra res.	DA28	grassland on limestone rocks	<i>Cru</i>	galls	3.xi.2001	
	Biedrzychów	FB54	grassland on loess	<i>LN</i>	galls	2.x.2001	
	Góry Pieprzowe res.	EB51	grassland on loess		1♂	29.v.1995	
					2♂	2.vi.2001	
				<i>Cru</i>	larvae	3.x.2002	1♂, 1♀, 1Chr, 6M*
Podlasie	Pińczów	DB60	grassland on gypsum	<i>LN</i>	galls	4.xi.2001	
					galls	10.x.2004	
	Wały res.	DA47	grassland on limestone rocks	<i>Cru</i>	galls	3.xi.2001	
	Ciesacín res.	FB49	neutral grassland on sand on the edge of peat-bog	<i>Cru</i>	old galls	9.v.2004	
	Dobrowoda	FD62	grassland on gravel	<i>Cra</i>	galls	2.x.2002	*
	Drohiczyn	FD10	grassland on gravel	<i>Cru</i>	galls	2.x.2002	*
	Grzybowce	FD89	grassland on gravel	<i>Cru</i>	galls	9.x.2004	
	Mielnik – Przedmieście	FC49	grassland on gravel	<i>Cru</i>	galls	7.x.2001	
	Oslowo	FC39	grassland on gravel	<i>Cru</i>	galls	2.x.2002	9M*
	Stare Stulno	FB89	neutral grassland on sand	<i>Cru</i>	galls	6.x.2001	
Roztocze	Witowo, in Białowieża Primeval Forest	FD63	grassland on gravel	<i>Cru</i>	galls	2.x.2002	*
	Zagórze	FC39	grassland on gravel	<i>Cru</i>	galls	7.x.2001	
	Biała Góra	FA79	grassland on limestone rocks	<i>Cru, LN</i>	galls	15.x.2001	
	Katy II	FB51	grassland on limestone rocks	<i>Cru, LN</i>	galls	8.x.2000	
	Korhynie	FA88	grassland on limestone rocks	<i>Cru, LN</i>	galls	14.x.2001	
Sandomierz Lowland	Machnów	FA88	grassland on limestone rocks	<i>Cru, LN</i>	galls	20.x.1998	
					galls	25.ix.2004	
	Radruż	FA66	grassland on limestone rocks	<i>Cru</i>	larva	24.ix.2004	
	Nowa Dęba	EA58	neutral grassland on sand		1♂	3.vi.1998	
Przemyśl Hills				<i>Cru</i>	galls	2.x.2001	
	Złota Góra	FA31	grassland on gravel	<i>Cru</i>	galls	1.x.2001	

**Tab. 2.** Records of *Trifurcula pallidella*, except for Poland. The countries are indicated by their ISO codes, and some provinces abbreviated. Hostplants: *Ca* = *Chamaecytisus austriacus*, *Ch* = *C. hirsutus*, *Cr* = *C. ratisbonensis*, *Ln* = *Lembotropis nigricans*. Abbreviations: AZL = A. & Z. Laštůvka, EvN = E. van Nieukerken. Collection codons follow Evenhuis & Samuelson (2004). Reliability: ! after collectioncodon: material seen by EvN; ? not impossible, fits in distribution and biology, but confirmation required, ?? questionable (but not completely rejected). The remaining literature references are either reliable (genitalia check, recent authors, galls or circumstantial evidence) or at least likely records.

**Key to sources:** 1. Anikin 2001, 2. Baudyš 1925, 3. Baudyš 1931, 4. Baudyš 1947, 5. Burmann & Huemer 1998, 6. Caradja 1899, 7. Caradja 1901, 8. Černík 1942, 9. Curó 1883, 10. Czekelius 1924, 11. Hartig 1964, 12. Hauder 1912, 13. Hauder 1924, 14. Herrich-Schäffer 1847-1855, 15. Hrubý 1964, 16. Issekutz 1972, 17. Kasy 1985, 18. Klimesch 1949, 19. Klimesch 1961, 20. Klimesch 1968, 21. Klimesch 1990, 22. Laštůvka 1994, 23. Z. & A. Laštůvka 1994, 24. A. & Z. Laštůvka in litt., 25. Mann 1862, 26. Mann 1866, 27. Mann 1869, 28. Mann 1885, 29. Mariani 1938, 30. Mariani 1943, 31. Müller-Rutz 1922, 32. Nickerl 1908, 33. van Nieukerken & Johansson 1986, 34. van Nieukerken et al. 2004, 35. Pável & Uhryk 1896, 36. Prohaska & Hoffmann 1929, 37. Pröse & Segerer 1999, 38. Puplesis 1994, 39. Rebel 1899, 40. Schmid 1886, 41. Skala 1933, 42. Szöcs 1965, 43. Tokár et al. 1999, 44. Toll 1936, 45. Toll 1938, 46. Vorbrodt 1932, 47. Zeller 1850, 48. Zimmermann 1944b, 49. Zimmermann 1944a.

Country, Locality	collector	stage, host	Source
AT: BU	Bruck an der Leitha, Spitalberg	Preissecker	1♂ 19, 49, NMW!
	Rechnitz, Donatikapella	Koschabek	1♂ 16, 19
	Schieferberg, Leithagebirge	Kasy	1♂ 33, NMW!
AT: KÄ	Villach	Preissecker	1♀ NHMB!
AT: NÖ	Fürbachwiesen, Gramatneusiedl	Kasy	4♂ 17, 33, NMW!, RMNH!
	Hundsheimer Berg	Kasy	1♂ 33, NMW!
	Jauerling	Preissecker	49
	Mödling [Goldene Stiege]		1♂ HNHM!
	Oberweiden	Preissecker	49
	Ostrong [Klimesch Haide]	Preissecker	1♂ 21, 49, NMW!
	Pfaffstätten		1♀ HNHM!
	Stein a. D., Goldburg	Preissecker	1♂ 49, NMW!
AT: OÖ	Ysper	Preissecker	1♂ 49, NMW!
	Berg (Alpenvorland)	Wolfschläger	12, 21, 41
	Bergham	Wolfschläger	13, 41
	Diessenleiten	Hauder	13, 21, 41
	Ebelsberg (Alpenvorland)	Klimesch	[1♀!] 21, ZSM!
	Koglerau, Waldblösse	Knitschke	1♂ 12, 21, 41, NMW!
	Linz, Linz kgl. [Koglerau]	Hauder	3♂ NHMB!, ZFMK!
AT: ST	Linz-Uhrfahrwänd	Klimesch	[1♂!] 18, 21, ZSM!
	Bruck a. Mur	Klimesch	19
	Graz; Graz, Plabutsch		9♂ 33, 36, HNHM!, NHMB!, NHRS!, NMW!
	Premstätten	Prohaska	1♂ 36, HNHM!
AT: WI	Weiz	Knitschke	2♂ NMW!
	Wien	Mann	4♂ 14, 33, MNHN (LT)!, BMNH!
	Laaerberger Remisen	Mann	28, 49
	Mauer	Mann	1♀ 19, 28, 49, NMW!
	Rodaun	Mann	19, 28, 49
	Tivoli	Mann	28, 49
CH	Wien, Dornbach		1♂ NMW!
	Ticino: Maroggia	Krüger	1♂ 31, 46, NHMB!
CZ:Boh	Praha (Prag-Krtsch)	Nickerl	23, 32, 48

<b>Country, Locality</b>	<b>collector</b>	<b>stage, host</b>	<b>Source</b>
<b>CZ: Mor.</b>	Brno	gall, <i>Ln</i>	2
	Bulhary	AZL	larvae, <i>Ca</i> 22, 23
	Čelechovice na Hané	AZL	23, AZL!
	Dětkovice	AZL	15♂ 23, RMNH!
	Dolní Dunajovice	AZL	larvae, <i>Ca</i> 23
	Hrušovany nad Jevišovkou	AZL	larvae, <i>Ca</i> 23
	Kvasice, Kvasická skála rock	Zavřel	gall, <i>Cr</i> 4
	Mikulov, Stráň u Sedlece	Picbauer, AZL	gall, <i>Ca</i> 4, 22,
	Mikulov, Pavlovské vrchy, Stolová h.	EvN	gall <i>Ca</i> RMNH!
	Myslejovice	AZL	1♂ 23, AZL!
<b>DE: BA</b>	Olomouc, Neboteiner Berg	Černík	<i>Cr, Ln</i> 8
	Pouzdřany	Picbauer	gall, <i>Ln</i> 2
	Tuřany		gall, <i>Ln</i> 2
	Vicov	AZL	1♂ 23
	Vranovice	AZL, EvN	7♂, 2♀, e.l. <i>Cr</i> 23, RMNH!
<b>FR- 2A</b>	Grafenwöhr	Klimesch	1♂ ZSM!
	Kelheim Donau: Auf der Brand	E. Jackh	1♂ USNM!
	Regensburg [Tegernheimerberge]		1♂ 14, 37, 40, ETHZ!
<b>FR- 2B</b>	St. Lucia di Porto Vecchio	Liška	1♂ 24
<b>FR- 2B</b>	15 km S. Bastia	Karsholt	1♂ 24, ZMUC
<b>GR-CR</b>	Iraklio: Potamies	Johansson	2♂ RMNH!, Johansson!
	Iraklio: Stalis, Chersonisos	Bengtsson	2♂ Bengtsson!
	Rethymno: Plakias	Johansson	1♂ Johansson!
	Rethymno	Johansson	1§ Johansson!
<b>GR</b>	Fthiotis: Delfoi	Selling	1♂, 1♀ ZMUC!
	Ioannina: Asprángeli	AZL	2♂ 24
	Lakonia: 5 km S Monemvasia	Skule, Christ.	1♂, 1♀ 33, ZMUC!
	Lakonia: 7 km SW Monemvasia	Skule	1♂ 33, ZMUC!
	Messinia: Messini	Horak	1♂ 33, ZMUC!
<b>HR</b>	Josipdol		35?
	Rijeka (Fiume) (Istria)		30, 35?
	Dalmatia [no detail]		27?
<b>HU</b>	Budakeszi, Hárzbokorhegy		42
	Nadap	Gozmany	1♂ 42, HNHM!
	Nyír, Kecskemét	Klimesch	1♂ ZSM!
<b>IT</b>	Bolzano	Hedemann	11, 30, 39
	Bolzano: Renon: Collalbo	Hartig	1♂ 11
	Catanzaro: Sta Catarina dello Ionio	AZL	1♀ 24
	Imperia: Pieve d.T., Monesi	Liška	2♂ 24
	Pisa	Mann	9, 14, 30, 47
	Trento: Pietramurata nei boschi	Jäckh	1♂ photograph in RMNH! [USNM]
	Verona: Mte Baldo, Ferrara + Pai	Burmann	5
<b>IT-SIC</b>	Messina: Francavilla di Sicilia	AZL	1♀ 24
	Palermo: Castelbuono	AZL	1♂ 24
	Palermitano (Sicilia)	Mann	9, 30
	Palermo: Vallecorta		29
	Palermo: San Martino delle Scale	Klimesch	3♂, 1♀ ZSM!
<b>MK</b>	Drenovo-Kav.	Klimesch	20
<b>RO</b>	Sibiu (as Hermannstadt)	Prall	10?
	Tulcea (as Tultscha)	Mann	6, 7?, 26
<b>RU</b>	Saratov	Anikin	1, 38
	Ul'yanovsk: Srednikovo, Povolzhye	Zolotuhin	1♂ 34, coll. Zolotuhin!
<b>SI</b>	Carniola	Mann	9?
<b>SK</b>	Čebovce	AZL	2♂ 23, AZL!
	Kočovce (Kocsócz)		15, 23?, 35
	Nové Mesto nad Váhom	Patočka	15, 23
	Plešivec	AZL, Patočka	1♂, <i>Ch</i> 23, 43
	Plešivec, Slov. Kras.: Hôrka	EvN	2♂, 2♀ e.l. <i>Ch</i> RMNH!
	Velký Šariš, Prešov	Suza	gall, <i>Ca</i> 3
	Zádiel, Slovensky Kras	AZL, EvN	3♂, 1♀, el <i>Ln</i> 23, AZL!, RMNH!
<b>TR</b>	Bursa (Brusza)	Mann	25??
<b>UA</b>	Borshchiv area, S: Babuchow	Toll	44, 45
	Zalishchyky area: Obizowa	Toll	44, 45

The old record from Turkey (Mann 1862) is questionable: in Turkey occurs also a large pale species of *Trifurcula* (*Glaucolepis*), which could easily be confused, but the occurrence of *T. pallidella* in Turkey is also to be expected.

The record of *T. pallidella* in September by Schmid (1886) is probably a misidentification for *T. beirnei*, likewise the August record from Macedonia (Klimesch 1968) is not accepted, but in both cases earlier records in the same area are considered as possibly correct *pallidella*. The record from 'Alpi Marittimi' (Mariani 1943) may refer to the incorrect record from the French department Alpes Maritimes (Cannes), but the recent record from Mónesi in the Alpi Marittimi very close to the French border also confirms its occurrence here in Italy. This also makes it more likely that real *T. pallidella* can after all be found in the French Department Alpes Maritimes.

In summary, *T. pallidella* is distributed from Southeast Germany (Bayern), east into Bohemia, Moravia, Slovakia, adjacent SE Poland, much of Austria, both along the Danube and East and South of the Alps, extending through the Balkans (with very few verified records) to southern Greece (Peloponnesus) and Crete, eastwards through Ukraine and Russia to the Volga, and one questionable record from Turkey. To the West it occurs in northern Italy almost reaching France, just extending into Switzerland (Ticino), and throughout Italy to Sicily and Corsica.

**Habitats.** In Poland and Slovakia the species was found in warm and dry grasslands or open woodland, often on calcareous soils, but also on sand or gravel (Figs. 1, 2, 4, Tab. 1). In one case (Brzeźno reserve, Fig. 3) it was found in a relatively wet habitat on the edge of a calcareous bog. Elsewhere in Europe it occurs in a variety of habitats, although usually on relatively warm places: often sunny slopes in river valleys, such as that of the Danube in Germany, Austria and Hungary. The hosts are frequently associated with *Quercus pubescens* woodland. The localities in Corsica, Greece and Crete are Mediterranean shrubland (garrigue, maquis or phrygana), often close to the sea. The altitudinal range is from sea level to 1000 m, with one record at 1600 m in Italy (Imperia, Mónesi, Alpi Marittimi).

In 2001 and 2002 the species was extremely abundant in Poland, and larvae could be found in large numbers in most places. In contrast to that, in October 2004 almost no larvae or galls could be found in several of the same localities. Most localities in Central Europe are in Nature reserves, and the species may be vulnerable to increasing pressure by agricultural development and manure.

## Discussion

**Distribution and hostplant relationships.** *Trifurcula pallidella* occurs commonly in many parts of the distribution area of its presumed hosts and is much more common than previously assumed (van Nieuwerkerken & Johansson 1986). However, it has as yet not been found in the more western part of the distribution area of species of *Chamaecytisus*, e.g. France, parts of Germany and Spain. On the other hand, the distribution of *T. pallidella* cannot be explained completely by the distribution of the known hosts in the southern part of its range: no species of *Lembotropis*, *Chamaecytisus* or *Cytisus* sect. *Corothamnus* are known from Corsica or Sicily, and

the relatively common occurrence of *T. pallidella* in southern Greece and Crete, contrasts with the sparse occurrence of just a few species of *Chamaecytisus*. In fact in this part of Greece and Crete only three species of brooms are frequent: *Calicotome villosa* (Poiret) Link, *Spartium junceum* L. and *Genista acanthoclada* DC. (Turland et al. 1993, EvN, personal observation in Peloponnesus).

Recent studies on the molecular phylogeny of the Genisteae (Cubas et al. 2002) support at least two large monophyletic clades: a *Genista* group and a *Cytisus* group. Käss & Wink (1995; 1997) find support for a division of the *Cytisus* groups in two clades using rbcL sequences and Neighbour Joining, in which *Chamaecytisus* is polyphyletic and dispersed over both clades, but all hostplants of *T. pallidella* in the analysis, group together with *Cytisus multiflorus*, *C. arboreus* and *Calicotome villosa*. However, in the ITS sequence and in MP analyses of both genes (Käss & Wink 1997), a monophyletic *Cytisus* group including *Calicotome* is supported. These authors therefore suggest to synonymise the genera *Lembotropis*, *Chamaecytisus* and *Calicotome* with *Cytisus*. Combining these results with the previous observations leads to the hypothesis that in the Mediterranean region *Calicotome* is a very likely alternative hostplant for *T. pallidella*, and that *T. pallidella* is specialised on a number of species (but not all) within the *Cytisus* clade.

There is an interesting parallel with the *Trifurcula immundella* species complex: this is a group of very closely related species, which feed on about the same hosts as *T. pallidella*, but the species in this complex are almost strictly monophagous: each feeds on only one or a few related hosts: *T. immundella* feeds on *Cytisus scoparius* (L.) Link, *T. moravica* Z. & A. Laštůvka, 1994 on *Lembotropis nigricans*, *T. corothamni* Z. & A. Laštůvka, 1994 on *C. procumbens*, *T. chamaecytisi* on *Chamaecytisus* and *T. calycotomella* A. & Z. & Laštůvka, 1997 on *Calicotome* spp. (Z. & A. Laštůvka 1994): all feeding on the *Cytisus* group and these species – apart from *T. immundella* – have combined the same host range as *T. pallidella*.

**Galling.** *Trifurcula pallidella* is the only species of the genus causing galls, the other species of the subgenus *Trifurcula* exclusively making stem-mines (van Nieuwerken 1990; Z. & A. Laštůvka 1994; A. & Z. Laštůvka 1997), and most representatives of the other subgenera (*Levarchama* and *Glaucolepis*) make leafmines, a few *Glaucolepis* also stem-mines. The galls of *T. pallidella* in a way resemble the stem-mines, but the larva seems to feed in deeper tissue than is usual for *Trifurcula* (s. str.) species. It places *T. pallidella* in a rather isolated position in the genus, and also morphologically it is not very close to any other species. There is no apparent explanation why this species changed the stemmining habit into galling: on the same hostplants stemmining species also occur as discussed above, so the plant itself cannot be the reason. The larva does not show special adaptations to the galling habit, it is only larger than most other Nepticulidae larvae, and the mandibular cusps are relatively strongly developed.

Galling has originated several times independently in the Nepticulidae: there are at least three other examples. The North American type species of the genus *Ectoedemia* (which means as much as an external swelling), *Ectoedemia populella* Busck, 1907 is causing globular galls in the petiole of *Populus* (Busck 1907). Related species in the

*populella* species group also cause thickened petioles by their feeding habit, but these are not considered real galls. Most likely *E. populella* evolved from a petiole mining species. *Ectoedemia (Zimmermannia) castaneae* Busck, 1913, another North-American species, makes galls on bark of *Castanea* (Busck 1913), and probably evolved from barkmining species. Further *Ectoedemia (Fomoria) nigrifasciata* (Walsingham, 1908) from the Canary Islands makes a combined gall and mine in the Apocynaceae *Periploca laevigata* Ait. (Klimesch 1972); its closest relatives are leafmining on *Euphorbia*.

For two species of *Trifurcula* the larval way of feeding is still unknown: *T. beirnei* and *T. squamatella* Stainton, 1849. Both are relatively large species (the largest in the genus), associated with resp. *Genista* species and *Cytisus scoparius*. It is well worth considering the possibility that they are also gall makers, when searching for their larvae.

### Acknowledgements

For the loan of material and information we are grateful to Daniel Bartsch (SMNS, Stuttgart, Germany), Bengt Å. Bengtsson (Färjestaden, Sweden), Michel Brancucci (NHMB, Basel, Switzerland), Ulf Buchsbaum (ZSMC, München, Germany), Don Davis (USNM, Washington, USA), Bert Gustafsson (NHRS, Stockholm, Sweden), Christoph Häuser (SMNS, Stuttgart, Germany), Peter Huemer (TLMF, Innsbruck, Austria), Roland Johansson (Växjö, Sweden), Ole Karsholt (ZMUC, Copenhagen, Denmark), the late F. Kasy (NMW, Wien, Austria), Gerard Luquet (MNHN, Paris, France), W. Sauter (ETHZ, Zürich, Switzerland), Dieter Stüning (ZFMK, Bonn, Germany), Kevin Tuck (BMNH, London, UK), András Vojnits (formerly HNHM, Budapest, Hungary) and Vadim V. Zolotuhin (Ul'yanovsk, Russia). We are particularly grateful to Aleš Laštůvka (Prostějov, Czech Republic) and Zdeněk Laštůvka (Brno, Czech Republic) for their generous hospitality to EvN during a fieldtrip in 1992, and lots of information and advise. Kees van den Berg (RMNH, Leiden) is acknowledged for assisting with rearing and preparation of material and Cees van Achterberg (RMNH, Leiden) for the identification of and information on the parasites. We acknowledge Jarosław Buszko (University of Toruń, Poland) and Tomasz Baran (University of Rzeszów, Poland) for their information and constructive advice. We are grateful to Beata Sielewicz (Nature Conservation Officer in Lublin, Poland) for permits to work in protected areas and to Marcin Sielezniew for the photographs of Polish habitats and J. Chobotow for photographing the moth. J. van Tol (RMNH, Leiden) and Peter Huemer (Innsbruck, Austria) kindly commented an earlier version of this manuscript.

### References

- Anikin, V. V. 2001. Cheshuekrylye (Lepidoptera) Inzhnega Povolzh'ya. – Izvestiya Saratovskogo Gosudarstvennogo Universiteta. Seriya Biologicheskaya **special issue**: 214–258.
- Baudyš, E. 1925. Čtvrtý příspěvek k zoocecidiologickému prozkoumání Moravy a Slezska. [4. Beitrag zur zoocecidiologischen Durchforschung von Mähren und Schlesiens]. – Sborník Klubu Přírodovědeckého v Brně **7**: 1–87.
- Baudyš, E. 1931. Třetí příspěvek k rozšíření hálek na Slovensku. (Dritter Beitrag zur Verbreitung der Zoocecidiien in der Slowakei). – Sborník Přírodovědecké Společnosti v Moravské Ostravě **6**: 361–376.
- Baudyš, E. 1947. Šestý příspěvek k zoocecidiologickému prozkoumání Moravy a Slezska [Sixth contribution to the zoocecidiological research of Moravia and Silesia.]. – Sborník Vysoké Školy Zemědělské v Brně **C37**: 1–55.
- Borkowski, A. 1975. Studien an Nepticuliden (Lepidoptera). Teil VI. Die Verbreitung der Nepticuliden in Polen. – Polskie Pismo entomologiczne **45**: 487–535.
- Buhr, H. 1964. Bestimmungstabellen der Gallen (Zoo- und Phytocecidiien) an Pflanzen Mittel- und Nordeuropas, I. – Fischer, Jena. 761 pp.
- Burmann, K. & P. Huemer 1998. Beitrag zur Kenntnis der Microlepidopterenfauna des Monte Baldo in Norditalien (Insecta, Lepidoptera). – Studi Trentini di Scienze Naturali Acta Biologica **73**: 5–53.
- Busck, A. 1907. New american Tineina. – Proceedings of the Entomological Society of Washington **8**: 86–99.

- Busck, A. 1913. Two microlepidoptera injurious to Chestnut. – Proceedings of the Entomological Society of Washington **15**: 102–104.
- Buszko, J. & J. Nowacki 2000. The Lepidoptera of Poland. A distributional checklist. – Polish Entomological Monographs **1**: 1–178.
- Caradja, A. 1899. Zusammenstellung der bisher in Rumänien beobachteten Microlepidopteren. – Deutsche Entomologische Zeitschrift Iris **12**: 171–217.
- Caradja, A. 1901. Die Microlepidopteren Rumäniens. – Buletinul Societății de Știinte din București-Romania **10**: 109–168.
- Černík, L. F. 1942. Krankheiten und teratologische Missbildungen (auch typische Verletzungen - Minen) an Pflanzen der Olmützer Flora. 14. Teil (Nr. 1801–2000). – Verhandlungen des Naturforschenden Vereines in Brünn **73**: 1–51.
- Cristofolini, G. 1991. Taxonomic revision of *Cytisus* Desf. Sect. *Tubocytisus* DC. (Fabaceae). – Webbia **45** (2): 187–219.
- Cubas, P., C. Pardo & H. Tahiri 2002. Molecular approach to the phylogeny and systematics of *Cytisus* (Leguminosae) and related genera based on nucleotide sequences of nrDNA (ITS region) and cpDNA (trnL-trnF intergenic spacer). – Plant Systematics and Evolution **233** (3–4): 223–242.
- Curò, A. 1883. Saggio di un catalogo dei Lepidotteri d'Italia. – Bulletino della Società Entomologica Italiana **15**: 1–124.
- Czekelius, D. 1924. Beiträge zur Schmetterlingsfauna Siebenbürgens, vii [recte viii]. – Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt **72–74**: 248–257 [sep. paginated as 1–10].
- Duponchel, P. A. J. [1842–1845]. Nocturnes, Supplément 4. Pp. li–xc, 9–554. – In: J.-B. Godart (ed.), Histoire naturelle des Lépidoptères ou papillons de France. – Méquignon-Marvis, Paris.
- Emmet, A. M. 1976. Nepticulidae. Pp. 171–267. – In: J. Heath (ed.), The moths and butterflies of Great Britain and Ireland, 1. Micropterigidae – Heliozelidae. The moths and butterflies of Great Britain and Ireland **1**. – Blackwell Scientific Publications Ltd, Oxford & London.
- Evenhuis, N. L. & G. A. Samuelson 2004. The insect and spider collections of the world website. Bishop Museum. – <http://hbs.bishopmuseum.org/codens/codens-r-us.html>. [Accessed 28 October 2004]
- Gustafsson, B. 1981. Characters of systematic importance in European Nepticulidae larvae (Lepidoptera). – Entomologica Scandinavica **12** (1): 109–116.
- Gustafsson, B. & E. J. van Nieuwerken 1990. Larvae of Nepticulidae. Pp. 323–356. – In: R. Johansson, E. S. Nielsen, E.J. van Nieuwerken & B. Gustafsson (eds.), The Nepticulidae and Opostegidae (Lepidoptera) of NW Europe. Fauna Entomologica Scandinavica **23**. – E. J. Brill, Leiden.
- Hartig, F. 1964. Microlepidotteri della Venezia Tridentina e delle regione adiacenti. – Studi Trentini di Scienze Naturali **41** (3): 3–296.
- Hauder, F. 1912. Beitrag zur Mikrolepidopteren-Fauna Oberösterreichs. – Jahresbericht des Museums Francisco-Carolinum, Linz **1912**: 1–321.
- Hauder, F. 1924. Nachtrag zur Mikrolepidopteren-Fauna Oberösterreichs. – Jahresbericht der Oberösterreichischen Musealvereines **80**: 265–294.
- Herrich-Schäffer, G. A. W. 1847–1855. Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text, Revision und Supplement zu Jakob Hübner's Sammlung europäischer Schmetterlinge, 5, Die Schaben und Federmotten. – G. J. Manz, Regensburg. 394 pp., plates Tineides 1–124, Pterophorides 1–7, Micropteryges 1.
- Heywood, V. H. & P. W. Ball 1968. Leguminosae. Pp. 80–187. – In: T. G. Tutin & et al. (eds.), Flora Europaea **2**. – University Press, Cambridge.
- Hoare, R. J. B. 2000. A new genus of primitive Nepticulidae (Lepidoptera) from eastern Australia, with a revised diagnosis of nepticulid subfamilies. – Zoological Journal of the Linnean Society **128** (3): 289–317.
- Hrubý, K. 1964. Prodromus Lepidopter Slovenska. – Slovenska Akadémia Vied, Bratislava. 962 pp.
- ILDIS 2003. ILDIS World Database of Legumes. International Legume Database & Information Service. – <http://www.ildis.org/LegumeWeb/>. [Accessed 28 June 2004.]
- Issekutz, L. 1972. Die Schmetterlingsfauna des südlichen Burgenlandes. II. Teil: Microlepidoptera. – Wissenschaftliche Arbeiten aus dem Burgenland **49**: 1–129, 5 pls.
- Käss, E. & M. Wink 1995. Molecular phylogeny of the Papilionoideae (Family Leguminosae): rbcL gene sequences versus chemical taxonomy. – Botanica Acta **108**: 149–162.
- Käss, E. & M. Wink 1997. Phylogenetic Relationships in the Papilionoideae (Family Leguminosae) Based on Nucleotide Sequences of cpDNA (rbcL) and ncDNA (ITS 1 and 2). – Molecular Phylogenetics and Evolution **8** (1): 65–88.

- Kasy, F. 1985. Die Schmetterlingsfauna des Naturschutzgebietes "Pischeldorf Fischawiesen", östliches Niederösterreich. – Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen **36 Supplement**: 1–27.
- Klimesch, J. 1949. Über einige bemerkenswerte Oberösterreichische Lepidopterenfunde. – Naturkundliche Mitteilungen aus Oberösterreich **1** (1): 17–18.
- Klimesch, J. 1961. Ordnung Lepidoptera. 1. Teil. Pyralidina, Tortricina, Tineina, Eriocraniina und Micropterygina. Pp. 481–789. – In: H. Franz (ed.), Die Nordost-Alpen im Spiegel ihrer Landtierwelt. **2**. – Universitätsverlag Wagner, Innsbruck.
- Klimesch, J. 1968. Die Lepidopterenfauna Mazedoniens, 4. Microlepidoptera. – Posebno Izdanje. Prirodonaucen Muzej Skopje **5**: 1–201.
- Klimesch, J. 1972. *Dechiria nigrifasciata* (Wlsm.) (Lep., Nepticulidae). Ein Beitrag zur Genitalmorphologie und -kologie. – Mitteilungen der Münchner Entomologischen Gesellschaft **62**: 1–12.
- Klimesch, J. 1990. Microlepidoptera (Kleinschmetterlinge), 1. Pp. 332. – In: K. Kusdas & E. R. Reichl (eds.), Die Schmetterlinge Oberösterreichs. **6**. – Entomologische Arbeitsgemeinschaft am OÖ. Landesmuseum zu Linz, Linz.
- Krause, A. 1871. Ueber einige Schmetterlinge der Umgegend von Altenburg. – Stettiner Entomologische Zeitung **32**: 296–304.
- Laštůvka, A. & Z. Laštůvka 1994. Zur Kenntnis der Tschechoslowakischen *Trifurcula*-Arten (Lepidoptera, Nepticulidae). – Acta Universitatis Agriculturae (Brno), Facultas Agronomica **40** (3–4): 197–220.
- Laštůvka, A. & Z. Laštůvka 1997. Nepticulidae Mitteleuropas. Ein illustrierter Begleiter (Lepidoptera). – Konvoj, Brno. 229 pp.
- Laštůvka, Z. 1994. Motýli rozšířeného území CHKO Pálava. Lepidoptera of the protected landscape area Pálava. – Faculty of Agronomy, University of Agriculture, Brno. 118 pp.
- Laštůvka, Z. & A. Laštůvka 1994. Drei neue Arten der *Trifurcula pallidella*-Gruppe aus Mitteleuropa (Lepidoptera: Nepticulidae). – Entomologia Generalis **18** (3–4): 201–212.
- Mann, J. 1862. Verzeichniss der im Jahre 1851 bei Brussa in Kleinasien gesammelten Schmetterlinge. – Wiener Entomologische Monatschrift **6**: 356–371, 373–409.
- Mann, J. 1866. Aufzählung der im Jahre 1865 in der Dobrudscha gesammelten Schmetterlinge. – Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien **16**: 321–360, pl. 1B.
- Mann, J. 1869. Lepidoptern gesammelt während dreier Reisen nach Dalmatien in den Jahren 1850, 1862, 1868. – Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien **19**: 371–388.
- Mann, J. 1885. Beiträge zur Kenntnis der Microlepidopteren-Fauna der Erzherzogthümer Oesterreich ob und unter der Enns und Salzburgs. 13. Fortsetzung und Schluss. – Wiener Entomologische Zeitung **4**: 265–273.
- Mariani, M. 1938. Fauna Lepidopterorum Siciliae. – Memorie della Società Entomologica Italiana **17**: 129–187.
- Mariani, M. 1943. Fauna Lepidopterorum Italiae. Parte 1. Catalogo ragionato dei Lepidotteri d'Italia. Fasc. 2 e 3. – Giornale di Scienze Naturali ed Economiche Palermo **42** (20): 81–236.
- Morton, A. 2000. DMAP for Windows. Version 7.0e. – Alan Morton, Winkfield, Windsor, Berkshire. Computer programme. <http://www.dmap.co.uk/welcome.htm>.
- Müller-Rutz, J. 1922. Die Schmetterlinge der Schweiz. 4. Nachtrag. Mit Beschreibung neuer Arten und Formen. – Mitteilungen der Schweizerischen Entomologischen Gesellschaft **13** (5): 217–259.
- Nickerl, O. 1908. Die Motten Böhmens (Tineen). Beiträge zur Insekten-Fauna Böhmens VI. – Verlag der Gesellschaft für Physiokratie, Prag. 161 pp.
- Nieuwerken, E. J. van 1990. The *Trifurcula subnitidella* group (Lepidoptera: Nepticulidae): taxonomy, distribution and biology. – Tijdschrift voor Entomologie **133** (2): 205–238.
- Nieuwerken, E. J. van 1996. Nepticulidae, Opostegidae. Pp. 21–27, 300. – In: O. Karsholt & J. Razowski (eds.), The Lepidoptera of Europe. A distributional checklist. – Apollo Books, Stenstrup.
- Nieuwerken, E. J. van & R. Johansson 1986. *Trifurcula pallidella* (Duponchel) and *T. beirnei* Puplesis (Lepidoptera: Nepticulidae): nomenclature, taxonomy and distribution. – Entomologica Scandinavica **17**: 271–278.
- Nieuwerken, E. J. van, A. Laštůvka & Z. Laštůvka 2004a. Annotated catalogue of the Nepticulidae and Opostegidae (Lepidoptera: Nepticuloidea) of the Iberian Peninsula. – SHILAP Revista de Lepidopterología **32** (127): 211–260.
- Nieuwerken, E. J. van, V. V. Zolotuhin & A. Mistchenko 2004b. Nepticulidae from the Volga and Ural region (Lepidoptera). – Nota lepidopterologica **27** (2/3): 125–157.

- Nieukerken, E. J. van, A. Laštuvka & Z. Laštuvka 2005. Annotated catalogue of the Nepticulidae and Opostegidae (Lepidoptera: Nepticuloidea) of France and Corsica. – Nota Lepidopterologica **in prep.**
- NIMA 2004. GEONet Names Server. National Imagery and Mapping Agency. – <http://gnswww.nima.mil/geonames/GNS/index.jsp>. [Accessed 28 June 2004.]
- Pável, J. & F. Uhryk 1896. Microlepidoptera. Pp. Lepidoptera 53–82 [order separately paginated]. – In: Fauni Regni Hungariae. Animalium Hungariae hucusque cognitorum enumeratio systematica, 3. Arthropoda. 3. – Regia Societatis Scientiarum naturalium Hungarica, Budapest.
- Prohaska, K. & F. Hoffmann 1929. Die Schmetterlinge Steiermarks, 10. A. Systematischer Teil (Schluss). Die Kleinschmetterlinge. – Mitteilungen des Naturwissenschaftlichen Vereins für Steiermark **64/65**: 272–321.
- Pröse, H. & A. H. Segerer 1999. Checkliste der “Kleinschmetterlinge” Bayerns (Insecta: Lepidoptera). – Beiträge zur bayerischen Entomofaunistik **3**: 3–90.
- Puplesis, R. K. 1984. Novye vidy Molei-malyutok (Lepidoptera, Nepticulidae) iz juzhnogo Primor'ya. (New species of nepticulids (Lepidoptera, Nepticulidae) from southern Primorye). – Entomologicheskoe Obozrenie **63**: 111–125.
- Puplesis, R. 1994. The Nepticulidae of eastern Europe and Asia. Western, central and eastern parts. – Backhuys Publishers, Leiden. 1–290 pp.
- Rebel, H. 1899. Zweiter Beitrag zur Lepidopteren-Fauna Südtirols. – Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien **49**: 158–185, pl. 4.
- Schmid, A. 1886. Die Lepidopteren-Fauna der Regensburger Umgegend mit Kelheim und Worth. – Korrespondenz-Blatt des Naturwissenschaftlichen Vereines in Regensburg **40**: 19–224.
- Segerer, A. H. 1997. Verifikation älterer und fraglicher Regensburger Lepidopterenmeldungen (Insecta: Lepidoptera). – Beiträge zur bayerischen Entomofaunistik **2**: 177–265.
- Shaw, M. R. & R. R. Askew 1976. Ichneumonoidea (Hymenoptera) parasitic upon leaf-mining insects of the orders Lepidoptera, Hymenoptera and Coleoptera. – Ecological Entomology **1**: 127–133.
- Skala, H. 1933. Die Nepticulidae Oberösterreichs. – Entomologischer Anzeiger **13**: 64–66, 99–101, 118–119, 129–132, 149–152.
- Szőcs, J. 1965. Molylepkék I. Microlepidoptera I. 7. Család: Nepticulidae-Törpemolyok. – Fauna Hungarica **76**: 48–104.
- Tokár, Z., F. Slamka & G. Pastorális 1999. New and interesting records of Lepidoptera from Slovakia in 1995–1997. – Entomofauna Carpathica **11**: 43–57.
- Toll, S. 1936. Badania nad narządami płciowymi u *Pyrausta purpuralis* L. i *P. ostrinalis* Hb., tudzież opisy 11 nowych gatunków Microlepidoptera. Untersuchung der Genitalien bei *Pyrausta purpuralis* L. und *P. ostrinalis* Hb., nebst Beschreibung 11 neuer Microlepidopteren-Arten. – Annales Musei Zoologiczi Polonici **11** (24): 401–413, pls. 47–49.
- Toll, S. 1938. Microlepidoptera zebrane w latach 1934–1937 w powiatach zaleszczyckim i borszczowskim na Podolu. (Verzeichnis der in den Jahren 1934–1937 in den Kreisen Zaleszczyki und Borszczów (Podolien) aufgefundenen Mikrolepidopteren). – Sprawozdania Komisji Fizyograficznej oraz Materyały do Fizyografii Kraju. Kraków **73**: 133–221.
- Turland, N. J., L. Chilton & J. R. Press 1993. Flora of the Cretan area, annotated checklist & atlas. – HMSO, London. xii + 439 pp.
- Vorbrodt, C. 1932. Tessiner und Misoxer Schmetterlinge. II. “Microlepidoptera”. – Deutsche Entomologische Zeitschrift Iris **46**: 66–92.
- Wocke, M. F. 1874. Verzeichniss der Falter Schlesiens, 2. Microlepidoptera. – Zeitschrift für Entomologie (Breslau) **4**: 1–107.
- Wörz, A. 1958. Die Lepidopterenfauna von Württemberg. II. Microlepidopteren. Kleinschmetterlinge. – Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg **113**: 253–312.
- Zajac, A. & M. Zajac 2001. Atlas rozmieszczenia roślin naczyniowych chronionych w Polsce. Distribution atlas of vascular plants in Poland. – Instytutu Botaniki, Uniwersytetu Jagiellońskiego, Kraków. 716 pp.
- Zeller, P. C. 1848. Die Gattungen der mit Augendeckeln versehenen blattminirenden Schaben. – Linnaea entomologica **3**: 248–344.
- Zeller, P. C. 1850. Verzeichniss der von Herrn Jos. Mann beobachteten Toscanischen Microlepidoptera. – Stettiner Entomologische Zeitung **11**: 195–212.
- Zimmermann, F. 1944a. Zur Kenntnis der Verbreitung der Nepticuliden in den Reichsgauen Wien und Niederdonau (Lepidopt.). – Zeitschrift der Wiener Entomologischen Gesellschaft **29**: 3–6, 60–64, 78–91, 107–122.
- Zimmermann, F. 1944b. Zur Nepticulidenfauna des böhmischen Raumes (Lep.). – Entomologische Zeitschrift **58**: 11–14, 21–24.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nota lepidopterologica](#)

Jahr/Year: 2004

Band/Volume: [27](#)

Autor(en)/Author(s): van Nieukerken Erik J., Mazurkiewicz Anna, Palka Krzysztof

Artikel/Article: [Trifurcula pallidella \(Duponchel, 1843\) \(Nepticulidae\): distribution, biology and immature stages, particularly in Poland 159-178](#)