

Genetic confirmation of *Aricia artaxerxes* (Fabricius, 1793) (Lepidoptera, Lycaenidae) in the Czech Republic, its conservation significance and biogeographic context

ALENA SUCHÁČKOVÁ BARTOŇOVÁ^{1,2}, JIŘÍ BENEŠ², ZDENĚK FALTÝNEK FRIC², MARTIN KONVIČKA^{1,2}

¹ University of South Bohemia in Ceske Budejovice, Faculty of Science, Branisovska 1760, Ceske Budejovice, Czech Republic

² Biology Centre, Czech Academy of Sciences, Institute of Entomology, Branisovska 1160/31, Ceske Budejovice, Czech Republic

<http://zoobank.org/5EB7C3A3-5BBE-468B-82DC-6D7D1676A679>

Received 6 August 2019; accepted 14 October 2019; published: 8 November 2019

Subject Editor: Maria Heikkilä.

Abstract. We report here the first molecular evidence for the occurrence of *Aricia artaxerxes* (Fabricius, 1793) (Lepidoptera: Lycaenidae) in the Czech Republic. In Central Europe, this species may co-occur with its more common sibling, *Aricia agestis* (Denis & Schiffermüller, 1775). We sequenced the cytochrome c oxidase subunit 1 of darkly-coloured, putative *A. artaxerxes* specimens in the Czech Republic. We confirmed *A. artaxerxes* only from a limestone area in South Bohemia (Vyšenské kopce National Nature Reserve), which is probably the only locality of the species in the Czech Republic. This area is located at ca. 550 m A.S.L., showing that the elevation overlap with *A. agestis* could be high in Central Europe. Other surveyed individuals were confirmed as *A. agestis*, with a minimum p-distance of 1.98% between the two species. The South Bohemian area of occurrence is probably highly isolated (approx. 190 km) from localities of the species in neighbouring countries, highlighting the conservation importance of the *A. artaxerxes* population and of the insular calcareous areas in the Šumava Mountains foothills. We used database sequences of *A. artaxerxes* to place the Czech population into a wider phylogeographic context. The Czech population is monomorphic, consisting of a single haplotype, which is present from Scandinavia through Germany to Central Asia.

Introduction

Aricia artaxerxes (Fabricius, 1793) is a Palaearctic species of lycaenid butterfly, occurring in Europe together with several cryptic siblings: the widely distributed *Aricia agestis* (Denis & Schiffermüller, 1775), the West Mediterranean *A. cramera* (Eschscholtz, 1821) and *A. montensis* Verity, 1928. *Aricia agestis* and *A. artaxerxes*, which are sympatric in Central Europe, differ in larval and pupal morphology, and rearing was traditionally used to distinguish them (Kames 1976; Lepidopterologen-Arbeitsgruppe 1987; Warecki 2010). Several studies combining morphology and molecular markers (Aagaard et al. 2002; Dincă et al. 2011; Sañudo-Restrepo et al. 2013) concluded that these taxa indeed represent true species and can be distinguished by allozyme profiles or the cytochrome c oxidase subunit 1 (COI, i.e., the standard DNA barcode for animals), but not by wing pattern or genital morphology. More specifically, adults from Scotland (nominotypical *A. artaxerxes artaxerxes* Fabricius, 1793) usually bear a white discoidal dot on the upper side of each brown fore wing. However, adults from mainland Europe (*A. artaxerxes allous* (Hübner, 1819), and other areas, cf. Sañudo-Restrepo et al. 2013) lack that trait, differing from adult *A. agestis* only

by a darker overall colouration, due to smaller or obscured orange spots on the upper sides of the fore wings, but also smaller spots on the hind wings (Tolman and Lewington 2008, p. 130).

Ecologically and biogeographically, the majority of *A. artaxerxes* records appear concentrated in calcareous short-turf grasslands at high latitudes or elevations (Lepidopterologen-Arbeitsgruppe 1987; Asher *et al.* 2001; Aagaard *et al.* 2002; Sañudo-Restrepo *et al.* 2013; Pecsénye *et al.* 2014), where the species has a single annual generation from June to August. Its confirmed larval host plants across the range include *Helianthemum* Mill. spp. (Cistaceae; *H. nummularium* (L.) Mill. in Britain: Asher *et al.* 2001) and large-flowered *Geranium* L. spp. (Geraniaceae; *G. sanguineum* L. in Poland: Warecki 2010; see also Tuzov 2000; Gorbunov and Kosterin 2003). The distribution of *A. agestis*, in contrast, includes most of the European continent except for the highest latitudes, and extends to the Tian Shan in Asia. It has two or three annual generations in the lowlands of Central Europe, and three or even more in more southerly regions (Hesselbarth *et al.* 1995). Its host plant range overlaps with *A. artaxerxes*, as it includes both many Geraniaceae (e.g., *Geranium dissectum* L., *G. pusillum* Burm. fil., *G. pratense* L. and *G. molle* L., and *Erodium* L'Her. ex Aiton spp.) and *Helianthemum* spp. in natural conditions (Thomas *et al.* 2001; Buckley *et al.* 2012). Reflecting the broad ecological range of the host plants used, *A. agestis* inhabits a broad range of biotopes, from xeric grasslands to mesic and even alluvial meadows (Bury 2016). In European mid-latitudes, the single generation of *A. artaxerxes* coincides with the gap between spring and summer generations of *A. agestis* (Beneš *et al.* 2002). *Aricia artaxerxes* is usually associated with higher elevations than *A. agestis* in Central and Southern Europe, but their occurrence could overlap at intermediate elevations, causing problems with identification in the potential contact zones (cf. Dincă *et al.* 2011).

Until the late 20th century, the two species were not distinguished in the lowlands of Central Europe, including in the Czech Republic. The first records of *A. artaxerxes* for the country are from 1964 (Králíček *et al.* 1970) and 1971 (Králíček and Gottwald 1980), from the warm region of South-Eastern Moravia (foothills of the White Carpathians Mts). These and later records were mainly identified based on the combination of flight period and a darker colouration than the more common *A. agestis*. The distribution atlas of Czech butterflies (Beneš *et al.* 2002) considered only the records of *A. artaxerxes* in a single atlas grid square (Vyšenské Kopce National Nature Reserve [= NNR] near Český Krumlov, South Bohemia) as recent and relatively reliable. The Czech Republic Butterflies and Moths Recording Database (Institute of Entomology, Biology Centre of the Czech Academy of Sciences) together with the Information System of Nature Conservation (Nature Conservation Agency of the Czech Republic) register possible, unreliable records of *A. artaxerxes* from 38 grid squares (as opposed to 427 for *A. agestis* – 63% of the country's area) (Fig. 1). The national Red list (Hejda *et al.* 2017) considers *A. artaxerxes* as critically endangered and *A. agestis* as of least concern.

The recent increase in the use of molecular markers for species identification, the emergence of national barcoding programmes (e.g., Dincă *et al.* 2011; Hausmann *et al.* 2011; Litman *et al.* 2018) and butterfly recording in European countries have significantly improved our knowledge of *A. artaxerxes* distribution. Besides the previously known distribution at high elevations, the species was reliably confirmed from relatively low elevations of Baden-Württemberg (at ca. 850 m A.S.L.), Saxony-Anhalt (ca. 430 m A.S.L.), and Thuringia (ca. 370 m A.S.L.) (Hausmann *et al.* 2011; and specimens used in Mutanen *et al.* 2016). It is also reported from central-northern Poland (Buszko and Masłowski 2008; Sielezniew and Dziekanska 2010), northern Hungary (800–850 m A.S.L.) (Pecsénye *et al.* 2014) and central-eastern Slovakia (600–800 m A.S.L.) (Reiprich and




Figure 1. Distribution map of *Aricia artaxerxes* and *A. agestis* in the Czech Republic. The map was created based on 8464 records both historical and recent (until 2018) of these two species from The Czech Republic Butterflies and Moths Recording Database (Institute of Entomology, Biology Centre of the Czech Academy of Sciences) and from the Information System of Nature Conservation (Nature Conservation Agency of the Czech Republic). Grey squares: *A. agestis*, crosses: probably darkly-coloured *A. agestis* (putative *A. artaxerxes* records), white points: barcoded individuals, black circle: confirmed *A. artaxerxes*.

Okáli 1988–1989; Pecsénye et al. 2014; L. Vítáz, personal communication), but as far as we know, these populations have not been studied using molecular sequencing.

In this contribution, we use DNA barcodes to clarify the status of *A. artaxerxes* in the Czech Republic. Specifically, we barcoded material from the putative *A. artaxerxes* population near Český Krumlov, and individuals with missing or reduced orange spots on upper sides of the wings (hereinafter “dark”) from several localities across the country with past *A. artaxerxes* reports, and a selection of typical *A. agestis* (with developed orange spots, hereinafter “light”). In addition to evaluating the current status of *A. artaxerxes* in the country, we place it into a wider biogeographic context and discuss its conservation significance.

Material and methods

We tested twelve individuals from the putative *Aricia artaxerxes* population from Vyšenské kopce NNR, South Bohemia (univoltine, dark, occurring between spring and summer broods of sympatric *A. agestis*; but also including reared individuals without reduced orange spots originating from dark females), five dark or intermediate individuals from other localities, and nine light individuals with developed orange spots (Table 1, Fig. 2). To place these individuals into a broader context, we further mined a total of 113 sequences from GenBank: *A. artaxerxes* (N=94), *A. agestis* (N=9), *A. anteros* (N=3), *A. cramer* (N=3) and *A. montensis* (N=4) (Suppl. materials 1, 3).

Table 1. Sampling sites of *Aricia artaxerxes* and *A. agestis* from the Czech Republic used for DNA-based identification.

Species	Voucher	Colouration	Locality	GenBank codes	Elevation [m A.S.L.]	Grid square
<i>A. artaxerxes</i>	6 specimens	dark	Vyšenské kopce National Nature Reserve, South Bohemia	MN107398	550	7151
	ZF-LY-001622 JB00138			–		
	ZF-LY-001774 to 1776, 1778			MN107403		
	AB24-1 to 6	light	Vyšenské kopce National Nature Reserve, South Bohemia – reared	MN107404 – MN107409	550	7151
<i>A. agestis</i>	JB00137	dark	Čepičná Nature Reserve, South Bohemia	MN107393	500	6747
	ZF-LY-001779	dark	Koněprusy, Čertovy schody quarry, Central Bohemia	MN107390	400	6050
	ZF-LY-001785	dark	Martinice, Velké Meziříčí, Vysočina region	MN107392	470	6662
	ArAg2	dark	Čimické údolí Nature Monument, Prague	MN107394	250	5852
	ArAg1	dark	Prokopské údolí Nature Reserve, Prague	MN107395	250	5952
	JB00124	light	Nerestský lom Nature Monument, South Bohemia	MN107396	470	6450
	ZF-LY-001784	light	Černousy, Frýdlant, Liberec region	MN107391	250	4956
	AB30-23	light	Nová Ves, Litovel, Olomouc region	MN107384	300	6368
	AB30-28	light	Mohelenská hadcová step National Nature Reserve, Vysočina region	MN107385	350	6863
	AB44-8	light	Hraniční hill, Město Albrechtice, Moravian-Silesian region	MN107386	500	5871
	AB44-9	light	Město Libavá, Olomouc region	MN107387	550	6271
	AB44-10	light	Chomýž, Krnov, Moravian-Silesian region	MN107388	350	5871
	AB44-11	light	Toužinské stráně Nature Monument, South Bohemia	MN107389	470	6958
	KA-2698	light	Havranické vřesoviště, Podyjí National Park, South Moravia	MN107397	330	7162

DNA was extracted from the legs using the Genomic DNA Mini Kit – Tissue (Geneaid) following the manufacturer's protocols. Using the Polymerase Chain Reaction (PCR), we amplified the mitochondrial gene cytochrome c oxidase subunit 1 gene (barcode/COI). We used the primer pair LCO/Nancy; or LCO/K699 and RON/HCO in case of fragmented DNA (primers: Monteiro and Pierce 2001; Wahlberg and Wheat 2008). We prepared the PCR mixture in 12.5 µl volume (6.25 µl Bioline 2× MyTaq HS Red Mix, 4 µl PCR H₂O, 0.625 + 0.625 µl primers; 1 µl DNA). The thermal cycling profile was 95 °C for 5 min; then 40 cycles of 94 °C for 30 s, 50 °C for 30 s, 72 °C for 90 s; and final extension 72 °C for 10 min. PCR products were cleaned with enzymes FastAP and ExoI (Thermofisher) and sequenced in one direction in Macrogen Inc. on ABI3730XL DNA analysers. Sequences were checked visually and aligned in GENEIOUS v. 8.0.5 (Kearse *et al.* 2012). Obtained sequences were submitted to the GenBank database (<https://www.ncbi.nlm.nih.gov/genbank/>; accession codes MN107385–MN107409, Suppl. material 1).




Figure 2. Examples of the sequenced specimens of *Aricia artaxerxes* and *A. agestis*. **a.** *Aricia artaxerxes*, (ZF-LY-001623, MN107399, Vyšenské kopce National Nature Reserve). **b, c.** *Aricia agestis*, darkly coloured individuals (**b.** JB00137, MN107393, Čepičná Nature Reserve. **c.** ZF-LY-001785, MN107392, Martinice, Velké Meziříčí). **d.** *Aricia agestis*, a typical, lightly coloured individual with developed orange spots (AB44-11, MN107389, Toužinské stráně Nature Monument).

The BLAST algorithm (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) with GenBank reference database assigned the sequences to either *A. artaxerxes* or *A. agestis*. MEGA7 (Kumar et al. 2016) was used for computing p-distances (i.e., proportion of nucleotide site differences) among sequences and groups of sequences (species). We mined 94 sequences of *A. artaxerxes* from GenBank (Suppl. materials 1, 3), which we used, together with the twelve sequences of Czech *A. artaxerxes*, to construct the TCS statistical parsimony haplotype network (Clement et al. 2000) in the program POPART (Leigh and Bryant 2015). Using one sample of each *A. artaxerxes* haplotype from GenBank, together with nine randomly chosen samples of European *A. agestis* and representatives of *A. anteros*, *A. cramera* and *A. montensis* (Suppl. materials 1, 2), we reconstructed the Maximum Likelihood tree of all tested specimens using the IQTREE web server (Nguyen et al. 2015; Trifinopoulos et al. 2016) with ultrafast bootstrap (Minh et al. 2013).

Results

All darkly coloured as well as the reared light putative *Aricia artaxerxes* individuals from Vyšenské Kopce NNR reserve (N=12) were unequivocally *A. artaxerxes* according to both BLAST and Maximum Likelihood analysis (Fig. 3). All other tested individuals, both darkly- and lightly-coloured,




Figure 3. Maximum Likelihood tree of the Czech Republic *Aricia* spp. butterflies. The tree is based on cytochrome c oxidase subunit I (DNA barcode) sequences obtained for this study (Czech material) and retrieved from GenBank. The sample AY556977 from Morocco is labelled as *Aricia artaxerxes* in GenBank, This specimen was renamed as *A. montensis*, which occurs in North Africa (Sañudo-Restrepo *et al.* 2013). Branch labels show bootstrap values.

were *A. agestis*. The minimum p-distance between Czech *A. artaxerxes* and *A. agestis* was 1.98% and the mean p-distance was 2.10%.

The Czech *Aricia artaxerxes* population is monomorphic, consisting of a single COI haplotype (Fig. 4). Comparing the published barcodes of *A. artaxerxes*, this haplotype (H1) is widely distributed in the Palearctic (Central Asia, Western Russia, Crimea, Central Germany, both Eastern and Western Alps and European North), and its derived haplotypes (Haplogroup A) are present in Greece, Baden-Württemberg, and reach the Far East. The other widespread haplotype (H2) and its derived haplotypes have a more southerly distribution – in the Balkans, Alps, and reach the United Kingdom. The highest haplotype diversity is in the Western Alps where five different haplotypes are present in twelve samples. However, as reported previously (Sañudo-Restrepo *et al.* 2013), the overall p-distance is low within this species (mean = 0.20%, max = 1.10%).




Figure 4. TCS haplotype network of *Aricia artaxerxes*. The network is based on cytochrome c oxidase subunit 1 (DNA barcode) sequences obtained for this study (Czech material) and retrieved from GenBank. Circles represent individual haplotypes, black dots represent putative haplotypes (not found among samples), and each connecting line represents one mutation.

Discussion

Using DNA barcoding, we confirmed that the putative *Aricia artaxerxes* population from Vyšenské Kopce NNR, Czech Republic, indeed belongs to this species, whereas other darkly-coloured samples from the country belong to *Aricia agestis* (Fig. 2). The fact that the reared individuals of *A. artaxerxes* resemble *A. agestis* suggests a phenotypic plasticity also in this species. Moreover, the Czech population is situated at a relatively low elevation (ca. 550 m A.S.L.). Relying on wing morphology or elevational records thus can lead to misidentifications between these two species. Our results highlight the need for DNA-based identification of these two species within their overlapping range (cf. Dincă et al. 2011).

Besides providing confirmation of *A. artaxerxes* distribution, our results indicate that its Czech population is isolated by ca. 190 km from the nearest genetically confirmed population of the species in Europe, possesses an extremely narrow distribution, and rightly deserves its critically endangered status (cf. Hejda et al. 2017).

This finding raises the national and even Central European importance of the Vyšenské Kopce NNR (area: 66.7 ha), plus two reserves within its closest environs (Výří vrch Nature Monument 12.9 ha, and Cvičák Nature Monument, 61.4 ha), for butterfly conservation. These three reserves, parts of the Blanský Les Landscape Protected Area, protect a system of (sub)xerophilous grasslands, shrublands and sparse woodlands on metamorphous limestones, a rare phenomenon within the prevailing base-poor, cold and damp north-eastern foothills of the Šumava Mts. These calcareous localities have been rescued from successional overgrowth by restoration activities, partly depending on volunteers, since the 1970s onwards (Albrecht 2003); the Cvičák Nature Monument was for a time used by the military (cf. Čížek *et al.* 2013). The current conservation management of these sites is exemplary, aiming at retaining a dynamic mosaic of coppiced woodlands, scrub and grasslands maintained by mowing and goat and cattle grazing of varying intensity. The reserves host a regionally outstanding butterfly diversity (Hanč 2005, 2011; Slámová *et al.* 2013). Within this system, *A. artaxerxes* inhabits short-sward, open-turf and flower-rich sites with dense growth of *Helianthemum grandiflorum obscurum* (most likely the local larval host), which cover an area not exceeding 20 ha in total (Fig. 5a).

The entire area belongs to a wider system of calcareous islets adjoining the Šumava Mts. in an approximately NW-SE direction, and supporting multiple range-restricted relict xerophilous plants and animals (Fig. 5b). These include species with southern distribution (e.g., the bush-cricket *Leptophyes bosci* Fieber, 1853 (Holuša *et al.* 2013)); south-western elements believed to have colonised Czech territory via south-western mountain passes (e.g., the lycaenid butterfly *Pseudophilotes baton* (Bergsträsser, 1779) (Konvička *et al.* 2008)); and alpine floral elements that likely descended to the Danube lowlands in colder periods of the Pleistocene (e.g., *Verbascum chaixii* subs. *austriacum* (R. *et* Sch.) Hayek (cf. Holub and Skalický 1959; Kaplan 2012)). The population of *A. artaxerxes* appears as another example of this “perialpine” element, a relict of a time when the currently high-elevation species descended to northern foothills of the Alps (Schmitt 2017). Our *Aricia* sp. samples included material from yet another calcareous islet of the Šumava Mts. foothills (Čepičná, see Table 1). Although the presence of *A. artaxerxes* has not yet been confirmed there, the possibility of its occurrence there or in other localities makes further surveys desirable.

On the other hand, we are sceptical regarding the earlier (1970s–1980s) records of *A. artaxerxes* from eastern parts of the Czech Republic (south-eastern Moravia). The records were not followed by subsequent observations, despite intensive recording in the region (e.g., Bělín and Gottwald 2001; Konvička *et al.* 2008; Spitzer and Beneš 2010; Uříčář *et al.* 2016). We have shown here that even *A. agestis* occasionally forms darker variants (Fig. 2). In addition, no *A. artaxerxes* records from adjoining Western Slovakia are known to Slovakian colleagues; the closest known populations of putative *A. artaxerxes* are from Central Slovakia (near Východná and Važec), more than 100 km distant from the East Moravian localities (L. Vířaz, personal communication).

In the wider geographical context, the overall COI genetic diversity within *A. artaxerxes* is low when compared to its sibling *A. agestis* (Sañudo-Restrepo *et al.* 2013; Vodá *et al.* 2015).

The mitochondrial haplotype network (Fig. 4) does not show pronounced genetic structure within *A. artaxerxes* and the following interpretation needs to be taken with caution. The network suggests the possible existence of two haplogroups, differing by a single mutation only, distributed in wide longitudinal belts across the Palaearctic. The first (haplogroup A) is distributed from the Far East to Scandinavia, including also the Eastern Alps and the German and Czech populations. The second (haplogroup B) has a more southerly distribution, in Romania and the Western Alps, but also




Figure 5. Calcareous localities adjoining the Šumava Mts. chain. **a.** Habitat of the Czech Republic population of *Aricia artaxerxes*, short-sward calcareous grassland near Český Krumlov, South Bohemia. **b.** Calcareous localities adjoining the Šumava Mts. chain in the wider geographic context. Although numerous, all the localities represent small and relatively isolated patches within predominately acidic bedrock, and the majority of their summed area is forested. The closest contiguous limestone areas are the Northern Alps, situated to the south, and the Franconian Jura, situated to the west. Blue field and coloured dots: calcareous areas. Yellow dot: Čepičná Nature Reserve, Red dot: Vyšenské kopce National Nature Reserve.

in northern Great Britain. Both haplogroups are present in the Western Alps, where several unique haplotypes also exist, and in Greece. The presence of unique haplotypes suggests that the species survived the glacial maxima in these topographically diverse regions, possibly on mountain foothills (cf. Schmitt 2017). Moreover, these areas could also represent a secondary contact zone for the two haplogroups. Samples from the Italian Peninsula, which are still lacking in the dataset, could shed more light on this problem. The Palaearctic-wide distribution of haplogroup A could either point to a rapid postglacial expansion from a refugium, or to a more continuous distribution during glacials, as detected also for other continental species (Marešová *et al.* 2019; Bartoňová *et al.* 2018). We incline towards the second option, as postglacial dispersal from a restricted refugium to the wide belt stretching from the Alps to the Altai Mts. seems unlikely. Unique haplotypes were detected in still more southerly areas (H7 – Uzbekistan, H11 – Greece, H12 – Armenia), but none was separated from the most frequent haplotypes (H1, H2) by more than three mutations. Their existence might indicate further structuring of populations in these areas, but more data are needed to ascertain this.

In any case, *Aricia artaxerxes* constitutes a boreomontane species with the European part of its range divided into a northern area where it frequents short-sward grasslands with the warmest microclimates, and a more southerly area where it inhabits localities of a similar character, but usually situated at high elevations. Adjoining the southern mountains, there exist isolated relict sites in relatively low elevations, potentially threatened by successional changes. This situation is remarkably similar to other butterfly species occurring both at high elevation and on piedmont grasslands, whose piedmont sites have been disappearing due to modern land use changes, whereas the high-elevation populations still hold on (e.g., *Lasiommata petropolitana* (Fabricius, 1787) (Spitzer *et al.* 2018) or *Parnassius apollo* (Linnaeus, 1758) (Todisco *et al.* 2010; Habel *et al.* 2012)).

Acknowledgements

We would like to thank Zdeněk Hanč, Tomáš Kadlec, Václav Krivan, Michal Plátek, Lukáš Spitzer, Bohumil Vodrlind, and Zdeněk Weidenhoffer for help with obtaining material, and to Jaroslav Buszko, Zdeněk Hanč, Petr Lepší, Pavel Marhoul and Lubomír Vítáz for valuable suggestions on the manuscript. We thank Jan Habel and Vlad Dincă for their reviews. We would like to thank the Nature Conservation Agency of the Czech Republic for the records compiled in this study. The study was supported by the University of South Bohemia in Ceske Budejovice (152/2016/P).

References

- Aagaard K, Hindar K., Pullin AS, James CH, Hammarstedt O, Balstad T, Hanssen O (2002) Phylogenetic relationships in brown argus butterflies (Lepidoptera: Lycaenidae: *Aricia*) from northwestern Europe. *Biological Journal of the Linnean Society* 75(1): 27–37. <https://doi.org/10.1046/j.1095-8312.2002.00004.x>
- Albrecht J (2003) Českobudějovicko. In: Mackovčín P, Sedláček M (Eds) *Chráněná území ČR VIII. Agentura Ochrany Přírody a Krajiny ČR and EkoCentrum Brno, Praha*, 808 pp.
- Asher J, Warren M, Fox R, Harding P, Jeffcoate G, Jeffcoate S (2001) *The Millennium Atlas of Butterflies in Britain and Ireland*. Oxford University Press, New York, 456 pp.
- Bartoňová A, Konvička M, Korb S, Kramp K, Schmitt T, Faltýnek Fric Z (2018) Range dynamics of Palaearctic steppe species under glacial cycles: the phylogeography of *Protorebia afra* (Lepidoptera: Nymphalidae: Satyrinae). *Biological Journal of the Linnean Society* 125(4) 867–884. <https://doi.org/10.1093/biolinnean/bly136>

- Bělin V, Gottwald A (2001) Motýli Bílých a Bielych Karpat. Přírodovědný klub v Uherském Hradišti, Uherské Hradiště, 153 pp.
- Beneš J, Konvička M, Dvořák J, Fric Z, Havelda Z, Pavlíčko A, Vrabec V, Weidenhoffer Z (2002) Motýli České republiky: Rozšíření a ochrana I, II. SOM, Praha, 857 pp.
- Buckley J, Butlin RK, Bridle JR (2011) Evidence for evolutionary change associated with the recent range expansion of the British butterfly, *Aricia agestis*, in response to climate change. *Molecular Ecology* 21(2): 267–280. <https://doi.org/10.1111/j.1365-294X.2011.05388.x>
- Bury J (2016) New data on *Aricia agestis* (Lepidoptera: Lycaenidae), its life history and occurrence in the Podkarpacie region of Poland. *Fragmenta Faunistica* 59(1): 29–37. <https://doi.org/10.3161/00159301FF2016.59.1.029>
- Buszko J, Maslowski J (2008) Motyle dzienne Polski (Lepidoptera: Hesperioidea, Papilionoidea). Koliber, Kraków, 274 pp.
- Čížek O, Vrba P, Beneš J, Hrázský Z, Koptík J, Kučera T, Marhoul P, Zámečník J, Konvička M (2013) Conservation potential of abandoned military areas matches that of established reserves: plants and butterflies in the Czech Republic. *PLOS ONE* 8(1): e53124. <https://doi.org/10.1371/journal.pone.0053124>
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology* 9(10): 1657–1659. <https://doi.org/10.1046/j.1365-294x.2000.01020.x>
- Dincă V, Zakharov EV, Hebert PDN, Vila R (2011) Complete DNA barcode reference library for a country's butterfly fauna reveals high performance for temperate Europe. *Proceedings of the Royal Society B: Biological Sciences* 278(1704): 347–355. <https://doi.org/10.1098/rspb.2010.1089>
- Gorbunov P, Kosterin O (2003) The butterflies of north Asia in nature (Hesperioidea and Papilionoidea) (Asian part of Russia), Volume 1. Publishers Koreschkov, Krylov, Moscow, 391 pp.
- Habel JC, Reuter M, Drees C, Pfaender J (2012) Does isolation affect phenotypic variability and fluctuating asymmetry in the endangered Red Apollo? *Journal of Insect Conservation* 16(4): 571–579. <https://doi.org/10.1007/s10841-011-9442-3>
- Hanč Z (2005) Denní motýli (Lepidoptera: Papilionoidea, Hesperioidea) Národní přírodní rezervace Vyšenské kopce. *Klapalekiana* 41: 33–42.
- Hanč Z (2011) Denní motýli a vřetenušky (Lepidoptera, Papilionoidea a Zygaenidae) Chráněné krajinné oblasti Blanský les. *Sborník Jihočeského Muzea v Českých Budějovicích, Přírodní vědy* 51: 101–117.
- Hanč Z, Beneš J, Faltýnek Fric Z, Pavlíčko A, Zapletal M (2019) Atlas denních motýlů a vřetenušek jižních Čech. Jihočeský kraj, České Budějovice, 437 pp.
- Hausmann A, Haszprunar G, Segerer AH, Speidel W, Behounek G, Hebert PDN (2011) Now DNA-barcoded: the butterflies and larger moths of Germany. *Spixiana* 34(1): 47–58.
- Hejda R, Farkač J, Chobot K (2017) Červený seznam ohrožených druhů České republiky. Bezobratlí. Agentura ochrany přírody a krajiny České republiky, Praha, 611 pp.
- Hesselbarth G, Oorschot van H, Wagener S (1995) Die Tagfalter der Türkei unter Berücksichtigung der angrenzenden Länder. Goecke & Evers, Keltern, 1354 pp.
- Holub J, Skalický V (1959) Floristicko-fytogeografické poznámky ke květeně jihovýchodní části Šumavy a přilehlé části Předšumaví. *Preslia* 31: 395–412.
- Holuša J, Kočárek P, Vlk R, Marhoul P (2013) Annotated checklist of the grasshoppers and crickets (Orthoptera) of the Czech Republic. *Zootaxa* 3616(5): 437–460. <https://doi.org/10.11646/zootaxa.3616.5.2>
- Kames P (1976) Die Aufklärung des Differenzierungsgrades und der Phylogenese der beiden *Aricia*-Arten *agestis* Den. et Schiff. und *artaxerxes* Fabr. (allous G.-Hb.) mit Hilfe von Eizuchten und Kreuzungsversuchen (Lep., Lycaenidae). *Mitteilungen der entomologischen Gesellschaft Basel, N. F.* 26: 7–13, 29–64.
- Kaplan Z (2012) Flóra a fytogeografie České republiky. *Preslia* 84: 505–573.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: an integrated and extend-

- able desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12): 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Konvička M, Dvořák L, Hanč Z, Pavlíčko A, Fric Z (2008) The Baton blue (*Pseudophilotes baton*) (Lepidoptera: Lycaenidae) in south-western Bohemia: iron curtain, military ranges and endangered butterfly. *Silva Gabreta* 14: 187–198.
- Králíček M, Gottwald A (1980) Motýli Jihovýchodní Moravy I. Muzeum J.A. Komenského Uherské Hradiště, 112 pp.
- Králíček M, Marek J, Povolný D (1970) Významné a nové nálezy Lepidopter z Moravy a Slovenska. *Ochrana fauny* (Bratislava) 4: 1–8.
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33(7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Leigh JW, Bryant D (2015) POPART: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6(9): 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- Lepidopterologen-Arbeitsgruppe (1987) Schmetterlinge und ihre Lebensräume, Band 1. Pro Natura – Schweizerischer Bund für Naturschutz, Basel, 516 pp.
- Litman J, Chittaro Y, Birrer S, Praz C, Wermeille E, Fluri M, Stalling T, Schmid S, Wyler S, Gonseth Y (2018) A DNA barcode reference library for Swiss butterflies and forester moths as a tool for species identification, systematics and conservation. *PLOS ONE* 13(12): e0208639. <https://doi.org/10.1371/journal.pone.0208639>
- Marešová J, Habel JC, Neve G, Sielezniew M, Bartoňová A, Kostro-Ambroziak A, Faltýnek Fric Z (2019) Cross-continental phylogeography of two Holarctic Nymphalid butterflies, *Boloria eunomia* and *Boloria selene*. *PLOS ONE* 14(3): e0214483. <https://doi.org/10.1371/journal.pone.0214483>
- Minh BQ, Nguyen MAT, Haeseler von A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- Monteiro A, Pierce NE (2001) Phylogeny of *Bicyclus* (Lepidoptera: Nymphalidae) inferred from COI, COII, and EF-1 α gene sequences. *Molecular Phylogenetics and Evolution* 18(2): 264–281. <https://doi.org/10.1006/mpev.2000.0872>
- Mutanen M, Kivelä SM, Vos RA, Doorenweerd C, Ratnasingham S, Hausmann A, Huemer P, Dincă V, van Nieukerken EJ, Lopez-Vaamonde C, Vila R, Aarvik L, Decaëns T, Efetov KA, Hebert PDN, Johnsen A, Karsholt O, Pentinsaari M, Rougerie R, Segerer A, Tarmann G, Zahir R, Godfray HCJ (2016) Species-level para- and polyphyly in DNA Barcode gene trees: strong operational bias in European Lepidoptera. *Systematic Biology* 65(6): 1024–1040. <https://doi.org/10.1093/sysbio/syw044>
- Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Pecsénye K, Rác R, Bereczki J, Batori E, Varga Z (2014) Loss of genetic variation in declining populations of *Aricia artaxerxes* in Northern Hungary. *Journal of Insect Conservation* 18(2): 233–243. <https://doi.org/10.1007/s10841-014-9634-8>
- Reiprich A, Okáli I (1988–1989) Dodatky k Prodrumu Lepidopter Slovenska. Biologické práce SAV, Bratislava, zv. 1 (1988), 135 pp., zv. 2 (1989), 106 pp., zv. 3 (1989), 139 pp.
- Sañudo-Restrepo CP, Dincă V, Talavera G, Vila R (2013) Biogeography and systematics of *Aricia* butterflies (Lepidoptera, Lycaenidae). *Molecular Phylogenetics and Evolution* 66(1): 369–379. <https://doi.org/10.1016/j.ympev.2012.10.010>

- Schmitt T (2017) Molecular biogeography of the high mountain systems of Europe: an overview. In: Catalan J, Ninot JM, Aniz MM (Eds) High Mountain Conservation in a Changing World. Springer International Publishing, Cham, 63–74. https://doi.org/10.1007/978-3-319-55982-7_3
- Sielezniew M, Dziekańska I (2010) Motyle dzienne. Multico, Warszawa, 336 pp.
- Slámová I, Klečka J, Konvička M (2013) Woodland and grassland mosaic from a butterfly perspective: habitat use by *Erebia aethiops* (Lepidoptera: Satyridae). Insect Conservation and Diversity 6(3): 243–254. <https://doi.org/10.1111/j.1752-4598.2012.00212.x>
- Spitzer L, Beneš J (2010) Nové a významné nálezy denních motýlů a vřetenuškovitých (Lepidoptera) na Valašsku (okres Vsetín, Česká republika). Acta Carpathica Occidentalis 1: 19–39.
- Spitzer L, Beneš J, Konvička M (2018) Extinction of *Lasiommata petropolitana* (Fabricius, 1787) (Lepidoptera: Nymphalidae) in the Czech Republic: a case of habitat loss at a range margin. Acta Musei Silesiae, Scientiae Naturales 66(3): 271–279. <https://doi.org/10.1515/cszma-2017-0030>
- Thomas CD, Bodsworth EJ, Wilson RJ, Simmons AD, Davies ZG, Musche M, Conradt L (2001) Ecological and evolutionary processes at expanding range margins. Nature 411(6837): 577–581. <https://doi.org/10.1038/35079066>
- Todisco V, Gratton P, Cesaroni D, Sbordoni V (2010) Phylogeography of *Parnassius apollo*: hints on taxonomy and conservation of a vulnerable glacial butterfly invader. Biological Journal of the Linnean Society 101(1): 169–183. <https://doi.org/10.1111/j.1095-8312.2010.01476.x>
- Tolman T, Lewington R (2008) Collins butterfly guide: the most complete field guide to the butterflies of Britain and Europe. Collins, London, 384 pp.
- Trifinopoulos J, Nguyen L-T, Haeseler von A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Research 44(W1): W232–5. <https://doi.org/10.1093/nar/gkw256>
- Tuzov VK (2000) Guide to the butterflies of Russia and adjacent territories (Lepidoptera, Rhopalocera), Volume 2: Libytheidae, Danaidae, Nymphalidae, Riodinidae, Lycaenidae. Pensoft, Sofia-Moscow, 581 pp.
- Uříčář J, Jongepierová I, Vondřejc TE (2016) Zásady péče o významné druhy motýlů Bílých Karpat. ZO ČSOP Bílé Karpaty, Veselí nad Moravou, 56 pp.
- Vodá R, Dapporto L, Dincă V, Vila R (2015) Why do cryptic species tend not to co-occur? A case study on two cryptic pairs of butterflies. PLOS ONE 10(2): e0117802. <https://doi.org/10.1371/journal.pone.0117802>
- Wahlberg N, Wheat CW (2008) Genomic outposts serve the phylogenomic pioneers: designing novel nuclear markers for genomic DNA extractions of lepidoptera. Systematic Biology 57(2): 231–242. <https://doi.org/10.1080/10635150802033006>
- Warecki A (2010) Motyle dzienne Polski. Atlas bionomii. Koliber, Nowy Sacz, 319 pp.

Supplementary material 1

The samples of *Aricia* spp. used for species identification in the Czech Republic and TCS haplotype network through its distributional range, including GenBank samples

Authors: Alena Sucháčková Bartoňová, Jiří Beneš, Zdeněk Faltýnek Fric, Martin Konvička

Data type: species data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/nl.42.38853.suppl1>

Supplementary material 2

NEXUS alignment of *Aricia* spp. cytochrome c oxidase subunit I sequences used for Maximum Likelihood analysis, covering samples from the Czech Republic and samples from GenBank

Authors: Alena Sucháčková Bartoňová, Jiří Beneš, Zdeněk Faltýnek Fric, Martin Konvička

Data type: molecular data

Copyright notice: This dataset is made available under the Open Database License (<http://open-datacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/nl.42.38853.suppl2>

Supplementary material 3

NEXUS alignment of *Aricia artaxerxes* cytochrome c oxidase subunit I sequences used for haplotype network analysis, covering samples from the Czech Republic and samples from GenBank

Authors: Alena Sucháčková Bartoňová, Jiří Beneš, Zdeněk Faltýnek Fric, Martin Konvička

Data type: molecular data

Copyright notice: This dataset is made available under the Open Database License (<http://open-datacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/nl.42.38853.suppl3>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nota lepidopterologica](#)

Jahr/Year: 2019

Band/Volume: [42](#)

Autor(en)/Author(s): Suchackova Bartonova Alena, Benes Jiri, Fric Zdenek Flatynek, Konvicka Martin

Artikel/Article: [Genetic confirmation of *Aricia artaxerxes* \(Fabricius, 1793\) \(Lepidoptera, Lycaenidae\) in the Czech Republic, its conservation significance and biogeographic context 163-176](#)