Nota Lepi. 43 2020: 1-14 | DOI 10.3897/nl.43.46354

# Systematics and evolution of the African butterfly genus *Mylothris* (Lepidoptera, Pieridae)

Haydon Warren-Gash<sup>1</sup>, Kwaku Aduse-Poku<sup>2</sup>, Leidys Murillo-Ramos<sup>3,4</sup>, Niklas Wahlberg<sup>3</sup>

1 98 Overstrand Mansions, Prince of Wales Drive, London SW11 4EU, UK

2 Biology Department, University of Richmond, 138 UR Drive, Richmond, VA 23173, USA

3 Department of Biology, Sölvegatan 37, Lund University, 223 62 Lund, Sweden

4 Grupo Biología Evolutiva, Department of Biology, Universidad de Sucre, Sincelejo, Sucre, Colombia

http://zoobank.org/CB00DD10-8657-4283-93BA-0FEFDEB922A7

Received 6 September 2019; accepted 6 January 2020; published: 11 February 2020 Subject Editor: Roger Vila.

Abstract. We study the systematics and evolutionary history of the Afrotropical butterfly genus *Mylothris* (Lepidoptera: Pieridae) based on six gene regions (COI, EF1a, GAPDH, MDH, RpS5 and wingless). We find that the genus can be placed into five species groups, termed the *jacksoni*, *elodina*, *rhodope*, *agathina* and *hila-ra* groups. Within these species groups, we find that many species show very little genetic differentiation based on the markers we sequenced, suggesting they have undergone rapid and recent speciation. Based on second-ary calibrations, we estimate the age of the crown group of *Mylothris* to be about 16 million years old, but that many of the species level divergences have happened in the Pleistocene. We infer that the clade has its origin in the forests of the Eastern part of Central Africa, and has spread out from there to other regions of Africa.

## Introduction

Studies on the systematics of Afrotropical butterflies using molecular data have been increasing in number in the past decade (Kodandaramaiah and Wahlberg 2007; Aduse-Poku et al. 2009; Nazari et al. 2011; Price et al. 2011; van Velzen et al. 2013; Aduse-Poku et al. 2017), giving us a better understanding of their evolutionary histories in one of the diversity hotspots of the planet. Many genera are endemic to the continent, although there are clear connections to the Oriental region (Kodandaramaiah and Wahlberg 2007; Aduse-Poku et al. 2009; Aduse-Poku et al. 2015; Sahoo et al. 2018; Toussaint et al. 2019) and even to the Neotropical region (Silva Brandão et al. 2008). Genera in the family Pieridae have not been the focus of many studies to date. So far only two pierid genera have been studied using molecular approaches in any detail, *Colotis* Hübner, 1819 (Nazari et al. 2011) and *Pseudopontia* Plötz, 1870 (Mitter et al. 2011). Here we investigate the systematics and evolutionary history of the Afrotropical pierid genus *Mylothris* Hübner, 1819.

The range of *Mylothris* includes most of the African continent south of the Sahara, extending to nearby islands such as Madagascar and the Comoros in the east, and to Bioko and São Tomé e Príncipe in the west. A single species flies alongside other pierid species belonging to the Afrotropical fauna in south west Arabia. It is speciose: a revision of the genus by one of the authors (HWG) is in progress and will list 105 species, a number of them new, many of them in turn divided into several subspecies (Warren-Gash in press).

A distinguishing feature of the genus, as with the related *Catasticta* Butler, 1870 in the Neotropics (Braby and Nishida 2010) and *Delias* Hübner, 1819 in the Australasian region (Braby 2006), is that the hostplants, with one known exception, are members of parasitic mistletoes belonging to the family Loranthaceae. This in turn determines where and how plentifully *Mylothris* species are found. Their foodplants are arboreal epiphytes and so they are insects of the canopies – unlike most Afrotropical pierids, which breed on low-growing shrubs such as *Capparis* Linnaeus, 1758, and many of which occur in more open, drier biotopes. Thus *Mylothris* species diversity is richest in the main African forest belt, with some species spreading into the forest-savannah mosaic and anthropogenic biomes including suburban gardens.

Two other consequences follow. The first is that, since Loranthaceae occur most frequently near the crown of the tree, many of the tropical forest species fly high, and some of them are seldom seen, accentuating their rarity in institutional collections. Secondly, they are particularly vulnerable to primary forest degradation.

Typically, *Mylothris* eggs are laid in a group on the underside of a leaf of the hostplant, and the larvae when they hatch are gregarious. The length of the lifecycle depends on the species and the biome or region. In southern Africa they follow the seasons in line with their foodplant, as they do in colder (usually montane) climates further north. In less seasonal biotopes, adults can be on the wing at any time of year (Warren-Gash in press).

In appearance *Mylothris* are a remarkably homogeneous group. In adults of the *M. jacksoni* species group, the hindwing is yellow and the forewing usually mostly white. The other clades typically show some orange (sometimes yellow) at the base of the forewing on one or both surfaces becoming white distally, with a greater or lesser black distal margin. There are exceptions, but they are similar enough to have baffled taxonomists over the years, leading to groups of species being lumped together under a single name in some cases and, no less frequently, names erected to describe what are no more than individual variations. If that were not enough, cryptic rings of different species that in adult morphology are externally almost indistinguishable, can be found flying together, most frequently in the two main areas of diversity: the Albertine Rift and central Cameroon (Warren-Gash in press).

The forthcoming revision (Warren-Gash in press) will look at all these issues in more detail based on morphological characters. Here we aim to investigate whether molecular data can shed light on species delimitations and their relationships. As yet undescribed species are here named nsp1 to nsp4. They will be formally described in Warren-Gash (in press).

#### Material and methods

We attempted to sample as many species as possible, and for selected species, as many populations as possible. We included 52 out of 105 species and a total of 235 individuals in our analyses (see Suppl. material 1: Table S1). Most specimens come from the private collection of HWG, with a few specimens taken from the collections of the African Butterfly Research Institute in Nairobi, Kenya. The specimens are part of a large taxonomic revision of *Mylothris* (Warren-Gash in press). In addition we included 23 outgroups from the subtribe Aporiina, taken from a previous publication (Wahlberg et al. 2014).

DNA was extracted from 2–3 legs of dried museum specimens using the Nucleospin Tissue Kit (Macherey-Nagel, Düren, Germany). From these specimens, we sequenced the mitochondrial COI gene and up to 5 nuclear genes (EF1a, GAPDH, MDH, RpS5 and wingless). Primers and laboratory protocols followed Wahlberg and Wheat (2008) in all cases. Successful PCR products were cleaned

Nota Lepi. 43: 1-14

enzymatically with exonuclease and FastAP (Thermo Fisher Scientific, Hampton, NH, USA) and sent to Macrogen Europe (Amsterdam, the Netherlands) for Sanger sequencing. The resulting chromatograms were examined by eye using BioEdit (Hall 1999). All six genes are protein-coding, thus alignments were trivial. The aligned sequences were curated and managed in VoSeq (Peña and Malm 2012).

Phylogenetic analyses were carried out using IQ-TREE 1.6.10 (Nguyen et al. 2015) in a maximum likelihood framework. The full dataset was analysed. The data were partitioned by gene and analysed with the partition finding (Chernomor et al. 2016) and model finding (Kalyaanamoorthy et al. 2017) algorithms of IQ-TREE (using the command MFP+MERGE). Robustness of the results were assessed using UFBoot2 (Hoang et al. 2018) and a SH-like approximate likelihood ratio test (Guindon et al. 2010). Analyses were run on the CIPRES server (Miller et al. 2010).

Timing of divergence was estimated using a reduced dataset. One specimen per species with the most DNA sequence data available was chosen for this analysis. The dataset was analysed using BEAST v1.8.3 (Drummond et al. 2012), partitioned according to the results of the IQ-TREE analysis. Since BEAST does not offer the possibility to use the TIM2 model, the model for COI was set to the GTR+G model, the other two partitions were assigned models according to the IQ-TREE results (see below). The nucleotide models and clock models were unlinked between partitions. The tree prior was set to the birth-death model and a lognormal relaxed clock was implemented. Monophyly of Aporiina was enforced. Four secondary calibration points were taken from Chazot et al. (2019) and were modeled with a normal prior. The root was set to 42 million years ago (Mya)  $\pm$  5 million years (My), the crown age of Aporiina to 36 Mya  $\pm$  4 My, the most common recent ancestor of *Mylothris* and *Aporia* Hübner, 1819 to 28 Mya  $\pm$  3.5 My, and the most recent common ancestor of *Catasticta* and *Melete* Swainson, 1831 to 20 Mya  $\pm$  3 My. The analyses were run four times independently at 10 million generations per run sampling at 10,000 generation intervals. Convergence was examined using Tracer v1.7 (Rambaut et al. 2014). The four runs were combined after a burnin of 1 million generations and a summary tree was generated using the maximum a posteriori tree and mean estimated times of divergence.

The historical biogeography of the group was inferred using the R package BioGeoBEARS v1.1.1 (Matzke 2014). The chronogram resulting from the BEAST analysis (see results section) with the outgroups pruned, was used for this analysis. Owing to the open conceptual debate on model comparisons in BioGeoBEARS (Ree and Sanmartín 2018), our ancestral range analysis were performed under the dispersal - extinction - cladogenesis (DEC) model (Ree et al. 2005; Ree and Smith 2008). DEC is a likelihood-based procedure that models range evolution as a discrete-valued process, specifying instantaneous transition rates between geographical ranges along the branches of a phylogenetic tree. Consequently, DEC requires each sampled taxon be assigned with one or more geographic range(s), reflecting its present distributional range. The geographic distributions of *Mylothris* were extrapolated from the literature and the degree of endemism of local species, which is highest in West Africa and Madagascar. The following regions were used in the analysis: A, West Africa; B, West Central Africa; C, East Central Africa; D, East Africa; E, South Africa; F, Malagasy Region (see inserted map in Figure 6 of Results section). B and C are closer than the others, corresponding roughly to the central forest belt, but with two main centres of endemism: the Cameroon highlands and the montane zone just west of the Albertine Rift. In all dispersal rate scaler matrices, dispersal events between adjacent areas were not scaled (dispersal d = 1.0). A scaler of 0.5 was applied for dispersal events between areas separated by an ocean barrier. A scaler of 0.25was applied for dispersal events between areas separated by a third region and a water barrier (see Suppl. material 2: Table S2). The maximum number of areas per ancestral state was fixed to four.

# Results

The dataset consisted of 5168 aligned base pairs. The IQ-TREE partition finding analysis combined EF1a, GAPDH, MDH, and RpS5 into one partition and assigned the GTR+I+G model to it. COI was kept as its own partition with TIM2+I+G model assigned to it, as was wingless with the K2P+G model.

We find *Mylothris* to be a well supported monophyletic group (UFB = 100, SH-like = 100), that is sister to a clade of Neotropical and Palaearctic/Oriental taxa (Figure 1). We identify five major clades that are strongly supported (the exception being the *hilara* clade with UFB = 87, SH-like = 90): the *jacksoni*, *elodina*, *rhodope*, *agathina* and *hilara* clades. We find the *jacksoni* clade to be sister to the rest of *Mylothris*, with *elodina* being the next lineage to diverge, followed by the *rhodope* clade, and finally the *agathina* and *hilara* clades being sister to each other.

Relationships within the *jacksoni* clade show four lineages (Figure 2), with *M. ducarmei* Hecq, 2001 being sister to the rest, followed by *M. crawshayi* Butler, 1896 and finally *M. trimenia* Butler,



**Figure 1.** Relationships of the five species groups in *Mylothris* based on a maximum likelihood analysis of 5 gene regions. Numbers below branches are branch support values (SH-like / Ultrafast Bootstrap).

1869 sister to a clade comprising five species with unresolved relationships. These five species, *M. jacksoni* Sharpe, 1891 *M. sagala*, Grose-Smith, 1886 *M. knutssoni* Aurivillius, 1891 *M. nagichota* Talbot, 1944 and *M. seminigra* d'Abrera, 1980 show very little genetic differentiation.

Only *M. elodina* Talbot, 1944 is found in the *elodina* clade (Figure 2). The eight individuals we sequenced show very little genetic differentiation, but a relatively long branch leads to them. The position of the lineage is not highly supported (UFB = 79, SH-like = 85.3), and there is a possibility that further data might place it as the sister to the rest of *Mylothris*.

The *rhodope* clade comprises 14 sampled species (Figure 3), which are genetically very close to each other. Indeed, *M. kiwuensis* Grunberg, 1910, *M. knoopi* Hecq, 2005, *M.* nsp2 and *M. jaopura* Karsch, 1893 are genetically identical at all the sequenced loci. *Mylothris uniformis* Talbot, 1944 and *M.* nsp3 are also intermixed and genetically extremely close to each other. Other species are reciprocally monophyletic, but genetically very close to their sister species, such as *M.* nsp4 + *M. rhodope* Fabricius, 1775 and *M. latimargo* Joicey & Talbot, 1921+ *M. zairiensis* Berger, 1981.



Figure 2. Relationships of sampled specimens within the *jacksoni* and *elodina* groups. Numbers below branches are branch support values (SH-like / Ultrafast Bootstrap).

#### HAYDON WARREN-GASH et al.: Mylothris phylogeny



Figure 3. Relationships of sampled specimens within the *rhodope* group. Numbers below branches are branch support values (SH-like / Ultrafast Bootstrap).

etas Europaea Lepidopterologica; download unter http://www.biodiversitylibrary.org/ und www.zobc



Figure 4. Relationships of sampled specimens within the *agathina* group. Numbers below branches are branch support values (SH-like / Ultrafast Bootstrap).

HAYDON WARREN-GASH et al.: Mylothris phylogeny



Figure 5. Relationships of sampled specimens within the *hilara* group. Numbers below branches are branch support values (SH-like / Ultrafast Bootstrap).



Figure 6. Timing of divergence and historical biogeographic analyses of *Mylothris*. Inset shows the biogeographical regions used for the latter.

The *agathina* clade has 16 sampled species in it, showing similar patterns of genetic differentiation as the previous clades (Figure 4). *Mylothris agathina* Cramer, 1779 and *M. chloris* Fabricius, 1775 are genetically indistinguishable (based on the markers used here), as are *M. sulphurea* Aurivillius, 1895, *M. holochroma* Talbot, 1944, *M. interposita* Joicey & Talbot, 1921 and *M. aurantiaca* Rebel, 1914. The Madagascan *M. phileris* Boisduval, 1833 and Comoro Islands *M. ngaziya* Oberthur, 1888 are found to be closely related to the *M. agathina/chloris* clade.

The *hilara* clade comprises 15 sampled species, and again similar patterns of little genetic differentiation are found among the species as in the other clades (Figure 5). The species *M. sjostedti* Aurivillius, 1895, *M. similis* Lathy, 1906, *M. poppea* Cramer, 1777 and *M. erlangeri* Pagenstecher, 1902 appear to be distinct from each other and the rest, while the rest of the species form a complex set of relationships with shared haplotypes and minor monophyletic groups (e.g. *M.* nsp1), which are very closely related genetically, even if morphological characters are clearly distinct.

Our timing of divergence analysis suggests that the crown clade of *Mylothris* began diversifying around 16 Mya (Figure 6) after diverging from their sister group some 10 My earlier. The five major clades diverged from each other between 16 and 8 Mya, during the mid to late Miocene, but almost all species level divergences have happened in the Pleistocene. Our ancestral range analysis recovers an origin of *Mylothris* in the East Central Africa. All the five major clades of the group are also inferred to have been derived from ancestors distributed in the East Central Africa. The ancestors of the extant species in Madagascar and Comoros islands are estimated to have range expanded from mainland Africa only recently (~2 Mya).

#### Discussion

*Mylothris* are a distinct and well-supported lineage of African pierids, not closely related to any other pierids on that continent. There are five clear species groups within the genus. Of the five groups, the *rhodope, agathina* and *hilara* species groups are more closely related to each other than to the other two groups. Based on our sampling of genes, it appears that the *jacksoni* group is sister to the rest, but with poor support. Indeed, in some preliminary analyses of our data, the *elodina* group came out as sister to the rest, and this may be the case with further gene and taxon sampling. The very different morphology of *elodina*, reflected also in the male genitalia, is a possible indicator that it is sister to the rest of *Mylothris*.

It is highly probable that we have uncovered all major species groups. The unsampled species are all, as far as we can tell, clearly morphologically related to already sampled species. The sampling process was conducted in the context of a revision of the genus (Warren-Gash in press), and large quantities of material were examined and dissected, including many specimens too old to be readily amenable to molecular sequencing. The unsampled species showed no differences which were important enough in terms of morphology or genitalia to suggest the likely existence of other species groups.

Regardless of the relationships of the major species groups, the biogeographic history of the deeper clades appears to be clear. We infer from our analyses that the group originated in East Central Africa, and from there spread to other parts of Africa. The fragmentation of the forests in the Miocene and Pleistocene may have contributed to the diversification of the group, perhaps by causing populations to be fragmented and isolated from each other. This appears to be a common pattern in forest dwelling butterflies of Africa (Aduse-Poku et al. 2009; van Velzen et al. 2013; Dhungel and Wahlberg 2018), but the mechanisms leading to their diversification are still poorly understood.

Within the species groups of *Mylothris* we find genetic variation between some of the species to be minimal, and indeed some species share the same haplotypes for the markers sequenced in this study. The infra-specific variation within *Mylothris* species is, however, considerable, and that has been given full weight both in the preparation of this paper and the forthcoming revision. A number of morphs previously considered as species have been synonymised in consequence (Warren-Gash in press). Nonetheless, such close genetic similarity needs explanation if the specific separation of these groups is to be justified.

There is no single answer to why some species are genetically identical for the markers we have sequenced. In one case – *Mylothris agathina* and *M. chloris* – they are sympatric with no evidence of interbreeding, and in addition the female genitalia of the two are clearly and consistently different. In the *M. jacksoni* group, the wing patterns vary, as does the biome in which they fly. Many of the scarcer species, both sampled and unsampled, are montane vicariants; others fly sympatrically without interbreeding and in several cases where the early stages have been observed, the larvae are more distinct than the adults and use different larval foodplants (e.g. *M. jacksoni* and *M. saga-la*). They not only look different but in their foodplant preferences behave differently.

In other groups, there are examples of species which fly together in at least a part of their range, which are morphologically similar in one sex (usually the male) but distinctively different in the other (e.g. *M. sulphurea basalis* Aurivillius, 1906 and *M. aurantiaca* Rebel, 1914, with overlapping ranges in the north-east of the DRC).

A group of more than a dozen species in the *hilara* group, which have in common a red basal patch at the base of the forewing and broadly similar (but not identical) male genitalia, is found across the width of the main forest belt. A few are very local. In the rugged terrain of the mountains west of the Albertine

#### Nota Lepi. 43: 1-14

Rift, they are separated more by terrain than geographical distance, but in those few instances where they overlap, they do not interbreed. At lower altitudes it is quite usual for several species in the group to fly together in varying combinations, without any evidence of cross-fertilisation. In the Albertine Rift widely varying female forms are found which occur nowhere else. Morphology shows up what appear to be widely separated vicariants of similar species in east and west Africa, more akin to each other than those flying in between. They almost certainly use different larval foodplants. Some are tied to strictly forest environments (e.g. *M. continua*); others prefer more open country (e.g. *M. rueppellii* Koch, 1865)

The case of the *jaopura* complex, with nine named and mostly allopatric species, offers a different challenge. Those sequenced are indistinguishable on that basis, and the genitalia, while distinctive within the complex, are very similar. However, morphology in terms of size, wingshape and markings is distinctive in each case, as very often is the biome in which they occur. Where they overlap (e.g. *M. kiwuensis* and *M.* nsp2), they do not interbreed. Again, and given the very different environments in which they occur, it is highly probable that their larval foodplants also differ. What they have in common is that in practically every case they mimic a *Mylothris* species in another group, sometimes so well that they can only be separated with certainty by dissection or sequencing.

Given the factors discussed above, the markers we have used are probably not sufficiently variable to differentiate the closely related species. It is likely that a genomic approach would shed further light on the species boundaries, as has been found in recent studies on nymphalid butterflies (Dinca et al. 2019; Campbell et al. 2020). Our results do however suggest that radiation within these groups is likely to be recent. Similar minimal genetic differentiation is found in the pierid genus *Colias* Fabricius, 1807 (Wheat and Watt 2008; Laiho and Ståhls 2013), but interestingly not in the closely related genus *Delias* (Braby et al. 2007; Müller et al. 2013). The reasons behind the close genetic relationships of several groups of species is not entirely clear, but may have to do with the repeated contraction and expansion of the forests in Africa, especially during the Miocene and Pleistocene. Similar patterns appear to be the case in other forest specialist genera of the family Nymphalidae: *Charaxes* Ochsenheimer, 1816 (Aduse-Poku et al. 2009), *Cymothoe* Hübner, 1819 (van Velzen et al. 2013) and *Euphaedra* Hübner, 1819 (Dhungel and Wahlberg 2018). *Mylothris* and the nymphalid genera cited appear to be going through rapid speciation in African forests and they require further study to understand the driving mechanisms

#### Acknowledgements

We thank Robin Pranter for help in the laboratory. We thank Steve Collins and ABRI, Kenya for providing us with much useful material and helpful insights. We also thank David Lees for extensive comments on the manuscript. NW acknowledges funding from the Swedish Research Council (grant number 2015-04441). We thank the Societas Europea Lepidopterologica (SEL) for support in publishing this article.

#### References

Aduse-Poku K, Brakefield PM, Wahlberg N, Brattström O (2017) Expanded molecular phylogeny of the genus *Bicyclus* shows the importance of increased sampling for detecting semi-cryptic species and highlights potentials for future studies. Systematics and Biodiversity 15: 115–130. https://doi.org/10.1080/1477200 0.2016.1226979

1	2
I	2

- Aduse-Poku K, Brattström O, Kodandaramaiah U, Lees DC, Brakefield P, Wahlberg N (2015) Systematics and historical biogeography of the Old World butterfly subtribe Mycalesina (Lepidoptera: Nymphalidae: Satyrinae). BMC Evolutionary Biology 15: 167. https://doi.org/10.1186/s12862-015-0449-3
- Aduse-Poku K, Vingerhoedt E, Wahlberg N (2009) Out-of-Africa again: a phylogenetic hypothesis of the genus *Charaxes* (Lepidoptera: Nymphalidae) based on 5 gene regions. Molecular Phylogenetics and Evolution 53: 463–478. https://doi.org/10.1016/j.ympev.2009.06.021
- Braby MF (2006) Evolution of larval food plant associations in *Delias* Hübner butterflies (Lepidoptera: Pieridae). Entomological Science 9: 383–398. https://doi.org/10.1111/j.1479-8298.2006.00185.x
- Braby MF, Nishida K (2010) The immature stages, larval food plants and biology of Neotropical mistletoe butterflies (Lepidoptera: Pieridae). II. The *Catasticta* group (Pierini: Aporiina). Journal of Natural History 44: 1831–1928. https://doi.org/10.1080/00222931003633227
- Braby MF, Pierce NE, Vila R (2007) Phylogeny and historical biogeography of the subtribe Aporiina (Lepidoptera : Pieridae): implications for the origin of Australian butterflies. Biological Journal of the Linnean Society 90: 413–440. https://doi.org/10.1111/j.1095-8312.2007.00732.x
- Campbell EO, Gage EV, Gage RV, Sperling FAH (2020) Single nucleotide polymorphism based species phylogeny of greater fritillary butterflies (Lepidoptera: Nymphalidae: Speyeria) demonstrates widespread mitonuclear discordance. Systematic Entomology 45: in press. https://doi.org/10.1111/syen.12393
- Chazot N, Wahlberg N, Freitas AVL, Mitter C, Labandeira C, Sohn J-C, Sahoo RK, Seraphim N, de Jong R, Heikkilä M (2019) Priors and posteriors in Bayesian timing of divergence analyses: the age of butterflies revisited. Systematic Biology 68: 797–813. https://doi.org/10.1093/sysbio/syz002
- Chernomor O, von Haeseler A, Minh BQ (2016) Terrace aware data structure for phylogenomic inference from supermatrices. Systematic Biology 65: 997–1008. https://doi.org/10.1093/sysbio/syw037
- Dhungel B, Wahlberg N (2018) Molecular systematics of the subfamily Limenitidinae (Lepidoptera: Nymphalidae). PeerJ 6: e4311. https://doi.org/10.7717/peerj.4311
- Dinca V, Lee KM, Vila R, Mutanen M (2019) The conundrum of species delimitation: a genomic perspective on a mitogenetically super-variable butterfly. Proceedings of the Royal Society of London B Biological Sciences 286: 20191311. https://doi.org/10.1098/rspb.2019.1311
- Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with BEAUTi and BEAST 1.7. Molecular Biology and Evolution 29: 1969–1973. https://doi.org/10.1093/molbev/mss075
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate Maximum-Likelihood phylogenies: Assessing the performance of PhyML 3.0. Systematic Biology 59: 307–321. https://doi.org/10.1093/sysbio/syq010
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucl. Acids. Symp. Ser. 41: 95–98.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the ultrafast bootstrap approximation. Molecular Biology and Evolution 35: 518–522. https://doi.org/10.1093/molbev/ msx281
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14: 587–589. https://doi.org/10.1038/ nmeth.4285
- Kodandaramaiah U, Wahlberg N (2007) Out-of-Africa origin and dispersal mediated diversification of the butterfly genus *Junonia* (Nymphalidae: Nymphalinae). Journal of Evolutionary Biology 20: 2181–2191. https://doi.org/10.1111/j.1420-9101.2007.01425.x
- Laiho J, Ståhls G (2013) DNA barcodes identify Central Asian *Colias* butterflies (Lepidoptera, Pieridae). Zookeys 365: 175–196. https://doi.org/10.3897/zookeys.365.5879
- Matzke NJ (2014) Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. Systematic Biology 63: 951–970. https://doi.org/10.1093/sysbio/syu056

Nota Lepi. 43: 1–14

- 13
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Proceedings of the Gateway Computing Environments Workshop (GCE), New Orleans, Louisiana. https://doi.org/10.1109/GCE.2010.5676129
- Mitter KT, Larsen TB, de Prins W, De Prins J, Collins S, Weghe GV, Safian S, Zakharov EV, Hawthorne DJ, Kawahara AY, Regier JC (2011) The butterfly subfamily Pseudopontiinae is not monobasic: marked genetic diversity and morphology reveal three new species of *Pseudopontia* (Lepidoptera: Pieridae). Systematic Entomology 36: 139–163. https://doi.org/10.1111/j.1365-3113.2010.00549.x
- Müller CJ, Matos-Maraví PF, Beheregaray LB (2013) Delving into *Delias* Hübner (Lepidoptera: Pieridae): fine-scale biogeography, phylogenetics and systematics of the world's largest butterfly genus. Journal of Biogeography 40: 881–893. https://doi.org/10.1111/jbi.12040
- Nazari V, Larsen TB, Lees DC, Brattström O, Bouyer T, Van de Poel G, Hebert PDN (2011) Phylogenetic systematics of *Colotis* and associated genera (Lepidoptera: Pieridae): evolutionary and taxonomic implications. Journal of Zoological Systematics and Evolutionary Research 49: 204–215. https://doi.org/10.1111/ j.1439-0469.2011.00620.x
- Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. Molecular Biology and Evolution 32: 268–274. https://doi.org/10.1093/molbev/msu300
- Peña C, Malm T (2012) VoSeq: a Voucher and DNA Sequence Web Application. PlosOne 7: e39071. https:// doi.org/10.1371/journal.pone.0039071
- Price BW, Villet MH, Walton SM, Barker NP (2011) Using molecules and morphology to infer the phylogenetic relationships and evolutionary history of the Dirini (Nymphalidae: Satyrinae), a tribe of butterflies endemic to Southern Africa. Systematic Entomology 36: 300–316. https://doi.org/10.1111/j.1365-3113.2010.00560.x
- Rambaut A, Suchard MA, Xie D, Drummond AJ (2014) Tracer v1.6. https://github.com/beast-dev/tracer/releases
- Ree RH, Moore BR, Webb CO, Donoghue MJ (2005) A likelihood framework for inferring the evolution of geographic range on phylogenetic trees. Evolution 59: 229–2311. https://doi.org/10.1111/j.0014-3820.2005. tb00940.x
- Ree RH, Sanmartín I (2018) Conceptual and statistical problems with the DEC+J model of founder-event speciation and its comparison with DEC via model selection. Journal of Biogeography in press. https://doi.org/10.1111/jbi.13173
- Ree RH, Smith SA (2008) Maximum Likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. Systematic Biology 57: 4–14. https://doi.org/10.1080/10635150701883881
- Sahoo RK, Lohman DJ, Wahlberg N, Müller CJ, Brattström O, Collins SC, Peggie D, Aduse-Poku K, Kodandaramaiah U (2018) Evolution of *Hypolimnas* butterflies (Nymphalidae): Out-of-Africa origin and *Wolbachia*-mediated introgression. Molecular Phylogenetics and Evolution 123: 50–58. https://doi. org/10.1016/j.ympev.2018.02.001
- Silva Brandão KL, Wahlberg N, Francini RB, Azeredo-Espin AML, Brown Jr. KS, Paluch M, Lees DC, Freitas AVL (2008) Phylogenetic relationships of butterflies of the tribe Acraeini (Lepidoptera, Nymphalidae, Heliconiinae) and the evolution of host plant use. Molecular Phylogenetics and Evolution 46: 515–531. https://doi.org/10.1016/j.ympev.2007.11.024
- Toussaint EFA, Vila R, Yago M, Chiba H, Warren AD, Aduse-Poku K, Storer C, Dexter KM, Maruyama K, Lohman DJ, Kawahara AY (2019) Out of the Orient: Post-Tethyan transoceanic and trans-Arabian routes fostered the spread of Baorini skippers in the Afrotropics. Systematic Entomology 44: 926–938. https:// doi.org/10.1111/syen.12365
- van Velzen R, Wahlberg N, Sosef MSM, Bakker FT (2013) Effects of changing climate and host plant association on species diversification rates in *Cymothoe* (Lepidoptera, Nymphalidae) tropical forest butterflies. Biological Journal of the Linnean Society 108: 546–564. https://doi.org/10.1111/bij.12012

- Wahlberg N, Rota J, Braby MF, Pierce NE, Wheat CW (2014) Revised systematics and higher classification of pierid butterflies (Lepidoptera: Pieridae) based on molecular data. Zoologica Scripta 43: 641–650. https:// doi.org/10.1111/zsc.12075
- Wahlberg N, Wheat CW (2008) Genomic outposts serve the phylogenomic pioneers: designing novel nuclear markers for genomic DNA extractions of Lepidoptera. Systematic Biology 57: 231–242. https://doi. org/10.1080/10635150802033006

Warren-Gash H (in press) Revision of the genus Mylothris. African Natural History Research Trust, London, UK.

Wheat CW, Watt WB (2008) A mitochondrial-DNA-based phylogeny for some evolutionary-genetic model species of *Colias* butterflies (Lepidoptera, Pieridae). Molecular Phylogenetics and Evolution 47: 893–902. https://doi.org/10.1016/j.ympev.2008.03.013

#### **Supplementary material 1**

## Table S1

Authors: Haydon Warren-Gash, Kwaku Aduse-Poku, Leidys Murillo-Ramos, Niklas Wahlberg Data type: List of specimens used in the study along with NCBI accession numbers.

- Explanation note: List of specimens used in this study, with species designations, NCBI accession numbers and collection localities. An "X" instead of an accession number indicates a sequence less than 200 bp (and thus unacceptable by NCBI).
- Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/nl.43.46354.suppl1

#### **Supplementary material 2**

#### Table S2

Authors: Haydon Warren-Gash, Kwaku Aduse-Poku, Leidys Murillo-Ramos, Niklas Wahlberg Data type: Scaler matrices used in BioGeoBEARS analyses.

Explanation note: Scaler matrices used in BioGeoBEARS analyses.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/nl.43.46354.suppl2

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Nota lepidopterologica

Jahr/Year: 2020

Band/Volume: 43

Autor(en)/Author(s): Warren-Gash Haydon, Aduse-Poku Kwaku, Murillo-Ramos Leidys, Wahlberg Niklas

Artikel/Article: <u>Systematics and evolution of the African butterfly genus Mylothris</u> (Lepidoptera, Pieridae) 1-14