

Proc.V.Congr.Eur.Lepid., Budapest 7-10.IV.1986

Nota lepid. Supplement No. 3 : 97-100 ; 30.IV.1992

ISSN 0342-7536

## Preliminary survey of the butterflies of Sierra de Gredos (Central Spain) (Lep. : Hesperioidea et Papilioidea)

J.L. VIEJO & J. MARTÍN, Universidad Autónoma de Madrid, Facultad de Ciencias, Departamento de Biología C-XV, Cantoblanco, E-28049 Madrid, Spain.

### Introduction and area description

The Sierra de Gredos is located in the Sistema Central, a mountain range which extends in a southwest-northeast direction, through the centre of the Iberian Peninsula. The Central portion of the Sierra de Gredos is about 60 km long, from Puerto del Pico (eastern limit) to Puerto de Tornavacas (western limit), being 20-30 km wide. This mountainous area rises from a flat region which is 800 m in altitude northwards and 500 m southwards ; the peaks themselves reach more than 2,500 m in their uppermost zones (Pico Almanzor, 2,592 m). Lithologically, Sierra de Gredos is composed of granitic rocks, mainly in the upper zones, resulting in acid soils. Both slopes are geomorphologically different ; the southern slope descends abruptly from the ridge to the southern plateau in some hundreds of metres, while the northern slope decreases smoothly and is several kilometres long.

Both slopes have therefore different climatic features. The southern slope has milder weather conditions, while the northern slope is colder, with longer, more severe winters and short cool summers. The southern slope has short winters with virtually no frosts and longer, hotter summers.

Vegetation ranges from an evergreen forest of *Quercus ilex* at low altitudes, to a grassland of *Festuca indigesta*, at the highest altitudes ; oak forests of *Quercus pyrenaica* and shrubs of broom (*Cytisus purgans*) occur in between (RIVAS MARTINEZ, 1975).

The lower zones of the southern slope are suitable for agriculture and grazing, and there are orchards of cherry trees, chestnuts, walnuts, etc. ; there are also herds, mainly of goats and sheep. On the northern slope there are low pastures and a few orchards.

Most of this area is given over to forestry, with pinewoods of *Pinus pinaster* on the southern slope, and *Pinus sylvestris* on the northern one. Some of these forests are 500 years old.

In spite of its proximity to Madrid (150 km from the nearest point), Sierra de Gredos has not been thoroughly surveyed from the Lepidoptera point of view, this being mainly due to the very poor network of roads.

Table I

Species abundance on both Sierra de Gredos slopes (sites above, 1,300 m have been considered in northern slope)

	N	S		N	S
<b>PAPILIONIDAE</b>			<b>Pandoriana pandora</b>	++	+++
<i>Papilio machaon</i>	++		<i>Argynnis paphia</i>	++	++
<i>Iphiclus podalirius</i>	+		<i>Fabriciana adippe</i>	++	++
<i>Zerynthia rumina</i>	+	+++	<i>Fabriciana niobe</i>	++	++
<b>PIERIDAE</b>			<i>Issoria lathonia</i>	+++	+++
<i>Aporia crataegi</i>	+		<i>Brenthis daphne</i>	+	+
<i>Pieris brassicae</i>		+	<i>Clossiana selene</i>	+++	
<i>Pieris rapae</i>	++	+++	<i>Mesocacidalia aglaja</i>	++	+
<i>Pieris napi</i>	++	+++	<i>Melitaea phoebe</i>	+	+++
<i>Pontia daplidice</i>	+	+++	<i>Melitaea didyma</i>		+++
<i>Euchloe ausonia</i>	+	+	<i>Melitaea trivia</i>	+	+
<i>Anthocharis cardamines</i>	+	++	<i>Melitaea athalia</i>	+	+
<i>Anthocharis belia</i>	+	++	<i>Melitaea deione</i>	+	+
<i>Colias croceus</i>	+++	+++	<i>Melitaea parthenoides</i>	+++	+
<i>Gonepteryx rhamni</i>	++	++	<i>Melitaea cinxia</i>	++	++
<i>Gonepteryx cleopatra</i>	+		<i>Euphydryas aurinia</i>		+++
<i>Leptidea sinapis</i>	+	+++	<b>LIBYTHEIDAE</b>		
			<i>Libythea celtis</i>		+
<b>LYCAENIDAE</b>			<b>SATYRIDAE</b>		
<i>Quercusia quercus</i>	+	++	<i>Melanargia lachesis</i>	+++	+++
<i>Laoeosipis roboris</i>	++	+	<i>Melanargia russiae</i>	++	
<i>Nordmannia ilicis</i>	+		<i>Melanargia ines</i>		+
<i>Nordmannia esculi</i>	+	+	<i>Hipparchia alcyone</i>	++	+++
<i>Strymonidia spini</i>	++		<i>Hipparchia semele</i>	+	+
<i>Callophrys rubi</i>	+		<i>Hipparchia statilinus</i>	+	++
<i>Lycaena phlaeas</i>	+++	+++	<i>Pseudotergumia fidia</i>	+	+
<i>Heodes virgaureae</i>	++		<i>Satyrus actaea</i>	+++	++
<i>Heodes tityrus</i>	+	+	<i>Brintesia circe</i>	++	+
<i>Heodes alciphron</i>	+++	+	<i>Arethusana arethusa</i>	++	+
<i>Lampides boeticus</i>	+	+	<i>Erebia meolans</i>	++	
<i>Syntarucus pirithous</i>	++	+	<i>Maniola jurtina</i>	+	+++
<i>Zizeeria knysna</i>	+		<i>Hyponephele lupina</i>	++	+
<i>Cupido minimus</i>	+		<i>Pyronia tithonus</i>	+++	+++
<i>Celastrina argiolus</i>	+	+++	<i>Pyronia cecilia</i>		++
<i>Glaucoopsyche alexis</i>	+	++	<i>Pyronia bathseba</i>	+	++
<i>Aricia cramera</i>	+++	+++	<i>Coenonympha pamphilus</i>	++	++
<i>Aricia morronensis</i>	+++		<i>Coenonympha iphioides</i>	+++	
<i>Eumedonia eumedon</i>	++		<i>Coenonympha arcania</i>	+	
<i>Cyaniris semiargus</i>	+++		<i>Pararge aegeria</i>	++	+++
<i>Lysandra bellargus</i>	+		<i>Lasiommata megera</i>	+++	+++
<i>Lysandra albicans</i>	+		<i>Lasiommata maera</i>	++	
<i>Philotes panoptes</i>	+		<b>HESPERIIDAE</b>		
<i>Polyommatus icarus</i>	++	++	<i>Pyrgus armoricanus</i>	+	
			<i>Pyrgus serratulae</i>	+++	
<b>NYMPHALIDAE</b>			<i>Spialia sertorius</i>		++
<i>Charaxes jasius</i>	+		<i>Carcharodus alceae</i>	+	+
<i>Limenitis reducta</i>	+	+	<i>Thymelicus acteon</i>		+++
<i>Inachis io</i>	+	+	<i>Thymelicus lineola</i>	+	+
<i>Vanessa atalanta</i>	+	+	<i>Thymelicus flavus</i>	+++	++
<i>Cynthia cardui</i>	++	+	<i>Hesperia comma</i>	++	+
<i>Aglais urticae</i>	+++	+	<i>Ochloides venatus</i>	+	+
<i>Polygonia c-album</i>	+				

N : northern slope ; S : southern slope ; + : 1 to 5 ; ++ : 6 to 15 ; +++ : more than 16 specimens (over 1,850 total specimens).

## Results

Since 1984 a faunal project has been under way in this mountainous area. This study involves (among other things) collecting butterflies and skippers at a dozen sampling sites, which constitute the main habitats of the Sierra.

The results of this preliminary study are shown in Table I, where the species abundances are given. Nomenclature follows HIGGINS & RILEY (1983), with modifications.

The number of species found thus far in the central portion of Sierra de Gredos amounts to 94 (77 on the northern slope and 77 on the southern). There is no difference in the number of species recorded on the two slopes. There are, however small differences in the faunistic composition and the relative abundance of some species.

These 94 species represent 44% of the Iberian butterflies and skippers (see Table II). Of the families, the Pieridae and Nymphalidae stand out, with 55% and 61%, respectively, of the Iberian total, while the Lycaenidae and Hesperiidae are the poorest represented, having 35% and 32%, respectively of the Iberian total.

Table II

Number of species of each family in comparison with total number of Iberian species

Family	N. spp.	S.Spp.	Total spp.	% Iberian spp.
Papilionidae	1	3	3	60%
Pieridae	10	11	12	54%
Lycaenidae	20	16	24	35%
Libytheidae		1	1	100%
Nymphalidae	19	22	23	60%
Satyridae	20	17	22	43%
Hesperiidae	7	7	9	32%
Total	77	77	94	44%

N.spp. : number of northern slope species ; S.spp. : number of southern slope species.

From the biogeographical point of view, Sierra de Gredos acts as a crossroads, where some butterfly and skipper species have their limits of distribution, converging as wet-climate species (for Iberian standards) like *Heodes virgaureae*, *H. tityrus*, *H. alciphron*, *Cupido minimus*, *Limenitis reducta*, *Argynnis paphia*, *Clossiana selene* and *Coenonympha arcania*, as typical Mediterranean species, like *Zerynthia rumina*, *Anthocharis belia*, *Laeosopis roboris*, *Nordmannia esculi*, *Lysandra albicans*, *Charaxes jasius*, *Melitaea deione*, *Libythea celtis*, *Pseudotergumia fidia*, *Saturys actaea*, *Hyponephele lupina*, etc., including the irregularly distributed species *Zizeeria knysna* (HIGGINS & RILEY, 1983 ; GOMEZ BUSTILLO & FERNANDEZ RUBIO, 1974).

Finally, it should be worth mentioning that some very local species have one of their few occurrences in the Sierra de Gredos, e.g. *Aricia morronensis*, *Eumedonia eumedon*, *Coenonympha iphioides* and *Erebia meolans* (EITSCHBERGER & STEINIGER, 1973 ; GARCIA DE VIEDMA & GOMEZ BUSTILLO, 1976 ; HIGGINS & RILEY, 1983 ; MUNGUIRA & MARTIN, 1986).

## References

- EITSCHBERGER, U. & STEINIGER, H., 1973. Die Verbreitung und die Beschreibung einer neuen Rasse von *Aricia morronensis* (RIBBE, 1910). *Atalanta* 4 (6) : 394-402.
- GARCIA DE VIEDMA, M. & GOMEZ BUSTILLO, M.R., 1976. *Libro rojo des los lepidópteros ibéricos*. ICONA, Madrid, 117 p.
- GOMEZ BUSTILLO, M.R. & FERNANDEZ RUBIO, F., 1974. *Mariposas de la Península Ibérica*. Ropalóceros. Tomo II. ICONA, Madrid, 258 pp.
- HIGGINS, L.G. & RILEY, N.D., 1983. *A Field Guide to the Butterflies of Britain and Europe*. Collins, London, 384 pp.
- MUNGUIRA, M.L. & MARTIN, J., 1992. Biology and distribution of the endangered lycaenid *Aricia morronensis* (RIBBE). *Proceedings of the Fifth European Congress of Lepidopterology, Budapest. Nota lepid.*, Suppl. 3 : 11-16.
- RIVAS MARTINEZ, S., 1975. Mapa de vegetación de la provincia de Avila. *Anal. Inst. Bot. Cavanilles* 32 (2) : 1493-1556.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nota lepidopterologica](#)

Jahr/Year: 1992

Band/Volume: [Supp\\_3](#)

Autor(en)/Author(s): Viejo Montesinos José Luis, Martin Jose

Artikel/Article: [Preliminary survey of the butterflies of Sierra de Gredos  
\(Central Spain\) \(Lep.: Hesperioidea et Papilioidea\) 97-100](#)