

ferne in arillum expansa ornatis, arillo sem. aequilongo latere sem. ventrali adpresso, in lobos 2 ovatos diffisso.

Brasilien, Plateau des Itatiaia; coll. II, 490.

Pavonia paraibica. Suffrutex erectus pauciramosus seabridus, caule hinc tomentello parce foliato; foliis spithameis ovato- vel obovato-oblongis producte acuminatis inferne nonnihil angustatis ipsa basi obtusis vel et retusis, crenato-serratis punctulis albis pubeque stellata parce obsitis; racemis aphyllis floriferis brevibus densis, fructiferis elongatis, pedunculis demum unciam longis strictis; involuci phyllis linearibus, cal. 4-plo superantibus, basi in patellam connatis; calyce obsolete repando 5-dentato; corolla rosea invol. haud superante extus pulverulenta; capsula 3—5-cocca, coccis seta iis aequi-longa glochidiata superatis, seminibus obtuse apiculatis.

Brasilien, Urwald von Juiz di fora; coll. II, 132.




Rosa Braunii n. sp.

Auctore J. B. Keller.

Frutex 4—5 pedalis, copiosissime subulato-aculeatus odorem terebinthinaceam conspicue redolens; ramis elongatis duris stricto-patulis, inferne denudatis bruneo-purpureis. Aculei dimorphi, aliis majoribus subulatis dense vel copiose dispositis, aliis in ramis annotinis et parte inferiore ramorum sparsis acutiusculis setaceo-glanduliferis. Foliola elliptica utrinque sensim rotundata supra pilosula subtus omnino villoso-rubiginosa, biserrata. Petioli villoso-glandulosi aculeati. Stipulae purpurascentes subtus villoso-rubiginosae breves, auriculis lanceolatis divergentibus. Pedunculi solitarii hispidae breviusculi (7—12 mm. longi) receptaculis purpurascensibus aequilongi aut paulo breviores. Receptaculum subglobosum totoque hispidum. Sepala dorso dense glanduloso-setosa, purpurea, breviuscule parceque pinnata, erecto-persistentia, apice haud dilatata. Styli e disco plano lanati. Petala . . . ? Fructus globosus setosus, sordide (?) -ruber, erectus.

Hab. In apricis collis Haglersberg prope Bruck a. d. Leitha solo schist. Legit dominus Henricus Braun in ejus honorem dicavi!

Species curiosissima inter Rubiginosas veras (Eurubiginosae Suavifoliae Aut.) distinctissima! Recedit ab affini *R. pimpinelloides* (Mey.) typo triplo majore, foliolis magis ellipticis, receptaculo et fructu majore subgloboso, sepalorum glandulis densis, serraturis argutioribus.



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Österreichische Botanische Zeitschrift = Plant Systematics and Evolution](#)

Jahr/Year: 1882

Band/Volume: [032](#)

Autor(en)/Author(s): Keller J. B.

Artikel/Article: [Rosa Braunii n. sp.. 39](#)