Infrageneric division of the genus *Conocybe* – a classical approach

ANTON HAUSKNECHT Sonndorferstraße 22 A-3712 Maissau, Austria Email: ahausknecht.oemg.@aon.at

IRMGARD KRISAI-GREILHUBER Institut für Botanik der Universität Wien Rennweg 14 A-1030 Wien, Austria Email: irmgard.greilhuber@univie.ac.at

Accepted 18. 9. 2006

Key words: Agaricales, Bolbitiaceae, Conocybe, Gastrocybe. - Infrageneric classification of the genus Conocybe. - New taxa, new combinations.

Abstract: An infrageneric concept of the genus Conocybe including all hitherto known taxa worldwide is presented. New sections, subsections and series are proposed along with listing all representatives in the respective categories. Gastrocybe is included in Conocybe sect. Candidae.

Zusammenfassung: Ein infragenerisches Konzept der Gattung Conocybe auf Basis aller bisher weltweit bekannten Taxa wird vorgestellt. Neue Sektionen, Subsektionen und Serien werden vorgeschlagen und die jeweiligen Vertreter diesen zugeordnet. Die Gattung Gastrocybe wird in Conocybe sect. Candidae eingeordnet.

While preparing a monographical study of the European taxa of the genus *Conocybe*, the first author has studied nearly all type specimens worldwide. Only very few type specimens, marked by (*) in the list, could not be examined microscopically so far. Subsequently, it is attempted to bring all resulting insights into a worldwide infrageneric concept of the genus.

Presently this infrageneric division can only rely upon macro- and microscopical characters and must be conservative in the basic concept in consideration of the existing genus classifications. In Conocybe up to know there is no comprehensive molecular biological study as in other genera. Within Bolbitiaceae MONCALVO & al. (2002) compared three taxa of Agrocybe, two of Bolbitius, one Conocybe, one Pholiotina and one Gastrocybe. They stated that the pair Conocybe-Gastrocybe builds a "conocyboid clade" and the pair Bolbitius-Pholiotina a "bolbitioid clade". Agrocybe is far apart and not necessarily closely related to Bolbitiaceae. However, the Panaeoloideae (Panaeolina, Panaeolus, Copelandia) are much closer to Bolbitiaceae than to Coprinaceae. WALTHER & al (2005) studied mycelia and anamorphs of more than 150 species morphologically and molecular phylogenetically and also included ten Conocybe species (incl. one Pholiotina), one Bolbitius, four Panaeolus and six Agrocybe spp. They found anamorphs in all Conocybe species investigated. This had already been observed by the second author in cultural studies several years ago (unpubl.). Bolbitius and the

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Conocybe species (incl. Pholiotina) appeared as a monophyletic clade, but Bolbitius and Pholiotina were nested in Conocybe. Panaeolus was the sister clade to Conocybe and Agrocybe again was far apart. As these studies are still based on few samples, they must be taken deliberately for a basic concept. They are not relevant for an infrageneric division of the genus Conocybe. HALLEN & al. (2003) examined molecular biologically members of sect. Candidae and compared materials from North America and Europe. Out of it two conclusions can be drawn which are used in the present paper: sect. Candidae is not related with Bolbitius, as stated by BON (1990, 1991) but belongs to Conocybe; and Gastrocybe cannot be maintained as a separate genus but forms a part of sect. Candidae.

In the history of Bolbitiaceae KÜHNER was the first to suggest an infrageneric division of the species known by him from Europe and North Africa, but without specifying valid hierarchical categories. SINGER (1951) for the first time brings a classification of the up to then worldwide known species and divides the genus into two subgenera, Euconocybe (SINGER 1947) with smooth or indistinctly rough spores of >6 µm length, and Ochromarasmius (SINGER 1947) with warty spores of <6 µm length. This concept does not fit any more for all taxa presently known and thus has to be abandoned. WATLING (1982) brings a systematic arrangement of the Bolbitiaceae of Great Britain, but subsumes also Pholiotina and Galerella under Conocybe. In the meantime the independence of *Pholiotina* and *Galerella* as separate genera has been accepted by most authors (at last ARNOLDS 2005). The infrageneric division suggested by us below is based in many details on WATLING (1982), who did not validly publish his "stirps". We publish them now as "series". ARNOLDS (2005) assigns the Dutch species only to sections and includes the rough-spored Conocybe dumetorum in sect. Candidae.

Alltogether, until now the genus Conocybe comprises 150 accepted species and 21 varieties and forms, i.e. 171 taxa.

Excluded or doubtful species

Not included in the 171 taxa are some which could not be examined microscopically so far and whereof a description is lacking or does not allow a classification in the infrageneric concept. These are: Conocybe microgranulosa BATYROVA (Asia), Conocybe morenoi RAITHELH. (South America), Conocybe nivea (MASSEE) WATLING (Europe), and Conocybe oculispora LOCQ. (Europe).

Taxonomical survey

Conocybe FAYOD 1889, Ann. Sci. Nat. Sér. VII, 9: 357.

Original diagnosis: Subhyménium très developpé, trame presque bilatérale. Cellules hyméniales (surtout les paraphyses) courtes, subcylindriques. Basides émergentes. Spores jaune-rouille (sous le microscope), ovoïdes-elliptiques ou plus ou moins lenticulaires, à pore germinatif distinct, sans dépression hilaire.

Type species: Conocybe tenera (SCHAEFF.: FR.) FAYOD, fixed by conservation (DONK 1949, see also HORAK 1968).

188

Österr. Z. Pilzk. 15 (2006)

Characters (predominantly following ARNOLDS 2005, amended on base of all species known worldwide): Basidiocarp mycenoid, rarely collybioid or tricholomatoid. Pileus hygrophanous, dry, greasy or glutinous, glabrous, pubescent or pruinose. Lamellae narrowly adnate or adnexed, sometimes free, close to distant, yellow, yellow-brown, rusty brown to fuscous brown. Stipe central, very rarely eccentric, slender, rarely stout, with equal, bulbous to distinctly radicant base. Partial veil usually absent, in some species present, fugacious or not, leaving small flocks adhering to margin of pileus, volvalike remains at base of stipe or a true membranous volva. Spore print yellow-brown, orange-brown to rusty brown, exceptionally ochre-yellow or blackish brown. Spores glabrous, very rarely nodulose or minutely to distinctly verrucose, thin- to thick-walled, generally yellow-brown to orange-brown, in some species nearly hyaline or blackish. Germ-pore present or not. Basidia clavate to suburniform, 4- or 2-spored. Clamp-connections present in most species. Cheilocystidia always lecythiform with thin, mostly short neck and globose capitulum. Pleurocystidia absent, except in one species. Hymenophoral trama made up of a narrow central strand of cylindrical hyphae surrounded by inflated elements. Stipitipellis a thin cutis, with clusters of lecythiform caulocystidia and/or non-lecythiform ones and long hairs. Pileipellis an epithelioid hymeniderm, made up of pyriform to spheropedunculate, in one species mixed with lageniform elements. Pileocystidia lecythiform or capilliform, sometimes absent. Development in most species paravelangiocarpic, sometimes gymnangiocarpic. Habitat on soil, litter, humus, dung and small pieces of wood, rarely on living wood, solitary or

on soil, litter, humus, dung and small pieces of wood, rarely on living wood, solitary or gregarious, rarely subcaespitose; usually on subneutral to basic substrates rich in nutrients. Widespread, with worldwide distribution.

Comments: The original diagnosis by FAYOD (1889) had to be amended in many respects. The genus description by ARNOLDS (2005) was retained and amended primarily for including deviating characters of all extra-European taxa, – see also SINGER (1986).

Section Conocybe

= Farinosae KÜHNER 1935: 51 inval., p. p.

Characters: basidiocarp mycenoid to collybioid, small to large, with equal, bulbous or radicant stipe. Stipe distinctly couloured, at least in older basidiocarps, more rarely pure white. Spore print ochre brownish, yellow-brown, rubiginous. Spores very small to very large, ellipsoidal, ovoid, limoniform, more rarely subhexagonal, lentiform or not, with very thin to thick wall. Germ-pore mostly present and distinct, in some species absent. Ammoniacal reaction negative to strongly positive. Cheilo- and caulocystidia small to voluminous, with small to large capitulum. Pleurocystidia and pseudoparaphyses absent. Stipitipellis mainly consisting of lecythiform caulocystidia, but in many species mixed with a few piliform, ellipsoidal, cylindrical to lageniform elements. Pileipellis hymeniform, lecythiform pileocystidia present or not. Meadows, grasslands, woods, bare soil, sand dunes, less oftenen dung, also in connection with roots of grasses and herbs.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Comments: KÜHNERS (1935) *Farinosae* also comprised the rough-spored *Conocybe laricina* KÜHNER [= *C. dumetorum* (VELEN.) SVRČEK], which was later placed by SIN-GER (1947) in a separate subgenus.

Series Tenera HAUSKN. & KRISAI, ser. nov.

190

= Stirps *Tenera* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval., p. p.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Stipes cylindricus vel bulbosus, nunquam radicans, nunquam pure albus, minime vetuste distincte coloratus. Sporae grandes usque ad valde grandes, crasse tunicatae, poro germinativo grandi. Reactio ammoniacalis plerumque valde positiva, in nonnullis speciebus infirma vel nulla.

Type species: Conocybe tenera (SCHAEFF.: FR.) FAYOD.

Characters: Basidiocarp mycenoid, small to large, with equal to distinctly bulbous, never radicating stipe. Stipe colour never pure white, soon distinctly coloured. Spore print yellow-brown, rubiginous. Spores large to very large, ellipsoidal, limoniform, in one species subhexagonal, lentiform or not, with thick to very thick wall and distinct germ-pore. Ammoniacal reaction often strongly positive, in some species weak or absent. Cheilo- and caulocystidia small to voluminous, capitulum small to large. Stipitipellis mainly consisting of lecythiform caulocystidia, but in many species mixed with piliform, ellipsoidal, cylindrical to lageniform elements. Pileipellis hymeniform, lecythiform pileocystidia present or not.

Representatives: C. macrocephala KÜHNER & WATLING (Europe, Asia), C. subpallida ENDERLE (Europe, Africa), C. subalpina (SINGER) SINGER & HAUSKN. (Europe), C. aurea (JUL. SCHÄFF.) HONGO (presumably worldwide), C. aurea var. hololeuca HAUSKN. (Europe), C. tenerrima SINGER (South America), C. tenera (SCHAEFF.: FR.) FAYOD (Europe, Asia, Africa, South America?), C. tenera f. excentrica SINGER inval. (South America, at least a good variety), C. subovalis KÜHNER & WATLING (Europe), C. semiglobata KÜHNER & WATLING (worldwide), C. cartilaginipes WATLING (Africa), C. subxerophytica SINGER & HAUSKN. (Europe, Asia, South America), C. subxerophytica var. brunnea HAUSKN. (EUROPE), C. mexicana (MURRILL) WATLING (North America, Europe), C. khasiensis (BERK.) WATLING (Asia).

Comments: Stirps *Tenera* of WATLING (1982) also comprised *Conocybe dunensis* WALLACE, which is assigned a separate series by us due to the specific habitat and different stipe base.

Series Leucopus HAUSKN. & KRISAI, ser. nov.

= Stirps *Leucopus* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium medium usque ad grande. Stipes cylindricus vel leve bulbosus, non radicans, longe albus. Reactio ammoniacalis nulla.

Type species: Conocybe leucopus KÜHNER & WATLING.

Characters: Basidiocarp mycenoid or collybioid, medium large to large. Stipe equal or bulbous, not radicant, white and very long time so. Spore print rubiginous. Spores large, ellipsoidal, ovoid, hardly lentiform, with thick wall and large germ-pore. Ammoniacal reaction negative. Cheilo- and caulocystidia small to medium large. Stipitipellis mainly consisting of lecythiform caulocystidia, non-lecythiform elements sometimes present in small number. Pileipellis hymeniform, lecythiform pileocystidia present or not. Forest soil rich in nitrate, meadows, dry grassland, bare soil (sand).

Representatives: C. ammophila M. LANGE (North America, Asia), C. leucopus KÜHNER & WATLING (Africa, South America), C. striatipes (SPEG.) SINGER (South America, Europe), C. striatipes var. sejuncta SINGER & HAUSKN.* (South America).

Series Dunensis HAUSKN. & KRISAI, ser. nov.

= Stirps *Tenera* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval., p. p.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium medium usque ad grande. Stipes coloratus, basi radicanti et profunde inserta arena. Reactio ammoniacalis positiva vel nulla.

Type species: Conocybe dunensis WALLACE.

Characters: Basidiocarp mycenoid, medium large to large. Stipe coloured, with base radicant to deeply fixed in sand. Spore print rubiginous. Spores medium large to large, ellipsoidal, not lentiform, to distinctly thick-walled, with large germ-pore. Ammonia-cal reaction positive or negative. Cheilo- and caulocystidia small, with small to medium large capitula. Stipitipellis mainly consisting of lecythiform elements, but also mixed with a few hairs and non-lecythiform elements. Pileipellis hymeniform, lecythiform pileocystidia present or not. Only sand soils, fixed dunes.

Representatives: *C. sabulicola* HAUSKN. & ENDERLE (Europe), *C. dunensis* WAL-LACE (Europe), *C. macrorhiza* (SPEG.) SINGER (South America).

Series Mesospora HAUSKN. & KRISAI, ser. nov.

= Stirps *Mesospora* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium medium usque grande, non connexum graminibus vel herbis; Stipes non pure albus, basi cylindrica, bulbosa, non radicanti. Sporae parvae ad mediae, ellipsoideae, limoniformes, tenuitunicatae, poro germinativo distincte. Reactio ammoniacalis (una specie excepto) nulla.

Type species: Conocybe mesospora KÜHNER & WATLING.

Characters: Basidiocarp mycenoid, small to medium large, not in connection with roots of grasses or herbs; Stipe not pure white, based equal to bulbous, not radicant. Spore print yellow-brown to rubiginous. Spores small to medium large, ellipsoidal, limoniform, thin-walled, with distinct germ-pore. Ammoniacal reaction (with one exception) negative. Cheilo- and caulocystidia small. Stipitipellis mainly consisting of lecythiform elements. Pileipellis hymeniform, lecythiform pileocystidia present or not.

Meadows, dry grassland, leaf or needle litter, up to the alpine area, rarely also on sanddunes.

Representatives: C. roberti SINGER & HAUSKN. (Europe), C. microsperma SIN-GER* (South America), C. microspora (VELEN.) DENNIS (Europe, Asia), C. microspora var. brunneola (KÜHNER & WATLING) SINGER & HAUSKN. (Europe, South America), C. brachypodii (VELEN.) HAUSKN. & SVRČEK (Europe), C. ochroalbida HAUSKN. (Europe), C. mesospora KÜHNER & WATLING (worldwide), C. xerophytica SINGER (South America), C. nigrodisca HAUSKN. & KRISAI (Europe), C. roseipes HAUSKN. (Europe).

Series Enderlei HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium valde parvum usque medium, basis stipitis aequalis vel bulbosa. Sporae ellipsoideae, guttiformes, parvae usque mediae, tenui- usque leve crassitunicatae, porus germinativus absens vel sole callus praesens.

Type species: Conocybe enderlei HAUSKN.

Characters: Basidiocarp mycenoid, very small to medium large, with equal or bulbous, never radicant stipe. Spore print yellow-brown to rubiginous. Spores ellipsoidal, drop-shaped, small to medium large at most, often of very different size within a species, thin- to moderately thick-walled. Germ-pore absent or only a callus present. Ammoniacal reaction negative. Stipitipellis mainly consisting of lecythiform elements. Pileipellis hymeniform, lecythiform pileocystidia present or not.

Representatives: C. dennisii HAUSKN. (South America), C. enderlei HAUSKN. (Europe, Asia), C. enderlei var. variispora HAUSKN. (Europe), C. minima SINGER & HAUSKN. (Europe), C. haglundii HAUSKN. (Europe).

Series Magnicapitata HAUSKN. & KRISAI, ser. nov.

= Stirps *Magnicapitata* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

= Stirps *Rickeniana* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: Stipitipellis pileipellisque ut in sectio *Conocybe*. Basidiocarpium parvum usque grande, basis stipitis aequalis usque bulbosa. Sporae ellipsoideae, parvae usque mediae, tenui- usque leve crassitunicatae, poro germinativo. Cheilo- et caulocystidia voluminosa, capitulo grande. Reactio ammoniacalis nulla.

Type species: Conocybe juniana (VELEN.) HAUSKN. & SVRČEK.

Characters: Basidiocarp mycenoid, with equal to bulbous stipe base. Spore print ochreyellow to rubiginous. Spores ellipsoidal, small to medium large, thin- to slightly thickwalled, with germ-pore. Ammoniacal reaction negative. Cheilo- and caulocystidia voluminous, with large capitulum. Stipitipellis mainly consisting of lecythiform caulocystidia, those being mostly more voluminous and with larger capitulum than the cheilocystidia. Pileipellis hymeniform, pileocystidia present or not. Meadows, woods. Österr. Z. Pilzk. 15 (2006)

Representatives: C. spiculoides KÜHNER & WATLING (Europe), C. echinata (VE-LEN.) SINGER (Europe, Asia), C. rickeniana P. D. ORTON (Europe), C. proxima SIN-GER (South America), C. juniana (VELEN.) HAUSKN. & SVRČEK (worldwide), C. juniana var. sordescens (P. D. ORTON) HAUSKN. (Europe), C. juniana var. subsejuncta HAUSKN. (Europe, Asia), C. amazonica SINGER* (South America).

Comments: WATLING (1982) created a separate stirps for *Conocybe rickeniana*; but the species with large, voluminous cystidia with large capitula microscopically are so similar that we refrained from further subdivision.

Series Graminis HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium parvum usque medium, stipes non pure albus, semper connexus radicibus graminum vel herbarum, basi radicanti vel non radicanti. Sporae ellipsoideae, mediae, poro germinativo. Pileocystidia nulla vel rarissima.

Type species: Conocybe graminis HAUSKN.

Characters: Basidiocarp mycenoid, small to medium large. Stipe not pure white, rootlike elongation present or not, always connected to roots of grasses or herbs. Spore print yellow-brown to rubiginous. Spores ellipsoidal, medium large, thin- to moderately thick-walled, with germ-pore. Ammoniacal reaction negative. Stipitipellis mainly consisting of lecythiform caulocystidia, in between sporadically non-lecythiform elements present. Pileipellis hymeniform, pileocystidia absent or very rare. Dry grassland, exposed loess slopes, steppes, semi-deserts.

Representatives: C. graminis HAUSKN. (Europe, Asia, Africa), C. herbarum HAUSKN. (Europe, Asia).

Series Antipus HAUSKN. & KRISAI, ser. nov.

= Stirps Antipus WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: Cystidia, stipitipellis pileipellisque ut in sectione *Conocybe*. Basidiocarpium medium usque grande, stipes (una specie excepto) longe radicans, juvenile saepe albus vel pallide coloratus. Sporae parvae usque grandes, ellipsoideae, ovoideae, submitriformes, angulatae usque hexagonae, lentiformes aut non lentiformes, plerumque crassitunicatae, poro germinativo distincte.

Type species: Conocybe antipus (LASCH: FR.) FAYOD.

Characters: Basidiocarp mycenoid, Stipe long radicant (with one exception), young often white or pale, old slightly stronger coloured. Spore print yellow-brown to rubiginous. Spores small to large, ellipsoidal, ovoid, submitriform, angular to hexagonal, lentiform or not, mostly thick-walled with distinct germ-pore. Ammoniacal reaction negative. Stipitipellis consisting of mainly lecythiform caulocystidia (these often with narrow ventre and small capitulum). Pileipellis hymeniform, lecythiform pileocystidia present or not. Grassland, meadows, fertilized soils, dung.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at

194

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Representatives: C. antipus (LASCH: FR.) FAYOD (Europe, North America), C. humicola (THIERS) HAUSKN., KRISAI & VOGLMAYR (North America), C. alboradicans ARNOLDS (Europe), C. pragensis HAUSKN. (Europe).

Section Mixtae SINGER 1962 a, Sydowia 15: 68

= Mixtae KÜHNER 1935: 85, inval.

Latin diagnosis: *Conocybe* sect. *Mixtae* KÜHN. (primitus subsectio!) ex SING. sect. nov. Dermatocystidiis stipitis et capitatis et non capitatis praesentibus; pilis stipitis etiam numerosis.

Type species: Conocybe cryptocystis (G. F. ATK.) SINGER.

Characters: Basidiocarp mycenoid, very small to large, with equal, distinctly bulbous to distinctly radicant stipe. Spore print ochre-yellow to rubiginous. Spores small to very large, ellipsoidal, naviculate, limoniform, in one species subhexagonal, barely lentiform, thin- to thick-walled, with distinct, often very large germ-pore. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of a mixture of lecythiform and non-lecythiform elements. Pileipellis hymeniform, often with capilliform, rarely lecythiform pileocystidia. Dung, compost, fertilised places, decomposing leaf- and needle litter, meadows, exceptionally wood.

Comments: The section *Mixtae* was created by SINGER (1962 a) for species with a mixture of lecythiform and non-lecythiform caulocystidia and in a very wide sense. WATLING (1986) and HAUSKNECHT (2003) only include species in this section which have a mixture ratio of ca. 1:5 to 5:1; species with only a very low proportion of lecythiform or non-lecythiform caulocystidia are excluded.

Subsection Pubescens HAUSKN. & KRISAI, subsect. nova

= Stirps *Pubescens* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: Basidiocarpium medium usque grande, stipes aequalis vel distincte bulbosus, non radicans. Sporae grandes usque valde grandes, ellipsoideae, poro germinativo grandi vel valde grandi. Stipitipellis caulocystidiis lecythiformibus et non lecythiformibus aequali numero consistens. Pileipellis hymeniformis, saepe pileocystidiis capilliformibus. In stercore, habitatione stercorata, acuis et foliis putridis.

Type species: C. pubescens (GILLET) KÜHNER.

Characters: Basidiocarp mycenoid, medium large to large, with equal or distinctly bulbous, not radicant stipe. Spore print yellow-brown to rubiginous. Spores large to very large, ellipsoidal, with large to very large, in one species eccentrical germ-pore. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of almost equivalent lecythiform and non-lecythiform caulocystidia. Pileipellis hymeniform, often with capilliform pileocystidia. Dung, compost, fertilised places, rotting leaf- and needle litter.

Representatives: C. subpubescens P. D. ORTON (worldwide), C. mixta SINGER (South America), C. pulchella (VELEN.) HAUSKN. & SVRČEK (Europe, Asia, Africa),

C. curta (G. F. ATK.) WATLING (North America), C. bulbifera (KAUFFMAN) ROMAGN. (North America), C. cryptocystis (G. F. ATK.) SINGER (North America), C. reticulatorugosa SINGER (South America), C. merdaria ARNOLDS & HAUSKN. (Europe, Asia, South America), C. pubescens (GILLET) KÜHNER (worldwide), C. acutoconica WAT-LING (Africa), C. macrospora (G. F. ATK.) HAUSKN. (worldwide).

Comments: In this very homogenous subsection all species are included with a mixture of almost equivalent lecythiform and non-lecythiform caulocystidia, it corresponds exactly stirps *Pubescens* (WATLING 1982).

Subsection Ambiguae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium valde parvum usque grande, stipes aequalis, bulbosus vel radicans. Sporae parvae usque grandes, naviculatae, limoniformes, rariore ellipsoideae, etiam lentiformes et distincte angulatae, tenuiter usque crasse tunicatae, poro germinativo aut nullo. Stipitipellis mixtura elementis lecythiformibus usque 80 % et non-lecythiformibus, vel caulocystidiis non-lecythiformibus maximam partem. Pileipellis hymeniformis, pileocystidiis (capilliformibus, lecythiformibus) aut nullis.

Type species: Conocybe ambigua WATLING.

Characters: Basidiocarp mycenoid, very small to large. Stipe with equal, bulbous or distinctly radicant base. Spore print ochre-yellow to rubiginous. Spores small to large, naviculate, limoniform, more rarely regularely ellipsoidal, in one species subhexagonal, hardly lentiform, thin- to thick-walled, with distinct germ-pore. Stipitipellis consisting of a mixture of lecythiform and non-lecythiform elements, either with up to 80 % lecythiform elements or with mainly non-lecythiform (capilliform, cylindrical, ellipsoidal to lageniform) caulocystidia. Pileipellis hymeniform, pileocystidia lecythiform, capilliform or absent. Meadows, forest soil, dung, rarely wood.

Type species: Conocybe ambigua WATLING.

Comments: This is a rather inhomogenous subsection. On the one hand it comprises species with mainly lecythiform caulocystidia (ser. *Cettoiana* and *Fragilis*), on the other hand such with mainly capilliform and non-lecythiform elements (ser. *Ambigua*). A further division into subsections does not seem to be meaningful, probably molecular biological examinations will provide a better insight into the relationships.

Series Ambigua

= Stirps *Ambigua* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Characters: Basidiocarp mycenoid, medium large to large. Stipe with equal to bulbous, never radicant base. Spore print ochre-yellow to rubiginous. Spores medium large to large, ellipsoidal, naviculate, limoniform, not to distinctly lentiform-hexagonal, thinto thick-walled. Basidia 2- or 4-spored. Stipitipellis consisting of a mixture of lecythiform and non-lecythiform elements. Pileipellis hymeniform, (capilliform, lecythiform) pileocystidia present or not. Meadows, forest soil, one species on wood of a living tree.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at 196 A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of *Conocybe*

Representatives: *C. ambigua* WATLING (Europe, Asia, North America), *C. alba* SINGER (South America), *C. acutoconica* WATLING (Africa), *C. siligineoides* R. HEIM (North America), *C. zuccherellii* HAUSKN. (Europe).

Comments: WATLING (1982) created this stirps for *Conocybe ambigua*; we include here all species with non-radicant stipe, in which capilliform and non-lecythiform caulocystidia, respectively, prevail and/or shape and size of spores deviates from subsection *Pubescens*.

Series Cettoiana HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Basidiocarpium medium usque grande, stipite distincte radicanti. Sporae mediae usque grandes, ellipsoideae, limoniformes, non lentiformes, tenuiter usque crasse tunicatae, poro germinativo. Stipitipellis consistens caulocystidiis lecythiformibus maximam partem, sed elementa capilliformia et alia elementa non-lecythiformia semper numerosa. Pileipellis hymeniformis, pileocystidia lecythiformia praesens aut non.

Type species: Conocybe cettoiana HAUSKN. & ENDERLE.

Characters: Basidiocarp mycenoid, medium large to large, with distinctly radicant stipe base. Spore print yellow-brown to rubiginous. Spores medium large to large, ellipsoidal, limoniform, not lentiform, thin- to thick-walled, with distinct germ-pore. Stipitipellis consisting of mainly lecythiform caulocystidia, but capilliform and other non-lecythiform elements always numerous (20-30 %). Pileipellis hymeniform, pileocystidia lecythiform or absent. Dung, compost, but also sand and forest soil.

Representatives: *C. izonetae* SINGER (South America), *C. cettoiana* HAUSKN. & ENDERLE (Europe), *C. solitaria* K. A. THOMAS, HAUSKN. & MANIM. (Asia), *C. fiorii* (D. SACC.) WATLING (Europe).

Series Fragilis HAUSKN. & KRISAI, ser. nov.

= Stirps *Fragilis* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval., p. p.

Latin diagnosis: Basidiocarpium valde parvum usque medium, basi stipitis aequali usque bulbosa. Sporae parvae usque mediae, ellipsoideae, ovoido-ellipsoideae, tenuiter usque leviter crasse tunicatae, poro germinativo praesens aut non. Stipitipellis consistens maximam partem caulocystidiis lecythiformibus, numerosa elementa non-lecythiformia immixta. Pileipellis hymeniformis, pileocystidia lecythiformia, capilliformia praesentia aut non.

Type species: Conocybe fragilis (PECK) SINGER.

Characters: Basidiocarp mycenoid, very small to medium large. Stipe not radicant, equal or bulbous. Spore print yellow-brown to rubiginous. Spores small to medium large, ellipsoidal, ovoid, not lentiform, thin- to thick-walled, germ-pore present or not. Stipitipellis consisting of mainly lecythiform caulocystidia, mixed with ca. 20-50 % non-lecythiform elements. Pileipellis hymeniform, pileocystidia lecythiform, capilliform or absent. Dry meadow, meadows, forest soil, also on wood.

Österr. Z. Pilzk. 15 (2006)

Representatives: *C. fragilis* (PECK) SINGER (North America), *C. xylophila* SINGER* (South America), *C. monicae* HAUSKN. (Europe, Africa), *C. tetrasporoides* HAUSKN. (North America, Europe), *C. tuxlaensis* SINGER (North- and South America, Europe), *C. lobauensis* SINGER & HAUSKN. (Europe).

Comments: This series also is not very homogenous, especially *C. tuxlaensis* and *C. lobauensis* have some characters deviating more from the other species in the series.

Subsection Mitrisporae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium medium, stipite leviter radicanti. Sporae parvae, distincte mitriformes, crasse tunicatae, poro germinativo truncato. Stipitipellis consistens caulocystidiis capilliformibus, cylindricis, ellipsoideis, lageniformibus et multis lecythiformibus. Pileipellis hymeniformis, pileocystidia absentia. Solo silvarum.

Type species: Conocybe mitrispora WATLING.

Characters: Basidiocarp mycenoid, medium large, with long, slightly radicant stipe base. Spore print colour unknown. Spores in front view distinctly mitriform, in side view ellipsoidal, lentiform, thick-walled with truncate germ-pore. Stipitipellis consisting of capilliform, cylindrical, ellipsoidal, lageniform and many lecythiform caulo-cystidia. Pileipellis hymeniform, pileocystidia absent. Forest soil.

Single representative: C. mitrispora WATLING (Asia).

Comments: WATLING (1994) states that within the genus *Conocybe* the exceptional form of spores is not correlated with the structure of the stipitipellis, there are species with similar form of spores also in sect. *Pilosellae*. Within sect. *Mixtae* there is no other species with similar spores.

Section Nodulososporae WATLING 1976 a, Kew Bull. 31: 594

Latin description: *Conocybe* section *nodulososporae* WATLING, sect. nov. Pileo epicute ex elementis pyriformibus vel elongato-clavatis; cheilocystidia lecythiformia; caulocystidia cylindrica vel clavata; basidiosporae ellipsoideae, nodulosae.

Type species: Conocybe nodulosospora (HONGO) WATLING.

Characters: Basidiocarp mycenoid, medium large to large, with equal, eventually radicant stipe base. Spore print colour unknown. Spores nodulous, cruciform, triangular, thick-walled, with distinct germ-pore. Stipitipellis consisting of non-lecythiform elements. Pileipellis hymeniform, pileocystidia present or not. Forest soil.

Representatives: *C. nodulosospora* (HONGO) WATLING* (Asia), *C. javanica* SINGER (Asia).

Comments: WATLING published the Latin description for this section in the year 1976 two times in different papers and journals (WATLING 1976 a, b). Due to the numbering of his "Observations" and according to some other hints the publication in Kew Bulletin may have been earlier, but the exact date of issuing is admittedly unknown to us.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at

198

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Section Pilosellae SINGER 1962 a, Sydowia 15: 68

= *Pilosellae* KÜHNER 1935: 91, inval.

Latin diagnosis: *Conocybe* sect. *Pilosellae* KÜHN. (primitus ut subsectio!) ex SING. sect. nov. Stipite haud toto albo; hyphis pilosis numerosis; dermatocystidiis capitatis in stipite nullis.

Type species: Conocybe pilosella (PERS.: FR.) KÜHNER.

Characters: Basidiocarp mycenoid, small to large. Stipe mostly coloured, at least when old, with equal, bulbous or radicant base. Pileus expanded, in one species elongated, cylindrical-ovoid. Spore print ochre-, yellow-brown to rubiginous. Spores very small to very large, ellipsoidal, ovoid, naviculate, limoniform or subcylindrical, lentiform or not, thin- to thick-walled, germ-pore present or not. Pleurocystidia and pseudoparaphyses absent. Stipitipellis mainly consisting of non-lecythiform elements, but in many species lecythiform caulocystidia present, at least near the stipe apex. Pileipellis hymeniform, consisting of sphaeropedunculate elements. Pileocystidia mostly capilliform, rarely absent or lecythiform. Meadows, woods, dung, decomposing plant debris, rarely also on wood.

Subsection Pilosellae

Characters: Basidiocarp mycenoid, small to large, Stipe with equal, bulbous or distinctly radicant base. Spore print yellow-brown to rubiginous. Spores very small to very large, ellipsoidal, ovoid, limoniform, lentiform or not, thin- to thick-walled, germ-pore present or not. Stipitipellis mainly consisting of non-lecythiform elements, in many species also lecythiform caulocystidia present, at least near the stipe apex. Pileipellis hymeniform, consisting of sphaeropedunculate elements. Pileocystidia mostly capilliform, rarely absent or also lecythiform. Meadows, woods, dung, decomposing plant debris, rarely also on wood.

Series Pilosella

= Stirps *Pilosella* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Characters: Stipitipellis and pileipellis as for the section. Basidiocarp mycenoid, medium large, with bulbous, not radicant stipe base. Pileus moist striate. Spore print yellow-brown to rubiginous. Spores very small to small, ellipsoidal, ovoid, thin-walled, hardly visible germ-pore present or not. Meadows, woods.

Representatives: *C. pilosella* (PERS.: FR.) KÜHNER (nearly worldwide), *C. pilosella* var. *brunneonigra* HAUSKN. & KRISAI (Europe), *C. pallidospora* KÜHNER & WATLING (Europe, Asia, North America).

Series Sienophylla HAUSKN. & KRISAI, ser. nov.

= Stirps *Siliginea* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval., p. p.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium parvum usque medium, base stipitis aequalis vel bulbosa, non radicanti. Pileus ochraceo-brunneus usque propre ater, plerumque distincte striatus. Sporae parvae usque mediae, ellipsoideae vel limoniformes, amygdaliformes, lentiformes aut non, tenuiter usque leviter crasse tunicatae, poro germinativo.

Type species: Conocybe sienophylla (BERK. & BROOME) SINGER.

Characters: Basidiocarp mycenoid, small to medium large, with equal or bulbous, never radicant stipe base. Pileus ochre, pale brown, dark brown to almost black, moist mostly distinctly striate. Spore print pale brown, yellow-brown to rubiginous. Spores small to medium large, ellipsoidal, lemon- or almond-shaped, lentiform or not, thin- to moderately thick-walled, with germ-pore. Stipitipellis mainly consisting of capilliform, cylindrical, ellipsoidal to lageniform elements, but lecythiform caulocystidia often present near the stipe apex. Pileipellis hymeniform, capilliform pileocystidia present or not. Meadows, woods, up to the alpine area, greenhouses.

Representatives: C. rostellata (VELEN.) HAUSKN. & SVRČEK (nearly worldwide), C. sienophylla (BERK. & BROOME) SINGER (nearly worldwide), C. halophila SINGER (Asia, Europe), C. ochrostriata HAUSKN. (nearly worldwide), C. ochrostriata var. favrei HAUSKN. (Europe), C. mazatecorum SINGER* (North America), C. nigrescens HAUSKN. & GUBITZ (Europe, South America), C. moseri WATLING (Europe, Asia, Africa), C. moseri var. bisporigera HAUSKN. & KRISAI (Europe), C. moseri var. robustior HAUSKN. & H.-J. HÜBNER (Europe).

Comments: WATLING (1982) included in his stirps *Siliginea* also the species around *Conocybe sienophylla* and *C. kuehneriana* (= *C. velutipes*). We retain the species with pale, non-striate pileus there and place all other species in two new series, mainly because many new taxa, especially in Europe, have been detected. The taxa with small spores belong now to ser. *Sienophylla*, *C. velutipes* with large spores is included in the following ser. *Anthracophila*.

Series Anthracophila HAUSKN. & KRISAI, ser. nov.

= Stirps *Anthracophila* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval.

= Stirps *Siliginea* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval., p. p.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium medium usque grande, base stipitis aequalis vel bulbosa, non radicanti. Pileus plerumque valde coloratus, umide fere distincte striatus. Sporae grandes, ellipsoideae, ovoideae, lentiformiter compressae aut non, crasse tunicatae, poro germinativo grandi.

Type species: Conocybe anthracophila KÜHNER & WATLING.

Characters: Basidiocarp mycenoid, medium large to large. Stipe not radicant, base equal or bulbous. Pileus mostly strongly coloured, moist mostly distinctly striate. Spore print yellow-brown to strongly rubiginous. Spores large, ellipsoidal or ovoid, partly lentiform compressed, thick-walled with large germ-pore. Stipitipellis consisting of non-lecythiform elements, lecythiform caulocystidia very rarely present near the

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at 200 A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of *Conocybe*

stipe apex. Pileipellis hymeniform, capilliforme pileocystidia present or not. Meadows, woods, fire places, also on fertilised soil.

Representatives: *C. anthracophila* KÜHNER & WATLING (Europe, Asia, Africa, South America), *C. anthracophila* var. *ovispora* HAUSKN. (Europe), *C. raphanacea* WATLING (Africa), *C. alachuana* (MURRILL) HESLER (North America), *C. caespitosa* (MURRILL) WATLING (North America), *C. velutipes* (VELEN.) HAUSKN. & SVRČEK (nearly worldwide), *C. peroxydata* (BERK.) D. A. REID (Africa), *C. ochraceodisca* WATLING (Africa).

Series Bispora HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium parvum usque medium, base stipitis aequalis vel bulbosa, non radicanti. Sporae naviculatae, limoniformes, subcylindricae, tenuiter usque crasse tunicatae, porus germinativus praesens aut non.

Type species: Conocybe bispora (SINGER) HAUSKN.

Characters: Basidiocarp mycenoid, small to medium large, with cylindrical, equal or slightly bulbous stipe, not radicant. Spore print yellow-brown to rubiginous. Spores naviculate, limoniform, subcylindrical, never markedly symmetrical ellipsoidal, germpore present or not, thin- to thick-walled. Basidia 2-, in one species also mostly 4-spored. Stipitipellis mainly consisting of non-lecythiform elements, sometimes mixed with a minor number of lecythiform caulocystidia. Pileipellis hymeniform. Pileocystidia absent, rarely capilliform and lecythiform. Meadows, forest soil.

Representatives: *C. phaeodropis* (BERK. & BROOME) PEGLER (Asia), *C. bispora* (SINGER) HAUSKN. (Europe, North America), *C. umbellula* (MONT.) SINGER (Southand North America, Asia).

Comments: Representatives of this series differ from the preceding mostly by spore shape and basidia mainly 2-spored.

Series Microrrhiza HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium parvum usque medium, stipite cylindrici, radicanti, basis stipitis numquam bulbosa. Sporae parvae usque grandes, naviculatae, limoniformes usque ellipsoideae, non-numquam lentiformes, tenuiter usque crasse tunicatae, poro germinativo.

Type species: Conocybe microrrhiza HAUSKN.

Characters: Basidiocarp mycenoid, small to medium large, with cylindrical, radicant stipe; Stipe base never bulbous. Spore print yellow-brown to rubiginous. Spores small to large, very variabel, naviculate, limoniform, ellipsoidal, in some species lentiform, with thin to thick wall, with germ-pore. Stipitipellis consisting of capilliform, cylindrical, ellipsoidal to lageniform elements, in some species with mixed with a few lecythiform caulocystidia. Pileipellis hymeniform, scarce lecythiform pileocystidia present or not.

Österr. Z. Pilzk. 15 (2006)

Representatives: C. leporina (VELEN.) SINGER (Europe, Asia), C. microrrhiza HAUSKN. (Europe), C. microrrhiza var. parvispora (HAUSKN.) HAUSKN. (Europe), C. microrrhiza var. tetraspora (SINGER & HAUSKN.) HAUSKN. (Europe), C. radicans K. A. THOMAS, HAUSKN. & MANIM. (Asia), C. myosura SINGER* (South America), C. incarnata (JUL. SCHÄFF.) HAUSKN. & ARNOLDS (Europe).

Series Inocybeoides HAUSKN. & KRISAI, ser. nov.

= Stirps *Inocybeoides* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium grande, stipiti albo, crasso et basi stipitis valde tumefacta usque bulbosa, non radicanti. Sporae grandes, ellipsoideae, leviter cylindricae usque irregulaes, crasse tunicatae, poro germinativo grandi.

Type species: Conocybe inocybeoides WATLING.

Characters: Basidiocarp mycenoid, large. Stipe white, old whitish, thick, with strongly inflated to broadly bulbous base, not radicant. Spore print yellow-brown to rubiginous. Spores large, ellipsoidal, slightly cylindrical to irregular, at most weakly lentiform, thick-walled with large germ-pore. Basidia 2- or 4-spored. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of non-lecythiform elements, lecythiform caulocystidia extremely rare near the stipe apex. Pileipellis hymeniform, with capilliform pileocystidia. Meadows, woods, disturbed sites, plant debris, compost.

Representatives: C. apala (FR.: FR.) ARNOLDS (Europe), C. inocybeoides WAT-LING (Europe).

Series Cylindracea HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium tenerum. Pileus cylindricus, haud expansus, semper altior ut latus. Sporae mediae, ellipsoideae, tenuiter tunicatae, distincte poro germinativo.

Type species: *Conocybe cylindracea* KÜHNER & WATLING.

Characters: Basidiocarp mycenoid, brittle. Pileus cylindrical to elongate ovoid, hardly expanding, always higher than wide. Stipe base slightly bulbous. Spore print yellow-brown. Spores medium large, ellipsoidal, thin-walled with distinct germ-pore. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of only capilliform, cylindrical to fusiform elements. Pileipellis hymeniform, with capilliform pileocystidia. Dry grassland, fire places.

Single representative: C. cylindracea KÜHNER & WATLING (Africa, Europe).

Subsection Siligineae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium medium usque grande, rare parvius, basis stipitis aequalis, bulbosa vel distincte radicans. Sporae grandes, ellipsoideae, lentiformes aut non, subhexagonae aut non, plerumque crasse tunicatae, poro germinativo grandi. A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Characters: Basidiocarp mycenoid, mostly medium large to large, rarely small. Stipe with equal, bulbous to distinctly radicant base. Spore print yellow-brown to rubiginous. Spores large, ellipsoidal, lentiform or not, subhexagonal or not, mostly thick-walled, with large germ-pore. Stipitipellis consisting of capilliform, cylindrical, ellipsoidal to slightly lageniform elements, lecythiform caulocystidia absent or present (in few species) in moderate number. Pileipellis hymeniform, capilliform pileocystidia present or not. Dung, fertilised soil, compost, few species also on meagre places in meadows and woods.

Comments: The mostly dung-inhabiting species of this subsection correspond to those of subsect. *Pubescens* within sect. *Mixtae*, but in the stipitipellis there are no or only exceptionally lecythiform caulocystidia. Especially representatives of ser. *Fimetaria* are macroscopically similar to the *C.-pubescens*-group.

Series Siliginea

= Stirps *Siliginea* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval., p. p.

Characters: Basidiocarp mycenoid, small to large, stipe base equal or bulbous. Pileus relatively pale coloured, moist never striate. Spore print yellow-brown to rubiginous. Spores large, ellipsoidal, never lentiform, thick-walled with large germ-pore. Basidia 2- or 4-spored. Stipitipellis mainly consisting of non-lecythiform elements, lecythiform caulocystidia sporadically present near the stipe apex. Pileipellis hymeniform, often with capilliform pileocystidia. Dung, rich and meagre meadows, grassland.

Representatives: C. fuscimarginata (MURRILL) SINGER (worldwide), C. crocospora (BERK. & M. A. CURTIS) KÜHNER inval. (North America), C. siliginea (FR.: FR.) KÜHNER (Europe, Asia, North America), C. capillaripes (PECK) WATLING (North America), C. rickenii (JUL. SCHÄFF.) KÜHNER (worldwide).

Series Fimetaria HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione *Pilosellae*. Basidiocarpium medium usque grande, basi stipitis aequali, numquam bulbosa, sed radicanti vel substrato profunde inserta. Sporae grandes, ellipsoideae, crasse tunicatae, poro germinativo grandi. In stercore.

Type species: Conocybe fimetaria WATLING.

Characters: Basidiocarp mycenoid, medium large to large, with equal, never bulbous stipe; stipe mostly distinctly radicant or deeply inserted in the substrate. Spore print rubiginous. Spores large, ellipsoidal, seldom weakly lentiform or indistinctly angular, with large germ-pore. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of non-lecythiform elements, lecythiform caulocystidia present in one species in low number. Pileipellis hymeniform, capilliform pileocystidia present or not. Dung.

Representatives: C. fimetaria WATLING (Europe, Africa, North America), C. diemii SINGER (South America), C. watlingii HAUSKN. (Europe).

202

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at Österr. Z. Pilzk. 15 (2006)

Series Murinacea HAUSKN. & KRISAI, ser. nov.

= Stirps Murinacea WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval.

Latin diagnosis: characteres ut in serie Fimetaria, sed basis stipitis aequalis usque distincte bulbosa, non radicans. In stercore, solo stercorato, una species acuis putridis. Type species: Conocybe murinacea WATLING.

Characters: Basidiocarp mycenoid, small to large, with equal to distinctly bulbous, never radicant stipe. Spore print yellow-brown to rubiginous. Spores large to very large, ellipsoidal, slightly lentiform or subhexagonal, with large, in one species eccentrical germ-pore. Stipitipellis consisting of non-lecythiform elements, very rarely mixed with lecythiform caulocystidia. Pileipellis hymeniform, capilliform pileocystidia present or not. Dung, compost, droppings, heavily fertilised soil, one species on needle litter.

Representatives: C. murinacea WATLING (Europe, Asia), C. farinacea WATLING (Europe), C. atkinsonii WATLING (North America), C. pseudopubescens K. A. THOMAS, HAUSKN. & MANIM. (Asia), C. singeriana HAUSKN. (Europe, Asia, Africa, Oceania), C. pinetorum WATLING, ESTEVE-RAV. & G. MORENO (Europe), C. magnispora (MUR-RILL) SINGER (North- and South America, Europe, Asia), C. gigasperma ENDERLE & HAUSKN. (Europe, Asia).

Series Lenticulospora HAUSKN. & KRISAI, ser. nov.

= Stirps *Fragilis* WATLING 1982, British Fungus Flora Agarics and Boleti 3: 41, inval., p. p.

Latin diagnosis: Stipitipellis et pileipellis ut in sectione Pilosellae. Basidiocarpium parvum usque medium, basi stipitis aequali usque leviter bulbosa, non radicanti. Sporae parvae usque grandes, semper distincte lentiformes et leviter angulatae usque subhexagonae, tenuiter usque crasse tunicatae, poro germinativo.

Characters: Basidiocarp mycenoid, small to medium large. Stipe cylindrical, equal or slightly bulbous, not radicant. Spore print yellow-brown to rubiginous. Spores small to large, always distinctly lentiform and in addition slightly angular to subhexagonal, thin- to thick-walled, with germ-pore. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of non-lecythiform elements. Pileipellis hymeniform, with capilliform pileocystidia. Dung, compost, meadows, woods, rarely on wood.

Representatives: C. lentispora SINGER (South America), C. hexagonospora HAUSKN. & ENDERLE (Europe, Asia), C. brunneoaurantiaca K. A. THOMAS, HAUSKN. & MA-NIM. (Asia, Australia), C. lenticulospora WATLING (Europe, Asia, North- and South America), C. pulchra (CLEM.) HAUSKN., KRISAI & VOGLMAYR (North- and South America, Europe).

Section Obscurae HAUSKN. & KRISAI, sect. nova

Latin diagnosis: Basidiocarpium validum, stipite cylindrico. Sporae parvae, ellipsoideae, poro germinativo distincte. Pleurocystidia et pseudoparaphyses nulla. Stipes 204

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

caulocystidiis cylindricis et grandibus late fusiformibus. Pileipellis hymeniformis, mixtura elementis sphaerico-pedunculatis et late fusiformibus.

Type species: Conocybe obscura WATLING.

Characters: Basidiocarp mycenoid, robust, with cylindrical stipe. Spore print pale cinnamon. Spores small, ellipsoidal, with small germ-pore. Pleurocystidia and pseudoparaphyses absent. Caulocystidia not lecythiform, cylindrical, mixed with large, broadly fusiform elements. Pileipellis hymeniform, consisting of a mixture of sphaeropedunculate and broadly fusiform, partly rostellate elements (similar to *Pluteus* subsekt. *Mixtini* SINGER, see SINGER 1958). Forest soil.

Single representative: C. obscura WATLING (Africa).

Comments: This is the only species in the genus *Conocybe* with a mixed pileipellis structure. WATLING (1974) states that this fact together with additional macroscopical informations gives a hint towards using *C. obscura* as type of a new section.

Section Conocybella (SINGER) HAUSKN. & KRISAI, comb. nova

Basionym: *Psathyrella* subg. *Conocybella* SINGER 1948, Sydowia **2**: 36. = *Conocybe* sect. *Conocybella* WATLING 1975, Notes Roy. Bot. Gard. Edinburgh **34**: 248, inval. (wrong basionym cited).

Latin diagnosis: *Psathyrella* subgenus *Conocybella* A. H. SMITH in litt. Cheilocystidiis vesiculosis vel conocyboideis, i. e. ventricosis et globulo parvo stipitato ad apicem praeditis; cystidiis ad latera lamellarum nullis; sporis exiguis (i. e. minoribus quam 10μ longitudine).

Type species: Conocybe michiganensis (A. H. SM.) WATLING.

Characters: Basidiocarp mycenoid. Lamellae mature "fuscous brown", Stipe stiff. Spore print "Benzo-brown". Spores ellipsoidal, with germ-pore, under 10 μ m long, nearly black in potassium hydroxide. Cheilocystidia lecythiform, pleurocystidia and pseudoparaphyses absent. On sawdust and pieces of wood.

Single representative: C. michiganensis (A. H. SM.) WATLING (North America).

Comments: Conocybe michiganensis was originally described as *Psathyrella* because of the colour of spores and spore print (SMITH 1941). WATLING (1975) made the new combination into *Conocybe* and erected the new sect. *Conocybella*, but invalid, as he cited a wrong basionym. Thus it proved to be necessary to validate sect. *Conocybella*.

Section Candidae SINGER 1962 a, Sydowia 15: 69

= Candidae KÜHNER 1935: 120 inval.

= Bolbitius sect. Candidi (KÜHNER ex SINGER) BON 1990, Doc. Mycol. 20/78: 39

Latin diagnosis: *Conocybe* sect. *Candidae* KÜHNER (primitus subsectio!). Stipite candido – dermatocystidiis stipitis nullis capitatis; hyphis piliformibus numerosis; pileo frequenter striato; pseudoparaphysibus frequenter numerosis. Typus sectionis: *C. lateritia* (FR.) KÜHNER sensu KÜHNER.

Type species: Conocybe albipes (G. H. OTTH) HAUSKN.

Characters: Basidiocarp mycenoid or gasteroid, small to large. Pileus dry or slimy, partly higher than wide, often with slightly crenulate margin. Lamellae, in one species also pileus deliquescent when mature. Stipe base equal to bulbous, stipe not radicant. Spore print ochre-yellow to rubiginous. Spores medium large to large, ellipsoidal, lentiform, in one species hexagonal-mitriform, thin- to thick-walled, with one exception with distinct germ-pore. Pseudoparaphyses present. Pleurocystidia absent. In meadows, grassland, woods, on dung, tropical species in Europe mostly in greenhouses.

Comments: BON (1990) included sect. *Candidae* in the genus *Bolbitius* due to the presence of pseudoparaphyses. HALLEN & al. (2003) provided molecular biological evidence that this section is clearly situated in the genus *Conocybe* and far apart from *Bolbitius*. *Bolbitius* rather is closer to *Pholiotina* than to *Conocybe*.

Subsection Candidae

Characters: Basidiocarp mycenoid, small to large, with often slightly crenulated pileus margin. Lamellae mature deliquescent. Stipe base equal to bulbous. Spore print ochre-yellow to rubiginous. Spores medium large to large, mostly ellipsoidal, sometimes lentiform, in one species hexagonal-mitriform. Pseudoparaphyses present. meadows, grassland, woods, on dung, in greenhouses.

Series Albipes HAUSKN. & KRISAI, ser. nov.

= Stirps *Lactea* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval.

Latin diagnosis: Sporae grandes, ellipsoideae, lentiformes aut non, numquam hexagono-mitriformes, crasse usque tenuiter tunicatae, porus germinativus praesens aut non. Pseudoparaphyses praesentes.

Type species: Conocybe albipes (G. H. OTTH) HAUSKN.

Characters: Basidiocarp mycenoid, medium large to large. Pileus whitish to orange brown, often with slightly crenulate margin. Lamellae mature deliquescent. Stipe base equal to slightly bulbous. Spore print ochre-yellow to rubiginous. Spores ellipsoidal, thick-walled, germ-pore present or not. Pseudoparaphyses present. In meadows, grassland, woods, on dung, in greenhouses.

Representatives: C. albipes (G. H. OTTH) HAUSKN. (worldwide), C. albipes var. crispa (LONGYEAR) HAUSKN. (North America), C. albipes var. rugata HAUSKN. (Europe), C. romagnesii HAUSKN. & G. MORENO (Europe), C. candida (COOKE & MASSEE) WATLING (Australia), C. fracticeps (PAT. & DEM.) Z. L. YANG (Asia), C. pseudocrispa (HAUSKN.) ARNOLDS (Europe), C. crispella (MURRILL) SINGER (worldwide), C. subcrispa (MURRILL) SINGER (North America), C. zeylanica (PETCH) BOEDIJN (worldwide), C. zeylanica var. marginata HAUSKN. (Africa).

Series Umbonata HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Sporae mediae, lentiformes, hexagono-mitriformes, crasse tunicatae, poro germinativo. Pseudoparaphyses praesentes.

Type species: Conocybe umbonata (MASSEE) WATLING.

Characters: Basidiocarp mycenoid, with bulbous stipe base. Pileus distinctly coloured, brown to dark brown, with crenulate margin. Stipe base distinctly bulbous. Spore print rubiginous. Spores medium large, lentiform, hexagonal-mitriform, thickwalled with distinct germ-pore. Pseudoparaphyses present. In greenhouses.

Single representative: C. umbonata (MASSEE) WATLING (Europe).

Comments: This species, rediscovered only quite recently (HAUSKNECHT & al. 2005), has completely different and much smaller spores as taxa of ser. *Albipes* and differs also macroscopically. The erection of a new series seemed to be adequate.

Subsection Deliquescens HAUSKN. & KRISAI, subsect. nova

= Genus Gastrocybe WATLING 1968, The Michigan Botanist 7: 19

Latin diagnosis: Pileus primo glandiformis vel ellipticus, vel cylindrico-ovoideus vix vel numquam expansus, unctus vel viscidus denique gelatinosus et informis. Stipes aequalis vel subaequalis, vulgo bulbillosus, albido-hyalinus dein pallidus, fragilis, gracilis, saepe elongatus. Lamellae adnatae vulgo ventricosae ad faciem venosae vel intervenosae, ferrugineae. Caro tenuis, albida vel pallida. Cellulae cuticulae pilei pyriformes vel sphaeropedunculatae. Basidia et sterigmata aliquantam crassitunicatae. Basidiosporae fere symmetrica, Cheilocystidia copiosa; pleurocystidia nulla. Type species: *G. lateritia*, by original designation.

Type species: Conocybe deliquescens HAUSKN. & KRISAI, nov. nom.

= Gastrocybe lateritia WATLING 1968, The Michigan Botanist 7: 20.

Characters: Basidiocarp gasteroid. Pileus glandiform to ovoid-cylindrical, hardly expanding, slimy. Stipe very long, with bulbous base. Lamellae anastomosing. Spore print rubiginous. Spores ellipsoidal, with germ-pore. Pleurocystidia absent, pseudo-paraphyses present. Meadows.

Single representative: C. deliquescens HAUSKN. & KRISAI (North America, Europe).

Comments: HALLEN & al. (2003) examined *Gastrocybe*, a genus with gasteroid basidiocarps erected by WATLING (1968), molecular biologically and recognized it as a good member of *Conocybe* sect. *Candidae*. The sliminess of the pileus is caused by a bacterium of the *Chryseobacterium gleum/indologenes*-group, so that HALLEN & al. (2003) were uncertain, if it would eventually be an already known species, disturbed in its growth by the bacteria. However, the combination of the microscopical data do not fit to any other species known of the section.

The epitheton lateritia already exists in the genus *Conocybe* [*Conocybe lateritia* (BATTARRA per FRIES) KÜHNER, Le Genre Galera 1935: 121], and thus cannot be used for the present species. Consequently, we propose the new name *Conocybe deliques-cens*.

Section Giganteae SINGER 1948, Sydowia 2: 36

Latin diagnosis: Sect. *Giganteae* SING. sectio nov. Stipite albo; dermatocystidiis pilisque praesentibus in habitu *Cortinarii*; carpophoris carnosis. Species typica: *Cono*-

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at

Österr. Z. Pilzk. 15 (2006)

cybe detrusa (PECK) SING. comb. nov. (*Cortinarius detrusus* PECK). **Type species:** *Conocybe intrusa* (PECK) SINGER.

Amended description by WATLING (1977: 293):

Pileus convex, expanding or not, thick-fleshy usually fairly brightly coloured in the range salmon buff or ochraceous buff, viscid under favourable conditions; pileipellis a well-developed palisadoderm but lacking distinctive dermatocystidia. Stipe robust, swollen at base, bulbous to clavate, white to yellowish (even primuline yellow), striate throughout or at apex only, with well-developed lecythiform caulocystidia. Cheilocystidia lecythiform. Basidiospores fairly to strongly pigmented, darkening in aqueous alkali solutions, fairly to strongly thick-walled with either a small narrow and indistinct germ-pore or a distinct thinning at the apical end giving a small hyaline patch. Basidio pyriform to clavate, normally 4-spored. Development typically gymnangiocarpic. Basidiocarp developing on base-rich and nitrogen-rich soil in some way connected with man.

Single representative: C. intrusa (PECK) SINGER (presumably worldwide).

Comments: SINGER (1948) states the type species as *Conocybe detrusa* (mis-spelt by SINGER, see SINGER 1950: 134, WATLING & GREGORY 1981: 112).

Section Ochromarasmius (SINGER) HAUSKN. & KRISAI, comb. & stat. nov. Basionym: Conocybe subgenus Ochromarasmius SINGER 1947, Mycologia **39**: 88

Latin diagnosis: 8. *Conocybe* FAYOD, subgenus *Ochromarasmius* SING., subgen. nov. A sectionibus generis *Conocybis* omnibus differt sporis minutis, distincte verrucosis stipiteque tenuissimo, cartilagineo. Ad truncos putridos caespitose in Brasilia.

Type species: Conocybe juruensis (HENN.) SINGER.

Characters: Basidiocarp small to medium large, mycenoid, with slightly bulbous, in one species additionally radicant stipe. Spore print yellow-brown to rubiginous. Spores s. m. mostly verrucous-rough, when smooth s. m., then in scanning electron micro-scope distinctly uneven, verrucous or crested. Pleurocystidia mostly absent, in one species present, Pseudoparaphyses absent. Stipitipellis consisting of lecythiform caulocystidia or a mixture of capilliform, lecythiform, ellipsoidal to lageniform elements. Pileipellis hymeniform, often with lecythiform pileocystidia, in one species heteromorph (elements with rostellate protuberances intermixed). Meadows, forest soil, on rotting wood.

Comments: SINGER (1947) erected a separate subgenus *Ochromarasmius* for species with rough spores. As he limited spore size in the original diagnosis to "smaller than 6 μ m", he later had to amend this diagnosis for the newly-discovered *Conocybe radicata* (SINGER 1962 b). Also the "distinctly warty" spores do not fit within the present concept, as in the meantime taxa were detected with spores seemingly smooth in the light microscope but distinctly uneven-rough in SEM. As in the present concept of the authors and also of other experts of *Bolbitiaceae* the spore surface is not any more that relevant, we suggest downgrading to the category of a section.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Subsection Dumetorae HAUSKN. & KRISAI, subsect. nova

= Stirps *Dumetorum* WATLING 1982, British Fungus Flora Agarics and Boleti **3**: 41, inval.

Latin diagnosis: Cystidia et pileipellis ut in sectione *Ochromarasmius*. Basidiocarpium parvum. Sporae per microscopum luminis visum leves usque subtile punctatae, per microscopum electronicum visum semper distincte asperae. Pleurocystidia et pseudoparaphyses nulla. Stipitipellis consistens elementis lecythiformibus vel mixtura elementis lecythiformibus et non-lecythiformibus.

Type species: Conocybe dumetorum (VELEN.) SVRČEK.

Characters: Basidiocarp mostly small, mycenoid, Stipe base often slightly bulbous, not radicant. Spore print yellow-brown to rubiginous. Spores tuberculous or rough, in one species smooth s. m., but distinctly rough in SEM. Cheilocystidia partly with large capitula. Pleurocystidia and pseudoparaphyses absent. Stipitipellis consisting of lecythiform elements or a mixture of lecythiform, capilliform, clavate-cylindrical or sphaerical elements. Pileipellis often with lecythiform pileocystidia. Forest soil, meadows, on rotting wood.

Representatives: C. dumetorum (VELEN.) SVRČEK (Europe), C. dumetorum var. laricina (KÜHNER) HAUSKN. (Europe), C. dumetorum var. phaeoleiospora HAUSKN. (Europe), C. abjecta (BERK. & BROOME) PEGLER (Asia), C. spinulosa HAUSKN. & KRISAI (Africa), C. stictospora SINGER (South America), C. misionum SINGER (South America, Australia), C. horakii WATLING (Oceania).

Subsection Juruensis HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium parvum. Sporae etiam per microscopum luminis visum distincte tuberculoso-asperae. Pleurocystidia et pseudoparaphyses nulla. Pileipellis hymeniformis, heteromorpha, elementis sphaerico-pedunculatis consistens, sub eis multis rostellatis.

Type species: Conocybe juruensis (HENN.) SINGER.

Characters: Basidiocarp mycenoid, small. Stipe base not radicant. Spore print rubiginous. Spores also s. m. distinctly tuberculous-rough. Cheilocystidia with medium large capitula, Pleurocystidia and pseudoparaphyses absent. Stipitipellis nearly exclusively consisting of lecythiform caulocystidia. Pileipellis hymeniform, heteromorph, consisting of sphaeropedunculate elements, often with rostellate protuberances. On rotting wood.

Single representative: C. juruensis (HENN.) SINGER (South America).

Comments: In the pileipellis of *Conocybe juruensis* there are many sphaerico-pedunculate elements with rostrate protuberances. These were also observed by SINGER (1969) and HORAK (pers. comm.). In the type material, investigated by the first author, the pileipellis elements were very much collapsed, but in another collection from Bolivia made by SINGER such elements could be found. This is a unique character within the genus and in our opinion justifies the erection of a separate subsection.

208

Subsection Pleurocystidiatae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium mycenoideum, stipite radicanti. Sporae valde verrucosae. Cheilo- et pleurocystidiis lecythiformibus.

Type species: Conocybe radicata SINGER.

Characters: Basidiocarp medium large, mycenoid, with strongly bulbous, distinctly radicant stipe base. Spore print rubiginous. Spores also s. m. distinctly tuberculous-rough. Pleurocystidia present, abundant, lecythiform, similar to cheilocystidia. Pseudoparaphyses absent. Stipitipellis consisting of lecythiform, capilliform and ellipsoidal-cylindrical elements. Pileipellis hymeniform, with lecythiform pileocystidia. On rotting wood.

Single representative: C. radicata SINGER (South America).

Comments: Pleurocystidia are not mentioned in the original description (SINGER 1953). The first author could examine the holotype (LIL) as well as the isotype (MICH) and found abundant pleurocystidia in both of them, not only near the lamellar edge, but also scattered over the whole area of the lamellae. Thus *Conocybe radicata* is the only representative of the genus with pleurocystidia.

Section Singerella WATLING 1979, Sydowia Beiheft 8: 408

Conocybe subgenus Conocybe Section Singerella WATLING sectio nova.

- ? Agaricus ser. Dermini sect. Acetabularia BERK. in J. Linn. Soc. (Bot.) London 18: 389, 1881
- = ? Acetabularia (BERK.) MASSEE, Brit. Fungi Fl. 2: 232, 1893 (non Acetabularia LAOUROUX 1912 (Dasycladaceae, Chlorophyta)
- =? Cyphellopus FAYOD, Ann. Sci. Nat. (Bot.) VII 9: 365, 1889
- = Conocybe subgenus Singerella (WATLING) BON 1991, Doc. Mycol. 21/83: 38

Latin diagnosis: Pileus hygrophanus, siccus, subinde paulum humidus, laevis vel subtiliter pubescens, propter pilocystidia. Annulus absens. Stipes cylindricus, laevis vel subtiliter pubescens. Pleurocystidia absentia; Cheilocystidia lecythiformia.

Type species: Conocybe corneri WATLING.

Characters: Basidiocarp mycenoid, medium large to large. Veil mostly a volva or a volva-like zone, rarely as fringes on the pileus margin and as girdle-like zones around the stipe. Stipe not radicant. Spore print yellow-brown, rubiginous or dark brown. Pseudoparaphyses absent or insignificant. Stipitipellis only consisting of non-lecythiform elements or a mixture of lecythiform and non-lecythiform caulocystidia. Dung, rotting plant and wood debris, fertilised soil, also in tropical deciduous woods.

Comments: For species of *Conocybe* with veil (fringes on the pileus margin or volvalike zone on the stipe) WATLING (1979) segregated the section *Singerella*. He critically studied all hitherto known species with veil or volva, respectively. He also discusses the eventual use of older names for the section (WATLING 1979 – see also above). A classification in the higher category of a subgenus, as suggested by BON (1991), does not seem to be appropriate in the present concept.

210

A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

Subsection Subvelatae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium cum velo, margo pilei velo fimbriatus, sine volva; base stipitis non radicanti. Pseudoparaphyses absentes vel non significantes, stipitipellis solum caulocystidiis non lecythiformibus consistens.

Type species: Conocybe subvelata SINGER.

Characters: Basidiocarp mycenoid; veil present as fine fringes on the pileus margin, stipe base volva or volva-like zone absent, not radicant. Spore print yellow-brown. Pseudoparaphyses absent or insignificant. Stipitipellis only consisting of clavate, lageniform and capilliform elements, lecythiform caulocystidia absent. Spores smooth, with germ-pore.

Single representative: C. subvelata SINGER (South America).

Subsection Vaginatae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium volva membranacea vel zona similis volvae, interdum etiam margo pilei velo fimbriatus; base stipitis non radicanti. Pseudoparaphyses absentes vel non significantes. Stipitipellis solum vel praecipue consistens caulocystidiis non lecythiformibus.

Type species: Conocybe vaginata WATLING.

Characters: Basidiocarp mycenoid; veil present as membranous volva or as volva-like zone, additionally sometimes girdle-like zones around the stipe and fringes on the pileus margin; stipe base not radicant. Spore print yellow-brown, rubiginous to dark brown. Pseudoparaphyses absent or insignificant. Stipitipellis consisting of capilliform, ellipsoidal, sphaerical or lageniform elements, lecythiform caulocystidia present in one species near the stipe apex. Spores smooth, thick-walled, often limoniform, lentiform to subhexagonal, with germ-pore. Dung, rotting plant and wood debris, fertilised soil, also in tropical deciduous woods.

Series Locellina HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Characters ut in subsectione *Vaginatae*, sed basidiocarpium zona similis volvae vel volva infirma non membranacea, interdum etiam margo pilei velo fimbriatus. Pulvis sporarum rubiginosus vel brunneus.

Type species: Conocybe locellina (MURRILL) WATLING.

Characters: Basidiocarp with *Mycena*-habit; veil present as vanishing volva-like zone or (when young) as weak, non-membranous volva, additionally sometimes with veilar fringes at the pileus margin or girdle-like zones around the stipe. Stipe base not radicant. Spore print rubiginous to dark brown. Pseudoparaphyses absent. Stipitipellis only consisting of capilliform, ellipsoidal, sphaerical to lageniform caulocystidia, lecythiform elements absent. Spores smooth, thick-walled, with germ-pore. Dung, rotting plant or wood debris.

©Österreichische Mykologische Gesellschaft, Austria, download unter www.biologiezentrum.at Österr. Z. Pilzk. 15 (2006) 211

Representatives: *C. locellina* (MURRILL) SINGER (North America), *C. hornana* SINGER & HAUSKN. (Europe), *C. hornana* var. *subcylindrospora* HAUSKN., VAURAS, KYTÖV. & OHENOJA (Europe).

Series Corneri HAUSKN. & KRISAI, ser. nov.

Latin diagnosis: Characters ut in subsectione *Vaginatae*, sed basidiocarpium volva membranacea persistente, stipes sine velo fimbriato. Pulvis sporarum luteo-brunneus usque rubiginosus.

Type species: Conocybe corneri WATLING.

Characters: Basidiocarp mycenoid; veil a membranous, persisting volva, veilar fringes at the pileus margin or girdle-like zones around the stipe absent. Stipe base not radicant. Spore print yellow-brown to rubiginous. Pseudoparaphyses absent. Stipitipellis consisting of capilliform, ellipsoidal, sphaerical to lageniform elements, in one species also with some lecythiform caulocystidia near the stipe apex. Spores smooth, thick-walled, often limoniform, lentiform to subhexagonal, with germ-pore. Dung, rotting plant or wood debris, forest soil.

Representatives: C. corneri WATLING (Asia, Ozeania), C. vaginata WATLING (Asia, Oceania), C. volvata K. A. THOMAS, HAUSKN. & MANIM. (Asia), C. volviornata E. HORAK, HAUSKN. & DESJARDIN (Asia), C. discorosea E. HORAK, HAUSKN. & DESJARDIN (Asia), C. anthurii WATLING & HAUSKN. (Africa).

Subsection Vinaceobrunneae HAUSKN. & KRISAI, subsect. nova

Latin diagnosis: Basidiocarpium tenuiter volva infirma furfuracea, stipes sine velo fimbriato, non radicans. Pseudoparaphyses absentes. Stipitipellis consistens elementis lecythiformibus et non lecythiformibus.

Type species: Conocybe vinaceobrunnea HAUSKN.

Characters: Basidiocarp mycenoid; volva weakly developed, young present as narrow scaly-mealy girdle, soon vanishing; veilar fringes at the pileus margin or girdle-like zones around the stipe absent; Stipe not radicant. Spore print rubiginous. Pseudopara-physes absent. Stipitipellis consisting of lecythiform, ellipsoidal, lageniform to capilliform caulocystidia. Spores smooth, with germ-pore. Dung, fertilised soil.

Single representative: C. vinaceobrunnea HAUSKN. (Oceania).

Comments: Conocybe vinaceobrunnea differs from all other species of the section in two characters: the veil is narrow and scaly-mealy, and the stipitipellis conforms to that of sect. *Mixtae* (HORAK & HAUSKNECHT 2002).

References

ARNOLDS, E., 2005: *Conocybe*. – In NOORDELOOS, M. E., KUYPER, T. W., VELLINGA, E. C., (Eds.): Flora Neerlandica 6. – Boca Raton: Taylor & Francis.

BON, M., 1990: Combinaisons nouvelles et validations. - Doc. Mycol. 20/78: 37-40.

— 1991: Espèces "galéro-naucoroïdes": Stat. et comb. nov. (Clé en préparation D. M. no 84). – Doc. Mycol. 21/83: 37-39. A. HAUSKNECHT & I. KRISAI-GREILHUBER: Infrageneric division of Conocybe

- DONK, M. A., 1949: Nomina generica conservanda and confusa for *Basidiomycetes* (Fungi). Bull. Bot. Gard. Buitenzorg **17**: 155-197.
- FAYOD, V., 1889: Prodrome d'une histoire naturelle des Agaricinées. Ann. Sci. Nat. Bot. VII, 9: 181-411.
- HALLEN, H. E., WATLING, R., ADAMS, G. C., 2003: Taxonomy and toxicity of *Conocybe lactea* and related species. – Mycol. Res. 107: 969-979.
- HAUSKNECHT, A., 2003: Beiträge zur Kenntnis der Bolbitiaceae 9. Conocybe Sekt. Mixtae. Österr. Z. Pilzk. 12: 41-83.
- VAURAS, J., KYTÖVUORI, I., OHENOJA, E., 2005: Die Gattung Conocybe in Finnland. Karstenia 45: 1-32.
- HORAK, E., 1968: Synopsis generum *Agaricalium* Die Gattungstypen der *Agaricales*. Beiträge zur Kryptogamenflora der Schweiz 13. Wabern, Bern: Büchler.
- HAUSKNECHT, A., 2002: Notes on extra-European taxa of *Bolbitiaceae (Agaricales, Basidiomy-cota)*. Österr. Z. Pilzk. 11: 213-264.
- KÜHNER, R., 1935: Le genre Galera. Paris: Lechevalier.
- MONCALVO, J.-M., VILGALYS, R., REDHEAD, S. A., JOHNSON, J. E., JAMES, T. Y., AIME, M. C., HOFSTETTER, V., VERDUIN, S. J. W., LASSON, E., BARONI, T. J., THORN, R. G., JACOBSSON, S., CLÉMENÇON, H., MILLER, JR., O. K., 2002: One hundred and seventeen clades of agarics. – Molec. Phylogenet. Evol. 23: 357-400.
- SINGER, R., 1947: New genera of Fungi III. Mycologia 39: 77-89.
- 1948: Diagnoses Fungorum novorum Agaricalium. Sydowia 2: 26-42.
- 1950: New and interesting species of Basidiomycetes III. Sydowia 4: 130-157.
- -, "1949" 1951: The Agaricales in modern taxonomy, 1st edn. Lilloa 22: 1-831.
- 1953: Quelques Agarics nouveaux de l'Argentine. Rev. Mycol. Paris 18: 3-23.
- 1958: Monographs of South American basidiomycetes I. Lloydia 21: 159-299.
- 1962 a: Diagnoses Fungorum novorum Agaricalium II. Sydowia 15: 45-83.
- 1962 b: The Agaricales in modern taxonomy, 2nd edn. Weinheim: Cramer.
- 1969: Mycoflora Australis. Nova Hedwigia 29: 1-405.
- 1986: The Agaricales in modern taxonomy, 4th edn. Koenigstein: Koeltz.
- SMITH, A. H., 1941: Studies of North American Agarics 1. Contrib. Univ. Mich. Herbarium 5: 1-73.
- WALTHER, G., GARNICA, S., WEISS, M., 2005: The systematic relevance of conidiogenesis modes in the gilled Agaricales. – Mycol. Res. 109: 525-544.
- WATLING, R., 1968: Observations on the *Bolbitiaceae*. IV. A new genus of gastromycetoid fungi. The Michigan Botanist 7: 19-24.
- 1974: Flore illustrée des champignons d'Afrique Centrale 3. *Bolbitiaceae*. Meise: Jardin Botanique de Belgique.
- 1975: Observations on the *Bolbitiaceae* 12: The affinities of two anomalous species. Notes Roy. Bot. Gard. Edinburgh 34: 245-251.
- 1976 a: Observations on the *Bolbitiaceae*, 13 & 14. Kew Bull. **31**: 587-594.
- 1976 b: Observations on the *Bolbitiaceae* XV. The taxonomic position of those species of *Conocybe* possessing ornamented basidiospores. Rev. Mycol. 40: 31-37.
- 1977: Observations on the *Bolbitiaceae* 16: *Conocybe* sect. *Giganteae*. Notes Roy. Bot. Gard. Edinburgh 35: 281-295.
- 1979: Observations on the *Bolbitiaceae* XVII. Volvate species of *Conocybe*. Sydowia, Beih. 8: 401-415.
- 1982: Bolbitiaceae: British Fungus Flora Agarics and Boleti 3: Agrocybe, Bolbitius & Conocybe. Edinburgh: Royal Botanic Garden.
- 1986: Observations on the *Bolbitiaceae*. 28. The *Conocybe pubescens* (C. GILLET) KÜHNER complex. *Galera neoantipus* and its various interpretations. Bol. Soc. Micol. Madrid 11: 91-96.
- 1994: Observations on Malaysian *Bolbitiaceae* with records from Solomon Islands. Garden's Bull. Singapore 45: 359-381.
- GREGORY, N., 1981: Census catalogue of world members of the *Bolbitiaceae*. Biblioth. Mycol. 82.

212

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Österreichische Zeitschrift für Pilzkunde

Jahr/Year: 2006

Band/Volume: 15

Autor(en)/Author(s): Hausknecht Anton, Krisai-Greilhuber Irmgard

Artikel/Article: Infrageneric division of the genus Conocybe - a classical approach. 187-212