

New data on the rare *Entoloma moserianum*

ELISEO BATTISTIN
Natural History Museum "Dr. Dal Lago"
Corso Italia, 63
I-36078 Valdagno, Italy
Email: eliseo_battistin@yahoo.it

NORBERTO RIGHETTO
Natural History Museum "Dr. Dal Lago"
Corso Italia, 63
I-36078 Valdagno, Italy
Email: righetto@interfree.it

Accepted 9. 10. 2009

Key words: *Entolomatales*, *Entoloma*, *E. moserianum*. – Stipitipellis, ecology, macrochemical reactions, new record, rare species. – Mycoflora of Italy.

Abstract: *Entoloma moserianum* is a very rare species described by the Dutch mycologist M. E. NOORDELOOS in 1983. It has been found in a few European countries like The Netherlands, Germany, Austria, and Italy. A recent finding in northern Italy has allowed us to analyse a lot of basidiomata and to notice some further characters not reported in the original diagnosis regarding the reactivity to several chemical compounds, microscopical and ecological features. Finally, a phenetic analysis of all the 11 species belonging to subg. *Entoloma* sect. *Entoloma*, based on 18 macro-, microscopical and ecological features and the principles of numerical taxonomy, has been carried out. Astonishing overall phenotypic relationships have been disclosed.

Zusammenfassung: *Entoloma moserianum* ist eine sehr seltene Art, die von dem niederländischen Mykologen M. E. NOORDELOOS im Jahr 1983 beschrieben wurde. Sie wurde bisher in einigen europäischen Ländern wie Italien, Österreich, Deutschland und den Niederlanden gefunden. Eine neue und reichliche Aufsammlung aus Norditalien erlaubte die Beobachtung von Merkmalen, die in der originalen Diagnose nicht beschrieben wurden, wie makrochemische Reaktionen, mikroskopische Merkmale und ökologische Angaben. Schließlich wurde eine phenotypische Analyse aller 11 Arten von subg. *Entoloma* sect. *Entoloma*, beruhend auf 18 mikro-, makroskopischen und ökologischen Merkmalen und auf den Prinzipien der numerischen Taxonomie, durchgeführt, die bemerkenswerte phänotypische Übereinstimmungen aufzeigte.

Riassunto: *Entoloma moserianum* è una specie piuttosto rara descritta dal micologo olandese M. E. NOORDELOOS nel 1983. Per quanto a nostra conoscenza è stata ritrovata in alcuni paesi europei quali l'Olanda, la Germania, l'Austria e l'Italia. Una recente e abbondante raccolta effettuata in Nord Italia ci ha permesso di notare delle caratteristiche non segnalate nella diagnosi originale concernenti la reattività a diversi composti chimici, la presenza di taluni elementi microscopici e annotazioni sull'ecologia. Infine viene riportata un'analisi di raggruppamento eseguita su tutte le 11 specie del sottogenere *Entoloma* sezione *Entoloma* basata su 18 caratteri macro-, microscopici ed ecologici, nonché sui principi della tassonomia numerica. Sorprendenti relazioni fenotipiche complessive vengono messe in evidenza.

Mycology has been undergoing an outstanding progress due to the application of molecular biology techniques: We look at these scientific developments with great enthusiasm.

siasm, nevertheless, we think many aspects concerning the morphological taxonomy still need further investigations. Morphological studies are still necessary, especially on rare or uncommon taxa like *E. moserianum*.

Table 1. Data matrix.

	Diameter of pileus (min.)	Diameter of pileus (MAX)	Length of stipe (min.)	Length of stipe (MAX.)	Diameter of stipe (min.)	Diameter of stipe (MAX)	Length of spore (min.)	Length of spore (MAX)	Breadth of the spores (min.)	Breadth of the spores (MAX.)	Pileipellis: transition to a trichoderm	Presence of cystidia	Yellowish lamellae in young specimens	Smell	Base of stipe differently coloured	Habitat: grasslands	Habitat: woods	
<i>Entoloma prunuloides</i>	20	70	30	80	3	12	6,5	9	6,5	8	1	0	c	0	0	0	1	0
<i>Entoloma sinuatum</i>	30	250	40	150	5	35	8	11	7	9,5	1	0	0	1	0	0	0	1
<i>Entoloma blixamii</i>	30	80	35	70	10	35	7	9	6,5	8	1	0	0	0	0	0	1	0
<i>Entoloma nitidum</i>	20	50	30	100	2	7	6,5	9	5,5	8	1	0	0	0	0	0	0	1
<i>Entoloma moserianum</i>	20	120	40	120	5	14	9	12	8	9,5	1	0	1	0	0	0	1	1
<i>Entoloma rubellum</i>	25	80	50	90	7	9	6,5	8	5,5	8	1	0	0	0	0	0	0	1
<i>Entoloma viridans</i>	30	40	40	60	6	9	7,5	9,5	6,5	8	1	0	0	0	0	0	0	1
<i>Entooma luteobasis</i>	25	50	60	80	8	14	6	8	6	7	1	0	0	0	0	1	0	1
<i>Entoloma olidum</i>	35	60	35	60	6	10	8,5	11	7	9	1	1	0	0	1	0	0	1
<i>Entoloma myochroum</i>	50	100	80	95	10	18	7	10	6,5	9	1	0	0	0	0	0	0	1
<i>Entoloma alcedicolor</i>	22	25	30	35	3	4	7	8,5	6	8	1	1	0	0	1	0	0	1

Material and methods


Photographs of the basidiomata were taken in the field by an Olympus FE-340 digital camera. Micro-photographs were taken by a Nikon Coolpix 5400 digital camera and a Nikon Eclipse E-200 LM. Fragments of fresh material for microscopical analysis were stained with Congo red (Titolchimica, Rovigo, Italy) or mounted directly in water, while dried material was previously hydrated in 5% KOH.

We followed the taxonomic arrangement of NOORDELOOS (1992, 2004, 2009). Abbreviations of the authors of fungal names and acronyms of herbaria follow KIRK (2008) and HOLMGREN & HOLMGREN (1998).


For the phenetic analysis two programs have been executed under Windows Vista: Microsoft Excel 2003 and ClustanGraphic (WISHART 2006).

The principles of numerical taxonomy we followed are clearly explained in ALDENDERFER & BLASHFIELD (1984), FIELDING (2007), ROMESBURG (2004), SNEATH & SOKAL (1973), SOKAL & SNEATH (1963). This kind of numerical analysis is composed of five consecutive steps:

- creation of a data matrix (Table 1) containing mixed data types (continuous and binary values), i.e., a table of n rows and m columns in Excel listing the species (rows) and the variables or attributes of those species (columns) obtained from NOORDELOOS' descriptions. We did not take into account the colour of the basidiomata because the software is not able to codify nominal variables.
- standardization of the continuous variables like the size of pileus, stipe, spore.
- calculus of a similarity matrix by Gower's similarity coefficient.



1



2



3

Figs. 1, 2. Macrochemical reactions on pileus surface. – Phot. ELISEO BATTISTIN. – Fig. 3. *Entoloma moserianum* in situ. – Phot. NORBERTO RIGHETTO.

– graphical representation by phenograms of the clusters' linkages got by several agglomerative hierarchical clustering methods like average linkage also known as UPGMA, probably the most frequently used clustering strategy (SNEATH & SOKAL 1973), weighted average linkage method and complete linkage method.

– estimate of the goodness of the results through the calculus of the cophenetic correlation coefficient (r), which assesses how much the phenogram faithfully portrays the similarities in the resemblance matrix, and the determination of the optimal number of clusters by a statistical method, a bootstrap validation test without replacement and 500 random trials.

We tested the following chemical compounds (Titolchimica, Rovigo, Italy) on the pileus surface of fresh basidiomata: potassium hydroxide 30%, sodium hydroxide 30%, nitric acid 65%, hydrochloric acid 10%, ferrous sulphate (powder), α -naphthol 4%, aniline, potassium dichromate 10%, ferrous chloride, toluene 2% and guaiac.

Collection examined: Italy: Veneto, Vicenza, Spagnago di Cornedo Vicentino, terricolous under deciduous trees, 23. 7. 2009, leg. ELISEO BATTISTIN & NORBERTO RIGHETTO (MCVE 24415).

Results and observations

One reagent, i.e., nitric acid produced a clearly positive, lemon-yellow macrochemical reaction after about 15 min from its application on fresh specimens; the other were inert on the whole (Figs. 1, 2).

In case of doubtful classification this simple macrochemical reaction can contribute significantly to get a certain identification of the species.

With regard to its habitat (Fig. 3) we collected it on a humus-rich calcareous soil in a definitely heliophilous broadleaved wood – it occurs in deciduous forest (*Quercus*) on relatively open, sunlit places according to NOORDELOOS (1992) – located at about 200 m s. m. consisting of *Ulmus minor* MILL., *Corylus avellana* L., *Acer campestre* L. and *Robinia pseudacacia* L. Many tufts of *Ruscus aculeatus* L. were all around the mushrooms.

According to our experience (BATTISTIN & RIGHETTO 1998) it appears in April, May, June, July and October, so it makes sense to reckon that it is neither a strictly thermophilous nor cryophilous taxon.

By handling some basidiomata we noticed under lens that the apex of many stipes was minutely pruinose: a microscopical check revealed bundles of hyphae perpendicular or almost so to the longitudinal axis of the stipe. The terminal elements are cylindrical and have an obtuse or subcapitulate apex, and are morphologically almost identical to cheilocystidia (Fig. 4). Their size is 35-64 \times 5-9 μm . This observation should be the first report of such a feature.

Finally, we performed an agglomerative hierarchical clustering analysis of the 11 species included in subg. *Entoloma* sect. *Entoloma* according to NOORDELOOS' taxonomy.

It is necessary to state first that cluster analysis is a type of multivariate statistical analysis whose aim is to divide a set of objects (species in this case) into relatively homogeneous subsets based on inter-object similarities (KACHIGAN 1991).

The outcome suggests that this section is phenotypically rather homogeneous: two clustering methods, average linkage ($r = 0.87$), the reference method, and weighted average linkage ($r = 0.83$) show that all the species belong to one cluster (Figs. 5, 6), while the third, i.e., complete linkage ($r = 0.78$) recognizes four clusters (Fig. 7). It is worth noting another element of homogeneity: The phenotypic relationship among species,

i.e., the topology of the phenograms, is identical (Figs. 5-7) in all of the three phenograms. It would be very interesting to acquire molecular data and compare them to the phenotypical ones.




Fig. 4. A cluster of terminal hyphae of the stipitipellis perpendicular to the longitudinal axis of the stipe. – Bar: 9 µm. – Phot. ELISEO BATTISTIN.

In any event, aside from the colour of the basidiomata, unexpected overall phenotypical resemblances among species were disclosed like those between *Entoloma nitidum* QUÉL. and *E. rubellum* (SCOP.) GILLET, *E. prunuloides* (FR.) QUÉL. and *E. bloxamii* (BERK. & BROOME) SACC., *E. oolidum* NOORDEL. & T. BORGEN and *E. alcedicolor* ARNOLDS & NOORDEL. or *E. moserianum* NOORDEL. itself and *E. sinuatum* (BULL.) P. KUMM.

With regard to *E. moserianum* and *E. sinuatum* they seem to be a little bit isolated within the phenograms and by comparing them to NOORDELOOS' taxonomy that couple of species could perhaps represent the subsect. *Sinuata* deprived of *Entoloma luteobasis* and *E. oolidum*. In our opinion it is a little difference which does not affect the substantial agreement between NOORDELOOS' taxonomic arrangement and the outcome of the numerical taxonomy.

We are indebted to Prof. GABRIELLA ZANROSSO (Valdagno, Italy) for improving the English text and Ing. ZILIO ZORDAN (Valdagno, Italy) for writing the Zusammenfassung.

References

- ALDENDERFER, M. S., BLASHFIELD, R. K., 1984: Cluster analysis. – Sage University Paper series on Quantitative Applications in the Social Science **44**. – Newbury Park: Sage Publications.

- BATTISTIN, E., RIGHETTO, N., 1998: Un *Entoloma* interessante: *E. moserianum* NOORDEL. – Riv. Mic. col. **41**: 43-46.
- BREITENBACH, J., KRÄNZLIN, F., 1995: Champignons de Suisse 4. – Luzern: Mykologia.
- FIELDING, A. H., 2007: Cluster and classification techniques for the biosciences. – New York: Cambridge University Press.
- HOLMGREN, P. K., HOLMGREN, N. H., 1998 [continuously updated]: Index Herbariorum: A global directory of public herbaria and associated staff. – New York Botanical Garden's Virtual Herbarium. [<http://sweetgum.nybg.org/ih/>].
- KACHIGAN, S. K., 1991: Multivariate statistical analysis. – New York: Radius Press.
- KIRK, P. M., 2008: Authors of fungal names. – Index Fungorum, the CABI Bioscience, CBS and Landcare Research database of fungal names. [www.indexfungorum.org, visited on 31. 7. 2009].
- NOORDELOOS, M. E., 1992: *Entoloma* s. l. – Saronno: Giovanna Biella.
- 2004: *Entoloma* s. l. Supplemento. – Alassio: Massimo Candusso.
- 2009: Machiels *Entoloma* pages. – [<http://www.entoloma.nl/html/entolomaeng.html>, visited on 4. 9. 2009]
- ROMESBURG, H. C., 2004: Cluster analysis for researchers. – Morrisville: Lulu Press.
- SNEATH, P. H. A., SOKAL, R. R., 1973: Numerical taxonomy. – San Francisco: Freeman.
- SOKAL, R. R., SNEATH, P. H. A., 1963: Principles of numerical taxonomy. – San Francisco: Freeman.
- WISHART, D., 2006: Clustan Graphics primer. A guide to cluster analysis. – Edinburgh: Clustan Limited.






Fig. 5. Average linkage phenogram. Cophenetic correlation coefficient $r = 0.87$. Clusters: 1 (all the 11 species included in section *Entoloma*).



⑦




Fig. 6. Weighted average linkage phenogram. Cophenetic correlation coefficient $r = 0.83$. Clusters: 1 (all the 11 species included in section *Entoloma*). – Fig. 7. Complete linkage phenogram. Cophenetic correlation coefficient $r = 0.78$. Clusters: 4 (*E. prunuloides* and *E. bloxamii*), (*E. nitidum*, *E. rubellum*, *E. viridans*, *E. luteobasis* and *E. myochroum*), (*E. olidum* and *E. alcedicolor*), (*E. sinuatum* and *E. moserianum*).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Österreichische Zeitschrift für Pilzkunde](#)

Jahr/Year: 2009

Band/Volume: [18](#)

Autor(en)/Author(s): Battistin Eliseo, Righetto Norberto

Artikel/Article: [New data on the rare Entoloma moserianum. 161-167](#)