

Chemistry of the earthy odour of basidiomata of *Cortinarius hinnuleus* (*Basidiomycota*, *Agaricales*)

NORBERT ARNOLD¹

GÖTZ PALFNER²

CHRISTINE KUHN¹

JÜRGEN SCHMIDT¹

Email: narnold@ipb-halle.de

¹Leibniz Institute of Plant Biochemistry

Department of Bioorganic Chemistry

Weinberg 3

06120 Halle (Saale), Germany

²Universidad de Concepción

Facultad de Ciencias Naturales y Oceanográficas

Departamento de Botánica

Casilla 160-C

Concepción, VIII Región, Chile

Accepted 13. January 2016

Key words: *Cortinarius hinnuleus*. – Geosmin, β -caryophyllene, β -barbatene, 1-octen-3-ol, volatile compounds.

Abstract: *Cortinarius hinnuleus* (Earthy Webcap), a common mycorrhizal mushroom in Central Europe, is characterized by a mouldy earthy odour. The relevant volatile compounds were detected by gas chromatography-mass spectrometry using headspace-solid phase microextraction technology and identified as geosmin, β -caryophyllene and β -barbatene together with the C₈-volatiles 1-octen-3-ol, 1-octen-3-one, octan-3-ol, octan-3-one, and 2-octen-1-ol.

Zusammenfassung: *Cortinarius hinnuleus* (Erdigriechender Gürtelfuß), ein in Mitteleuropa häufiger Mykorrhiza-Pilz, zeichnet sich durch schimmelig-erdigen Geruch aus. Die relevanten flüchtigen Verbindungen wurden durch Gaschromatographie-Massenspektrometrie (GC-MS) unter Verwendung der „headspace-solid phase microextraction“ (HS-SPME)-Technologie ermittelt und als Geosmin, β -Caryophyllen und β -Barbaten zusammen mit den flüchtigen C₈-Verbindungen 1-Octen-3-ol, 1-Octen-3-on, Octan-3-ol, Octan-3-on und 2-Octen-1-ol identifiziert.

Cortinarius (PERS.) GRAY (*Cortinariaceae*, *Agaricales*, *Basidiomycota*) is the most diverse genus of ectomycorrhizal fungi with about 5000 epithets listed in the database www.indexfungorum.org. *Cortinarius hinnuleus* FR. (Earthy Webcap) is especially frequent in Central Europe, where it occurs in mixed and deciduous forests and parks on rich, acidic to calcareous soil, mainly associated with *Quercus*, *Fagus*, *Tilia*, *Carpinus*, but also near *Betula*, *Populus* and *Corylus* (GMINDER 2010, NISKANEN & KYTÖVUORI 2012). However, *C. hinnuleus* is not cultivable on artificial media so far. The basidiomata appear between late summer and late autumn and the species is common in nemoral to hemiboreal zones in Europe, therefore less common in northern Europe (BRANDRUD & al. 1990, GMINDER 2010, NISKANEN & KYTÖVUORI 2012). This is one of the classical European *Cortinarii* developing reddish brown to ochra-

ceous brown carpophores, defined by a universal veil forming a white ring and incomplete girdles on the stem, distant pale yellow to red brown gills and, most conspicuously, by a pungent, unpleasant earthy odour. We analysed its chemical composition from collected basidiomata using headspace-solid phase microextraction gas chromatography-mass spectrometry (HS-SPME/GC-MS) for the identification of the compounds responsible for the strong earthy aroma of *C. hinnuleus*.

Materials and methods

Experimental: The GC-EIMS measurements were performed on a QP-2010 Ultra (Shimadzu Corporation, Kyoto, Japan) by using the following conditions: electron energy 70 eV, detected mass range m/z 40–500; source temperature 200 °C; column: ZB-5MS (Phenomenex, 30 m x 0.25 mm, 0.25 μm film thickness), interface temperature 270 °C, carrier gas Helium, column flow 1.02 ml/min, constant flow mode, column temperature program: 40 °C for 2 min, then raised to 260 °C at a rate of 10 °C min^{-1} and then held on 260 °C for 5 min.

The following operating conditions for the HS-SPME (headspace-solid phase microextraction) were used: SPME fibre: 50/30 μm divinylbenzene(DVB)/carboxen/polydimethylsiloxane(PDMS), adsorption temperature: 65 °C, adsorption time: 5 min; split injection (split ratio 1:5), injector port temperature: 250 °C, desorption time: 2 min. The volatiles were identified by comparing their EI mass spectra with the databases NIST 11 (National Institute of Standard and Technology), FFNSC 2 (Shimadzu) and MassFinder 4.0 (Dr. Hochmuth Scientific Consulting, Hamburg, Germany). The relative abundances of components **1-10** were calculated from the peak areas of the total ion chromatogram (TIC).

Fungal material: *Cortinarius hinnuleus*: Germany, Saxony-Anhalt, near Freyburg (Unstrut), 18. October 2012, leg. & det. N. Arnold & G. Palfner, near *Quercus robur*, voucher deposited at the Leibniz Institute of Plant Biochemistry (Halle/Saale).














Fig. 1. Gas chromatogram of main volatile compounds of *Cortinarius hinnuleus*.

Results and discussion

Headspace-solid phase microextraction (HS-SPME) analysis of volatiles emitted by the cap of *C. hinnuleus* revealed several aliphatic alcohols and ketones, viz.: hexanal (**1**), a mixture of the C₈-volatiles 1-octen-3-one (**2**), 1-octen-3-ol (**3**), octan-3-one (**4**), octan-3-ol (**5**), 2-octen-1-ol (**6**) and the monoterpene linalool (**7**) (Tab. 1, Fig. 1). Besides these compounds, geosmin (**8**, 3% of identified compounds, RT 14.67 min) as well as the sesquiterpenes β -caryophyllene (**9**, 0.6 % of identified compounds, RT 14.69 min) and β -barbatene (**10**, 0.3% of identified compounds, RT 15.17 min) could be identified.

Tab. 1. GC data of main volatile compounds of *Cortinarius hinnuleus*, obtained by GC-MS measurements.

No.	RT (min)	Structure	Compound	Relative Composition (%)
1	4.55		Hexanal	0.2
2	7.77		1-Octen-3-one	1.2
3	7.80		1-Octen-3-ol	35.7
4	7.91		Octan-3-one	49.4
5	8.10		Octan-3-ol	5.4
6	9.28		2-Octen-1-ol	3.9
7	9.81		Linalool	0.2
8	14.67		Geosmin	3.0
9	14.69		β -Caryophyllene	0.6
10	15.17		β -Barbatene	0.3
			Others (not identified)	0.1

Compounds **1–8** represent 96 % of total volatiles in *C. hinnuleus*. In particular, the volatiles **2–6** are responsible for the typical mushroom smell of fungi and have been known for some time (e.g. BELTRAN-GARCÍA & al. 1997, CHO & al. 2008, COMB & al. 2006, MALHEIRO & al. 2013). The occurrence of the alkyl aldehyde hexanal (**1**, 0.2 % of identified compounds, RT 4.55 min) could be interpreted as a degradation product by oxidation of fatty acids or a biosynthetic side product during formation of 1-octen-3-ol (**3**). Hexanal (**1**) has been shown to inhibit spore germination of different fungi and to act against microorganisms (SPITELLER 2002). The eight-carbon volatiles **2–6** seem to possess two biological functions. On one side, they play an important role in attracting mycophilous diptera, which are likely to disseminate spores (CHIRON & MICHELOT 2005). On the other side, it has been demonstrated that 1-octen-3-ol (**3**) inhibits germ tube formation in spores of parasitic fungi like *Penicillium paneum* and exhib-

its bactericidal effects at the same time (COMB & al. 2006), protecting the fungal basidioma against ubiquitous moulds and bacteria in its natural habitat.

The terpene linalool (7) together with other constituents could be partially related to the characteristic odours of some fungi (BREHER & al. 1997, MALHEIRO & al. 2013). Linalool (7) shows several biological activities, among them also bactericidal and fungicidal effects (PATNAIK & al. 1997).

The natural bicyclic sesquiterpene β -caryophyllene (9), occurs in more than 50 % of all plant families, representing a very common constituent of essential oils of various aromatic plants such as *Syzygium aromaticum*, *Cannabis sativa* and *Rosmarinus officinalis* (KNUDSEN & al. 2006). Recently, β -caryophyllene could also be identified as VOC (Volatile Organic Compounds) from a fungal strain of *Talaromyces wortmannii* (YAMAGIWA & al. 2011) and *Cladosporium cladosporioides* (DIBY & PARK 2013). We can therefore conclude that β -caryophyllene (9) could also be a naturally occurring constituent of the basidioma of *C. hinnuleus*. Derivatives of β -caryophyllene (9) have so far been reported from the culture broth of the basidiomycetes *Hebeloma longicaudum* (WICHLACZ & al. 1999) and *Hypholoma fasciculare* (SHIONO & al. 2004).

The sesquiterpene β -barbatene (10) was so far only known as VOC of *Fomitopsis pinicola* (FÄLDT & al. 1999) and could be also detected after inoculation experiments of *Trametes versicolor* on *Fagus sylvatica* (THAKEOW & al. 2006). The authors suggested that two beetle species might use this compound to discriminate the host odours of chopped *F. pinicola* and *Fomes fomentarius* (FÄLDT & al. 1999).

The unpleasant, musty, earthy odour of *C. hinnuleus* is caused by geosmin (8) and was previously reported as the key compound also responsible for the musty-earthly aroma of *Cortinarius herculeus*, *Cystoderma amianthinum* and *Cy. carcharias* (BREHER & al. 1999). Geosmin (8), normally accompanied by 2-methylisoborneol, is common in nature, being produced by blue-green algae (Cyanobacteria), some vascular plants, mosses and protozoan symbionts (WATSON 2003), but is mainly synthesized by terrestrial actinomycetes (genus *Streptomyces*) and some filamentous fungi such as *Penicillium* or *Aspergillus* spp., giving soil its characteristic smell (BUTTERY & GARBALDI 1976). The geosmin (8) biosynthesis investigated by feeding experiments to *Streptomyces* and the liverwort *Fossombronina pusilla* demonstrates that it is synthesized by cyclization of farnesyl diphosphate (SPITELLER & al. 2012). The odour threshold concentrations (OTCs; minimum detectable concentrations) is about 10 ng l⁻¹ or less (WATSON & al. 2000) and its presence in drinking water creates consumer complaints. Objectionable soily taste and odour in drinking water mainly caused by geosmin (8) and 2-methylisoborneol is a common problem for water suppliers using surface water sources (BAO & al. 1997). Furthermore, geosmin (8) is also resistant to conventional water treatment, and its bioaccumulation in fish and shellfish causes off-flavours in farmed and wild stocks (WATSON 2003). The occurrence of geosmin (8) also indicates mould-related spoilage (SCHNÜRER & al. 1999).

It is not surprising that in a large genus like *Cortinarius*, several species are known to produce conspicuous and diagnostic odours, although only very few of them have been chemically analysed, such as the anise-like odour of *C. odorifer* which was very recently identified as methyl p-anisate (KLEOFAS & al. 2015) or the naphthalene-like smell of *C. lebre* from Chile which is caused by indol (ARNOLD & al. 2012). Other scents like that of fruits in *C. traganus*, or of *Pelargonium* leaves in *C. flexipes*, among

others are used as characters in species discrimination, but remain to be chemically identified.

In general, VOCs play an important role for fungi in their natural environments. They act as signals between fungi and plants, arthropods, bacteria, and other fungi (BENNETT & al. 2012). Such fungal VOCs can further developed for use in biotechnological applications (MORATH & al. 2012). But it has to be taken in account, that for this purposes the fungal organism must be cultivable on artificial media, which is not the case in highly specialized ectomycorrhizal species such as *C. hinnuleus*.

G. PALFNER and N. ARNOLD are grateful for support by CONICYT-BMBF [bilateral cooperation project CONICYT (no. PCCI 2011-609) – German Ministry of Education and Research and (no. 01DN12107)].

References

- ARNOLD, N., PALFNER, G., SCHMIDT, J., KUHN, C., BECERRA, J., 2012: Chemistry of the aroma bouquet of the edible mushroom “lebre” (*Cortinarius lebre*, Basidiomycota, Agaricales) from Chile. – J. Chilean Chem. Soc. **58**: 1333–1335.
- BAO, M.-L., BARBIERI, K., BURRINI, D., GRIFFINI, O., PANTANI, F., 1997: Determination of trace levels of taste and odor compounds in water by microextraction and gas chromatography-ion-trap detection-mass spectrometry. – Water Res. **31**: 1719–1727.
- BELTRAN-GARCÍA, M. J., ESPINOSA-ESTARRON, M., OGURA, T., 1997: Volatile compounds secreted by the oyster mushroom (*Pleurotus ostreatus*) and their antibacterial activities. – J. Agric. Food Chem. **45**: 4049–4052.
- BENNETT, J. W., HUNG, R., LEE, S., PADHI, S., 2012: Fungal and bacterial volatile organic compounds: an overview and their role as ecological signaling agents. – In: Hock, B. (Ed.): Fungal associations. pp. 373–393. – Berlin: Springer.
- BRANDRUD, T. E., LINDSTROM, H., MARKLUND, H., MELOT, J., MUSKOS, S., 1990: *Cortinarius* Flora Photographica. – Matfors: Fotoflora.
- BREHERET, S., TALOU, T., RAPIOR, S., BESSIÈRE, J.-M., 1997: Monoterpenes in the aromas of fresh wild mushrooms (*Basidiomycetes*). – J. Agric. Food Chem. **45**: 831–836.
- BREHERET, S., TALOU, T., RAPIOR, S., BESSIÈRE, J.-M., 1999: Geosmin, a sesquiterpenoid compound responsible for the musty-earthly odor of *Cortinarius herculeus*, *Cystoderma amianthinum*, and *Cy. carcharias*. – Mycologia **91**: 117–120.
- BUTTERY, R., GARIBALDI, J., 1976: Geosmin and methylisoborneol in garden soil. – J. Agric. Food Chem. **24**: 1246–1247.
- CHIRON, N., MICHELOT, D., 2005: Mushrooms odors, chemistry and role in the biotic interactions: a review. – Cryptogamie Mycol. **26**: 299–364.
- CHO, I. H., NAMGUNG, H.-J., CHOI, H.-K., KIM, Y.-S., 2008: Volatiles and key odorants in the pileus and stipe of pine-mushroom (*Tricholoma matsutake* SING.). – Food Chemistry **106**: 71–76.
- COMBET, E., HENDERSON, J., EASTWOOD, D. C., BURTON, K. S., 2006: Eight-carbon volatiles in mushrooms and fungi: properties, analysis, and biosynthesis. – Mycoscience **47**: 317–326.
- DIBY, P., PARK, K. S., 2013: Identification of volatiles produced by *Cladosporium cladosporioides* CL-1, a fungal biocontrol agent that promotes plant growth. – Sensors **13**: 13969–13977.
- FÄLDT, J., JONSELL, M., NORDLANDER, G., BORG-KARLSON, A. K., 1999: Volatiles of bracket fungi *Fomitopsis pinicola* and *Fomes fomentarius* and their functions as insect attractants. – J. Chem. Ecol. **25**: 567–590.
- GMINDER, A., 2010: Die Grosspilze Baden-Württembergs **5**. – Stuttgart: Ulmer.
- INDEX FUNGORUM: <http://www.indexfungorum.org>. [Accessed January,12, 2014].
- KLEOFAS, V., POPA, F., FRAATZ, M. A., RÜHL, M., KOST, G. ZORN, H., 2015: Aroma profile of the anise-like odour mushroom *Cortinarius odorifer*. – Flavour Fragr. J. **30**: 381–386.
- KNUDSEN, J. T., ERIKSSON, R., GERSHENZON, J., STÄHL, B., 2006: Diversity and distribution of floral scent. – Bot. Rev. **72**: 1–120.

- MALHEIRO, R., DE PINHO, P.G., SOARES, S., DA SILVA FERREIRA, A. C., BAPTISTA, P., 2013: Volatile biomarkers for wild mushrooms species discrimination. – *Food Res. Internat.* **54**: 186–194.
- MORATH, S. U., HUNG, R., BENNETT, J. W., 2012: Fungal volatile organic compounds: a review with emphasis on their biotechnological potential. – *Fungal Biol. Rev.* **26**: 73–83.
- NISKANEN, T., KYTÖVUORI, I., 2012: Subgen. *Telamonia* sects. *Hinnulei* MELOT and *Safranopedes* LIIMAT., KYTÖV. & NISKANEN. – In KNUDSEN, H., VESTERHOLT, J., (Eds.): *Funga Nordica. Agaricoid boletoid, clavarioid, cyphelloid and gastroid genera.* – Copenhagen: Nordsvamp.
- PATTHAIK, S., SUBRAMANYAM, V. R., BAPAJI, M., KOLE, C. R. 1997: Antibacterial and antifungal activity of aromatic constituents of essential oils. – *Microbios.* **89**: 39–46.
- SCHNÜRER, J., OLSSON, J., BÖRJESSON, T., 1999: Fungal volatiles as indicators of food and feeds spoilage. – *Fungal Gen. Biol.* **27**: 209–217.
- SHIONO, Y., MATSUZAKA, R., WAKAMATSU, H., MUNETA, K., MURAYAMA, T., IKEDA, M., 2004: Fascicularones A and B from a mycelial culture of *Naematoloma fasciculare*. – *Phytochemistry* **65**: 491–496.
- SPIPELLER, G., 2002: Chemical responses to plant injury and plant aging – In: RAHMAN, A.-U., (Ed.) *Studies in natural product chemistry 27, bioactive natural products (part H)*, pp. 59–102. – Amsterdam: Elsevier.
- SPIPELLER, D., JUX, A., PIEL, J., BOLAND, W., 2012: Feeding of [5,5-²H₂]-1-desoxy-d-xylulose and [4,4,6,6,6-²H₅]-mevalolactone to a geosmin-producing *Streptomyces* sp. and *Fossombronia pusilla*. – *Phytochemistry* **61**: 827–834.
- THAKEOW, P., WEIßBECKER, B., SCHÜTZ, S., 2006: Volatile organic compounds emitted from fungal rotting beech (*Fagus sylvatica*). – *Mitt. Deutsch. Ges. Allg. Angew. Entomol* **15**: 157–160.
- WATSON, S. B. 2003: Cyanobacterial and eukaryotic algal odour compounds: signals or by-products? A review of their biological activity. – *Phycologia* **42**: 332–350.
- WATSON, S. B., BROWNLEE, B., SATCHWILL, T., HARGESHEIMER, E., 2000: Quantitative analysis of trace levels of geosmin and MIB in source and drinking water using headspace SPME. – *Water Res.* **34**: 2818–2828.
- WICHLACZ, M., AYER, W. A., TRIFONOV, L. S., CHAKRAVARTY, P., KHASA, D., 1999: A caryophyllene-related sesquiterpene and two 6, 7-seco-caryo-phyllenes from liquid cultures of *Hebeloma longicaudum*. – *J. Nat. Prod.* **62**: 484–486.
- YAMAGIWA, Y., INAGAKI, Y., ICHINOSE, Y., TOYODA, K., HYAKUMACHI, M., SHIRAIISHI, T., 2011: *Talaromyces wortmannii* FS2 emits β-caryophyllene, which promotes plant growth and induces resistance. – *J. Gen. Pl. Pathol.* **77**: 336–341.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Österreichische Zeitschrift für Pilzkunde](#)

Jahr/Year: 2016

Band/Volume: [25](#)

Autor(en)/Author(s): Arnold Norbert, Palfner Götz, Kuhnt Christine, Schmidt Jürgen

Artikel/Article: [Chemistry of the earthy odour of basidiomata of Cortinarius hinnuleus \(Basidiomycota, Agaricales\) 5-10](#)