

Key to the *Byssoloma* species (lichenised Ascomycota, Pilocarpaceae) known from Macaronesia

OTTHMAR BREUSS

Naturhistorisches Museum Wien

Botanische Abteilung (Kryptogamenherbar)

Burgring 7

1010 Wien, Österreich

Email: obreuss@bg9.at

Accepted 3. October 2016

Key words: Lichenised Ascomycota. – Taxonomy. – Mycobiota of Macaronesia, distribution.

Abstract: A key to the 11 species of *Byssoloma* known from Macaronesia is provided. Short notes on substrata, distribution in Macaronesia and worldwide, as well as relevant literature references are added.

Zusammenfassung: Die 11 bisher aus Makaronesien bekannten *Byssoloma*-Arten werden geschlüsselt. Kurze Angaben über die Substrate, die Verbreitung in Makaronesien und weltweit, sowie relevante Literaturangaben werden beigefügt.

The genus *Byssoloma* is characterised by a ± byssoid exciple, an amyloid tubular structure in the ascus apex (*Byssoloma*-type sensu HAFELLNER 1984), and mostly pyri-form conidia (LÜCKING 2008). It comprises 59 species (LÜCKING, pers.com.), most of which occur on leaves in the tropics.

During examination of *Byssoloma* collections from the island of Madeira (BREUSS 2012, 2013, 2014) species descriptions and keys widely distributed in the literature have been used. As a result, a compilation of Macaronesian species is herewith presented.

Currently 11 species of *Byssoloma* are known from Macaronesia, two of which (*Byssoloma kalbii* SÉRUS. and *B. laurisilvae* BREUSS) seem to be endemic to Madeira because they had not been recorded from anywhere else.

Key to the *Byssoloma* species of Macaronesia

- 1 Spores 11–20-septate, 40–70 µm long, tapering towards one end; apothecial margin smooth or finely pulverulent and shortly byssoid 2
- 1* Spores 3–7-septate, shorter 3

- 2 Apothecial discs brownish grey to dark brown or blackish, often with olive or bluish tinge; hypothecium brownish to orange brown, 100–160 µm thick; pycnidia pale orange or bluish to blackish blue ***B. kakouettae***
- 2* Apothecial discs yellowish to ochre; hypothecium 35–50 µm thick, colourless; pycnidia white ***B. laurisilvae***

- 3 Apothecia very small, usually less than 0.3 mm diam., apothecial margin byssoid but indistinct and disappearing in old apothecia 4
- 3* Apothecia larger, 0.3–0.6(–0.8) mm diam., apothecial margin byssoid or smooth, usually distinct 6
- 4 Apothecia tiny, 0.05–0.1 mm; margin very thin; pycnidia abundant, globose, whitish to bluish grey, N+ red; mature ascospores unknown ***B. diederichii***
- 4* Apothecia 0.1–0.3(–0.4) mm 5
- 5 Apothecia 0.15–0.2(–0.3) mm; discs almost white when young, becoming brownish when old; excipulum thin, almost absent in old apothecia; ascospores 3-septate; pycnidia abundant, globose, sessile, white or pale brownish ***B. kalbii***
- 5* Apothecia 0.1–0.3(–0.4) mm; discs at first chamois or rose-orange, soon brown orange to dark brown, with violet tinge; margin thin, disappearing in old apothecia, with colourless or brown crystals; ascospores 3-septate; pycnidia rare, globose, sessile, brown ***B. croceum***
- 6 Apothecial margin smooth or shortly byssoid 7
- 6* Apothecial margin distinctly byssoid (composed of free hyphae), but sometimes thin or becoming excluded in old apothecia 8
- 7 Apothecial margin with a compact appearance due to a gelatinous matrix embedding the hyphae, without crystals, thin but distinct, pale grey; discs dark brown and often with a purplish tinge; ascospores 3-septate ***B. marginatum***
- 7* Apothecial margin smooth or shortly byssoid; excipulum composed of short hyphae at the margin, inner part paraplectenchymatous (composed of polyhedral brown cells); apothecial discs variable in colour (bluish grey to blackish when young, becoming brown or greenish grey when mature, sometimes with a piebald mixture of these colours), margin usually paler than the disc in old apothecia; ascospores 3–5(–7)-septate ***B. ilimonae***
- 8 Apothecial margin comparatively thin but distinct, barely spreading laterally over the thallus surface; excipulum and epihymenium inspersed with crystals ***B. maderense***
- 8* Apothecial margin thick, spreading laterally over the thallus surface; excipulum and epihymenium without crystals 9
- 9 Apothecial discs pure black; thallus at least marginally dispersed into rounded patches; apothecial margin pure white ***B. subdiscordans***
- 9* Apothecial discs brownish grey to dark brown; thallus continuous or effuse; apothecial margin chamois-coloured 10

- 10 Ascospores 3-septate, 10–18 × 2.5–4 µm..... *B. leucoblepharum*
 10* Ascospores 5(–7)-septate, 18–28 × 3.5–5.5 µm..... *B. vezdanum*

Species list

Byssoloma croceum SÉRUS. & PUNTILLO

Substrata: Leaves, bark. – Distribution in Macaronesia: Canary Islands, Madeira. – World distribution: Macaronesia, Italy (Calabria). – Reference: SÉRUSIAUX (1998).

Byssoloma diederichii SÉRUS.

Substratum: Leaves. – Distribution in Macaronesia: Madeira. – World distribution: Macaronesia, France. – Reference: SÉRUSIAUX (1998).

Byssoloma kakouettae (SÉRUS.) LÜCKING & SÉRUS.

≡ *Bapalmia kakouettae* SÉRUS.,

= *Byssoloma aptrootii* SÉRUS.

Substrata: Leaves, bark. – Distribution in Macaronesia: Canary Islands, Madeira. – World distribution: Macaronesia, France (Pyrenees), Italy (Campania). – References: SÉRUSIAUX (1993, 1996, 1998), ETAYO (1998), SÉRUSIAUX et al. (2002), BREUSS (2014).

Byssoloma kalpii SÉRUS.

Substratum: Fern leaves. – Distribution in Macaronesia: Madeira. – World distribution: Madeira. – Reference: SÉRUSIAUX (1996).

Byssoloma laurisilvae BREUSS

Substratum: Leaves. – Distribution in Macaronesia: Madeira. – World distribution: Madeira. – References: BREUSS (2013, 2014).

Byssoloma leucoblepharum (NYL.) VAIN.

Substrata: Leaves, bark. – Distribution in Macaronesia: Azores, Canary Islands, Madeira. – World distribution: pantropical, subtropical-temperate. – References: TAVARES (1952), SANTESSON (1952), HERNÁNDEZ-PADRÓN et al. (1987), APTROOT (1989), ARVIDSSON (1990), ETAYO (1996, 1998), SÉRUSIAUX (1996), PIŠÚT (2009), SCHUMM & APTROOT (2013) (belongs to *B. maderense*), BREUSS (2014, 2016).

Byssoloma llimonae SÉRUS., GÓMEZ-BOLEA, LONGÁN & LÜCKING

Substrata: Bark, leaves. – Distribution in Macaronesia: Canary Islands, Madeira. – World distribution: Macaronesia, Spain. – References: SÉRUSIAUX et al. (2002), BREUSS (2014).

Byssoloma maderense BREUSS

Substrata: Leaves, bark. – Distribution in Macaronesia: Azores, Canary Islands, Madeira.

World distribution: Macaronesia, southeastern USA, mediterranean and suboceanic Eurasia. – References: BREUSS (2014, 2016).

***Byssoloma marginatum* (ARNOLD) SÉRUS.**

= *Tapellaria similis* KALB

Substrata: Leaves, bark. – Distribution in Macaronesia: Azores, Canary Islands, Madeira. – World distribution: cosmopolitan. – References: KALB & HAFELLNER (1992), ETAYO (1996, 1998), SÉRUSIAUX (1996), BERGER & APTROOT (2002), BOOM & ERTZ (2002).

***Byssoloma subdiscordans* (NYL.) P. JAMES**

= *B. rotuliforme* (MÜLL.ARG.) R. SANT.

Substrata: Leaves, rarely bark. – Distribution in Macaronesia: Azores, Canary Islands, Madeira. – World distribution: cosmopolitan. – References: TAVARES (1952), SANTESSON (1952), CHAMPION (1976), FOLLMANN & HERNÁNDEZ-PADRÓN (1978), PURVIS & JAMES (1993), ETAYO (1998), SÉRUSIAUX (1996), PIŠÚT (2009), SCHUMM & APTROOT (2013).

***Byssoloma vezdanum* SÉRUS.**

Substrata: Bark, mosses over bark. – Distribution in Macaronesia: Madeira. – World distribution: pantropical. – References: BREUSS (2012, 2014).

The author thanks ROBERT LÜCKING, Berlin, for valuable information.

References

- APTROOT, A., 1989: Contribution to the Azores lichen flora. – *Lichenologist* **21**: 59–65.
 ARVIDSSON, L., 1990: Additions to the lichen flora of the Azores. – *Bibl. Lichenol.* **38**: 13–27.
 BERGER, F., APTROOT, A., 2002: Further contributions to the flora of lichens and lichenicolous fungi of the Azores. – *Arquipélago. Life and Marine Sciences* **19A**: 1–12.
 BOOM, P. VAN DEN, ERTZ, D., 2012: Lichens and lichenicolous fungi from El Hierro (Canary Islands), a survey, including five new species. – *Cryptog. Mycol.* **33**: 59–97.
 BREUSS, O., 2012: Flechtenfunde auf Madeira. – *Stapfia* **97**: 47–52.
 BREUSS, O., 2013: *Byssoloma laurisilvae* und *Thelotrema lueckingii*, zwei neue Flechtenarten aus Madeira. – *Österr. Z. Pilzk.* **22**: 99–105.
 BREUSS, O., 2014: Weitere Flechtenfunde aus Madeira. – *Stapfia* **101**: 47–51.
 BREUSS, O., 2016: *Byssoloma maderense* is not endemic to Macaronesia. – *Evansia* **33**(2): 54–62.
 CHAMPION, C. L., 1976: Algunos líquenes nuevos para las Islas Canarias. – *Vieraea* **6**: 25–32.
 ETAYO, J., 1996: Contribution to a lichen flora of the Canary Islands. II. Epiphytic lichens from La Palma. – *Österr. Z. Pilzk.* **5**: 149–159.
 ETAYO, J., 1998: Aportación a la flora liquénica de las Islas Canarias. IV. Líquenes epífitos de La Gomera (Islas Canarias). – *Tropical Bryology* **14**: 85–107.
 FOLLMANN, G., HERNÁNDEZ-PADRÓN, C., 1978: Zur Kenntnis der Flechtenflora und Flechtenvegetation der Kanarischen Inseln. II. Über einige Neufunde, besonders von der Insel Hierro. – *Philippia* **3**: 360–378.
 HAFELLNER, J., 1984: Studien in Richtung einer natürlicheren Gliederung der Sammelfamilien *Lecanoraceae* und *Lecideaceae*. – *Nova Hedwigia Beih.* **79**: 241–371.
 HERNÁNDEZ-PADRÓN, C., SANCHEZ-PINTO, L., FOLLMANN, G., 1987: Zur Kenntnis der Flechtenflora und Flechtenvegetation der Kanarischen Inseln. VII. Arealytypen und Verbreitungsmuster einiger Neufunde. – *Cour. Forsch.-Inst. Senckenberg* **95**: 189–199.

- KALB, K., HAFELLNER, J., 1992: Bemerkenswerte Flechten und lichenicole Pilze von der Insel Madeira. – *Herzogia* **9**: 45–102.
- LÜCKING, R., 2008: Foliicolous lichenized fungi. – *Flora Neotropica Monograph* **103**.
- PÍŠUT, I., 2009: Some lichens from the vicinity of Ribeiro Frio (Madeira, Macaronesia). – *Acta Mycol.* **44**: 179–184.
- PURVIS, O.W., JAMES, P.W., 1993: Studies on the lichens of the Azores. Part 1 – Caldeira do Faial. – Arquipélago. Life and Marine Sciences **11A**: 1–15.
- SANTESSON, R., 1952: Foliicolous lichens I. A revision of the taxonomy of the obligately foliicolous, lichenized fungi. – *Symb. Bot. Upsal.* **12**(1): 1–590.
- SCHUMM, F., APTROOT, A., 2013: Flechten Madeiras, der Kanaren und Azoren 2 (Ergänzungsband). – Norderstedt: Books on Demand.
- SÉRUSIAUX, E., 1993: New taxa of foliicolous lichens from Western Europe and Macaronesia. – Nord. J. Bot. **13**: 447–461.
- SÉRUSIAUX, E., 1996: Foliicolous lichens from Madeira, with the description of a new genus and two new species and a world-wide key of foliicolous *Fellhanera*. – *Lichenologist* **28**: 197–227.
- SÉRUSIAUX, E., 1998: Deux nouvelles espèces de *Byssoloma* TREVIS. (lichens, *Pilocarpaceae*) d'Europe occidentale et Macaronésie. – *Cryptog. Bryol. Lichénol.* **19**: 197–209.
- SÉRUSIAUX, E., GÓMEZ-BOLEA, A., LONGÁN, A., LÜCKING, R., 2002: *Byssoloma llimonae* sp. nov., from continental Spain, Madeira and the Canary Islands. – *Lichenologist* **34**: 183–188.
- TAVARES, C. N., 1952: Contribution to the lichen flora of Macaronesia I. Lichens from Madeira. – *Portugaliae Acta Biol. (B)* **3**(3): 308–391.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Österreichische Zeitschrift für Pilzkunde](#)

Jahr/Year: 2016

Band/Volume: [25](#)

Autor(en)/Author(s): Breuss Othmar

Artikel/Article: [Key to the Byssoloma species \(lichenised Ascomycota, Pilocarpaceae\) known from Macaronesia 95-99](#)