

Phyton (Horn, Austria)	Vol. 38	Fasc. 2	239–250	29. 12. 1998
------------------------	---------	---------	---------	--------------

## Comparison between Growth Responses of Autotrophic and Heterotrophic Populations of Lichen Photobiont *Trebouxia irregularis* (Chlorophyta) on Cu, Hg and Cd Chlorides Treatment

By

Martin BAČKOR\*\*), Ján HUDÁK\*) and Miriam BAČKOROVÁ\*)

With 7 Figures

Received February 5, 1998

Accepted March 17, 1998

Key words: *Trebouxia*, photobiont, algae, lichens, growth, heavy metals, copper, mercury, cadmium.

### Summary

BAČKOR M., HUDÁK J. & BAČKOROVÁ M. 1998. Comparison between growth responses of autotrophic and heterotrophic populations of lichen photobiont *Trebouxia irregularis* (Chlorophyta) on Cu, Hg and Cd chlorides treatment. – *Phyton* (Horn, Austria) 38 (2): 239–250, 7 figures. – English with German summary.

Comparison between growth responses of autotrophic and heterotrophic populations of lichen photobiont *Trebouxia irregularis* (isolated from heavy metal tolerant population of lichen *Cladonia mitis*) on Cu, Hg and Cd chlorides treatment were studied. Heavy metal content in the *C. mitis* thalli, relative toxicity of single metal chlorides, effect of the concentration, differences between toxicity to autotrophic and heterotrophic strains and effective concentrations are discussed.

### Zusammenfassung

BAČKOR M., HUDÁK J. & BAČKOROVÁ M. 1998. Wirkungen der Schwermetallchloride (Cu, Hg und Cd) auf das Wachstum autotropher und heterotropher Popu-

\*) M. BAČKOR, J. HUDÁK and M. BAČKOROVÁ, Department of Plant Physiology, Faculty of Natural Sciences, Comenius University, Mlynská dolina B-2, SK-842 15 Bratislava, Slovak Republic.

\*\*) M. BAČKOR, Department of Experimental Botany and Genetics, Faculty of Natural Sciences, Šafárik University, Mánesova 23, SK-041 54 Košice, Slovak Republic (permanent address). E-mail: mbackor@kosice.upjs.sk

lationen des Photobionten *Trebouxia irregularis* (Chlorophyta) – ein Vergleich. – Phytion (Horn, Austria) 38 (2): 239–250, mit 7 Abbildungen. – Englisch mit deutscher Zusammenfassung.

Es wurde die Wirkung der Schwermetalle – Cu, Hg und Cd (als Chloride) auf den Wuchs auto- und heterotropher Populationen des Photobionten *Trebouxia irregularis* (isoliert von einer schwermetall-toleranten Population der Flechte *Cladonia mitis*) untersucht. Der Schwermetall-Gehalt in Thalli von *C. mitis*, die relative Toxizität der einzelnen Metallchloride, der Einfluß der Konzentration und die unterschiedliche Empfindlichkeit der auto- und heterotrophen Stämme wird diskutiert.

## Introduction

The lichens which are, in general, sensitive to different pollutants, play an important role in numerous studies focused on air pollution (for review see: AHMADJIAN 1993, GALUN & RONEN 1988, HAWKSWORTH 1990, NASH 1988, NASH & GRIES 1991, RICHARDSON 1988, 1991, SEAWARD 1989, TÜRK 1988).

The absorption, accumulation and toxicity of heavy metals in lichens are also well documented (AHMADJIAN 1993, BROWN 1991, GARTY 1993, NASH 1989, PURVIS & HALLS 1996, RICHARDSON 1995, SEAWARD & RICHARDSON 1989, TYLER 1989, TYLER & al. 1989). The main source of the heavy metals accumulation in lichens is air, however, in lichens growing on metalliferous rocks and derived soils from mine and smelter wastes (PURVIS & HALLS 1996) the substrate may also play an important role in their absorption.

Heavy metals affect both the mycobionts and photobionts associated in lichen thallus. Similarly, in non-lichenized algae (GENTER 1996) the changes in photosynthesis, respiration, chlorophyll content and in cyanolichens (e.g. *Peltigera* sp.) nitrogen fixation are observed as a result of the air pollution and heavy metal influence (BOONPRAGOB & NASH 1991, DERUELLE & PETIT 1983, FIELDS 1988, GARTY & al. 1988, 1992, RICHARDSON & NIEBOER 1983, SHOWMAN 1972, VON ARB & al. 1990). However, only few investigations are given on poisoning of pollutants including heavy metals on isolated lichen bionts, although photobionts appear to be the key element in lichen sensitivity to air pollution (AHMADJIAN 1993). Effect of ozone on photobiont *Trebouxia* (from *Cladonia stellaris*) was observed by ROSEN-TRETTNER & AHMADJIAN 1977. EWALD & SCHLEE 1983 showed the effect of sulfite on the proline content decrease in *Trebouxia* (influence on proline dehydrogenase as the last enzyme of proline synthesis). SCHLEE & al. 1984 studied the effect of sulphite on the activity of aminopeptidases in the lichen alga *Trebouxia* sp. Similarly, photobionts isolated from *Usnea* sp., *Hypogymnia physodes* and *Lecanora conizaeoides* (the latter lichen is known as very tolerant to air pollution) were sensitive to low concentrations of sulphite (HILL 1974). MARTI 1983, 1985 exposed cultured photobionts to aqueous solutions of sulfite, nitrite, sulfate and nitrate and

shown that they were distinctly sensitive to sulfite. Nitrite and zinc were far less toxic, and it is interesting, sulfate and nitrate in a few cases increased the growth rates. NASH 1975 found that zinc and cadmium (at concentration  $5 \times 10^{-5}$  M) inhibited growth of two cultured photobionts (*Trebouxia gelatinosa*, *T. erici*).

Most species of *Trebouxia* can grow heterotrophically in the dark on organic media. Evidence for existence of heterotrophy in situ (in lichen thallus) comes from studies of Antarctic lichens where they are exposed to light for less than 2 months per year during the summer, and during the winter they grow up to 3 m under the snow (AHMADJIAN 1993).

## Material and Methods

### 1. Isolation

Photobiont *Trebouxia irregularis* was isolated from lichen *Cladina mitis* (heavy metal tolerant population from copper ore depositions at Špania dolina, Central Slovakia) by both, centrifugation and micropipette methods (ASCASO 1980, AHMADJIAN 1993).

### 2. Influence of heavy metals

For cultivation, 1N *Trebouxia* medium (AHMADJIAN 1993) containing glucose (20g/l) and caseine (10g/l) was used. The final pH of the media was adjusted to 7. Medium (50 ml) was pipetted into Erlenmayer flasks and autoclaved. The chlorides of heavy metals (Cu, Cd, Hg) were added after the sterilisation of the media in final concentrations 30  $\mu$ M and 300  $\mu$ M (for Cu, Cd, Hg) and 3mM (for Cu).

Algae from culture were homogenised by gentle stirring on magnetic stirrer for 1 hour. Approximately  $6 \times 10^5$  (final concentration  $0,12 \times 10^5$  in 1 ml of cultivation medium) cells of photobiont was transferred to each flask as an inoculum. The presence and homogeneity of algae were examined microscopically. Growth of cultures was calculated using standard hemocytometer chamber every 5 days (up to 30 days) as number of cells per 1 ml of medium. Growth of heterotrophic cultures was calculated under green light.

### 3. Incubation

The autotrophic cultures were maintained at 25 °C under a 16-h photoperiod and 24  $\mu$ M . m<sup>-2</sup> . s<sup>-1</sup> artificial irradiance. Heterotrophic cultures were kept in the dark at 25 °C.

## Results and Discussion

### 1. Heavy metal content in lichen *Cladina mitis*

The mean content of metals in thallus of *Cladina mitis* was the following: 28,18  $\mu$ g . g<sup>-1</sup> for Cu, 0,88  $\mu$ g . g<sup>-1</sup> for Hg and 1,85  $\mu$ g . g<sup>-1</sup> for Cd (unpublished data). These concentrations are lower or higher in comparison with other studies (BARGAGLI & BARGHIGIANI 1991, GALUN & al. 1984, GARTY & AMMANN 1987, GARTY & al. 1986, NASH 1975, PAKARINEN 1985, RAO & al. 1977).

## 2. Comparison between autotrophic and heterotrophic growth of control

Fig. 1 shows the differences between autotrophic (in the light) and heterotrophic (in the dark) growth of strains of *Trebouxia irregularis* on organic trebouxia medium. Although most species of *Trebouxia* can grow heterotrophically in the dark on organic media (AHMADJIAN 1993), our results suggest that great differences in quantity of the cells between autotrophic and heterotrophic populations (after 30 days of cultivation approximately 50 times more cells in the light cultivation) exist.




Fig. 1. The growth of *Trebouxia irregularis* calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).

## 3. Influence of single metals on autotrophic growth

The figures 2–4 represent autotrophic growth of photobiont cultures after the Hg, Cd and Cu chloride treatment in different final concentrations (0,03 mM, 0,3 mM and for Cu 3 mM). After 30 day cultivation, the relative toxicity of metals within the given concentrations were as follows: for 0,03 mM:  $\text{Cu}^{2+} \ll \text{control} \leq \text{Cd}^{2+} \leq \text{Hg}^{2+}$ , for 0,3 mM:  $\text{control} \leq \text{Cu}^{2+} < \text{Cd}^{2+} < \text{Hg}^{2+}$  and for 3 mM:  $\text{control} < \text{Cu}^{2+}$ . The concentrations used in our study are much more higher (for Hg, Cd) than the metal content in *C. mitis* thallus. For Cu due to its higher concentration in the thallus as well as chemical nature of substrate (dumps of metal ore depositions) three concentrations were used. The results suggest that only Cu at 0,03 mM concentration stimulates the growth of cultures.






Fig. 2. The autotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).




Fig. 3. The autotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).




Fig. 4. The autotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).

In our study we have used only chlorides of metals, and therefore a toxicity of different salts was not tested. LEE & al. 1993 observed no difference in toxicity between  $\text{CuCl}_2$  and  $\text{CuSO}_4$  in the growth studies of *Anacystis nidulans*. All metals, even those that are nutrients at low concentrations, are toxic to algae at high concentrations (RAI & al. 1981) and inhibition effect depends on concentrations of metals, where higher concentrations decrease growth of the cultures.

The effective concentrations ( $\text{EC}_{50}$ , what means that growth is inhibited for 50%) for 5 day or 30 day autotrophic cultivation are calculated as follows: for Cd 5 days  $\text{EC}_{50} = 0,9$  mM and for 30 days  $\text{EC}_{50} = 0,88$  mM, for Cu 5 days  $\text{EC}_{50} = 2,66$  mM and for 30 days 1,6 mM, for Hg 5 days  $\text{EC}_{50} = 0,23$  mM and for 30 days 0,11 mM growth inhibition. The results indicate that during 5 days the  $\text{EC}_{50}$  indexes are higher and result from biosorption, probably accumulation in cell wall components, which does not cause high toxicity.

#### 4. Influence of single metals on heterotrophic growth

The figures 5–7 represent heterotrophic growth photobiont cultures after treatment of chlorides in different final concentrations, the same as for autotrophic growth. After 30 days cultivation the required sequences of metals toxicity were as follows: (from the best growth to less) for 0,03 mM




Fig. 5. The heterotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).




Fig. 6. The heterotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).




Fig. 7. The heterotrophic growth of *Trebouxia irregularis* cultures calculated as number of cells for 1 ml of medium ( $\times 10^5$ ).

are:  $\text{Cu}^{2+} < \text{Cd}^{2+} \leq \text{Hg}^{2+} \leq \text{control}$ , for 0,3 mM are:  $\text{control} \leq \text{Cu}^{2+} < < \text{Cd}^{2+} < < < \text{Hg}^{2+}$ . The results indicate, that probably exist different mechanisms of detoxications of metals in the light and dark. Similarly, the influence of single metals is different and depends on their final concentrations.

There are differences in toxicity of single chlorides of metals in medium between autotrophe and heterotrophe growth. There are two phases of heavy metals uptake (RAI & al. 1981). The metabolism independent phase (biosorption) involves the accumulation of inorganic chemicals to cell wall components. The metabolism dependent phase is often slower (lasting hours or days) and is inhibited by low temperatures, absence of energy sources (light) or metabolic inhibitors (GADD 1988). Absence of energy sources (light in our case) is resulted in lower toxicity of metals in concentration 0,03 mM at heterotrophe growth. At 0,3 mM concentrations of metals the situation is different and metals are toxic in order:  $\text{control} \leq \text{Cu}^{2+} < < \text{Cd}^{2+} < < < \text{Hg}^{2+}$ .

Following conclusions and explanations can be drawn from the above presented results. First, inorganic stressors almost always occur as mixtures in nature (GENTER 1996). RAI & al. 1981 found that the same algae may behave synergistically to one metal combination and antagonistically to another metal combination and, therefore different metal concentrations must be tested. Second, another important factor is metal concentration. Our results indicate relatively high tolerance to three metals at pH 7 (optimum for this strain) but pH which is an important factor of the



heavy metal toxicity (PAWLIK & al. 1993, PETERSON & al. 1984) should also be tested. Third, toxicity of heavy metals depends on temperature (FLORENCE & al. 1994). Temperature used in our study (25 °C) is relatively high for *Trebouxia*. Algae at lower temperatures can tolerate higher concentrations of heavy metals. Fourth, relative toxicity of chemical mixtures can differ between clones of the same species (BRAEK & al. 1980). EDTA, and other organic substances, can bind to metals and reduce their toxicity (GENTER 1996).

Dumps of copper ore depositions in Špania dolina are relatively old and possibilities for genetic tolerance are an interesting question. TWISS & al. 1993 showed possibilities of the selection of a Cu-resistant strain from a Cu-sensitive strain of *Scenedesmus* by repeated culturing under sublethal concentrations of Cu at laboratory conditions. Probably, there are more mechanisms to tolerate heavy metals, but similar selection can occur in nature, in case of photobionts from heavy metal tolerant populations of lichens growing hundreds of years on heavy metal enriched substrates.

#### Acknowledgement

This work was financially supported by the following grants: grant 1/4098/97 from Slovak Ministry of Education to J. HUDÁK, M. BAČKOR, and grant 1573 from Rector of Comenius University for M. BAČKOR and M. BAČKOROVÁ.

#### References

- AHMADJIAN V. 1993. The lichen symbiosis. – John Wiley & sons, Inc., 250 pp.
- ASCASO C. 1980. A rapid method for the quantitative isolation of green algae from lichens. – Ann. Bot. 45: 483.
- BARGAGLI R. & BARGHIGIANI C. 1991. Lichen biomonitoring of mercury emission and deposition in mining, geothermal and volcanic areas of Italy. – Environmental Monitoring and Assessment 16: 265–275.
- BOONPRAGOB K. & NASH T. H. 1991. Physiological responses of the lichen *Ramalina menziesii* Tayl. to the Los Angeles urban environment. – Environmental and Experimental Botany 31: 229–238.
- BRAEK G. S., MALNES D. & JENSEN A. 1980. Heavy metal tolerance of marine phytoplankton. IV. Combined effect of zinc and cadmium on growth and uptake in some marine diatoms. – J. Exp. Mar. Biol. Ecol. 42: 39–54.
- BROWN D. H. 1991. Lichen mineral studies – currently clarified or confused. – Symbiosis 11: 207–223.
- DERUELLE S. & PETIT P. J. X. 1983. Preliminary studies on the net photosynthesis and respiration responses of some lichens to automobile pollution. – Cryptogamie, Bryologie et Lichenologie 4: 269–278.
- EWALD D. & SCHLEE D. 1983. Biochemical effects of sulphur dioxide on proline metabolism in the alga *Trebouxia*. – New Phytologist 94: 235–240.
- FIELDS R. F. 1988. Physiological responses of lichens to air pollutant fumigations. – In: NASH T. H. & WIRTH V. (Eds.), Lichens, bryophytes and air quality. Bibliotheca Lichenologica, Vol. 30, pp. 175–200, J. Cramer, Berlin-Stuttgart.

- FLORENCE T. M., STAUBER J. L. & AHSANULLAH M. 1994. Toxicity of nickel ores to marine organisms. – *Sci. Total Environ.* 148: 139–155.
- GADD G. M. 1988. Accumulation of metals by microorganisms and algae. – In: REHM H. J. (Ed.), *Biotechnology*, Vol. 60, pp. 401–434, VCH, Weinheim.
- GALUN M. & RONEN R. 1988. Interactions of lichens and pollutants. – In: GALUN M. (Ed.), *CRC Handbook of lichenology*. Volume III, pp. 55–72, CRC Press, Inc., Boca Raton.
- , GARTY J. & RONEN R. 1984. Lichens as bioindicators of air pollution. – *Webbia* 38: 371–383.
- GARTY J. 1993. Lichens as biomonitors for heavy metal pollution. – In: MARKERT B. (Ed.), *Plants as biomonitors. Indicators for heavy metals in the terrestrial environment*, pp. 193–263, VCH, Weinheim.
- & AMMANN K. 1987. The amounts of Ni, Cr, Zn, Pb, Cu, Fe and Mn in some lichens growing in Switzerland. – *Environmental and Experimental Botany* 27: 127–138.
- , GALUN M. & HOCHBERG Y. 1986. The accumulation of metals in *Caloplaca aurantia* growing on concrete roof tiles. – *Lichenologist* 18: 257–263.
- , KARARY Y. & HAREL J. 1992. Effect of low pH, heavy metals and anions on chlorophyll degradation in the lichen *Ramalina duriaei* (De Not.) Bagl. – *Environmental and Experimental Botany* 32: 229–241.
- , KARDISH N., HAGEMEYER J. & RONEN R. 1988. Correlations between the concentration of adenosine triphosphate, chlorophyll degradation and the amounts of airborne heavy metals and sulphur in a transplanted lichen. – *Archives of Environmental Contamination and Toxicology* 17: 601–611.
- GENTER R. B. 1996. Ecotoxicology of inorganic chemical stress to algae. – In: STEVENSON R. J., BOTHWELL M. L. & LOWE R. L. (Eds.), *Algal ecology*, pp. 403–468, Academic Press, San Diego-Toronto.
- HAWKSWORTH D. L. 1990. The long-term effects of air pollution on lichen communities in Europe and North America. – In: WOODWELL G. M. (Ed.), *The earth in transition: patterns and processes of biotic impoverishment*, pp. 45–64, Cambridge University Press, Cambridge.
- HILL D. J. 1974. Some effects of sulphite on photosynthesis in lichens. – *New Phytologist* 73: 1193–1205.
- LEE L. H., LUSTIGMAN B. & MACCARI J. 1993. Effect of copper on the growth of *Anacytis nidulans*. – *Bull. Environ. Contam. Toxicol.* 50: 600–607.
- MARTI J. 1983. Sensitivity of lichen phycobionts to dissolved air pollutants. – *Canadian Journal of Botany* 61: 1647–1653.
- 1985. Die Toxizität von Zink, Schwefel- und Stickstoffverbindungen auf Flechten-Symbionten. – *Bibliotheca Lichenologica* 21, 129 pp, J. Cramer, Vaduz.
- NASH T. H. III. 1975. Influence of effluents from a zinc factory on lichens. – *Ecological Monographs* 45: 183–198.
- 1988. Correlating fumigation studies with field effects. – In: NASH T. H. III. & WIRTH V. (Eds.), *Lichens, bryophytes and air quality*. *Bibliotheca Lichenologica* No. 30, pp. 201–216, J. Cramer, Berlin-Stuttgart.
- 1989. Metal tolerance in lichens. – In: SHAW A. J. (Ed.), *Heavy metal tolerance in plants: Evolutionary aspects*, pp. 119–131, CRC Press, Boca Raton.

- & GRIES C. 1991. Lichens as indicators of air pollution. – In: HUTZINGER O. (Ed.), *The handbook of environmental chemistry*, pp. 1–29, Springer-Verlag, Heidelberg, New York.
- PAKARINEN P. 1985. Mineral element accumulation in bog lichens. – In: BROWN D. H. (Ed.), *Lichen physiology and cell biology*, pp. 185–192, Plenum Press, New York and London.
- PAWLIK B., SKOWROŃSKI T., RAMAZANOW Z., GARDESTROM P. & SAMUELSSON G. 1993. pH-dependent cadmium transport inhibits photosynthesis in the cyanobacterium *Synechocystis aquatilis*. – *Environ. Exp. Bot.* 33: 331–337.
- PETERSON H. G., HEALEY F. P. & WAGEMANN R. 1984. Metal toxicity to algae: A highly pH dependent phenomenon. – *Can. J. Fish. Aquat. Sci.* 41: 974–979.
- PURVIS O. W. & HALLS C. 1996. A review of lichens in metal-enriched environments. – *Lichenologist* 28: 571–601.
- RAI L. C., GAUR J. P. & KUMAR H. D. 1981. Phycology and heavy-metal pollution. – *Biol. Rev. Cambridge Philos. Soc.* 56: 99–151.
- RAO D. N., ROBITAILLE G. & LEBLANC F. 1977. Influence of heavy metal pollution on lichens and bryophytes. – *Jour. Hattori Bot. Lab.* 42: 213–239.
- RICHARDSON D. H. S. 1988. Understanding the pollution sensitivity of lichens. – *Botanical Journal of the Linnean Society* 96: 31–43.
- 1991. Lichens as biological indicators – recent developments. – In: JEFFREY D. W. & MADDEN B. (Eds.), *Bioindicators and environmental management*, pp. 263–272, Academic Press, London – San Diego.
- 1995. Metal uptake in lichens. – *Symbiosis* 18: 119–127.
- & NIEBOER E. 1983. Ecophysiological responses of lichens to sulphur dioxide. – *Journal of the Hattori Botanical Laboratory* 54: 331–351.
- ROSENTRETTER R. & AHMADJIAN V. 1977. Effect of ozone on the lichen *Cladonia arbuscula* and the *Trebouxia* phycobiont of *Cladonia stellaris*. – *The Bryologist* 80: 600–605.
- SCHLEE D., EWALD D. & ALBRECHT R. 1984. Effect of sulphite on the activity of aminopeptidases in the lichen alga *Trebouxia* sp. – *Biologia (Bratislava)* 39: 1199–1207.
- SEAWARD M. R. D. 1989. Lichens as monitors of recent changes in air pollution. – *Plants Today* 2: 64–69.
- & RICHARDSON D. H. S. 1989. Atmospheric sources of metal pollution and effects on vegetation. – In: SHAW A. J. (Ed.), *Heavy metal tolerance in plants: Evolutionary aspects*, pp. 75–92, CRC Press, Boca Raton.
- SHOWMAN R. E. 1972. Residual effects of sulfur dioxide on the net photosynthesis and respiratory rates of lichen thalli and cultured lichen symbionts. – *The Bryologist* 75: 335–341.
- TÜRK R. 1988. Bioindikation von Luftverunreinigungen mittels Flechten. – In: GRILL D. & GUTTENBERGER H. (Eds.), *Ökophysiologische Probleme durch Luftverunreinigungen*, pp. 13–27, Institut für Pflanzenphysiologie, Karl-Franzens-Universität, Graz.
- TWISS M. R., WELBOURN P. M. & SCHWARTZEL E. 1993. Laboratory selection for copper tolerance in *Scenedesmus acutus* (*Chlorophyceae*). – *Can. J. Bot.* 71: 333–338.
- TYLER G. 1989. Uptake, retention, and toxicity of heavy metals in lichens. – *Water, Air, and Soil Pollution* 47: 321–333.

- , PAHLSSON A. M. B., BENGTSSON G., BAATH E. & TRANVIK L. 1989. Heavy metal ecology of terrestrial plants, microorganisms and invertebrates. A review. – Water, Air, and Soil Pollution 47: 189–215.
- VON ARB C., MUELLER C., AMMANN K. & BRUNOLD C. 1990. Lichen physiology and air pollution. II. Statistical analysis of the correlation between SO<sub>2</sub>, NO<sub>2</sub>, NO and O<sub>3</sub>, and chlorophyll content, net photosynthesis, sulphate uptake and protein synthesis of *Parmelia sulcata* Taylor. – New Phytologist 115: 431–437.

Phyton (Horn, Austria) 38 (2): 250 (1998)

## Recensio

**Flora of North America North of Mexico. Volume 3. Magnoliophyta: Magnoliidae and Hamamelididae 1997.** Edited by Flora of North America Editorial Committee. – Lex. 8°, XXIV + 590 Seiten, 100 Abbildungen, 1470 Verbreitungskarten. – Oxford University Press, New York, Oxford. – £ 65,-. – ISBN 0-19-511246-6.

Der dritte Band der neuen Nord-Amerika-Flora enthält mit den Unterklassen *Magnoliidae* (angenehm bzw. praktisch, daß diese nicht weiter gesplittet sind) und *Hamamelididae* (sensu CRONQUIST) die Ordnungen *Magnoliales*, *Laurales*, *Piperales*, *Aristolochiales*, *Illiciales*, *Nymphaeales*, *Ranunculales*, *Papaverales*, *Hamamelidales*, *Urticales*, *Leitneriales*, *Juglandales*, *Myricales*, *Fagales* und *Casuarinales*. Dem Konzept der Flora entsprechend gibt es für alle Taxa, von der Familie an abwärts, ausführliche, sorgfältige Beschreibungen und Bestimmungsschlüssel. Auf den angenommenen Species-Namen mit Zitat folgt(-en) meist (ein) Volksname(n). Gegebenenfalls folgt ein Gefährdungshinweis. Ein E keinnzeichnet Arten, die nur im Gebiet der Flora vorkommen. Wichtige Synonyme sind angeführt. Nach der Beschreibung folgen Hinweise auf Blütezeit, Standort, Höhenverbreitung und geographische Verbreitung. Schließlich gibt es kurze Diskussionen nomenklatorisch-taxonomischer Fragen, von Variabilität, ökologischer Ansprüche, von Nutzung (Kultur, Heilpflanze), Unkrauteigenschaften oder anderes. Für jedes Taxon gibt es eine kleine, schematisierte Verbreitungskarte. Da der Band 741 Arten enthält und die Zahl der Karten im Verlagsprospekt mit 1470 angegeben wird, müssen außer den Typussippen zusätzlich noch 729 infraspezifische Taxa berücksichtigt sein. Welche Autoren welche Familie oder Gattungen bearbeitet haben, ist stets angegeben.

Bei der Erstellung des Konzeptes für die Flora, dürfte manche Inspiration von seiten der Flora Europaea eingeflossen sein, aber die Flora of North America ist in der Qualität weit über diese hinausgewachsen. Eine ganz exzellente Flora! Hoffentlich lassen die ausständigen Bände (die Flora ist auf 30 Bände angelegt) nicht zu lange auf sich warten; der Rezensent würde den Abschluß noch gerne erleben.

Der vorliegende Band hat auch dadurch erhebliche praktische Bedeutung, daß er viele vegetationsprägende Gehölze enthält, z.B. *Fagaceae* [alleine 90 *Quercus*-Arten, *Castanea* (3) mit ausführlicher Diskussion der Schäden durch *Cryphonectria parasitica*], *Betulaceae* s.l. [*Alnus* (5), *Betula* (18)], *Juglandaceae* [*Carya* (11), *Juglans* (6)], *Ulmaceae*, *Maclura*, *Platanus*, *Magnolia* u.a. Die größte behandelte Familie sind die *Ranunculaceae*.



# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 1998

Band/Volume: [38\\_2](#)

Autor(en)/Author(s): Backor Martin, Hudak Jan, Backorova Miriam

Artikel/Article: [Comparison between Growth Responses of Autotrophic and Heterotrophic Populations of Lichen Photobiont \*Trebouxia irregularis\* \(Chlorophyta\) on Cu, Hg and Cd Chlorides Treatment. 239-250](#)