

Phyton (Austria) Special issue: "Global change"	Vol. 42	Fasc. 3	(55)-(63)	1.10.2002
---	---------	---------	-----------	-----------

The Impact of Atmospheric H₂S on Growth and Sulfur Metabolism of *Allium cepa* L.

By

M. DURENKAMP¹⁾ & L. J. DE KOK¹⁾

Key words: *Allium cepa*, onion, hydrogen sulfide, sulfur metabolism, nitrogen metabolism, H₂S deposition, alliins.

Summary

DURENKAMP M. & DE KOK L. J. 2002. The impact of atmospheric H₂S on growth and sulfur metabolism of *Allium cepa* L. - *Phyton* (Horn, Austria) 42 (3): (55) - (63).

The impact of atmospheric H₂S deposition on growth and sulfur metabolism has been studied in onion (*Allium cepa* L.). The H₂S uptake followed saturation kinetics with respect to the H₂S concentration. The maximum H₂S uptake rate (JH₂S_{max}) was approx. 1 µmol g⁻¹ FW h⁻¹ and the KH₂S (H₂S concentration at which ½JH₂S_{max} was reached) was approx. 1.5 µl l⁻¹, which demonstrated that onion had a rather high H₂S uptake rate when compared with other species. Upon exposure of onion to 0, 0.075, 0.15, 0.225 and 0.3 µl l⁻¹ H₂S for two weeks, growth was only slightly reduced at 0.3 µl l⁻¹ H₂S. H₂S exposure resulted in an increased content of sulfate, total thiols and total sulfur in the shoot, while that in the roots was unaltered. There was a slight increase in total nitrogen content in the shoot, whereas the content of nitrate and amino acids was hardly affected. There was a substantial increase in the organic sulfur fraction in the shoot upon H₂S fumigation, which was probably due to an increase in the alliin content.

Introduction

Atmospheric H₂S has a paradoxical impact on plants. It may negatively affect growth at atmospheric levels of 0.03 µl l⁻¹ and higher and may even cause visible injury and defoliation at ≥ 0.3 µl l⁻¹, however, there is a wide variation in susceptibility between species to H₂S (DE KOK & al. 1998, 2000, 2002). On the other hand, foliarly absorbed H₂S may be utilized as sulfur source for growth and may be beneficial if the pedospheric sulfur supply is limited (DE KOK & al. 2000, 2002, WESTERMAN & al. 2000). H₂S is taken up via the stomates, metabolized with high

¹⁾ Laboratory of Plant Physiology, University of Groningen, P.O.Box 14, 9750 AA Haren, The Netherlands.

affinity into cysteine and subsequently into other sulfur compounds (DE KOK & al. 1998, STUIVER & DE KOK 2001). In *Brassica oleracea* L. there was a direct interaction between foliar H_2S deposition and the uptake and metabolism of pedospheric sulfate (DE KOK & al. 2000, WESTERMAN & al. 2000, 2001a,b).

Allium cepa L. (onion) was possibly one of the first domesticated vegetables by man and it was already cultivated by the ancient Egyptians. *Allium cepa* appears to be a species with a high sink capacity for reduced sulfur. This species belongs to the genus *Allium* (about 400 species), which derives its name from an allyl group in its secondary sulfur compounds. These compounds, specific for the family Alliaceae, are known as alliiins and consist of different stereoisomers of *S*-alk(en)ylcysteine sulfoxides. Cysteine is the major precursor for the synthesis of alliiins (BLOCK 1992). The content of alliiins in the leaves can directly be related to the sulfur status and can be affected by the sulfate supply to the roots (RANDLE & al. 1993, 1999, HANEKLAUS & al. 1997). The alliiins may account for up to 80% of the total organic sulfur fraction in onion and they are metabolically inert end-products, which can not be re-metabolized as sulfur source (SCHNUG 1993).

The aim of our study is to get insight into the interaction between atmospheric and pedospheric sulfur nutrition in onion and the significance of secondary sulfur compounds as sink for foliarly absorbed sulfur gases. The present study is focused on the kinetics of H_2S uptake by onion and on the impact of H_2S fumigation on growth, the sulfur content and its distribution in both shoot and roots.

Material and Methods

Seeds of onion (*Allium cepa* L. cv. Nerato F1; Nickerson-Zwaan, Made, The Netherlands) were germinated in vermiculite in a climate-controlled room. Day and night temperatures were 20 and 17 °C, respectively, with a relative humidity of 60-70%. The photoperiod was 14 hours at a photon fluence rate of 250-300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR 400-700 nm).

For H_2S deposition measurements two-week-old seedlings were transferred to 30 l tanks (12 plants per set, 20 sets per tank) containing a 25% Hoagland nutrient solution, pH 5.9, and grown for 10 days. H_2S uptake kinetics were measured as described by STUIVER & DE KOK 2001. Plants were taken from the nutrient solution, transferred to plastic pots filled with fine perlite, 12 plants per pot, and placed in 3 l glass cuvettes. The air flow through the cuvette was approximately 100 l h^{-1} , the temperature was 23 ± 1 °C, the relative humidity was about $40 \pm 10\%$ and the photon fluence rate was $190 \pm 10 \mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR 400-700 nm). The H_2S uptake rate (JH_2S) and the transpiration rate (JH_2O) were derived from the difference in concentration of H_2S and that of H_2O vapor between the inlet and outlet port of the cuvette and the total shoot fresh weight of the exposed plants, and calculated according to DE KOK & al. 1991. The apparent maximum H_2S uptake rate ($\text{JH}_2\text{S}_{\text{max}}$) and the atmospheric H_2S concentration at which $\frac{1}{2}\text{JH}_2\text{S}_{\text{max}}$ was reached (KH_2S) were calculated and fitted for Michaelis-Menten kinetics by Enzfitter (a Data Analysis Program by R.J. Leatherbarrow, Elsevier-BIOSOFT, Cambridge, U.K.). The ratio of H_2S deposition velocity to aqueous vapor efflux ($\text{gH}_2\text{S}/\text{gH}_2\text{O}$) was calculated and values were multiplied by 1.369 ($\sqrt{(\text{molecular weight H}_2\text{S})/(\text{molecular weight H}_2\text{O})}$), in order to correct for the differences in diffusion velocity between H_2S and H_2O (DE KOK & al. 1991, 1998).

For the assessment of the impact of H_2S fumigation on growth and metabolite content, two-week-old seedlings were transferred to stainless steel containers (19.5 x 15.0 x 45 cm; 6 plants per set, 10 sets per container) filled with a 25% Hoagland nutrient solution, pH 5.9. Containers were

placed in 150 l cylindrical stainless steel cabinets (0.6 m diameter) with a polycarbonate top, as described by STUIVER & al. 1992 and exposed to 0, 0.075, 0.15, 0.225 and 0.3 $\mu\text{l l}^{-1}$ H_2S for two weeks. Day and night temperatures were 24 and 18 $^{\circ}\text{C}$ (± 1 $^{\circ}\text{C}$) respectively, relative humidity was 40-50% and the photoperiod was 14 hours at a photon fluence rate of 300-350 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (PAR 400-700 nm) at plant height, with Philips HPL(R)N (400 W) as light source. Temperature was controlled by adjusting the cabinet wall temperature; the air exchange was 40 l min^{-1} and the air inside the cabinets was stirred continuously by a ventilator. Pressurized H_2S diluted with N_2 (1 ml l^{-1}) was injected into the incoming airstream and adjusted to the desired level by ASM electronic mass flow controllers (Bilthoven, The Netherlands). H_2S levels in the cabinets were controlled by an SO_2 analyzer (model 9850) equipped with a H_2S converter (model 8770, Monitor Labs, Measurement Controls Corporation, Englewood, CO 80112, USA).

Anions were determined according to MAAS & al. 1986. Shoot and roots of powdered dry material (dried at 120 $^{\circ}\text{C}$ for 18 h) were extracted for anion determination by incubating them overnight in demineralized water (50 mg in 5 ml) at 40 $^{\circ}\text{C}$ (TAUSZ & al. 1996). Homogenates were filtered through one layer of Miracloth and the filtrate was centrifuged at 30,000 g for 15 min (0 $^{\circ}\text{C}$). Sulfate and nitrate in the supernatant were separated on an Ionosphere 5A anion exchange column with guard column (250 x 4.6 and 30 x 3.0 mm, respectively, Varian/Chrompack Benelux, Bergen op Zoom, The Netherlands). The HPLC apparatus further consisted of a Separations high precision pump, model 300 (H.I. Ambacht, The Netherlands), provided with a Rheodyne sample injector, model 7175 (loop volume 20 μl ; Cotati, CA 94928, USA), and a Knauer differential refractometer, model 98.00 (Bad Homburg, Germany). Potassium biphthalate (25 mM, pH 4.3), containing 0.02% NaN_3 , was used as a mobile phase. The flow rate was 1 ml min^{-1} ; detector temperature was kept at 25 $^{\circ}\text{C}$ by a waterbath. The HPLC was connected with a Shimadzu Chromatopac C-R8A data processor (Kyoto, Japan).

Free amino acid content was determined according to ROSEN 1957. Shoot and roots of frozen fresh material were homogenized in demineralized water at 0 $^{\circ}\text{C}$ with an Ultra Turrax for 15 s (1 g in 10 ml; STUIVER & al. 1992). Homogenates were filtered through one layer of Miracloth, incubated in a boiling waterbath for 10 min and the filtrate was centrifuged at 30,000 g for 15 min (0 $^{\circ}\text{C}$). Free amino acid content of the supernatant was measured by ninhydrin colorimetric determination at 579 nm.

Analysis of the total sulfur content of shoot and roots was performed using a modification of the method as described by JONES 1995. Samples were dried at 120 $^{\circ}\text{C}$ for 18 h, pulverized by a Retsch microdismembrator (type MM2; Haan, Germany) and 50-100 mg of the samples was weighed into porcelain ashing trays. 50% $\text{Mg}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O}$ was added to saturate the material, after which it was dried in an oven at 100 $^{\circ}\text{C}$ overnight. The samples were ashed in an oven at 650 $^{\circ}\text{C}$ for 12 h. The residue was dissolved in 10 ml 20% aqua regia (50 ml HNO_3 and 150 ml HCl in 1 l distilled water) and the volume was adjusted to 100 ml with distilled water. One SulfaVer[®] 4 Reagent Powder Pillow (HACH, Permachem[®] reagents, Loveland, USA) containing BaCl_2 was added to 25 ml of the mixture and the turbidity was measured on a spectrophotometer (HACH DR/400V, Loveland, USA) at 450 nm.

Total nitrogen content of shoot and roots was determined with the Kjeldahl method according to BARNEIX & al. 1988. All nitrogen was reduced to NH_4^+ by boiling the tissue in concentrated H_2SO_4 in the presence of a catalyzer. NH_4^+ was determined colorimetrically at 410 nm after reaction with Nessler's reagent.

The organic N and S content was calculated by subtracting the inorganic N and S content from the total N and S content.

For measurements of the total thiol content, shoot and roots were extracted after STUIVER & al. 1992 and the content of total water-soluble non-protein sulfhydryl compounds was measured according to DE KOK & al. 1988. The relative growth rate (RGR) was calculated from the ln-transformed plant fresh weights (HUNT 1982).

Data were statistically analyzed using an unpaired Student's t-test and a F-test for linearity.

Results and Discussion

The uptake of gases by plant shoots is predominantly determined by both the diffusive conductance of the stomates and the mesophyll (internal) resistance (DE KOK & TAUSZ 2001). The uptake of H_2S by plant shoots appears to be largely dependent on the mesophyll resistance, *viz.* the rate of H_2S metabolism into cysteine and subsequently into other sulfur compounds (DE KOK & al. 1998, 2000, 2002, DE KOK & TAUSZ 2001). The maximum uptake rates of H_2S differ considerably among species (DE KOK & al. 2002).




Fig. 1. Kinetics of H_2S uptake in shoots of onion. Two-week-old seedlings were grown on a 25% Hoagland nutrient solution at 24°C for 10 days after which the H_2S uptake (JH_2S) was measured at various atmospheric H_2S levels. The mean transpiration rate (JH_2O) was $18.6 \pm 0.8 \text{ mmol g}^{-1} \text{ fresh weight h}^{-1}$. KH_2S and $\text{JH}_2\text{S}_{\text{max}}$ are expressed as $\mu\text{l l}^{-1}$ and $\mu\text{mol g}^{-1} \text{ fresh weight h}^{-1}$, respectively. Data represent the mean of 4 measurements with 12 plants in each ($\pm \text{SD}$). Repeating the experiment gave approx. the same results (1.38 and 1.04 for KH_2S and $\text{JH}_2\text{S}_{\text{max}}$, respectively).

Similar to observations with other species, the H_2S uptake by shoots of onion followed saturation kinetics with respect to the H_2S concentration and the saturation kinetics fitted well with the Michaelis-Menten equation. The maximum H_2S uptake rate ($\text{JH}_2\text{S}_{\text{max}}$) was $1.1 \mu\text{mol g}^{-1} \text{FW h}^{-1}$ and the H_2S concentration at which $\frac{1}{2}\text{JH}_2\text{S}_{\text{max}}$ was reached (KH_2S) was $1.5 \mu\text{l l}^{-1} \text{H}_2\text{S}$ (Fig. 1), which showed that onion had a rather high capacity for H_2S uptake and a higher KH_2S value when compared with other species (DE KOK & al. 2002). The ratio of H_2S deposition velocity to aqueous vapor efflux ($g(\text{H}_2\text{S})/g(\text{H}_2\text{O})$) was calculated to obtain insight into the factors determining H_2S uptake in the shoots of onion. The $g(\text{H}_2\text{S})/g(\text{H}_2\text{O})$ ratios remained constant and close to 1 at levels up to $0.3 \mu\text{l l}^{-1} \text{H}_2\text{S}$, which indicated that uptake of H_2S was limited by its diffusion through the stomates only and that the mesophyll (internal) resistance was close to zero (Fig. 1).

Onion is a rather slow growing species with a RGR of approx. 9.5% day⁻¹ (on a plant fresh weight basis at a day/night temperature of 24/18 °C) and appeared to be resistant to relatively high levels of H₂S (Table 1). The plant biomass production was hardly affected upon a two-week exposure to atmospheric H₂S levels up to 0.3 µl l⁻¹. Shoot/root ratio and dry matter content in shoot and roots were not affected upon H₂S fumigation (Table 1).

Table 1. The impact of H₂S exposure on growth of onion. Two-week-old seedlings were grown on a 25% Hoagland nutrient solution at 24 °C for two weeks and exposed to 0, 0.075, 0.15, 0.225 and 0.3 µl l⁻¹ H₂S. Data on fresh weight (g) and dry matter content (DMC) represent the mean of 6 and 3 measurements, respectively, with 6 plants in each (± SD). RGR (% day⁻¹) was calculated using the ln-transformed plant fresh weight over a two-week interval as described by HUNT 1982. There were no significant differences between treatments (P<0.05, Student's *t*-test).

	0 µl l ⁻¹ H ₂ S	0.075 µl l ⁻¹ H ₂ S	0.15 µl l ⁻¹ H ₂ S	0.225 µl l ⁻¹ H ₂ S	0.3 µl l ⁻¹ H ₂ S
Fresh weight					
Shoot	0.31± 0.07	0.32 ± 0.07	0.34 ± 0.05	0.31 ± 0.07	0.26 ± 0.05
Roots	0.12 ± 0.02	0.13 ± 0.03	0.13 ± 0.02	0.12 ± 0.02	0.11 ± 0.02
S/R ratio	2.55 ± 0.30	2.55 ± 0.32	2.55 ± 0.31	2.48 ± 0.21	2.36 ± 0.24
RGR					
Plant	9.5 ± 1.0	9.9 ± 1.4	10.5 ± 1.5	9.7 ± 1.2	8.7 ± 1.0
DMC					
Shoot	6.59 ± 0.65	6.39 ± 0.54	6.65 ± 0.80	6.51 ± 0.48	6.65 ± 0.58
Roots	4.46 ± 0.49	4.21 ± 0.35	4.19 ± 0.55	4.22 ± 0.50	3.99 ± 0.36

Exposure of onion seedlings to atmospheric H₂S resulted in an increased content of sulfate, thiols and total sulfur in the shoots. There was a slight increase in total nitrogen content of the shoot, whereas that of nitrate and amino acids was hardly affected. Metabolite content of the roots was not affected upon H₂S exposure. At 0.3 µl l⁻¹ H₂S sulfate and total sulfur contents in the shoot were increased with 6.9 and 17.7 µmol g⁻¹ FW, respectively, whereas the total nitrogen content in the shoot increased with 29 µmol g⁻¹ FW and the nitrate content was not significantly affected (Fig. 2). This demonstrated that in particular for sulfur the greater proportion of the increase in its content was due to an increase in the organic sulfur fraction. In unexposed plants the organic N/S ratio was around 41, predominantly representing the molar ratio of N/S in the proteins. However, upon H₂S exposure the organic N/S ratio decreased with the H₂S concentration, which likely has to be attributed to an increase in non-protein organic sulfur compounds (Fig. 3). It has been observed that onion has a sink capacity for sulfur which is metabolized into secondary sulfur compounds viz. alliins (RANDLE & al. 1993, 1999, HANEKLAUS & al. 1997). The molar ratio of N/S in alliins and its precursors is ≤ 2. The present data provide circumstantial evidence that a great part of the increase in total sulfur content upon H₂S exposure may be attributed to an accumulation of alliins or its precursors. The impact of H₂S exposure on alliin

synthesis will further be investigated. The experimental techniques will have to be optimized, since alliin compounds are rather unstable during extraction and subject to degradation.




Fig. 2. The impact of H_2S exposure on the sulfate, nitrate, thiol, amino acid, total sulfur and total nitrogen content of onion. Two-week-old seedlings were grown on a 25% Hoagland nutrient solution at 24 °C for two weeks and exposed to 0, 0.075, 0.15, 0.225 and 0.3 $\mu l\ l^{-1}$ H_2S . The sulfate content (a), the nitrate content (b), the total thiol content (c), the free amino acid content (d), the total sulfur content (e) and the total nitrogen content (f) of shoot (closed circles) and roots (open circles) are expressed in $\mu mol\ g^{-1}\ FW$. Data on sulfate, nitrate, free amino acid, total sulfur and total nitrogen represent the mean of 3 measurements with 36 plants in each (\pm SD). Total nitrogen and total sulfur contents increased linearly with the H_2S concentration ($P < 0.05$, F-test). Data on the total thiol content represent the mean of 4 measurements with 6 plants in each (\pm SD).




Fig. 3. The impact of H₂S exposure on the organic and inorganic N/S ratio of onion. The organic (closed circles) and inorganic (open circles) N/S ratio's (\pm SD) were derived from data presented in Fig. 2.

H₂S had a strong impact on thiol levels of the shoot. At 0.3 µl l⁻¹ H₂S the thiol content in the shoot was 2.8 times higher than that in the shoot of the control plants but at 0.075 µl l⁻¹ H₂S it was only 1.1 times higher, indicating that at low levels of H₂S the incorporation of cysteine (and glutathione) into other metabolic compounds, possibly alliin, was high and prevented a substantial accumulation of thiols (Fig. 2c). Future studies will include measurements on the composition of the thiol pool, since H₂S exposure may not only affect the content but also the composition of the thiol pool (BUWALDA & al. 1993, POORTINGA & DE KOK 1997, TAUSZ & al. 1998, WESTERMAN & al. 2000).

In *Brassica oleracea* there was a strong interaction between the uptake and metabolism of atmospheric H₂S and pedospheric sulfate. Upon H₂S exposure the sulfate uptake was downregulated, whereas the total sulfur content was hardly affected (DE KOK & al. 2000, WESTERMAN & al. 2001a). It appeared from our experiments that the sulfur metabolism in onion was differently regulated since the total sulfur content increased linearly with the H₂S concentration.

The interaction between atmospheric and pedospheric sulfur nutrition and its consequences for alliin synthesis will further be evaluated.

References

- BARNEIX A. J., COOPER H. D., STULEN I. & LAMBERS H. 1988. Metabolism and translocation of nitrogen in two *Lolium perenne* populations with contrasting rates of mature leaf respiration and yield. - *Physiol. Plant.* 72: 631 - 636.

- BUWALDA F., DE KOK L. J. & STULEN I. 1993. Effects of atmospheric H₂S on thiol composition of crop plants. - J. Plant Physiol. 142: 281 - 285.
- BLOCK E. 1992. The organosulfur chemistry of the genus *Allium* - implications for the organic chemistry of sulfur. - Angew. Chem. Int. Ed. Eng. 31: 1135 - 1178.
- DE KOK L. J. & TAUSZ M. 2001. The role of glutathione in plant reaction and adaptation to air pollutants. - In: GRILL D., TAUSZ M. & DE KOK L. J. (Eds.), Significance of glutathione in plant adaptation to the environment, pp. 185 - 200. - Kluwer Academic Publishers, Dordrecht.
- , BUWALDA F. & BOSMA B. 1988. Determination of cysteine and its accumulation in spinach leaf tissue upon exposure to excess sulfur. - J. Plant Physiol. 133: 502 - 505.
- , RENNENBERG H. & KUIPER P. J. C. 1991. The internal resistance in spinach leaves to atmospheric H₂S deposition is determined by metabolic processes. - Plant Physiol. Biochem. 29: 463 - 470.
- , STUIVER C. E. E. & STULEN I. 1998. Impact of atmospheric H₂S on plants. - In: DE KOK L. J. & STULEN I. (Eds.), Responses of plant metabolism to air pollution and global change, pp. 51 - 63. - Backhuys Publishers, Leiden.
- , WESTERMAN S., STUIVER C. E. E. & STULEN I. 2000. Atmospheric H₂S as plant sulfur source: interaction with pedospheric sulfur nutrition - a case study with *Brassica oleracea* L. - In: BRUNOLD C., RENNENBERG H., DE KOK L. J., STULEN I. & DAVIDIAN J.-C. (Eds.), Sulfur nutrition and sulfur assimilation in higher plants: molecular, biochemical and physiological aspects, pp. 41 - 56. - Paul Haupt Verlag, Bern.
- , STUIVER C. E. E., WESTERMAN S. & STULEN I. 2002. Elevated levels of hydrogen sulfide in the plant environment: nutrient or toxin. - In: OMASA K., SAJI H., YOUSSEFIAN S. & KONDO N. (Eds.), Air pollution and plant biotechnology, pp. 201 - 213. - Springer-Verlag, Tokyo.
- HANEKLAUS S., HOPPE L., BAHADIR M. & SCHNUG E. 1997. Sulphur nutrition and alliin concentrations in *Allium* species. - In: CRAM W. J., DE KOK L. J., STULEN I., BRUNOLD C. & RENNENBERG H. (Eds.), Sulphur metabolism in higher plants, pp. 331 - 334. - Backhuys Publishers, Leiden.
- HUNT R. 1982. Plant growth curves. - In: The functional approach to plant growth analysis, pp. 5 - 33. - E. Arnold Publishers, London.
- JONES J. B. Jr. 1995. Determining total sulphur in plant tissue using the HACH kit spectrophotometer technique. - Sulphur in Agriculture 19: 58 - 62.
- MAAS F. M., HOFFMANN I., VAN HARMELEN M. J. & DE KOK L. J. 1986. Refractometric determination of sulphate and other anions in plants separated by high-performance liquid chromatography. - Plant and Soil 91: 129 - 132.
- POORTINGA A. M. & DE KOK L. J. 1997. Uptake of atmospheric H₂S by *Spinacia oleracea* L. and consequences for thiol content and composition in shoots and roots. - In: CRAM W. J., DE KOK L. J., STULEN I., BRUNOLD C. & RENNENBERG H. (Eds.), Sulphur metabolism in higher plants, pp. 285 - 288. - Backhuys Publishers, Leiden.
- RANDLE W. M., BUSSARD M. L. & WARNOCK D. F. 1993. Ontogeny and sulfur fertility affect leaf sulfur in short-day onions. - J. Amer. Soc. Hort. Sci. 118: 762 - 765.
- , KOPSELL D. E., KOPSELL D. A. & SNYDER R. L. 1999. Total sulfur and sulfate accumulation in onion is affected by sulfur fertility. - J. Plant Nutr. 22: 45 - 51.
- ROSEN H. 1957. A modified ninhydrin colorimetric analysis for amino acids. - Arch. Biochem. Biophys. 67: 10 - 15.
- SCHNUG E. 1993. Physiological functions and environmental relevance of sulfur-containing secondary metabolites. - In: DE KOK L. J., STULEN I., RENNENBERG H., BRUNOLD C. & RAUSER W. E. (Eds.), Sulfur nutrition and assimilation in higher plants - regulatory, agricultural and environmental aspects, pp. 179 - 190. - SPB Academic Publishing, The Hague.
- STUIVER C. E. E. & DE KOK L. J. 2001. Atmospheric H₂S as sulfur source for *Brassica oleracea*: kinetics of H₂S uptake and activity of O-acetylserine (thiol)lyase as affected by sulfur nutrition. - Environ. Exp. Bot. 46: 29 - 36.

- , — & KUIPER P. J. C. 1992. Freezing tolerance and biochemical changes in wheat shoots as affected by H₂S fumigation. - Plant Physiol. Biochem. 30: 47 - 55.
- TAUSZ M., DE KOK L. J., STULEN I. & GRILL D. 1996. Physiological responses of Norway spruce trees to elevated CO₂ and SO₂. - J. Plant Physiol. 148: 362 - 367.
- , VAN DER KOOIJ T. A. W., MÜLLER M., DE KOK L. J. & GRILL D. 1998. Uptake and metabolism of oxidized and reduced sulfur pollutants by spruce trees. - In: DE KOK L. J. & STULEN I. (Eds.), Responses of plant metabolism to air pollution and global change, pp. 455 - 458. - Backhuys Publishers, Leiden.
- WESTERMAN S., DE KOK L. J., STUIVER C. E. E. & STULEN I. 2000. Interaction between metabolism of atmospheric H₂S in the shoot and sulfate uptake by the roots of curly kale (*Brassica oleracea*). - Physiol. Plant. 109: 443 - 449.
- , BLAKE-KALFF M. M. A., DE KOK L. J. & STULEN I. 2001a. Sulfate uptake and utilization by two different varieties of *Brassica oleracea* with different sulfur need as affected by atmospheric H₂S. - Phyton 41: 49 - 62.
- , STULEN I., SUTER M., BRUNOLD C. & DE KOK L. J. 2001b. Atmospheric H₂S as sulphur source for *Brassica oleracea*: consequences for the activity of the enzymes of the assimilatory sulphate reduction pathway. - Plant. Physiol. Biochem. 39: 1 - 8.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 2002

Band/Volume: [42_3](#)

Autor(en)/Author(s): Durenkamp M., De Kok Luit J.

Artikel/Article: [The Impact of Atmospheric H₂S on Growth and Sulfur Metabolism of *Allium cepa* L. 55-63](#)