Phyton (Austria) Special issue:	Vol. 42	Fasc. 3	(83)-(88)	
"Global change"				

A Comparison of Impacts of N Deposition Applied as NH₃ or as NH₄Cl on Ombrotrophic Mire Vegetation

By

I. D. LEITH¹⁾, C. E. R. PITCAIRN¹⁾, L. J. SHEPPARD¹⁾, P. W. HILL¹⁾, J. N. CAPE¹⁾, D. FOWLER¹⁾, S. TANG¹⁾, R. I. SMITH¹⁾ & J. A. PARRINGTON²⁾

K e y w o r d s : Ammonia, ammonium, dry N deposition, foliar nutrition, moorland vegetation, wet N deposition.

Summary

LEITH I. D., PITCAIRN C. E. R., SHEPPARD L. J., HILL P. W., CAPE J. N., FOWLER D., TANG S., SMITH R.I. & PARRINGTON J. A. 2002. A comparison of impacts of N deposition applied as NH_3 or as NH_4Cl on ombrotrophic mire vegetation - Phyton (Horn, Austria) 42 (3): (83)-(88).

The impacts of increased N deposition (wet; NH_4^+ and dry; NH_3) on ombrotrophic mire vegetation were compared in a two-year dose-response study. Vegetation from an ombrotrophic mire in SE Scotland was exposed in open-top chambers to a range of wet N deposition at 0 (deionised water), 8, 16, 32, 64 & 128 kg N ha⁻¹ y⁻¹ applied as NH₄Cl and dry N deposition as gaseous NH₃ at concentrations of 2, 6, 20, 50 & 90 μ g m⁻³. N concentrations in the foliage of all species showed a significant linear increase in response to dose when N was applied as wet deposited NH₄⁺ in both 1999 and 2000. Comparison of the linear regression for foliar N versus NH₄⁺ and NH₃ in the dose-response relationships showed significant differences (p<0.05) between the form of N applied and %N concentration for two species in 1999 (*C. vulgaris* and *P. commune*) but not in 2000. There was a significant increase in *C. vulgaris* shoot growth in the NH₃ treatments compared to NH₄⁺ treatments when expressed per unit deposited N. Effects of nitrogen deposited as gaseous NH₃ are generally greater than those of wet deposited NH₄⁺ on moorland species, when expressed per unit deposited NH₄⁺ on moorland species, when expressed per unit deposited NH₄⁺ on moorland species.

Introduction

Increases in both wet and dry N deposition have been reported throughout Europe in the past 30 years ('UNITED KINGDOM REVIEW' 1994). Enhanced N input

¹⁾Centre for Ecology and Hydrology Edinburgh, Bush Estate, Penicuik, Midlothian, EH26 0QB, U.K.

²⁾ Centre for Ecology and Hydrology Merlewood, Grange-over Sands, Cumbria LA11 6JU, U.K.

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at (84)

to semi-natural lowland heaths has caused changes in species composition, stimulation of secondary environmental stresses such as drought, frost and insect infestations (AERTS 1989, CARROLL & al. 1999, POWER & al. 1998). Studies of enhanced N deposition on semi-natural communities such as ombrotrophic mires and moorland have generally reported effects of wet deposition as NH_4^+ and NO_3^- (GORDON & al. 1999, CARROLL & al. 1999). Although critical loads for N, set to protect these sensitive communities are 5-10 and 10-20 kg N ha⁻¹ y⁻¹ respectively, the effects of increased dry N deposition as NH_3 or a comparison of NH_4^+ and NH_3 have not been widely reported (LEITH & al. 2001a).

In order to compare directly the effects of increased wet NH₄⁺ deposition (expressed as kg N ha⁻¹ y⁻¹) and dry NH₃ concentrations (μ g m⁻³) the deposition rates (kg N ha⁻¹ y⁻¹) of NH₃ must be determined. To estimate NH₃ deposition, deposition velocities, canopy resistances and their dependence on ambient NH₃ concentrations must be known. Effects research to date has generally assumed that the dry deposition rate of NH₃ increases linearly with ambient NH₃ concentration, i.e. that deposition velocity is constant with increasing concentration. FLECHARD & FOWLER 1998 have shown that deposition velocity decreases as NH₃ concentration increases in the range 0.1-2.0 μ g m⁻³, due to chemical processes at the vegetation surface. However, this concentration range is not sufficiently wide to cover the range of NH₃ conditions in the UK where ambient concentrations close to point sources often exceed 10 μ g m⁻³.

In a 2-year open-top chamber (OTC) study, the effects of N form (NH₃ versus NH₄⁺) on nutrient-limited, upland ombrotrophic mire vegetation (7 species) were compared in N dose-response experiments. The vegetation was exposed to a range of wet N deposition treatments (0-128 kg N ha⁻¹ y⁻¹) applied as NH₄Cl and dry deposition as gaseous NH₃ at a range of concentrations (2-90 μ g m⁻³) for 2 field seasons (1999 and 2000). These concentrations are typical of gaseous NH₃ concentrations found in close proximity to intensive livestock units (FOWLER & al. 1998). Rates of NH₃ dry deposition were measured over a range of concentrations in a flux chamber to quantify inputs to the vegetation and the dependence of deposition rate on the NH₃ concentration.

Material and Methods

Measurements of NH₃ deposition velocities were made over ombrotrophic mire vegetation (identical to that used in the dose response experiment), using an additional OTC with a purpose built flux chamber. A full description of methods and results is given in LEITH & al. 2001b. This enabled deposition rates (kg N ha⁻¹ y⁻¹) at a range of ambient NH₃ concentrations to be calculated. By applying the calculated deposition velocity to the measured NH₃ treatment concentrations the NH₃ deposition could be determined.

In the summer of 1998, a central square $(1.75 \text{ m x } 1.75 \text{ m}; \text{ area} = 3.06 \text{ m}^2)$ in each of eleven OTC's was excavated to a depth of 50 cm. The excavated areas were then back-filled with peat and individual plants of seven species, *Calluna vulgaris* (L.) Hull, *Narthecium ossifragum* (L.) Hudson, *Potentilla erecta* (L.) Rauschel, *Deschampsia flexuosa* (L.) Beauv., *Eriophorum vaginatum* (L.), *Polytrichum commune* Hedw. and *Molinia caerulea* (L.) Moench from an ombrotrophic mire (30

(85)

km SW of Edinburgh). All chambers were watered with de-ionised water using either an underground drip system or by overhead misting.

During the field season 1998, a dose-response experiment, exposing ombrotrophic mire vegetation to either a range of NH₃ concentrations or to NH₄⁺ (applied as NH₄Cl solutions) was established using eleven OTCs at the Centre for Ecology and Hydrology- Edinburgh. Six wet and five dry N deposition treatments were applied throughout the 1999 and 2000 field seasons. Six wet deposition mist treatments (de-ionised water, 8, 16, 32, 64, 128 kg N ha⁻¹ y⁻¹ as NH₄Cl) were applied over a 25-week period during both the 1999 and 2000 growing seasons at a rate of 12 mm per week (3 applications per week) from individual pressurized vessels via a spray droplet generator. The concentrations of the 6 wet deposition treatments were 0, 0.19, 0.38, 0.76, 1.52 and 2.85 mM NH₄Cl respectively. A full description of the application system is given in FOWLER & al. 1989. Four gaseous NH₃ treatments (6, 20, 50, 90 µg m⁻³) were generated by volatilization of a pumped 1 % aqueous NH₃ solution into the air stream of the OTC manifold. NH₃ treatments were applied continuously throughout the growing seasons of 1999 and 2000. NH₃ treatment chambers received background ambient NH₃ concentrations of approximately 1-2 µg m⁻³ throughout the winter nontreatment period. NH3 concentrations in each treatment chamber were continuously monitored using passive ammonia diffusion samplers (3 replicate samplers per chamber) changed every 4 weeks (SUTTON & al. 1998). The calculated depositions for 1999 and 2000 were 4, 9, 14, 34 and 48 kg N ha⁻¹ y⁻¹ and 4, 9, 30, 65, 80 kg N ha⁻¹ y⁻¹ respectively for mean measured treatment concentrations of 1.6, 4.0, 13, 30 and 73 in 1999 and 2, 8, 28, 55 and 93 µg NH₃ m⁻³ in 2000.

Above ground biomass of *N. ossifragum, P. erecta, D. flexuosa, E. vaginatum* and *M. caerulea* was determined by destructive harvest in both October 1999 and 2000. All above ground foliage was harvested and split into tillers/leaves and flowers. The tillers/leaves were then thoroughly washed with de-ionized water, to remove surface N deposition, dried at 80 $^{\circ}$ C then weighed. Sub-samples were ground, digested and analyzed for NH₄-N by the indophenol-blue method and for phosphate-P by the molybdenum-blue reaction (GRIMSHAW & al. 1989).

Ten individual shoots of *Calluna vulgaris* per treatment were randomly selected prior to shoot extension in March 1999 and 2000. Each shoot was marked with a waterproof solution, approximately 1 cm below the growing shoot tip. At regular intervals throughout the season the extension (mm) of each shoot was measured.

Results and Discussion

Although the higher N depositions in both the NH₃ and NH₄⁺ treatments exceeded the critical load for moorland vegetation of 10-20 kg N ha⁻¹ y⁻¹ (BOBBINK & al. 1996) there was no loss of species or visible foliar injury in the two years of the experiment. This could be due to the absence in this study of the additional secondary stress factors such as heather beetle and drought reported to be required to initiate species composition change (POWER & al. 1998).

For a given amount of N deposited the effects on moorland vegetation of NH₃ appear to be greater than for $\rm NH_4^+$. This is consistent with VAN DER EERDEN & al. 1990, who suggest that foliar NH₃ uptake is quicker than foliar and root assimilation of NH₄⁺ or NO₃⁻. It is suspected that the uptake pathway for NH₃ is through the stomata and not through the leaf surface, as DUECK & al. 1991 found little or no NH₃ absorbed to the leaf surface of either graminoid or ericaceous species.

There was a significant linear dose response relationship (p<0.05) between foliar N content in all plant species and NH₄-N applied in both 1999 and 2000. The N concentrations in *C. vulgaris* ranged from 1.70 % to 2.25 % and 1.4 % to 1.93 %,

(86)

going from the lowest and highest NH_4^+ treatments in 1999 and 2000 respectively. The decrease in % N concentration in the second year of treatment could be due to

Table 1. Linear regression analysis of foliar N with applied N as NH_4^+ or NH_3 in 1999 and 2000. The difference in response to NH_3 and NH_4^+ (*Prob.) was tested using a generalized linear regression model, where the slope and r^2 refer to mean values of 5(6) points. Data marked with an asterisk (*) are statistically significant at p < 0.05.

%N concentration	NH_4^+ (kg N ha ⁻¹ y ⁻¹)		NH_3 (kg N ha ⁻¹ y ⁻¹)		Prob. NH4 ⁺	
	Slope	r^2	Slope	r^2	Versus NH ₃	
1999						
C. vulgaris	0.004*	0.98	0.013*	0.96	0.003*	
M. caerulea	0.006*	0.79	0.090	0.70	0.572	
E. vaginatum	0.006*	0.91	0.001	0.01	0.314	
P. erecta	0.011*	0.76	0.017	0.61	0.505	
D. flexuosa	0.011*	0.90	0.022	0.61	0.116	
P. commune	0.010*	0.93	0.033	0.86	0.007*	
N. ossifragum	0.005*	0.78	0.008	0.43	0.536	
2000						
C. vulgaris	0.004*	0.98	0.006*	0.89	0.106	
M. caerulea	0.017*	0.88	0.004*	0.87	0.562	
E. vaginatum	0.006*	0.96	0.005*	0.95	0.167	
P. erecta	0.006*	0.71	0.008	0.91	0.530	
D. flexuosa	0.011*	0.88	0.012*	0.96	0.716	
N. ossifragum	0.007*	0.92	0.008*	0.93	0.542	
S. capillifolium	0.010*	0.94	0.027*	0.91	0.068	

growth dilution. PRINS & al. 1991 also found an increase in % N in *C. vulgaris* in response to increased NH_4^+ deposition. In the NH₃ treatment, only *C. vulgaris*, and *P. commune* showed a significant response to increasing N additions and also to the form of N applied (Table 1) in 1999. Both species exhibited greater N uptake when N was supplied as NH₃ rather than as NH₄⁺ on a per unit N basis. A linear dose response relationship was also found by PITCAIRN & al. 1995, 1998 for a number of species including *C. vulgaris*. This dose response relationship found for both N forms is further support for the use of foliar N concentration as a bio-indicator of N deposition.

A significant relationship between % N and deposition was found for all species except *P. erecta* in the NH_3 treatments in 2000. This increase in foliar N concentration with increased NH_3 concentrations has been found in a number of species (FANGMEIER & al. 1994). It is not clear why this significant linear relationship was not found in 1999, but was present in the second year of treatment (2000) for *M. caerulea, E. vaginatum, D. flexuosa* and *N. ossifragum*. In contrast to 1999, none of the species in 2000 showed significant differences between the form of N applied and % N in the foliage.

Enhanced N deposition increased shoot extension in *C. vulgaris* in each year of N treatment. This is consistent with the results of CARROLL & al. 1999. Analysis of individual N treatment growth data found that Gompertz curves could be fitted to all individual treatment shoot elongation curves. There was a significant difference (p<0.001) between the two N forms with NH₃ treatments increasing



Fig. 1. Shoot extension of *C. vulgaris* shoot extension from April to September 2000. Numbers in legend refer to N deposition rate (kg N ha⁻¹ y⁻¹).

shoot elongation compared to NH_4^+ treatments. This difference could be increased by excluding the high NH_3 treatment (61 kg N ha⁻¹ y⁻¹) and was not included as a separate component in the analysis, because shoot elongation was significantly decreased in this high NH_3 treatment. This could be attributed to changes in shoot morphology with increased lateral branching of the terminal shoot in the highest NH_3 treatments compared to other N treatments. Comparison of the dry weights of 3 cm length of *C. vulgaris* shoots showed a 35 % increase in weight in the highest NH_3 treatment compared to lowest. Generally, on an N input basis, N applied as dry NH_3 increased shoot extension more than wet NH_4^+ . The de-ionized water treatment had the smallest shoot extension compared to all the other N treatments irrespective of N form. This could be attributed to N deficiency as the foliar % N concentration decreased from 1.75 % in 1999 to 1.4 % in 2000.

This study has shown that NH_3 uptake is greater on a per unit of N deposited than NH_4^+ uptake and that the form of N applied influencec the assimilation of N and growth rates in *C. vulgaris*.

Acknowledgements

The UK Department for Environment, Food and Rural Affairs (Contract no. 1/3/94) and the Natural Environment Research Council are thanked for their financial support.

References

AERTS R. 1989. Aboveground biomass and nutrient dynamics of *Calluna vulgaris* and *Molinia caerulea* in a dry heathland - Oikos 56: 31 - 38.

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at (88)

- BOBBINK R. HORNUNG M. & ROELOFS J. G. M. 1996. In: Manual of methodologies and criteria for mapping critical levels/loads and geographical areas where they are exceeded. - UN-ECE Convention on Long-range Transboundary Air Pollution. - Federal Environmental Agency, Berlin.
- CARROLL J. A., CAPORN S. J. M., CAWLEY L., READ D. J. & LEE J. A. 1999. The effect of increased deposition of atmospheric nitrogen on *Calluna vulgaris* in upland Britain.- New Phytol.141: 423 - 431.
- FANGMEIER A., FANGMEIER A. H., VAN DER EERDEN L. & JÄGER H.-J. 1994. Effects of atmospheric ammonia on vegetation - A review. - Environ. Pollut. 86: 43 - 82.
- DUECK TH. A., VAN DER EERDEN L. J. & BEEMSTERBOER B. 1991. Nitrogen uptake and allocation by *Calluna vulgaris* (L.) Hull and *Deschampsia flexuosa* (L.) Trin. Exposed to ¹⁵NH₃. -Acta Bot. Neerl. 40: 257 - 267.
- FLECHARD C. R. & FOWLER D. 1998. Atmospheric ammonia at a moorland site. I: the meteorological control of ambient ammonia concentrations and the influence of local sources. -Quartery Journal Royal Meteorological Society. 124: 733 - 757.
- FOWLER D., PITCAIRN C. E. R., SUTTON M. A., FLECHARD C., LOUBET B., CAPE J. N. & MUNRO R. C. 1998. The mass budget of atmospheric ammonia in woodland within 1 km of livestock buildings. - Environ. Pollut. 102: S1, 343 - 348.
 - , CAPE J. N., DEANS J. D., LEITH I. D., MURRAY M. B., SMITH R. I. SHEPPARD L. J. & UNSWORTH M. H. 1989. Effects of acid mist on the frost hardiness of red spruce seedlings. - New Phytol. 113: 321 - 335.
- GORDON C, WOODIN S. J., ALEXANDER I. J. & MULLINS C. E. 1999. Effects of increased temperature, drought and nitrogen supply on two upland perennials of contrasting functional type: *Calluna vulgaris* and *Pteridium aquilinum*. - New Phytol. 142: 243 - 258
- GRIMSHAW H. M., ALLEN S. E. & PARKINSON J. A. 1989. Nutrient elements. In: ALLEN S. E. (Ed.), Chemical analysis of ecological materials (2nd ed.), pp. 81 - 159. - Blackwell Scientific Publications. Oxford.
- LEITH I. D., SHEPPARD L. J., PITCAIRN C. E. R, CAPE J. N. HILL P. W., KENNEDY V. H., SIM Y. T. SMITH R. I. & FOWLER D. 2001a. Comparison of efects of N applied as NH₃ or as NH₄Cl (aq) to moorland vegetation growing in open-top chambers. - Final report to department for environment, food and rural affairs. London, UK.
 - , , , , , , , & 2001b. Comparison of the effects of N deposition (NH₄Cl) and dry deposition (NH₃) ON UK moorland vegetation. -Water Air Soil Pollut. 130: 1043 - 1048.
- PITCAIRN C. E. R., FOWLER D. & GRACE J. 1995. Deposition of fixed atmospheric nitrogen and foliar nitrogen content of bryphytes and *Calluna vulgaris* (L.) Hull. - Environ. Pollut. 88: 193 - 205.
 - , LEITH I. D., SHEPPARD L. J. SUTTON M. A., FOWLER D., MUNRO R. C., TANG. S. & WISON D. 1998. The relationship between nitrogen deposition, species composition and foliar nitrogen concentrations in woodland flora in the vicinity of livestock farms. Environ. Pollut. 102 S1: 41 48.
- POWER S. A., ASHMORE M. R., COUSINS D. A. 1998. Effects of nitrogen additions on the stress sensitivity of *Calluna vulgaris*. - New Phytol. 138: 663 - 673.
- PRINS A. H., BERDOWSKI J. J. M. & LATUHIHIN M. J. 1991. Effect of NH₄-fertilisation on the maintenance of *C. vulgaris* vegetation. Acta Bot. Neerl. 40: 269 279.
- SUTTON M. A. TANG Y. S., MINERS B. P., COYLE M., SMITH R. I. & FOWLER D. 1998. Final report to DETR. - Results of national ammonia monitoring network. London, UK.
- UNITED KINGDOM REVIEW GROUP ON IMPACTS OF ATMOSPHERIC NITROGEN. 1994. Impacts of nitrogen deposition on terrestrial ecosystems. - London, UK; Department of the Environment.
- VAN DER EERDEN L. J., DUECK TH., ELDERSON J., VAN DODDEN H. F., BERDOWSKI J. J. M. & LATUHIHIN M. J. 1990. Effects of NH₃ and (NH₄)₂SO₄ deposition on terrestrial seminatural ecosystems on nutrient-poor sandy soils. - Report 90/06 (IPO) and 90/20 (RIN) Dutch Priority Programme on Acidification.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Phyton, Annales Rei Botanicae, Horn

Jahr/Year: 2002

Band/Volume: 42_3

Autor(en)/Author(s): Leith I. D., Pitcairn C. E. R., Sheppard L. J., Hill P. W., Cape J. N., Fowler D., Tang S., Smith R. I., Parrington J.

Artikel/Article: <u>A Comparison of Impacts of N Deposition Applied as NH3</u> or as NH 4Cl on Ombrotrophic Mire Vegetation. 83-88