

## First confirmed record of *Golovinomyces neosalviae* from Argentina

Uwe BRAUN & Maria HAVRYLENKO

**Abstract:** Braun, U. & Havrylenko, M. 2018: First confirmed record of *Golovinomyces neosalviae* from Argentina. Schlechtendalia **34**: 101–102.

In the 1990s, powdery mildew was collected on *Salvia officinalis* in Argentina and assigned to *Erysiphe biocellata* (= *Golovinomyces biocellatus*). The material concerned has been re-examined and identified as *Golovinomyces neosalviae*. This is the first confirmed record of this species from Argentina and possibly the first South American record at all.

**Zusammenfassung:** Braun, U. & Havrylenko, M. 2018: Erster bestätigter Fund von *Golovinomyces neosalviae* aus Argentinien. Schlechtendalia **34**: 101–102.

In den 1990iger Jahren wurde in Argentinien Echter Mehltau auf *Salvia officinalis* gefunden und zu *Erysiphe biocellata* (= *Golovinomyces biocellatus*) gestellt. Das Material wurde neu untersucht und als *Golovinomyces neosalviae* identifiziert. Dabei handelt es sich um den ersten bestätigten Nachweis dieser Art für Argentinien und wahrscheinlich auch für Südamerika überhaupt.

**Key words:** Powdery mildew, *Golovinomyces*, *Salvia officinalis*, new record, South America.

Published online 07 Aug. 2018

*Salvia officinalis*, native to the Mediterranean region, is a popular, frequently cultivated plant, used as ornamental garden plant and for culinary and medical purposes (Clebsch & Barner 2002). Powdery mildew on this host was previously assigned to *Erysiphe biocellata* Ehrenb. (= *Golovinomyces biocellatus* (Ehrenb.) Heluta) or *Erysiphe salviae* (Jacz.) S. Blumer (Amano 1986, Braun 1995, Braun & Cook 2012). Scholler et al. (2016) published a comprehensive revision of *Golovinomyces biocellatus* s. lat. and split it into *G. biocellatus* s. str., *G. monardae* (G.S. Nagy) M. Scholler, U. Braun & Anke Schmidt, *G. salviae* (Jacz.) M. Scholler, U. Braun & Anke Schmidt, and the new species *G. neosalviae* M. Scholler, U. Braun & Anke Schmidt with *Salvia officinalis* as type host and *S. officinalis* subsp. *lavandulifolia* and *S. fruticosa* as additional hosts. *G. neosalviae* is genetically clearly separated from *G. salviae* and morphologically readily distinguishable by its characteristic, quite distinct conidiophores, which are very long, often with a single or several short basal cells followed by a much longer cell. The basal septum is almost always distinctly elevated above the junction with the supporting hypha. Götz et al. (2018) described a new unusual branching pattern in conidiophores of *G. neosalviae*. A Mediterranean origin of this species can be postulated. Scholler et al. (2016) examined collections on *Salvia officinalis* from Germany and Switzerland and included several German collections in phylogenetic analyses. *Golovinomyces* on *Salvia officinalis* has been recorded from several European countries [Bulgaria, Czech Republic, France, Hungary, Italy, Switzerland (Amano 1986, Braun 1995, Bolay 2005), Croatia (Radisek et al. 2012)]. *Golovinomyces* infections on *S. officinalis* in Germany are undoubtedly not rare (Jage et al. 2010, Scholler et al. 2016, Götz et al. 2018). Non-European powdery mildew records on *S. officinalis* are from Argentina (Havrylenko 1997, Cabrera et al. 2010), New Zealand (Amano 1986) and South Africa (Gorter 1993). Voytyuk et al. (2009) reported *Golovinomyces biocellatus* on *Salvia fruticosa* from Israel. Most collections of *Golovinomyces* on *S. officinalis* and *S. fruticosa* seem to belong to *G. neosalviae*, but they need to be re-examined and confirmed.

Havrylenko (1997: 94–96, fig. 15 A–D) described and illustrated *Erysiphe* aff. *biocellata* on *Salvia officinalis* from Argentina characterized by having conidiophores with long foot-cells, up to 140 µm, and ellipsoid-doliiform conidia, 28–40 × 16–21 µm. A duplicate of this material has been re-examined [Argentina, Provincia de Río Negro, Bariloche, San Carlos de Bariloche, garden, on *Salvia officinalis*, May 1996, M. Havrylenko 540 (HAL 544 F)] and turned out to be identical with *G. neosalviae*. The conidiophores are either composed of a very long foot-cell followed by 1–3 short cells or shorter basal cells followed by a much longer cell and 1–3 shorter cells, and the basal septum is always elevated, 5–25(–30) µm. This is the first confirmation of

*Golovinomyces neosalviae* for Argentina and probably for South America in general. The description and illustration of *Golovinomyces biocellatus* on *Salvia officinalis* found in Corrientes, Argentina (Cabrera et al. 2010) does not agree with *G. neosalviae* (conidiophores consistently composed of a long foot-cell, on average  $66.5 \times 11.25 \mu\text{m}$ , followed by short cells, basal septum at the junction with the mother cell or only slightly elevated, conidia smaller, on average  $33.2 \times 16.5 \mu\text{m}$ ). The identity of this collection is unclear and requires molecular examinations. It might belong to *Golovinomyces salviae* or the infection could be caused by a plurivorous species of *Golovinomyces*.

## Literature

- Amano (Hirata), K. 1986: Host Range and Geographical Distribution of the Powdery Mildew Fungi. Japan Scientific Societies Press, Tokyo.
- Bolay, A. 2005: Les Oïdiums de Suisse (Erysiphacées). Cryptogamica Helvetica **20**: 1–173.
- Braun, U. 1995: The Powdery Mildews (*Erysiphales*) of Europe. G. Fischer Verlag, Jena.
- Braun, U. & Cook, R. T. A. 2012: Taxonomic Manual of the *Erysiphales* (Powdery Mildews). CBS Biodiversity Series **11**: 1–707.
- Cabrera, M. G., Vobis, G. & Alvarez, R. E. 2010: Powdery mildew on *Salvia officinalis* in Corrientes, Argentina. Mycosphere **1**(4): 289–291.
- Clebsch, B. & Barner, C. D. 2003: The New Book of Salvias: Sages for Every Garden. Timber Press, Portland.
- Gorter, G. J. M. A. 1993: A revised list of South African *Erysiphaceae* (powdery mildews) and their host plants. South African Journal of Botany **59**: 566–568.
- Götz, M., Itczak, E., Richert-Pöggeler, K. & Braun, U. 2018: A new morphotype of *Golovinomyces neosalviae* infecting *Salvia officinalis*. Plant Pathology & Quarantine **8**(2): 131–139.
- Havrylenko, M. 1997: *Erysiphales* de la region Andino-Patagonica. Thesis, Centro Regional Universitario Bariloche, Universidad Nacional del Comahue.
- Jage, H., Klenke, F. & Kummer, V. 2010: Neufunde und bemerkenswerte Bestätigungen von phytoparasitischen Kleinpilzen in Deutschland – *Erysiphales* (Echte Mehltäupilze). Schlechtendalia **21**: 1–140.
- Radisek, S., Ferant, N., Jakse, J. & Javornik, B. 2012: First Report of Powdery Mildew Caused by *Golovinomyces biocellatus* on Common Sage (*Salvia officinalis*) in Slovenia. Plant Disease **96**: 1065.
- Scholler, M., Schmidt, A., Siahaan, S. A. S., Takamatsu, S. & Braun, U. 2016: A taxonomic and phylogenetic study of the *Golovinomyces biocellatus* complex (*Erysiphales*, *Ascomycota*) using asexual state morphology and rDNA sequence data. Mycological Progress **15**(56): 1–13.
- Voytyuk, O., Heluta, V. P., Wasser, S. P. & Nevo, E. 2009: Biodiversity of the powdery mildew fungi (*Erysiphales*, *Ascomycota*) of Israel. Ganter Verlag, Ruggell.

## Addresses of the authors

Uwe Braun, Martin-Luther-Universität, Institut für Biologie, Bereich Geobotanik und Botanischer Garten, Neuwerk 21, 06099 Halle (Saale), Germany.  
(E-mail: uwe.braun@botanik.uni-halle.de)

Maria Havrylenko, Calle Francisco de Biedma 80, 8400 San Carlos de Bariloche, Río Negro, Argentina.  
(E-mail: mhavryl@gmail.com)

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Schlechtendalia](#)

Jahr/Year: 2018

Band/Volume: [34](#)

Autor(en)/Author(s): Braun Uwe, Havrylenko Maria

Artikel/Article: [First confirmed record of Golovinomyces neosalviae from Argentina](#)  
[101-102](#)