

# Societas entomologica.

„Societas Entomologica“ gegründet 1886 von *Fritz Rühl*, fortgeführt von seinen Erben unter Mitwirkung bedeutender Entomologen und ausgewiesener Fachmänner, erscheint monatlich zweimal im Kommissionsverlage von *Felix L. Dames*, Steglitz-Berlin.

## Journal de la Société entomologique internationale.

Toute la correspondance scientifique et les contributions originales doivent être envoyées aux héritiers de M. Fritz Rühl à Zurich V. Pour toutes les autres communications, annonces, collations &c. s'adresser à M. Felix L. Dames à Steglitz-Berlin.

Jährlicher Beitrag für Mitglieder 8 Mark = 8 Shillings = 10 Francs = 10 Kronen S. W. — Durch den Buchhandel bezogen 10 Mark. — Das Vereinsblatt erscheint zweimal im Monat (am 1. und 15.). — Insertionspreis für die 5-gespaltenen Peitscheln oder deren Raum 20 Pf. — Mitglieder haben in jedem Jahre 125 Zeilen frei für Inserate entomologischen Inhalts, Mehrzeilen werden mit 5 Pf. berechnet.

## Organ für den internationalen Entomologen-Verein.

Alle wissenschaftlichen Mitteilungen und Originalbeiträge sind an Herrn *Fritz Rühl's* Erben in Zürich V. zu richten. Alle geschäftlichen Mitteilungen, Inserate, Zahlungen etc. sind an *Felix L. Dames*, Steglitz-Berlin einzusenden.

## Journal of the International Entomological Society.

Any scientific correspondence and original contributions to be addressed to Mr. *Fritz Rühl's* Heirs, Zurich V. All other communications, insertions, payments &c. to be sent to Mr. *Felix L. Dames*, Sieglitz-Berlin.

### 57.92 Stictometeorus (68.7)

#### Description of a New Genus and Species of Meteorinae (Braconidae) from Cape Colony, South Africa.

By P. Cameron.

##### Stictometeorus gen. nov.

Pterostigma large, semicircular, the radius issuing from shortly behind its middle; 1<sup>st</sup> abscissa of radius straight, not oblique, twice the length of the 2<sup>nd</sup>, the radius not reaching to the middle of the space between the stigma and apex of the wing; transverse median nervure received beyond the transverse basal, the recurrent nervure in the base of the 2<sup>nd</sup> cubital cellule, which is short, twice wider at the base than at the apex; the 2<sup>nd</sup> abscissa of cubitus roundly curved backwards. Marginal cellule in hind wings narrowed towards the apex, undivided. Abdominal petiole slender, very slightly dilated towards the apex, half the length of the rest of the abdomen; the 2<sup>nd</sup> segment greatly enlarged, almost as long as the thorax, the other segments minute. Thorax stout, steeply sloped at base and apex, transverse, but with the sides rounded; parapsidal furrows absent. Antennae 18-jointed, placed above the middle of the face. Clypeus separated from the face by a furrow. Malar space large, but not so long as the eyes. Mandibles with a long, upper, sharply-pointed tooth. Occiput margined. Below the middle of the metanotum is a transverse keel.

The Genera of Meteorinae are reduced by Szpligeti to 2 — *Meteorus* and *Zemiotics*, distinguished by the latter having a deviated radial cellule in the hind wings. In this respect the present genus agrees with *Meteorus*, but it is readily separated from it by the steeply sloped base and apex of thorax, by the absence of parapsidal furrows, by the greatly enlarged 2<sup>nd</sup> ab-

dominal segment, by the basal abscissa of radius being longer than the 2<sup>nd</sup>, and by the much smaller 2<sup>nd</sup> cubital cellule, narrowed towards the apex. The type of the Genus looks more like one of the *Euphorinae* than of the *Meteorinae*.

##### Stictometeorus rufus sp. nov.

Rufous, the flagellum of antennae blackish above on the basal half, the lower part and the apex brownish, the mesonotum black; the legs and abdominal petiole paler, more yellowish coloured than the body; wings hyaline, the stigma and stigma blackish, the costa and nervures testaceous. ♀. Length 4 mm.

Face, front and vertex closely, rather strongly punctured, the temples more finely and weakly punctured, the clypeus still more finely punctured. Mesonotum and scutellum coarsely rugosely punctured; the metanotum still more coarsely punctured, almost reticulated, as are also the whole of the pleurae. Tegulae rufo-testaceous. Abdomen very smooth and shining.

### 57.83 (47.3)

#### Kleine Beiträge zur Kenntnis der Lepidopterenfauna Krestowkas (Gouv. Orel [spr.: Arjol]).

von B. Slevogt, Bathen.  
(Schluss.)

28. *Acherontia atropos* L. In einem Exemplare gesehen, aber nicht gefangen. 29. *Dilina tiliae* L. War 1908 dort so häufig, dass mitunter 10—15 Stück gleichzeitig die Lampe umschwärmen. Unterscheidet sich nicht unbedeutend von der kurfürstlichen Form. Während die hiesigen Tiere bei rotbrauner Grundfarbe der Vorderflügel immer grüne Flecke besitzen, die bei libauischen Exemplaren sehr ausgedehnt sind und fast die Grundfarbe verdriingen, haben die Falter aus

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Societas entomologica](#)

Jahr/Year: 1909

Band/Volume: [24](#)

Autor(en)/Author(s): Cameron Peter

Artikel/Article: [Description of a New Genus and Spezies of Meteorinae \(Braconidae\) from Cape Colony, South Africa. 9](#)