

fältig gefasst und in Wasserbehälter, in denen sich nebenbei bemerkt *Hyla meridionalis* Böttger, der südliche Laubfrosch, tummelt, geleitet, von welchen aus die Ländereien zu bestimmten Tageszeiten abwechselnd nach Gemeindeordnung berieselten werden. So kommt es, dass bereits hier die beiden östlich und westlich den Ort tangierenden Barrancos um diese Jahreszeit fast vollkommen ausgetrocknet waren und nur eine halbe Stunde weiter aufwärts beide einen kleinen dünnen Wasserfall aufweisen, der zwar bald zwischen Geröll verschwindet, aber doch eine üppige Vegetation, namentlich mit Polstern von Cinerarien, hervorzaubert. Derartige Stellen sind es vor allem, die den, der die landschaftlichen Reize einer Gegend nur nach seinen natürlichen, ungekünstelten Naturschönheiten einschätzt, für mancherlei Enttäuschungen auf den Canaren zu entschädigen vermögen.

Unsere Fonda hatte eine nach Nordosten freiliegende Veranda, von der man über den kleinen davorliegenden Garten, über ein Kartoffelfeld weit hinweg bis zu dem sterilen Höhenzug im Osten blicken konnte. Wenn auch ab und zu eine frisch geschlüpfte *Vanessa indica* v. *vulcanica* God., die wir schon in Las Palmas an Blumenbeeten bemerkt hatten, uns beim Frühstück störte und infolge der zu schonenden Anpflanzungen nur zu einer vergeblichen Jagd verleitete, so war es doch anderseits erstaunlich zu bemerken, dass abends an die von der Veranda weithinleuchtende Lampe, so gut wie kein Anflug von Insekten stattfand.

(Fortsetzung folgt.)

57.92 Braconidae (91.1)

On some New Bornean Species of Braconidae.

By P. Cameron, New Mills by Stockport, England.
(Continuation.)

Iphiaulax obesculus, sp. nov.

Entirely black, except the basal ventral segments, which are obscure white; wings hyaline, slightly obscured with fuscous, the nervures and stigma black; the 2nd transverse cubital nervure is very obliquely sloped; the recurrent nervure is received distinctly in front of the transverse cubital. Abdomen broad, ovate; closely, finely, but distinctly punctured; the 2nd segment with out an area or keel, the 3rd, 4th and 5th with a transverse furrow at the base. Head and thorax smooth and shining, covered with a short white pile, as are also the legs. ♀.

Length 3 mm, terebra 2 mm, longer than the abdomen.

Kuching, Borneo, July (John Hewitt, B. A.)

Hind legs thickened; the metatarsus thicker than the other joints, as long as the following 2 joints united; tibial spurs short, the longer as

long at the 3rd tarsal joint. Temples wide, obliquely narrowed. Front very slightly depressed; a narrow, shallow furrow down its centre. The pubescence is denser on the pleurae than it is on the mesonotum. Parapsidal furrows narrow but distinct. There is no transverse furrow ('suturiform articulation') on the 2nd abdominal segment.

Iphiaulax varireticulata, sp. nov.

Black, the prothorax except the lower part of the pleurae, the mesonotum, scutellum and the mesopleurae above the oblique basal furrow; red; the rest of the mesopleurae and the base of metapleurae tinged with red; wings hyaline, tinged with fulvous, the stigma and costa at apex, pale ochraceous, the radius, cubitus and the anal nervure from near the base of a paler ochraceous colour; the costa behind the stigma and the basal nervures black: the recurrent nervure received distinctly before the transverse cubital nervure. ♀.

Length 9 mm, terebra 5 mm.

Kuching, Borneo (John Hewitt, B. A.)

Face finely rugose, opaque; there is shallow furrow, widened below on its lower two-thirds; the sides of the face below are more strongly and distinctly punctured. Palpi black, densely covered with white pubescence. Front deeply furrowed down the middle, the furrow deepest in the middle, with the sides oblique. Metanotum opaque, densely covered with white pubescence; on the sides it is finely closely reticulated; the apex is depressed and bears some, more or less oblique, striae. Pro- and mesopleurae smooth, the metapleurae closely shagreened and densely pilose. Basal half of centre of 1st abdominal segment closely, rugosely punctured, the apex with a straight central and 2 or 3 irregular stout keels on either side; the base laterally with some oblique striae. Basal plate on 2nd segment opaque, closely longitudinally rugose; the keel extends shortly beyond the middle; the part bordering it stoutly widely reticulated, bordered on the outerside by an oblique keel; outside this keel it is obliquely striated, stoutly at the base, more weakly and closely towards the apex. The furrow is crenulated, curved and is without an apical branch; the other segments are closely longitudinally striated, more or less punctured and do not have furrows.

Iphiaulax saitis, sp. nov.

Black, the thorax rufous, broadly fuscous on the apex of metanotum laterally, the wings hyaline, a little tinged with fuscous, the nervures and stigma black; mandibles yellow, tinged with rufous above, the teeth black; palpi blackish, the apical 2 joints of maxillary pale fuscous. Head, metanotum, sternum and legs densely

covered with white pubescence; the pro- and mesonotum and their pleurae sparsely marked with shorter, white pile. Abdomen opaque, closely punctured, the central part of 1st abdominal segment irregularly rugosely punctured, more or less reticulated, the apex in the middle with a narrow longitudinal keel, with a smooth dilated part at the apex; the 2nd segment has the basal plate comparatively large, smooth, obliquely widened from the base to the middle, and obliquely narrowed from the middle to the apex; the keel extends close to the apex of the segment; the central part is closely irregularly reticulated and bounded by a slightly oblique keel, which is smooth and a little dilated at the apex; outside this keel the puncturation is closer than it is in the middle. The lateral furrow on the 1st segment is closely, obliquely striated. There is only one furrow, that between the 2nd and 3rd segments; it is closely, somewhat strongly striated; the apices of the 3rd and 4th segments have a narrow striated band; the apical 2 segments are pallid yellow. ♀ and ♂.

Length 5 mm; terebra 2 mm.

Kuching, Borneo (John Hewitt, B. A.)

The face is opaque and finely, closely rugosely punctured. The middle lobe of mesonotum is clearly separated at the base; the metanotum and metapleurae may be for the greater part black. The basal ventral abdominal segments are for the greater part pale yellow. There is a clearly defined furrow on the top half of the front commencing at the ocelli. Temples broad, distinctly, obliquely narrowed.

(to be continued.)

Der Traum des Entomologen.

Humoristisches Märchen in 3 Bildern von Ernst Breit,
cand. theol. et phil.¹⁾.

I.

Eines heissen Sommertages
Lag in schönster Mittagsstunde
Ein Entomologe schnarchend
Auf dem grünen Wiesengrunde.

Seiner müden Hand entfallen
War das Netz, das weissgewebte,
Das im Sicherheitsgefühl
Nun ein Weissling frech umschwebte.

Heimchen zirpten lust'ge Lieder
Zu der bunten Falter Tanze,
Und von Blum zu Blume schwirrten
Bienen rings im Sonnenglanze.

¹⁾ Anmerkung der Redaktion: Ganz ausnahmsweise wird einmal der Poesie in den Spalten der „Soc. entom.“ Raum gewährt. Der Verfasser ist eines Entomologen Sohn, nämlich des bei unseren Lesern in guter Erinnerung stehenden Herrn Julius Breit in Düsseldorf.

Doch es sah der müde Forscher
Nicht, was um ihn vorgegangen,
Denn den Schmetterlingsverderber
Hielt ein tiefer Traum umfangen.

Ihm erschien ein holdes Mädchen
Mit den Flügeln der Libelle,
Die in ihrem blauen Kleide
Schwebt ob der krystall'nellen Quelle.

Und sie sprach: „Komm mit, mein lieber,
Will dir etwas Schönes zeigen,
Doch du darfst es nicht verscheuchen,
Darum musst du stille schweigen.

Und vor allem auch dich hüten
Mit dem Netze dreinzuhauen,
Wie es die Entomologen
Immer tun auf Wiesenauen.

Diesmal musst du dich beherrschen,
Sonst kann es dir schlimm ergehen
Schauen, ohne zu begehrn,
Lerne, bester Freund, verstehen!“

Und der Forscher zupft am Barte,
Lächelt leis und voller Tücke,
Hm, da könnte es passieren,
Dass ein guter Fang mir glücke.

„Topp! Ich folge, Kind!“ So sprach er,
Packt zusammen seine Dosen,
Und in hoffender Erwartung
Folgte er der Ahnungslosen.

Schon war's dunkle Nacht geworden,
Als sie an die Wiese kamen,
Die von allen Seiten Wasser,
Wälder und Gebüsche umrahmen.

Und sie sprach: „Hier bleibe sitzen,
Schön's soll dein Auge schauen,
Aber hüte dich, du Mörder,
Mit dem Netze dreinzuhauen.“

II.

Elfen geben auf der Wiese
Heute einen grossen Ball,
Wollen fröhlich sein und tanzen
In des Mondes bassem Strahl.

Allerhand Insektenlarven
Haben sie sich angelegt,
Fühler, Flügel, drei paar Beine
Heut ein jedes an sich trägt.

Nun schlägt's 12. Aus düstren Wolken
Bricht ein Meer von mildem Glanz,
Und vom Wiesengrunde hebt es
Leise sich zum Geistertanz. —

Hier ein Elf als Raupertöter,
Schillernd goldig, grün und blau,
Lädt zum Tanz ein blondes Mädchen,
Kostümiert als Gelblingsfrau.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Societas entomologica](#)

Jahr/Year: 1909

Band/Volume: [24](#)

Autor(en)/Author(s): Cameron Peter

Artikel/Article: [On some New Bornean Species of Braconidae. 138-139](#)