

# Diplosmittia recisus spec. nov. from Peru

(Diptera, Chironomidae)

By Ole A. Sæther

## Abstract

The male of *Diplosmittia recisus* spec. nov. is described from Peru. It differs from *D. harrisoni* Sæther from the Eastern Lesser Antillean Islands, and *D. carinata* Sæther from Michigan, the two previously known members of the genus, in having only 9 flagellomeres and a strongly extended costa.

## Introduction

SÆTHER (1981) described four new genera from the British West Indies. Three of these genera later were described also from the southeastern parts of the U.S.A. (SÆTHER 1982). The remaining genus, *Diplosmittia*, recently was described from Michigan (SÆTHER 1985). Recently Dr. F. Reiss, Zoologische Staatssammlung, Munich, sent me a male imago collected in Peru. It proved to belong to a new species of *Diplosmittia* closely related to the previously described species.

## Methods and morphology

The general terminology follows SÆTHER (1980) with the exception that the apical spine of the gonostylus is termed the megaseta. In the figure of the male hypopygium the dorsal aspect is shown to the left, the ventral aspect and the apodemes to the right.

## *Diplosmittia* Sæther, 1981: 29, emended

Antenna with 9–13 flagellomeres, flagellomeres 2–8 or possibly sometimes only 2–5, and ultimate with sensilla chaetica. Costa not or barely to relatively strongly extended, with or without additional false vein extending beyond costa nearly to wing tip. Otherwise as in SÆTHER (1981: 29).

*Diplosmittia recisus* spec. nov.  
(Fig. 1)

Type material: Holotype, male, Station Koepke Panguana, about 260 m a. s. l., 9°37'S, 74°56'W, Peru, 17.2.1974, W. Kurz (Nr. 10), in the collection of Zoologische Staatssammlung, Munich.

### Diagnostic characters

Nine flagellomeres, costal extension moderately long (about 68 µm), LR<sub>1</sub> 0.36, LR<sub>2</sub> 0.36, LR<sub>3</sub> 0.43.




Fig. 1. *Diplosmittia recisus* spec. nov., male imago. — A. Wing. — B. Hypopygium.

**Etymology:** From Latin, *recisus*, cut back, cut short, referring to the reduced number of flagellomeres.

#### Description

##### Male imago ( $n = 1$ )

Total length 1.33 mm. Wing length 0.79 mm. Total length/wing length 1.68. Wing length/length of profemur 2.98. Coloration brown.

Head (as in *D. harrisoni*, SÆTHER 1981 fig. 13 A). Nine flagellomeres, ultimate 244  $\mu\text{m}$  long. AR 0.80. Temporal setae apparently consisting of 1 inner vertical. Clypeus apparently with 4 setae. Tentorium and stipes not measurable. Palp segments length (micrometers): 15, 19, 41, 45, 45.

Thorax (as in *D. harrisoni*, SÆTHER 1981 fig. 13 B). Antepronotum apparently with 1 lateral seta only. Dorsocentrals 4, prealars 3. Scutellum with 2 setae.

Wing (Fig. 1 A). VR 1.44. C extension 68  $\mu\text{m}$  long. Brachiolum with 1 seta.

Legs. Spur of front tibia 26  $\mu\text{m}$  long, spurs of middle tibia 15  $\mu\text{m}$  and 11  $\mu\text{m}$  long, of hind tibia 32  $\mu\text{m}$  and 15  $\mu\text{m}$  long. Width at apex of front tibia 26  $\mu\text{m}$ , of middle tibia 23  $\mu\text{m}$ , of hind tibia 23  $\mu\text{m}$ . Comb with 8 setae, 17–23  $\mu\text{m}$  long. Lengths (micrometers) and proportions of legs:

	fe	ti	ta <sub>1</sub>	ta <sub>2</sub>	ta <sub>3</sub>	ta <sub>4</sub>	ta <sub>5</sub>	LR	BV	SV	BR
P <sub>1</sub>	266	341	124	79	56	30	30	0.36	3.75	4.91	2.3
P <sub>2</sub>	323	356	129	79	58	30	32	0.36	4.07	5.25	2.7
P <sub>3</sub>	293	328	143	90	94	24	30	0.43	3.08	4.36	3.3

Hypopygium (Fig. 1B). Anal point with 6 lateral and basal setae, laterosternite IX with 2 setae. Phallapodeme 63  $\mu\text{m}$  long, transverse sternapodeme 54  $\mu\text{m}$  long. Virga 38  $\mu\text{m}$  long. Gonocoxite 92  $\mu\text{m}$  long. Gonostylus 49  $\mu\text{m}$  long, with lobe 57  $\mu\text{m}$  long, apical megaseta 6  $\mu\text{m}$  long. HR 1.91, HV 2.72.

### Acknowledgement

I am much indebted to Dr. F. Reiss, Zoologische Staatssammlung, Munich, for the type material.

### Literature

- SÆTHER, O. A. 1980: Glossary of chironomid morphology terminology (Diptera: Chironomidae). — Ent. scand. Suppl. 14: 1–51
- 1981: Orthocladiinae (Diptera: chironomidae) from the British West Indies, with descriptions of *Antillocladius* n. gen., *Lipurometriocnemus* n. gen., *Compterosmittia* n. gen. and *Diplosmittia* n. gen. — Ent. scand. Suppl. 16: 1–46
- 1982: Orthocladiinae (Diptera: Chironomidae) from SE U.S.A., with descriptions of *Pl budsonia*, *Unniella* and *Platysmittia* n. genera and *Atelopodella* n. subgen. — Ent. scand. 13: 465–510

Prof. O. Sæther  
Museum of Zoology  
University of Bergen  
N-5007 Bergen, Norway

# ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie, Supplement](#)

Jahr/Year: 1988

Band/Volume: [014](#)

Autor(en)/Author(s): Saether Ole A.

Artikel/Article: [Diplosmittia recisus spec. nov. from Peru \(Diptera,  
Chironomidae\) 45-47](#)