SPIXIANA S	Supplement 14 57-	-64 Münche	en, 15. Juli 1988	ISSN 0177-7424
------------	-------------------	------------	-------------------	----------------

Heterotrissocladius brundini spec. nov. from Norway

(Diptera, Chironomidae)

By Ole A. Sæther and Øyvind A. Schnell

Abstract

Heterotrissocladius brundini spec. nov. is described in all stages and both sexes. It apparently form the plesiomorphic sister species of *H. maeaeri* Brundin.

Introduction

LANGTON (1984: 88, fig. 28 a) illustrated a pupa of *Heterotrissocladius* with imbedded spines on segments VI-VIII and very large frontal warts. During an investigation of some acidified lakes in the southernmost part of Norway the junior author found all stages of apparently the same species. Later it was found also in western and northwestern Norway.

Methods and morphology

Morphological nomenclature follows SÆTHER (1980) with the exception that the apical spine of the gonostylus is termed the megaseta. The measurements are given as ranges followed by a mean when four or more measurements are made, followed by the number measured in parentheses (n).

The type material is kept at the Museum of Zoology, University of Bergen (ZMB).

Heterotrissocladius Spärck 1922, emended

Imago – Eyes completely naked, with a few microtrichia between marginal ommatids or pubescent, i. e. with microtrichia between all ommatids, with reticulation between central ommatids. Otherwise as in SÆTHER (1975: 3).

Pupa – Frontal warts weak to strongly developed. Frontal setae on frontal apotome (most species) or on prefrons (at least *H. latilaminus* Sæther). Caudolateral corners of segments VI-VIII occasionally with inner sclerotization resembling imbedded spines (*H. brundini* spec. nov), these strong on VII, weaker on VI, and only indicated on VIII. Otherwise as in SÆTHER (1975: 4).

Larva – Anterior parapods with apical claws grading over into basal hair-like spines anteriorly and on a common base. Otherwise as in $S_{\text{#THER}}$ (1975: 4).

Heterotrissocladius brundini spec. nov. (Figs 1-3)

Heterotrissocladius Pe 1, Langton 1984: 88

Type locality: Norway, Aust-Agder, Birkenes, Lake Repstadvatn.

Type material: Holotype, male, Lake Repstadvatn, Birkenes, Aust-Agder, Norway, 19/9/86, leg. H. Sægrov & Ø. A. Schnell, in coll. Mus. Zool. Univ. of Bergen (ZMB No. 115).

Paratypes: 5 males, 4 females, 2 male pupae, 147 pupal exuviae, 6 larval exuviae, as holotype; 5 pupal exuviae, Lake Store Hovvatn, Birkenes, Aust-Agder, Norway, 6/9/86, Ø. A. Schnell mature female pupa, Lake Jølster, Jølster, Sogn & Fjordane, Norway, 18/8/87, H. Sægrov; mature male pupa, Lake Litlebøvatn, Volda, Møre & Romsdal, Norway 26/7/86, Ø. A. Schnell; Other material (?): Male, Nordseter, Lillehammer, Oppland, Norway, 31/8/ 86, R. Bergo.

Diagnostic characters

The normal clypeus and cibarial pump combined with the moderately short acrostichals starting some distance from the scutal projection and the few setae on the wing membrane will separate the species from other *Heterotrissocladius* except *H. maeaeri* Brundin. However, the stipes is reduced, the wing slightly darker, the AR lower (0.7–1.1), the LR₁ higher (0.76–0.78) and the number of setae in cell r_{4+5} usually lower (7–18, except in one mature male pupa apparently with about 50 setae and another possible specimen with 45 setae, see remarks) in *H. brundini* spec. nov. The pupa can be separated from other members of the genus by means of the large frontal warts and the imbedded spines on segments VI-VIII. The larva can be distinguished by the narrowly separated two median teeth of the mentum with distinct lateral notches, the brownish black submentum which is conspicously darker than the surrounding areas of the head capsule, and the VM ratio of 1.1–1.5.

Etymology: Named in honour of Prof. emer. Lars Z. Brundin to his 80 year birthday and to signify his eminent contribution to chironomidology and lake typology where *Heterotrissocladius* has occupied a central position.

Description

Male imago (n = 7-9, except when otherwise stated).

Total length 3.93-4.22, 4.03 mm (6). Wing length 2.10-2.19, 2.14 mm (4). Total length/wing length 1.82-1.96, 1.90 (4). Wing length/length of profemur 2.44-2.55, 2.50 (4). Coloration brown with central parts of scutellum conspicuously pale. Wing pale yellowish brown.

Head. Eye with weak microtrichia between all ommatids. AR 0.92-1.05, 0.98 with one antenna of one specimen with an AR of 0.68. Ultimate flagellomere 424-484. 460 μ m long with one antenna of one specimen with ultimate flagellomere 381 μ m long. Pedicel 129-160, 139 μ m wide. Temporal setae 10-17, 14, including 3-5, 4 inner verticals; 2-5, 4 outer verticals; and 5-8, 7 postorbitals. Clypeus 123-138, 132 μ m wide; with 6-13, 10 setae. Cibarial pump, tentorium and stipes as in Fig. 1A, of doubtful specimen from Nordseter as in Fig. 1B. Tentorium 150-176, 162 μ m long, 35-55, 42 μ m wide. Stipes with reduced sclerotization in basal half and reduced median plate; 105-139, 120 μ m long; 19-38, 26 μ m wide (normal in possible specimen from Nordseter, see remarks). Palp lengths (micrometers): 29-45, 32; 41-62, 51; 109-149, 134; 86-103, 98; 106-138, 125. Third palpal segment with 2-4, 3(6) sensilla clavata at apex.

Thorax (Fig. 1 C). Antepronotum with 6–14, 11 setae. Dorsocentrals 15–25, 18; acrostichals 5–9, 7, starting some distance from scutal projection ranging from $\frac{1}{3}-\frac{1}{2}$ the length of scutum, longest acrostichal 38–56, 45 μ m long; prealars 7–9,8. Scutellum with 16–23, 19 setae.

Wing (Fig. 1D). VR 1.13–1.19, 1.16 (4). Brachiolum with 1–2, 2 setae; R with 11–15, 13 (6): R_1 with 4–11, 7 (6); R_{4+5} with 7–10, 9 (6); M_{1+2} with 0–8, 2 setae; other veins bare. Cell r_{4+5} with 7–18, 14 (5) setae, apparently with about 50 setae in mature male pupa from Volda; cell m_{1+2} with 0–3, 1 seta; cell an with 0–1, 0 setae. Squama with 18–21, 19 setae.

Legs. Spur of front tibia 47–56, 50 μ m long; spurs of middle tibia 35–38, 36 μ m and 21–34, 25 μ m long; of hind tibia 50–67, 62 μ m and 18–26, 22 μ m long.

Width at apex of front tibia 44–50, 47 μ m (4); of middle tibia 47–53, 49 μ m (4); of hind tibia 56–59, 56 μ m (4). Comb with 11–12, 11 setae; shortest seta 21–29, 24 μ m long; longest seta 53–70, 61 μ m long. Sensilla chaetica absent. Lengths (micrometers) and proportions of legs (n = 4):

	fe	ti	tal	ta ₂	ta ₃	ta ₄	ta ₅	LR	BV	SV	BR
Pl	849 - 860	88 4 - 920	681-717	341-382	245-263	179 -1 85	126 - 132	0.76-0.78	2.58-2.70	2.47-2.56	2.6-3.0
	857	905	696	360	258	181	127	0.77	2.66	2.53	2.8
Р ₂	801-860 837	777 - 848 822	406-418 412	227	167-173 170	131	114-120 116	0.49-0.52 0.50	3.10-3.27 3.22	3.88-4.12 4.02	2.0-2.8 2.4
^Р З	884 - 956	920 - 1004	568-6.21	311-335	251-263	155-167	126-132	0.60-0.62	2.80-2.94	3.15-3.28	3.8-4.8
	935	980	599	323	260	158	131	0.61	2.89	3.20	4.2



Fig. 1. Heterotrissocladius spp, male imago: A, C-E. H. brundini spec. nov; B, F. H. brundini or maeaeri (Brundin); A, B. Cibarial pump, tentorium and stipes; C. Thorax; D. Wing; E-F. Hypopygium.

Hypopygium (Fig. 1E). Anal point 53–70, 63 μ m long; tergite IX including anal point with 20–27, 23 setae; laterosternite IX with 4–8, 7 setae. Phallapodeme 117–139, 125 μ m long; transverse sternapodeme 106–116, 111 μ m (6) long. Virga very weak and inconspicuous, consisting of one or

two clusters of all together 6–7 spines; 11–15, 13 μ m (5) long. Gonocoxite 255–278, 268 μ m long; inferior volsella rounded, with long setae on distal side. Gonostylus 117–141, 127 μ m long; outer margin rounded; crista dorsalis long, relatively pronounced, rounded near apex; megaseta 10–15, 12 μ m long. HR 1.96–2.28, 2.12; HV 2.95–3.46, 3.18.

Female imago (n = 9-11, except where otherwise stated).

Total length 3.28-4.00, 3.61 mm. Wing length 2.11-2.33, 2.18 mm. Total length/wing length 1.51-1.80, 1.66. Wing length/length of profemur 3.01-3.24, 3.12 (8). Coloration as in male.

Head. Flagellomere lengths (micrometers): 83-109, 98; 45-60, 55; 53-64, 59; 56-69, 64; 120-156, 136. Pedicel 68-82, $73 \ \mu$ m wide. AR 0.44-0.55, 0.50. Temporals 9-14, 12; including 4-6, 5 inner verticals; 2-5, 3 outer verticals; and 2-5, 3 postorbitals. Clypeus 138-161, $151 \ \mu$ m wide; 1.85-2.22, 2.06 times as wide as pedicel; with 7-15, 11 setae. Tentorium 138-154, $146 \ \mu$ m (8) long; 18-26, $22 \ \mu$ m (8) wide. Stipes 86-149, $118 \ \mu$ m (8) long; 9-23, $16 \ \mu$ m (8) wide. Palp segments length (micrometers): 26-35, 31; 38-49, 43; 82-103, 94; 68-88, 78 (7); 98-135, 118 (7). Two specimens with fourth and fifth palp segments fused; $120-135 \ \mu$ m long. Third palpal segment with 3-5, 4 sensilla clavata at apex. Coronal suture incomplete; 64-98, $82 \ \mu$ m long.

Thorax. Antepronotum with 8-17, 11 lateral setae. Dorsocentrals 14-29, 19; acrostichals 4-7, 5; prealars 6-10, 8. Scutellum with 16-24, 19 setae.

Wing. VR 1.11–1.16, 1.13. Brachiolum with 2 setae; R with 12–25, 17; R_1 with 5–13, 8; R_{4+5} with 12–33, 24; M_{1+2} with 0–2, 1 setae; other veins bare. Cell r_{4+5} with 23–54, 36; cell m_{1+2} with 4–17, 13 (8) setae; cell an with 2–14, 6 (8) setae. Squama with 15–25, 21 setae.

Legs. Spur on front tibia 41–56, 50 μ m long; spurs on middle tibia 30–41, 37 μ m and 23–34, 28 μ m long; on hind tibia 60–71, 66 μ m and 18–26, 24 μ m long. Width at apex of front tibia 38–47, 43 μ m; of middle tibia 41–53, 48 μ m; of hind tibia 49–59, 53 μ m. Comb of 7–10, 8 setae; shortest seta 23–34, 29 μ m long; longest seta 49–67, 56 μ m long. Sensilla chaetica 1–7, 4 at 0.20–0.30, 0.28 (8) to 0.40–0.60, 0.53 on ta₁ of hind leg; apparently absent in mid leg. Lengths (micrometers) and proportions of legs (n = 7–9):

	fe	ti	tal	ta ₂	ta ₃	ta ₄	ta ₅	LR	BV	SV	BR
P ₁	669 - 753	753 - 848	548-624	274-321	194 - 217	137-151	99 - 131	0.72-0.76	2.68-2.92	2.47-2.63	2.1-3.1
	697	793	582	291	200	143	106	0.74	2.87	2.54	2.9
P2	690 - 807	717 - 807	359 - 384	191 - 227	142 - 167	95 - 120	85 - 114	0.49 - 0.51	3.23 - 3.47	3.76-4.10	2.1-3.0
	716	749	377	203	149	106	99	0.50	3.31	3.89	2.5
^Р з	784-896	884-992	548-624	293 - 335	227 - 263	135 - 161	99 - 134	0.61-0.65	2.78-2.98	2.88-3.10	2.9 - 3.9
	817	921	576	303	242	144	114	0.63	2.88	3.02	3.3

Abdomen. Number of setae on tergites I-VIII as: 33–53, 42; 38–46, 42; 26–40, 25; 24–44, 34; 24–38, 31; 19–35, 28; 19–35, 24; 9–26, 16. Number of setae on sternites I-VIII as: 0; 3–7, 5; 5–10, 7; 8–17, 11; 12–20, 15; 14–24, 17; 15–23, 18; 43–71, 56.

Genitalia (Fig. 2). Gonocoxite with 13–20, 16 setae. Tergite IX well divided, with 20–31, 25 setae. Cercus 114–142, 133 μ m long. Seminal capsule 76–101, 86 μ m (8) long; excluding 11–26, 17 μ m long neck; 64–79, 71 μ m wide. Notum 105–161, 132 μ m long.

Pupa (n = 11, except when otherwise stated)

Total length 3.76–4.78, 4.31 mm (9). Length of thoracic horn/length of anal macrosetae 1.10–1.50, 1.34. Thorax of exuviae pale brownish grey, abdomen nearly transparent.

Cephalothorax. Frontal warts (Fig. 3 A) conspicuous, 82-117, $97 \ \mu m$ high; 35-53, $40 \ \mu m$ (10) wide at base. Frontal setae on frontal apotome; 59-83, $73 \ \mu m$ (8) long. Postorbitals 59-73, $66 \ \mu m$ (4) and $41-56 \ \mu m$ (2) long. Median antepronotals 117-147, $140 \ \mu m$ (9) and 103-132, $120 \ \mu m$ long; lateral antepronotals 73-88, $80 \ \mu m$ and 0-44, $9 \ \mu m$ long (when absent reduced to setal mark). Thoracic horn (Fig. 3 B) 290-396, $349 \ \mu m$ long; 44-70, $54 \ \mu m$ (9) wide; 5.44-8.00, 6.57 (9) times as long as wide. Anterior dorsocentral 44-88, $65 \ \mu m$ long; $Dc_2 \ 53-117$, $88 \ \mu m$ long; $Dc_3 \ 32-129$, $68 \ \mu m$ long;

©Zoologische Staatssammlung München;download: http://www.biodiversitylibrary.org/; www.biologiezentrum.at



Fig. 2. *Heterotrissocladius brundini* spec. nov, female genitalia: A. Dorsal view; B. Ventral view; C. Lateral view; D. Lobes of gonapophysis VIII (DmL dorsomesal lobe; VlL, ventrolateral lobe; ApL, apodeme lobe).

Dc₄ 64-88, 78 μ m long. Distance between Dc₁ and Dc₂ 18-67, 42 μ m; between Dc₂ and Dc₃ 15-53, 28 μ m; between Dc₃ and Dc₄ 23-82, 58 μ m. Prealar observed in one specimen, 18 μ m long. Wing sheath with fine marginal lines.

Abdomen (Fig. 3 C, D). Shagreen absent on tergite I (T I) and T IX, weak median on T II, more extensive and stronger posterior on TIII–VI, weak anteriomedian on T VII–VIII. Sternites I (S I) and IX bare, S II–III with weak anterior and median shagreen, S IV–VII with weak anterior shagreen and a few posterior spinules as S VI. Pedes spurii A present on sternites IV-VII and occasionally VIII (3 individuals). Pedes spurii B wider than high. Caudal hooklets 20-40, 32 on T II. Sternite VIII of male with 19-33, 27 (7) posterior spines. Caudolateral corners of segments VI–VIII with imbedded spines, distinct on VII, only indicated on VI and VIII. L setae on segments I–VIII as: 1-2 (4): 3:3:3:3:3-4, 3:5;3 lamelliform on VII, 4 lamelliform on VIII, occasionally also fifth L seta on VIII slightly lamelliform. Anal lobe 269-337, $302 \,\mu$ m long; with 19-25, 22 setae in fringe; anal macrosetae 234-293, $261 \,\mu$ m long. Genital sac of male overreaching anal lobe by 88-124, $109 \,\mu$ m (4).

Larva (n = 6, except when otherwise stated)

Total length about 6 mm (larval exuviae). Head capsule length $447-502 \ \mu m$ (3).

Head. Antenna as in Fig. 3 E. Lengths of antennal segments (micrometers): 82-88, 86; 35-41, 36; 3-4, 4; 18-22, 19 (4); 6-9, 7 (4); 4-6, 4 (4). AR 1.07-1.16, 1.12 (4). Basal antennal segment 18-21, 20 μ m wide; 4.05-4.32, 4.24 times as long as wide; distance from base to annular organ 9-18, $15 \ \mu$ m; to distal mark of seta 21-35, $31 \ \mu$ m; blade at apex 50-64, 50 μ m long; accessory blade 7-9, $9 \ \mu$ m long. Subapical style of second segment 6-9, $7 \ \mu$ m (5) long. Labrum and epipharyngeal areas as in



Fig. 3. *Heterotrissocladius brundini* spec. nov, pupa: A. Frontal apotome; B. Thoracic horn; C. Tergites of male; D. Sternites of male. – larva: E. Antenna; F. Labrum and epipharyngeal area; G. Mandible; H. Maxilla; I. Mentum.

Fig. 3 F. S I broad, finely plumose. Labral lamellae rounded apically. Median spine of pecten epipharyngis narrower than lateral spines. Premandible 79–85, 83 μ m (5) long. Mandible (Fig. 3G) 158–176, 168 μ m (4) long. Maxilla as in Fig. 3 H. Mentum (Fig. 3 I) with two close median teeth each with distinct lateral accessory tooth, width of one median tooth including accessory tooth 18–21, 19 μ m; ventromental plates 21–26, 25 μ m (4) wide; 1.08–1.50, 1.32 times as wide as one median tooth including accessory tooth. Postmentum distinctly darker than surrounding areas of head capsule; 208–229, 222 μ m long.

Abdomen. Procercus 38–47, 43 μ m high; 29–32, 31 μ m wide; anal setae 647–747, 706 μ m (5) long. Supraanal seta 234–264, 242 μ m long; 0.32–0.35, 0.34 (5) times as long as anal setae. Anal tubules and posterior parapods not measurable.

Remarks

A male imago from Nordseter near Lillehammer is not included in the type material. Judging from the leg proportions and the antennal ratio the specimen belong to *H. brundini*. The normally sclerotized stipes (132 μ m long, Fig. 1B) and the more numerous setae on the wing membrane (45 setae in cell r₄₊₅, 7 setae in each of m₁₊₂ and m₃₊₄), however, indicate that it belongs to *H. maeaeri*. The hypopygium (Fig. 1F) could belong to any of the two species. Associated pupal material is needed to decide whether the stipes and the wing setation is more variable within *H. brundini* or whether the specimen represents *H. maeaeri* or an additional new species.

The pupa described by Langton (1984: 88) do not quite fit *H. brundini*. The thoracic horn of *Heterotrissocladius* Pe I of Langton is wider (length/width 4.0-5.1), and there are more numerous setae in the anal lobe fringe (26-30). These variations, however, probably is not more than can be expected between various populations.

Ecology and distribution

H. brundini has been found in two acidified lakes in Southern Norway, two lakes in the provinces of Møre & Romsdal and Sogn & Fjordane, and in Lake Assynt in the Sutherland District of Scotland. It could conceivably be an indicator species of oligotrophic acid lakes.

Systematics

SÆTHER (1975: 57-62, fig. 15) does a phylogenetic analysis of *Heterotrissocladius* and related genera. In the scheme of argumentation *H. brundini* will be synapomorphous for trends 57, 58, 59, 60; 9, 10, 11; and 51, 52, showing that the species is a good *Heterotrissocladius*. It is symplesiomorphous for trends 41 and 44, apomorphous for trends 42 (reduced acrostichals) and 43 (reduced number of setae on the wing membrane). The last two trends show parallelisms between the *subpilosus* and the *maeaeri* groups. However, there are no pearl rows on the pupal wing sheaths (trend 37), only 4 filamentous setae on segment VIII of the pupa (trend 39), and no setae dorsad of the genital sac (trend 40), showing that the species belong with the *maeaeri* and *marcidus* groups. However, the pedes spurii A which may occur on sternite VIII and the fifth L seta which in some specimen is somewhat filamentous, indicate a plesiomorphic position within these groups.

The species is synapomorphous with *H. maeaeri* for trends 32 (short acrostichals) and 35 (distribution of L setae in pupa). It is plesiomorphous for trends 33 (PSB), 34 (PSA), 36 (median teeth of mentum), and apomorphous for trends 30 (setae on male tergite IX) and 31 (AR of larva), indicating a placement in the *marcidus* group. However, trends 30 and 31 are not very significant, representing continuous variation and probably good only as specific autapomorphies. Also several additional synapomorphies between *H. maeaeri* and *H. brundini* can be found in the male hypopygium in the shape of the inferior volsella, the inner margin of the gonocoxite and in the shape of the gonostylus. We feel relatively certain about the phylogenetic position of the species as the sister species of *H. maeaeri*. There are few problems fitting the imagines and the pupa within the concept of the *maeaeri* group (SÆTHER 1975: 22); the stipes of the imagines and the pedes spurii B of the pupa have to be deleted as distinguishing characters. However, the larva will key to *H. marcidus* (Walker) in the larval key (SÆTHER 1975: 9) because of the two median teeth of the mentum. The median teeth, however, are closer together than in other described species except for *Heterotrissocladius* sp. E from Lake Superior (SÆTHER 1975: 55) which is mentioned as intermediate between the *marcidus* and *maeaeri* groups. *Heterotrissocladius* sp E almost certainly is close to *H. brundini*. A redefinition of the *maeaeri* group to include larvae with median teeth close together and distinct accessory teeth will make a separation between the *maeaeri* and *marcidus* groups still feasible.

The imbedded spines found in the pupa is nearly unique within the orthoclads, otherwise found in some members of Zalutschia Lipina only. SÆTHER (1976) did not attempt to classify Oliveridia Sæther, Hydrobaenus Fries, Trissocladius Kieffer and Zalutschia combined as a monophyletic group. According to SÆTHER (1977: 82), however, the genera apparently form a monophyletic unit based on the shape of the female gonapophysis VIII. This trend, however, is rather insecure as similar gonapophyses occur elsewhere including in the Heterotrissocladius group of genera. It thus is possible that Zalutschia is more closely related to the Heterotrissocladius group than to Hydrobaenus, and that the imbedded spines represent an underlying synapomorphy occurring in some, bot not all members of both Zalutschia and Heterotrissocladius. We have observed indication of imbedded spines also in some exuviae of H. marcidus. On the other hand the imbedded spines merely represent a strengthening of the paratergital margin and could easily be a result of parallel selection.

The larvae of *Heterotrissocladius* all have hair-like spines at base of the anterior parapods similar to those illustrated by STRENZKE (1950 fig. 11) for *Paraphaenocladius* Thienemann, but not found in *Parametriocnemus* Goetghebuer or apparently in the same form in other orthoclads. However, other synapomorphies seem to confirm the relationship shown by SÆTHER (1975 fig. 15) and these spines must be regarded either as secondarily reduced in *Parametriocnemus* or as an underlying synapomorphy for the whole group.

Acknowledgements

We would like to thank Laboratory for Freshwater Ecology and Inland Fisheries Bergen for financial support. We also are indebted to Mr. Tien van Trieu for doing the illustrations and to Mrs. N. Waagsberg for typing the manuscript.

Literature

LANGTON, P. H. 1984: A key to pupal exuviae of British Chironomidae. – P. H. Langton, March, Cambridgeshire, 324 pp

SÆTHER, O. A. 1975: Nearctic and Palaearctic *Heterotrissocladius* (Diptera: Chironomidae). – Bull. Fish. Res. Bd Can. 193: 1–67

- 1976: Revision of Hydrobaenus, Trissocladius, Zalutschia, Paratrissocladius, and some related genera (Diptera: Chironomidae).
 Bull. Fish. Res. Bd Can. 195: 1–287
- -- 1977: Female genitalia in Chironomidae and other Nematocera: morphology, phylogenies, keys. Bull. Fish. Res. Bd Can. 197: 1–209
- -- 1980: A glossary of chironomid morphology terminology (Chironomidae: Diptera). Ent. scand. Suppl. 15: 1–51

STRENZKE, K. 1950: Systematik, Morphologie und Ökologie der terrestrichen Chironomiden. – Arch. Hydrobiol. Suppl. 18(2): 207–414

> Prof. Ole A. Sæther Øyvind Schnell Museum of Zoology University of Bergen N-5007 Bergen, Norway

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Spixiana, Zeitschrift für Zoologie, Supplement

Jahr/Year: 1988

Band/Volume: 014

Autor(en)/Author(s): Saether Ole A., Schnell Oyvind A.

Artikel/Article: <u>Heterotrissocladius brundini spec. nov. from Norway</u> (Diptera, Chironomidae) 57-64