

| | | | | | |
|----------|---|---|---------|-----------------------|----------------|
| SPIXIANA | 7 | 2 | 105–113 | München, 1. Juli 1984 | ISSN 0341-8391 |
|----------|---|---|---------|-----------------------|----------------|

Brachyura of the coast of Southern Spain

(Crustacea, Decapoda)

By J. Enrique García Raso

Dept. Zoology, Fac. Science, Univ. Málaga, Spain

Abstract

During a study of the Crustacea Brachyura in the 0 to 310 m depth range on the coast of Southern Spain, from 1978 to 1981, eighty three species were taken. Data is given in this paper on the vertical distribution of each species and type of substrate where each species was found. Information is included on the presence of ovigerous females and other biological data. The record of *Palicus caronii*, *Euchirograpsus liguricus*, *Heterocrypta maltzami*, *Asthenognathus atlanticus* and *Pisa carinimana* are of particular interest while the capture of *Brachynotus atlanticus* represent the first record of this species from the Mediterranean Sea and Europe.

Introduction

The Brachyura are perhaps the best known Decapodan group on African and European coasts as evidenced by the huge number of works already published about these areas, whether in the form of monographic studies, such as those of BARNARD (1950), CAPART (1951), MONOD (1956), MANNING and HOLTHUIS (1981), NOBRE (1936), INGLE (1980) and CHRISTIANSEN (1969) on the Eastern Atlantic; or those of PESTA (1918), BACESCU (1967) and ZARIQUIEY (1968) on the Mediterranean Sea; or by the large number of smaller but none the less important works expeditions and faunal inventories too numerous to mention here.

The Southern coast of Spain, however, in spite of its important geographical location between Africa and Europe has not been investigated in detail and references to species captured in this area are very few.

The present study was undertaken to remedy this state of affairs. It represents the first faunal study of the area and can also be considered a complement to works such as MANNING and HOLTHUIS (1981) concerning West African Brachyuran crabs.

Area of Investigation and Methods

The coastal region of the province of Málaga, extending 129 Km. westwards from Maro to San Pedro de Alcántara and located on the northern margin of the Sea of Alboran in the Western Mediterranean, was chosen for this investigation (Fig. 1). Samples were collected from the shore to 310 metres depth but the greater part of the sampling was carried out between 0 and 100 m. In addition a few samples were taken at greater depths.

Specimens were obtained by methods varying according to the type of substrate and depth. In general trawling was employed for soft substrates, fish traps or free diving for hard substrates, while supplementary set netting was also carried out on the sea floor.

The duration of the study was four years, from 1978 to 1981 inclusive.




Fig. 1. Map of study area.

Results

The results of this study are given in tabular format for a better and more rapid comprehension. The table gives the list of species found in the Málaga region together with a record of the depth of occurrence and substrate type where each species was found. Data on the presence of ovigerous females in samples collected in this study is given on a monthly basis for a annual cycle.

Included in the table is information, for comparative purposes, on the vertical distribution range and presence of ovigerous females in Spanish waters. This information has been compiled from data of some authors who are designated by numbers in brackets in the table.

The numbers shown in brackets correspond to the following authors: (1) SANTAELLA (1974); (2) FOREST (1974); (3) ZARIQUIEY (1968); (4) FOREST (1965); (5) PASTORE (1972); (6) MANNING and HOLTHUIS (1981); (7) WILLIAMS and WIGLEY (1977); (8) MIRANDA y RIVERA (1921, 1933); (9) FOREST et GUINOT (1966); (10) PESTA (1918); (11) TÜRKAY (1976a); (12) POWERS (1977); (13) CRESPO, REY & CAMIÑAS (1976); (14) TÜRKAY (1976b); (15) MAURIN (1968) taken of MANNING and HOLTHUIS (1981); (16) BOUVIER (1940); (17) MONOD (1956); (18) LAGARDERE (1973); (19) INGLE (1980); (20) CHRISTIANSEN (1969); (21) CAPART (1951); (22) FOREST (1967); (23) GONZALEZ G. (1978); (24) CROSNIER (1970); (25) O. DE BUEN (1925); (26) FOREST et ZARIQUIEY (1964); (27) FOREST (1978); (28) MASSUTI (1967); (29) RAMADAN and DOWIDAR (1972); (30) FOREST et GUINOT (1958); (31) DRACH et FOREST (1953); (32) FOREST et GANTES (1960); (33) SCHMITT, McCAIN and DAVIDSON (1973); (34) TÜRKAY (1975); (35) ŠTEVČIĆ (1973); (36) LEWINSOHN (1976); (37) FROGLIA and MANNING (1982).


The signs employed to describe the type of substrate are: S-sand, V-mud, SV-muddy sand, C-shell, R-rocky, G-caves and fissures, F-phanerogams, A-algae, Co-coralliferous, SR-sand and rocks, SA-sand and algae, SVR-muddy sand and rocks, SG-coarse sand, SGA-coarse sand and algae, SF-fine sand.

The data on ovigerous females in Spanish waters compiled, except in the one case referred to in the table, from ZARIQUIEY (1968) and FOREST (1965) is represented by broken lines. The information on egg carrying females collected in this study is represented by solid lines.

| SPECIES | General | VERTICAL RANGE Spain | Málaga | TYPES OF SUBSTRATES Málaga | OVIGEROUS FEMALES IN SPAIN | | | | | | | | | |
|---|--|-----------------------------|-------------------------|-------------------------------|----------------------------|---|---|---|---|---|---|---|---|---|
| | | | | | J | F | M | A | M | J | J | A | S | O |
| Fam. DROMIOIDAE | | | | | | | | | | | | | | |
| De Haan, 1833 | | | | | | | | | | | | | | |
| Dromia personata (Linnaeus, 1759) | 2(1)(5)-300(2) | 10(3)-48(4) | 2-82 | R,S,G. | | | | | | | | | | |
| Fam. HOMOLIOIDAE | | | | | | | | | | | | | | |
| De Haan, 1839 | | | | | | | | | | | | | | |
| Homola barbata (Fabricius, 1793) | 10-30(6)-685(7) | 40(3)-250(8) | 50-110 250(8) | VR,V,SV,?R | | | | | | | | | | |
| Fam. ORIPIPOIDAE | | | | | | | | | | | | | | |
| Mac Leay, 1838 | | | | | | | | | | | | | | |
| Medorippe lanata (Linnaeus, 1767) | 10(9)(10)-1038(11) | 40-100(3) | 11-67 200(8) | V | | | | | | | | | | |
| Ethusa mascaronne (Herbst, 1785) | Littoral-82(12) 2-650(16) | Shallow water- 75(3) | 10-81 Max.18-30 | V,SV,S | | | | | | | | | | |
| Fam. CALAPPIODAE | | | | | | | | | | | | | | |
| Da Haan, 1833 | | | | | | | | | | | | | | |
| Calappa granulata (Linnaeus, 1758) | 13(14)-400-700(15) | 30(3)-150(8)(3) | 3-102 150(8) | S,SV,V,C | | | | | | | | | | |
| Fam. LEUCOSIIDAE | | | | | | | | | | | | | | |
| Samouelle, 1819 | | | | | | | | | | | | | | |
| Ilia nucleus (Linnaeus, 1758) | 4(3)-162(16)(17) | 4(3)-48(4) | 31-34 | SV,S | | | | | | | | | | |
| Ebalia tuberosa (Pennant, 1777) | Sublittoral(3)-200(18) | 40-98(4) | 26-111 | SV,V | | | | | | | | | | |
| Ebalia cranchii Leach, 1817 | 7(19)-550(3)-1000(17) | | In fishing nets | | | | | | | | | | | |
| Ebalia granulosa H. Milne Edwards, 1837 | Littoral-445(3)-2983?(19) | | 35-102 | SV,V | | | | | | | | | | |
| Ebalia deshayesi Lucas, 1846 | 12(14)-100(3) | 5-50(3) | 18-72 | S,SV | | | | | | | | | | |
| Ebalia tumefacta (Montagu, 1808) | 2-155(19) | 20-54(3) | 35-42 | SV | | | | | | | | | | |
| Ebalia edwardsi Costa, 1838 | Sea shore-190(3) | Sea shore(3)- 5-7(4) | Sea shore | S | | | | | | | | | | |
| Fam. CORYSTIDAE | | | | | | | | | | | | | | |
| Samouelle, 1819 | | | | | | | | | | | | | | |
| Corystes cassivelaunus (Pennant, 1777) | 0-50(18)-90(20) | 9-20(3) | Sea shore-15 | SF,S | | | | | | | | | | |
| Fam. ATELECYCLIOIDAE | | | | | | | | | | | | | | |
| Ortmann, 1893 | | | | | | | | | | | | | | |
| Atelocyclus rotundatus (Olivier, 1792) | Littoral(17)-300(6)- 748(16)(17) | 20-90(3) | 11-78 Max.18-29 | SV,V,S | | | | | | | | | | |
| Atelocyclus undecimdentatus (Herbst, 1783) | Sublittoral-30(21)(17) | To 30(3) | 1,5-51 Max.2-15 | S | | | | | | | | | | |
| Fam. THIIDAE | | | | | | | | | | | | | | |
| Dana, 1852 | | | | | | | | | | | | | | |
| Thia scutellata (Fabricius, 1793) | 0-50(18) | 10-20(3) | 2-7 | S | | | | | | | | | | |
| Fam. PIRIMELIOIDAE | | | | | | | | | | | | | | |
| Alcock, 1899 | | | | | | | | | | | | | | |
| Pirimela denticulata (Monatagu, 1808) | 0,3(3)-200(16) | 0,3-1(3) | 0,1-6 | SR,SA,F | | | | | | | | | | |
| Fam. PORTUNIDAE | | | | | | | | | | | | | | |
| Rafinesque, 1815 | | | | | | | | | | | | | | |
| Carcinus aestuarii Nardo, 1847 | Sea shore-6(22) | Sea shore | 0-1 | SVR | | | | | | | | | | |
| Carcinus maenas (Linnaeus, 1758) | 0-62(17)-rare at 200 (20)(7) | Shallow water | 0-1 | SVR,SR | | | | | | | | | | |
| Portunus latipes (Pennant, 1777) | 0(3)-28(17) | Shallow water | 0-10 Max. 2-4 | S | | | | | | | | | | |
| Xaiva biguttata (Risso, 1816) | Littoral-Some half a score of meters (17) | Few meters(3) | 0-6 | S,SAF | | | | | | | | | | |
| Macropodus tuberculatus (Roux, 1830) | 80(19)-500(16) | Common at 300-400(3) | 90-310 | V,SV | | | | | | | | | | |
| Liocarcinus arcuatus (Leach, 1814) | Littoral(17)-108(6) | Shallow water (3)- 19(4) | 0,1-44 | S,SAF | | | | | | | | | | |
| Liocarcinus puber (Linnaeus, 1767) | Littoral-80(19) | Littoral | 1,5-15 | R | | | | | | | | | | |
| Liocarcinus corrugatus (Pennant, 1777) | 10(3)-90(14) | 10-50(3) | 2-44 Max. 25-30 | S,SAF | | | | | | | | | | |
| Liocarcinus maculatus (Risso, 1827) | S-73(37) | 20(3)-50(4) | 14-109 Max. 25-30 | S,SV,SG,G | | | | | | | | | | |
| Liocarcinus zariqueyi (Gordon, 1968) | S(3)-60(37) | S-30(3) | 27-31 | S | | | | | | | | | | |
| Liocarcinus depurator (Linnaeus, 1758) | 1-450(20) | Shallow water-300(3) | 1-310-450 Max. 36-90 | SV,V,SG,C,... | | | | | | | | | | |
| Liocarcinus marmoratus (Leach, 1816) | 0(18)-85(20)-200(19) | | 2-6 | S | | | | | | | | | | |
| Liocarcinus vernalis (Risso, 1816) | Sea shore-20(17) | Sea shore-20(3) | 0-7 | S | | | | | | | | | | |
| Polybius henslowii (Leach, 1820) | To 375(11) | To 250(25) | 2-15 | Pelagic | | | | | | | | | | |
| Sathyneutes maravigna (Prestandrea, 1839) | 100-1455(3)(12) | 100-1455(3) | 60-300 | SV,V | | | | | | | | | | |
| Sathyneutes longipes (Risso, 1816) | 20(3)-91(19) | 20-90(3) | ? | ? | | | | | | | | | | |
| Portunus hastatus (Linnaeus, 1767) | 1(1)-25(5) | | 2-6 | S | | | | | | | | | | |
| Fam. GERYONIOIDAE | | | | | | | | | | | | | | |
| Colosi, 1923 | | | | | | | | | | | | | | |
| Geryon longipes A. Milne Edwards, 1881 | 250(22)-1430(11) | 300(3)-1000(4) | 300-310 890(28) | V | | | | | | | | | | |

| | | | | | J F M A M J J A S O N D |
|----------------------------|---------------------------------|------------------------|-------------|-----------------------|-------------------------|
| Fam. XANTHICAE | | | | | |
| MacLeay,1838 | | | | | |
| Pilumnus spinifer | Sublittoral-100(17)-180(29) | 40-100(3) | 1-89 | SV,50,S,V,C,... | |
| H. Milne Edwards,1834 | | | Max. 18-37 | | |
| Pilumnus villosoissimus | 3(3)-70(29) | 3-4(3) | 2-6 43? | SR,R | |
| (Rafinesque,1814) | | | | | |
| Pilumnus hirtellus | 1(30)-77(4) | Shallow water(3)-77(4) | 1-6 | Co,SF,SR | |
| (Linnaeus,1761) | | | | | |
| Eriphia verrucosa | A few meters | A few meters | 0-1 | R | |
| (Forskal,1775) | | | | | |
| Xantho poressa | First meters(31)-0(8)(22)-5(4) | 0(8)-5(4) | 0-1 | SR | |
| (Olivier,1792) | | | | | |
| Xantho pilipes | To 100(31)-110(19) | Shallow water(3)-70(4) | 0-45 | S,SVR,SGA | |
| A. Milne Edwards,1867 | | | | | |
| Xantho incisus incisus | Littoral-40-100(31)(3) | Littoral-40(3) | 0-1,5 | SR | |
| (Leach,1814) | | | | | |
| Xantho incisus granularius | 0-100(3) | 0,5-10(3) | 0-1 | SR | |
| (Forest,1953) | | | | | |
| Monodaeus couchii | 7(19)-60(3)(32)-1300(11) | 60-1000(3) | 85-300 | SV,V | |
| (Couch,1851) | | | | | |
| Fam. PINNOTHERIOAE | | | | | |
| De Haan,1833 | | | | | |
| Pinnotheres pisum | Littoral(21)-50(4) | 47-50(4) | 0-48 | Commensal in Bivalves | |
| (Linnaeus,1767) | | | | | |
| Pinnotheres pinnotheres | 16-94(33) | | 6-65 | Commensal in Bivalves | |
| (Linnaeus,1758) | | | | | |
| Asthenognathus atlanticus | 8-70(33) | | 0-16 | S,?Commensal? | |
| Monod,1933 | | | | | |
| Fam. PALCIDAE | | | | | |
| Rathbun,1898 | | | | | |
| Palicus caronii | 23(9)-100(9)(17) | 52-91(3) | 46-50 | SAC | |
| (Roux,1830) | | | | | |
| Fam. GONEPLACIOAE | | | | | |
| MacLeay,1838 | | | | | |
| Goneplax rhomboides | 4(8)-760(9)-1478(19) | 4(8)-550(4) | 0-310 | SV,V,S | |
| (Linnaeus,1758) | | | Max. 37-74 | | |
| Fam. GRAPSIDAE | | | | | |
| MacLeay,1838 | | | | | |
| Pachygrapsus marmoratus | 0-5(4) | Sea shore(3)-5(4) | 0-5 | SR,R | |
| (Fabricius,1787) | | | Max. 0-0,5 | | |
| Pachygrapsus maurus | Intertidal | Intertidal(3) | 0-3 | R,SR | |
| (Lucas,1846) | | | | | |
| Pachygrapsus transversus | 0-10(9) | Intertidal(3) | 0-0,5 | SR,R | |
| (Gibbes,1850) | | | | | |
| Planes minutus | On floating objects | | | | |
| (Linnaeus,1758) | | | | | |
| Euchirograpsus liguricus | 10(32)-350(34) | 54-350(3) | ? | ? | |
| H. Milne Edwards,1853 | | | | | |
| Brachynotus sexdentatus | Littoral | 1-2(3) | 0-1 | SR,SVR,R | |
| (Risso,1827) | | | | | |
| Brachynotus atlanticus | Littoral | | 0,1 | SR | |
| Forest,1957 | | | | | |
| Fam. PARTHENOPIDAE | | | | | |
| MacLeay,1838 | | | | | |
| Heterocrypta maltzami | 5-550(9)-900(16) | 110(8) | 22-50 | SF | |
| Miers,1881 | | | | | |
| Parthenope massena | 3(3)(9)-500(16)(17) | 3-100(3) | 3-55 | S,SC,SA,SG | |
| (Roux,1830) | | | Max. 14-27 | | |
| Parthenope angulifrons | 10(3)-80(5) | 10-40(3) | 2-30 | S,SA | |
| Latreille,1825 | | | | | |
| Parthenope macrochelos | 80(5)-1478?(16) | Quite deep(3) | 55-74 | SV | |
| (Herbst,1790) | | | | | |
| Fam. MAJIDAE | | | | | |
| Samouelle,1819 | | | | | |
| Maja squinado | 2(35)(1)(19)-75(9) | To 50(3) | 3-55 | S,SV,R | |
| (Herbst,1788) | | | | | |
| Maja crispata | 0(3)(36)-50(29) | Sea shore-40(3) | 0,1-40 | SR,R,SA | |
| Risso,1827 | | | | | |
| Acanthonyx lunulatus | 0-0,1(30)(36)-10(9) | 0(3)-5(4) | 0-2 | SA,RA,A | |
| (Risso,1816) | | | | | |
| Eury nome aspera | 10-550(16)(17)(21)-684(19) | To 75(3) | 14-96 | S,SA,SC,SV | |
| (Pennant,1777) | | | | | |
| Herbstia condylata | 2(6)-3(22)-54(3) | To 54(3) | 1-20 | R | |
| (Fabricius,1787) | | | | | |
| Lissae chiragra | 20(3)-60(16) | 20-40(3) | 1-27 | R,S,Co | |
| (Fabricius,1775) | | | | | |
| Anamathia rissoana | 30-1000(16) | 400(3)-775(13) | 650-775(13) | | |
| (Roux,1828) | | | | | |
| Pisa tetraodon | Littoral-100(16)(17)(21) | 0,3-more than 50(3) | 0-44 | R,RA,SR | |
| (Pennant,1777) | | | | | |
| Pisa muscosa | 4(3)-70(29) | 4-40(3) | 14-33 | S,SV | |
| (Linnaeus,1758) | | | | | |
| Pisa carinimana | 2(21)-100(6) | | 4-44 | SA | |
| Miers,1879 | | | | | |
| Pisa armata | Littoral(17)-162(24) | To 50 or more(3) | 18-102 | Mainly SV | |
| (Latreille,1803) | | | Max. 74-78 | | |
| Pisa nodipes | Sublittoral-75(17) | To 75(3) | 9-44 | S,SRA | |
| (Leach,1815) | | | | | |
| Achaetus gracilis | 0,5-20(3) | 0,5-20(3) | 0,1-43 | RA,SA,S,F | |
| (Costa,1839) | | | | | |
| Achaetus cranchii | Littoral-68(6)-70(19) | 20(3)-50(4)(3) | 22-40 | SA | |
| Leach,1817 | | | | | |
| Inachus communissimus | 15(3)-48(4) | 15(3)-48(4) | 5-78 | S,SV,V,C | |
| Rizza,1839 | | | Max. 15-42 | | |
| Inachus dorsettensis | A few meters(6)-168(11)-256(19) | 30(3)-98(4) | 5-109 | V,SV,S | |
| (Pennant,1777) | | | Max. 20-43 | | |
| Inachus phalangium | 4(3)-150(6) | 4-40(3) | 6-15 | SR | |
| (Fabricius,1775) | | | | | |

| | | | | | |
|--|----------------------|---------------------------|----------------------|-----------------|--|
| <i>Inachus thoracicus</i> | 25(21)-200(29) | 30(4)-80(3) | 14-78 | SA,S,SV,R | |
| Roux, 1830 | | | | | |
| <i>Inachus aguiarii</i> | 20(16)-200(6) | 20-100(3) <i>Melillia</i> | 14-66 | SA,S,SV,R | |
| Brito Capello, 1876 | | | | | |
| <i>Macropodia rostrata</i> (Linnaeus, 1775) | 0-50-150(26)(27) | A few meters-70(3) | 5-94 Max. 18-42 | S,SV,C,SR,SA... | |
| <i>Macropodia linaresi</i> | 20-80(26)(27)-86(19) | 30-80(3) | 18-55 | S | |
| Forest & Zariquey, 1964 | | | | | |
| <i>Macropodia czerniavskii</i> (Brandt, 1880) | 10-80(26) | 10(3)-37(4) | 0,3-2 | RA,F | |
| <i>Macropodia longirostris</i> (Fabricius, 1775) | Some meters-130(26) | 4-50(3) | 18-55 Max. 18-36 | SA,S | |
| <i>Macropodia longipes</i> (A. Milne Edwards & Bouvier 1899) | 50-445(27) | 70-100(3) | 9-310 Max. 40-270 | SV,V,S | |



Comments on some species

A record of the capture the *Cancer pagurus* Linnaeus, 1758, in the port of Málaga and the report of a sighting of another at 10 meters depth in the littoral region in the eastern region of Málaga by a free diver cannot be confirmed since this species was not found or seen during the current study.

A variety of sponges (*Euspongia* sp., *Suberites* sp., etc) were observed on the dorsal surface of *Dromia personata* (L.) where they are attached near the fifth legs, while on occasion specimens of *Alcyonium palmatum* Pallas were sometimes found in the same position.

Although *Homola barbata* (Fab.) is cited as being found beyond 50 m in this study, one specimen was taken on the harbour walls of Caleta de Vélez, perhaps dislodged during the cleaning of a fishing boat.

Bathymetric distribution data on *Ebalia granulosa* (H. Milne Edw.) and *E. cranchii* (Leach) has not previously been given for Spanish waters (ZARIQUEY, 1968). One specimen of this genus, possessing intermediate characters, could not be clearly assigned to either species.

Thia scutellata (Fab.) isn't frequently encountered. It appears in the samples in association with *Albunea carabus* (L.) and *Atelecyclus undecimdentatus* (Herbst) in sandy littoral substrates.

The width/length relationship of specimens of the genus *Carcinus* taken in this study in comparison with data of ALMAÇA (1972) and ZARIQUEY (1968) is:

| | Data obtained by author | ALMAÇA | ZARIQUEY |
|---------------------|----------------------------|-----------|-----------|
| <i>C. maenas</i> | 1.29-1.35 | 1.21-1.37 | 1.22-1.35 |
| <i>C. aestuarii</i> | 1.25-1.27 | 1.17-1.27 | — |

Pilumnus spinifer H. Milne Edw. is quite frequently encountered. Two anomalies were observed, in one specimen the carpus of the fifth leg has two spines while in another specimen the left anterolateral margin has only three teeth instead of the usual five. Small individuals of this species were occasionally found in the roots of phanerogams but this was not the case with the adults.

Pilumnus villosissimus (Raf.) is a species less frequently found in comparison to the previous one. In observations made by free diving it has been established that it quite commonly occurs on rocky substrates beneath the tentacles of large anemones together with several specimens of *Scyllarus arctus* (L., 1758). *Inachus phalangium* (Fab.), on the other hand, appears also in the middle of the anemone tentacles, the only place where this species has been found.

The species *Paractaea rufopunctata* (H. Milne Edw., 1834), not found in the study area, appears very occasionally in the Granada coastal region at the Punta de la Mona (GARCIA RASO & BARRAJON, 1982) which is very close to the study zone. It is possible that it could occur on rocky substrates with phanerogams in the extreme east of the study area.

Both species of the genus *Pinnotheres* Bosc, 1802, *P. pisum* (L.) and *P. pinnotheres* (L.) have been found principally inside the bivalve *Atrina pectinata* (L., 1767). *P. pisum*, was also encountered, but less frequently, in *Mytilus edulis* (L., 1758) and only rarely in *Donax trunculus* L., 1758.




Fig. 2. *Brachynotus atlanticus* Forest, ♀: a, carapace; b, left cheliped; c, third maxilliped; d, second pereiopod; e, fifth pereiopod.

Specimens of *Asthenognathus atlanticus* Monod have been taken free-living on the soft substrates or found thrown up on the beach, while one specimen was found in the stomach of *Raja asterias* Delaroche 1809. The only records from the Mediterranean of this species, known to the author, are those of NOEL et AMOUROUX (1977) and GARCIA RASO & JIMENEZ MILLAN (1981). The average dimensions given by them are:

| | Width | Length |
|-----|----------|---------|
| ♂ ♂ | 11.78 mm | 7.50 mm |
| ♀ ♀ | 13.33 mm | 7.96 mm |

Goneplax rhomboides (L.) is one of most abundant Brachyuran species in the study zone, together with *Liocarcinus depurator* (L.) and *Pachygrapsus marmoratus* (Fab.). The capture of two specimens at 0 m. was very unusual. The smaller ovigerous female found had a width of 13.4 mm.

The only Brachyuran taken with the parasite *Sacculina* were *Inachus communissimus* Rizza, *I. dorsettensis* (Pennant) and *Monodaeus couchi* (Couch). The former species, in a number of cases, even carried two of these parasites while the latter was observed to do so in one case only.

A number of interesting species were taken during this study. *Bathynectes longipes* (Risso), is a rare species and only one specimen was encountered. *Liocarcinus zariqueyi* (Gordon), a species with a wide geographical distribution and known from England (ZARIQUIEY, 1968; INGLE, 1980) to Canary Islands (ZARIQUIEY, 1968; MANNING & HOLTHUIS 1981) and Mediterranean is considered quite a rare species and was previously known only from the Catalan coast in Spain. Similarly only two specimens of *Palicus caronii* (Roux), captured in 1950 at Cadaqués were reported by ZARIQUIEY (1968). *Euchirograpsus liguricus* H. Milne Edw. is not a common species in the Mediterranean Sea and only two specimens were taken during this study. ZARIQUIEY (1968) did not record *Heterocripta maltzami* Miers from Spain and considered it very rare based in the works of MIRANDA & RIVERA (1921, 1933). Representatives of this species, moreover, are absent from the reference collections in the Oceanographic Institute (I.E.O.) of Málaga. The presence of *Pisa carinimana* Miers in the study zone represents for the European Mediterranean the single area where this species is reported. Other species of rare occurrence are *Inachus aguiarii* Britto Capello, *Macropodia linarensis* Forest et Zariquey and *M. czerniavskii* (Brand). According to available data this record of the latter species is its most westerly reported occurrence to date.

Finally, the capture of two specimens of *Brachynotus atlanticus* Forest is of special note. These conform exactly to the descriptions given by FOREST (1957) concerning the line of frontal granulations, the internal margin of the third maxillipeds, the position of the anterolateral teeth and the keels of the chelipeds (Fig. 2). However the length/width ratio of the propodite of the P2 and P5 legs is similar to it of *B. sexdentatus*. The capture of these specimens represents the first record of this species in the Mediterranean Sea. It extends northwards its known distribution since the previous most northerly records of the species are from the Atlantic coast of Morocco at 34° 53' latitud.

References

- ALMAÇA, C., 1972: Preliminary note on some ratios between carapace dimensions of *Carcinus maenas* (L.) and *C. mediterraneus* Czerniavski. – Thalassia Jugoslavica 8(1): 49–57
- BACESCU, M., 1967: Decapoda. – In: Fauna Republicii Socialiste România. Crustacea. Bucharest, 4(9): 1–351
- BARNARD, K. H., 1950: Descriptive Catalogue of South African Decapod Crustacea (Crabs and Shrimps). – Annals of South African Mus., 38: 1–837
- BOUVIER, E. L., 1940: Décapodes Marcheurs. – Faune de France, Paris, 37: 1–404
- BUEN, O. DE, 1925: Sur les essaims de *Polybius beslowi* Leach. – Extrait Bulletin de la Société Zoologique France, 50: 90–92
- CAPART, A., 1951: Crustacés Décapodes Brachyures. – In: Expédition Océanographique Belge dans les Eaux Côtières Africaines de l'Atlantique Sud (1948–1949). Résultats Scientifiques. 3(1): 11–205
- CHRISTIANSEN, M. E., 1969: Crustacea Decapoda Brachyura. – Marine invertebrates of Scandinavia, Oslo, 2: 1–143
- CRESPO, J.; J. C. REY & J. A. CAMINAS, 1976: *Trachyrhynchus trachyrhynchus* (Risso, 1810) del Mar de Alborán. Pesca y rendimiento. Areas de distribución. Biología y biometría. – Bol. Instituto Español de Oceanografía, 218: 1–21
- CROSNIER, A., 1970: Crustacés Décapodes Brachyures et Macrures recueillis par l’“Undaunted” au sud de l’Angola: Description de *Scyllarus subarctus* sp. nov. – Bull. Mus. National d’Histoire naturelle, ser. 2, (1969), 41(5): 1214–1227
- DRACH, P. & J. FOREST, 1953: Description et répartition des *Xantho* des mers d’Europe. – Archives de Zoologie Expérimentale et Générale, 90: 1–35

- ©Zoologische Staatssammlung München; download: <http://www.biodiversitylibrary.org/>; www.biologiezentrum.at
- FOREST, J., 1957: Mise au point sur les *Brachynotus* de la Méditerranée et d'Afrique occidentale: *Brachynotus sex-dentatus* Riso et *Brachynotus atlanticus* nov. sp. – Bull. Inst. Français d'Afrique Noire, ser. A, 19(2): 501–510
- — 1965: Crustacés Décapodes: Campagnes du «Professeur Lacaze-Duthiers» aux Baleares, Juin 1953 et Aout 1954. – Vie et Milieu, 16(1B): 325–413
- — 1967: Sur une collection de Crustacés Décapodes de Porto Cesareo: Description de *Portumnus pestai* sp. nov. – Thalassia Salentina 2: 1–29
- — 1974: Les Dromies de l'Atlantique Oriental: Description de *Sternodromia* gen. nov. et deux espèces nouvelles du genre *Dromia* Weber (Crustacea Decapoda Dromiidae). – Annales de l'Institut Océanographique, new ser., 50(1): 71–123
- — 1978: Le genre *Macropodia* Leach dans les eaux Atlantiques européennes (Crustacea Brachyura Majidae). – Cahiers de Biologie Marine, 19: 323–342
- — & GANTES, 1960: Sur une collection des Crustacés Décapodes marcheurs du Maroc. – Bull. Mus. National d'Histoire naturelle, ser. 2, 32(4): 346–358
- — & D. GUINOT, 1958: Sur une collection de Crustacés Décapodes des côtes d'Israël. – Bull. Sea Fisheries Research Station Haifa 15: 4–16
- — 1966: Crustacés Décapodes Brachyures. – In: Campagne de la Calypso dans le Golfe Guinée et aux îles Principe, São Tomé et Annobon (1956). 16. – Annales d'Institut Océanographique, 44: 23–124
- — & R. ZARIQUIEY, 1965: Le genre *Macropodia* Leach en Méditerranée, I: Description et étude comparative des espèces (Crustacea Brachyura Majidae). – Bul. Mus. National d'Histoire naturelle, ser. 2, 36(2): 222–244
- FROGLIA, C. & R. B. MANNING, 1982: Notes on *Liocarcinus pusillus* (Leach) and related species. – Quad. Lab. Tecnol. Pesca, 3(2–5): 257–266
- GARCIA RASO, J. E. & A. BARRAJON, 1982: Contribución al conocimiento de los Xanthidae Mac-Leay (Crustacea, Decapoda, Brachyura) del Sur de España. – Monografías y Trabajos Depto. Zoología Málaga, (1981–1982), 3–4: 3–14
- — & F. JIMENEZ MILLAN, 1981: Algunas especies de Decápodos (Crust.) del litoral malagueño. – Bol. Asociación Española Entomología, 4(1980): 19–24
- GONZALEZ GURRIARAN, E., 1978: Introducción al estudio de una población de *Macropipus puber* (L.) (Decapoda Brachyura) en la Ría de Arosa (Galicia, España). – Bol. Instituto Español de Oceanografía, 4, 241: 73–80
- INGLE, R. W., 1980: British Crabs. – British Museum (Natural History) London, 1–222
- KOCATAS, A., 1981: Liste préliminaire et répartition des Crustacés Décapodes des eaux turques. – Rapports Comm. International Mer Méditerranéen, 27(2): 161–162
- LAGARDERE, J. P., 1973: Distribution des Décapodes dans le Sud du Golfe de Gascogne. – Revue des Travaux de l'Institut des Pêches Maritimes, 37(1): 77–95
- LEWINSOHN, Ch., 1976: Crustacea Decapoda von der Insel Rhodos, Griechenland. – Zoologische Mededelingen, 49(17): 237–254
- MANNING, R. B. & L. B. HOLTHUIS, 1981: West African Brachyuran Crabs (Crustacea: Decapoda). – Smithsonian Contributions to Zoology, 306: 1–379
- MASSUTI, M., 1967: Carta de pesca de la región sur-mediterránea española (desde Estepona a Adra). – Bol. Instituto Español de Oceanografía, 33: 1–24
- MAURIN, C., 1968a: Ecologie ichthyologique des fonds chatulables Atlantiques (de la Baie Ibéromarocaine à la Mauritanie) et de la Méditerranée occidentale. – Rev. Travaux d'Institut des Pêches maritimes, 32(1): 3–147
- — 1968b: Les Crustacés capturés par la «Thalassa» au large des côtes nord-ouest africaines. – Rev. Romaine de Biologie, ser. Zoologie, 13(6): 479–493
- MIRANDA Y RIVERA, A., 1921: Algunos crustáceos de la colección del laboratorio biológico de Málaga. – Boletín de Pescas, 6: 179–205
- — 1933: Ensayo de un catálogo de los Crustáceos Decápodos marinos de España y Marruecos español. – Notas y Resúmenes. Bol. Instituto Español de Oceanografía, ser. 2, 67: 1–72
- MONOD, TH., 1956: Hippidea et Brachyura uest-africains. – Mémoires de l'Institut Français d'Afrique Noire, 45: 1–674
- NOBRE, A., 1936: Crustáceos Decápodos e Stomatópodes Marinhos de Portugal. – In: Fauna Marinha de Portugal, 2a ed., 4: 1–213
- NOËL, P. & J. M. AMOUROUX, 1977: Sur la présence d'*Asthenognathus atlanticus* Monod, 1932, (Crustacea Brachyura) dans la région de Banyuls-sur-Mer (Méditerranée). – Vie Milieu ser. A, 27(1): 135–136

- © Zoologische Staatssammlung München download: <http://www.biodiversitylibrary.org/>; www.biologiezentrum.at
- PASTORE, M., 1972: Decapod Crustacea in Taranto's and Catania's Seas and a discussion on a new species of Dromiidae (Decapoda Brachyura) in the Mediterranean Sea. – *Thalassia Jugoslavica*, 8(1): 105–117
- PESTA, O., 1918: Die Decapodenfauna der Adria. – Versuch einer Monographie, Leipzig and Wien, 1–500
- POWERS, L. W., 1977: A catalogue and bibliography to the crabs of the Gulf of Mexico. – Contributions in Marine Science (suppl.) 20: 1–190.
- RAMADAN, SH. E. & N. M. DOWIDAR, 1972: Brachyura (Decapoda Crustacea) from the Mediterranean waters of Egypt. – *Thalassia Jugoslavica*, 8(1): 127–139
- SANTAELLA, E., 1974: Crustáceos Decápodos del Archipiélago Canario. I. Contribución al estudio de las especies de Braquiuros (Crustacea Decapoda Brachyura). – *Bol. Instituto Español de Oceanografía*, 182: 1–31
- SCHMITT, W. L.; J. C. McCAIN & E. S. DAVIDSON, 1973: Decapoda, I: Brachyura, I: Fam. Pinnotheridae. – In: H.-E. GRUNER and L. B. HOLTHUIS (eds.), *Crustaceorum Catalogus*, 3: 1–160
- ŠTEVČIC, Z., 1973: Les migrations de l'Araignée de Mer. – *Rapp. Comm. International Mer Méditerranéen*, 21(9): 597–598
- TURKAY, M., 1975: Zur Kenntnis der Gattung *Euchiro grapsus* mit Bemerkungen zu *Brachygrapsus* und *Litochir* (Crustacea: Decapoda). – *Senckenbergiana Biologica*, 56(1/3): 103–132
- — 1976a: Decapoda Reptantia von der portugiesischen und marokkanischen Küste. Auswertung der Fahrten 8, 9c (1967), 19 (1970), 23 (1971) und 36 (1975) von F. S. „Meteor“. „Meteor“-Forschungsergebnisse, Reihe D: Biologie 23: 23–44
- — 1976b: Die Madeirensischen Brachyuren des Museu Municipal do Funchal und des Forschungs-Instituts Senckenberg. I. Familien Dromiidae, Homolidae, Calappidae, Leucosiidae, Cancridae, Portunidae, Xanthidae, Geryonidae, Gonoplacidae und Palicidae (Crustacea Decapoda). – *Bol. Mus. Municipal do Funchal* 30(art. 133): 57–74
- WILLIAMS, A. B. and R. L. WIGLEY, 1977: Distribution of Decapod Crustacea off Northeastern United States based on specimens at the Northeast Fisheries Center, Woods Hole, Massachusetts. – U. S. Dept of Commerce, N.O.A.A. Technical report N.M.F.S., 407: 1–44
- ZARIQUIEY, A., R., 1968: Crustáceos Decápodos Ibéricos. – *Investigaciones Pesqueras*, 32: 1–510

Author's address:

Dr. J. Enrique García Raso,
Dpto. Zoology, Fac. Science,
Univ. Málaga, Spain.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 1984

Band/Volume: [007](#)

Autor(en)/Author(s): Garcia Raso J. Enrique

Artikel/Article: [Brachyura of the coast of Southern Spain \(Crustacea, Decapoda\)](#)
[105-113](#)