

SPIXIANA	16	2	171–172	München, 1. Juli 1993	ISSN 0341–8391
----------	----	---	---------	-----------------------	----------------

***Doronomyrmex pacis* Kutter 1945,
a socially parasitic ant new to Germany**

(Insecta, Hymenoptera, Formicidae)

By J. Heinze, S. Kauffmann & B. Hülsen

Heinze, J., Kauffmann, S. & B. Hülsen (1993): *Doronomyrmex pacis* Kutter 1945, a socially parasitic ant new to Germany (Insecta, Hymenoptera, Formicidae). - Spixiana 16/2: 171-172

The parasitic ant *Doronomyrmex pacis* is recorded for the first time from Germany: three colonies were found at an elevation of 1660m at Jenner, Berchtesgaden National Park.

J. Heinze, S. Kauffmann, Theodor-Boveri-Institut für Biowissenschaften, Am Hubland, W-8700 Würzburg, Germany

B. Hülsen, Max-Planck-Institut für Verhaltensphysiologie, W-8130 Seewiesen, Germany

Introduction

Of the approximately 10.000 described species of ants some 200 are social parasites, i.e. they temporarily or permanently depend on the help of workers from another ant species (e. g., Hölldobler & Wilson 1990). Workerless parasites (“inquilines”) are among the rarest species of ants. Many are known only from their type localities, whereas others are discontinuously distributed over large areas. The knowledge on the geographical range of social parasites might help to understand the evolution of social parasitism (Baroni-Urbani 1967) and is fundamental for the protection of these rare ant species (Buschinger 1987). Each new record is therefore of considerable interest.

The holarctic ant, *Leptothorax* (s.str.) *acervorum* is host of at least four parasites: the slave-maker *Harpagoxenus sublaevis*, and the workerless *Doronomyrmex kutteri*, *D. goesswaldi*, and *D. pacis*. *Doronomyrmex pacis*, Kutter 1945 was described from a single queen collected in Binntal, Valais, Switzerland. Since then it has repeatedly been rediscovered in various parts of the Alps: Switzerland (Saas Fee, Zermatt, Ofenpaß etc., Kutter 1969, Buschinger 1971), northern Italy (Schludersbach, Buschinger 1971; Toblach, Buschinger et al. 1981; Schlanders, Buschinger, pers. comm.), Austria (Tennengebirge, Winter 1972), France (Maurienne, Buschinger et al. 1981; Briançon, Heinze & Buschinger, unpubl.), and Slovenia (Triglav, Buschinger pers. comm., Buschinger & Douwes in press). We here report the first finding of *Doronomyrmex pacis* from Germany and document some life history traits of this species.

Results and Discussion

Some thirty complete colonies of *Leptothorax* (s.str.) *acervorum* were collected on July 20 and 21, 1991 near “Schneibsteinhaus”, Jenner, Berchtesgaden National Park, southern Germany, at an elevation of appr. 1660 m. Nests were found in rotting sticks and roots of dwarf pine, *Pinus mugo*, on the edge of stands of coniferous trees amidst alpine meadows. In two colonies a *D. pacis* queen was found, and in a third nest, *D. pacis* sexuals enclosed in the laboratory shortly after collection. *D. pacis* queens are easily distinguished from host queens: whereas the latter are brownish and coarsely sculptured, *D. pacis*

queens are shiny and black. Furthermore, parasite queens are much smaller than host queens.

All three *D. pacis* colonies from Jenner contained alate sexuals of *L. acervorum*, one or more *L. acervorum* queens, and between appr. 20 and 150 workers. Five *L. acervorum* queens were dissected: their ovaries were strongly elongated and contained corpora lutea, but no developing or mature oocytes. Whether this condition reflects a regular reduction in fertility in midsummer or reproductive inhibition through the parasite queen, as suggested by Buschinger (1990), is not yet clear. Many social parasites, which were thought to tolerate fertile host queens, interfere with egg-laying or even kill host queens (e.g., Allies et al. 1986; Buschinger & Klump 1988). Openly aggressive or ritualized dominance interactions of queens, which might probably cause ovary degeneration (e.g., Heinze & Smith 1990), were not observed.

According to laboratory observations, queens of *D. pacis* exhibit sexual calling behavior (e.g., Buschinger 1972) and mate in the immediate neighborhood of the maternal nest. As yet it is not known whether mated queens may seek readoption into the maternal nest instead of invading a new host colony. Winter (1972) found a colony with two dealate female sexuals of *D. pacis*, however, their reproductive status was not examined. After eclosion of sexuals we observed heavy fighting between two dealate gynes of *D. pacis* in a single nest, probably between mother and one of her offspring. Both tumbled about the nest, pulling each other's legs and antennae with their mandibles and extruding their stings. One individual left the nest about half an hour later. Aggression between parasite queens is known from several species, e.g. *Harpagoxenus sublaevis*, *Doronomyrmex goesswaldi* (Buschinger & Klump 1988), *Leptothorax wilsoni* (Heinze 1989), and *Epimyrma algeriana* (Buschinger et al. 1990), all of which also attack fertile host queens. More observations are needed to clarify the social organization of *D. pacis* colonies.

Acknowledgements

Studies on *Leptothorax* in Berchtesgaden National Park were made possible by a permit of the Park authorities (520 NA 01-604/91). A. Buschinger, TH Darmstadt, provided specimens for comparison.

References

- Allies, A. B., Bourke, A. F. G. & N. R. Franks 1986. Propaganda substances in the cuckoo ant *Leptothorax kutteri* and the slavemaker *Harpagoxenus sublaevis*. - J. chem. Ecol. 18: 1285-1293
- Baroni-Urbani, C. 1967. Le distribuzioni geografiche discontinue dei Formicidi mirmecobiotici. - Arch. Botan. Biogeograf. Ital. 43: 355-364
- Buschinger, A. 1971. Zur Verbreitung der Sozialparasiten von *Leptothorax acervorum* (Fabr.) (Hym., Formicidae). - Bonner zool. Beitr. 22: 322-331
- 1972. Kreuzung zweier sozialparasitischer Ameisenarten, *Doronomyrmex pacis* Kutter und *Leptothorax kutteri* Buschinger (Hym., Formicidae). - Zool. Anz. 189: 169-179
- 1987. *Teleutomyrmex schneideri* Kutter 1950 and other parasitic ants found in the Pyrenees. - Spixiana 10: 81-83
- 1990. Sympatric speciation and radiative evolution of socially parasitic ants - Heretic hypotheses and their factual background. - Z. zool. Syst. Evolut.-forsch. 88: 241-260
- Buschinger, A. & P. Douwes (in press). Socially parasitic ants of Greece. - Biol. gallo-hellenica.
- Buschinger, A. & B. Klump 1988. Novel strategy of host-colony exploitation in a permanently parasitic ant, *Doronomyrmex goesswaldi*. - Naturwissenschaften 75: 577-578
- Buschinger, A., Ehrhardt, W. & K. Fischer 1981. *Doronomyrmex pacis*, *Epimyrma stumperi*, und *E. goesswaldi* (Hym., Formicidae) neu für Frankreich. - Ins. Soc. 28: 67-70
- Buschinger, A., Jessen, K. & H. Cagniant 1990. The life history of *Epimyrma algeriana*, a slave-making ant with facultative polygyny (Hymenoptera, Formicidae). - Zool. Beitr. N.F. 33: 23-49
- Heinze, J. 1989. *Leptothorax wilsoni* n. sp., a new parasitic ant from eastern North America (Hymenoptera: Formicidae). - Psyche 96: 49-61
- Heinze, J. & T. A. Smith 1990. Dominance and fertility in a functionally monogynous ant. - Behav. Ecol. and Sociobiol. 27: 1-10
- Hölldobler, B. & E. O. Wilson 1990. The ants. - Harvard University Press, Cambridge, Mass. 732p.
- Kutter, H. 1969. Die sozialparasitischen Ameisen der Schweiz. - Neujahrsblatt d. Naturforsch. Ges. Zürich 171: 1-62
- Winter, U. 1972. Sozialparasiten der *Leptothorax* Gruppe (Hym.; Formicidae) aus der Umgebung des Tennen-Gebirges (Österreich). - Zeitschr. der Arbeitgem. Österr. Entomologen 84: 124-126

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 1993

Band/Volume: [016](#)

Autor(en)/Author(s): Heinze J., Kauffmann S., Hülsen B.

Artikel/Article: [Doronomyrmex pacis Kutter 1945, a socially parasitican new to Germany \(Insecta, Hymenoptera, Formicidae\) 171-172](#)