

Tanytarsus usambarae, spec. nov.
from West Usambara Mts., Tanzania, East Africa

(Insecta, Diptera, Chironomidae)

Elisabeth Stur & Torbjørn Ekrem

Stur, E. & T. Ekrem (2000): *Tanytarsus usambarae*, spec. nov. from West Usambara Mts., Tanzania, East Africa (Insecta, Diptera, Chironomidae). – In: Baehr, M. & M. Spies (eds): Contributions to chironomid research in memory of Dr. Friedrich Reiss. – Spixiana 23/2: 219–223.

Tanytarsus usambarae, spec. nov. from West Usambara Mts. is described and figured as male imago. In the Usambaras, the species co-exists with the closely related *Tanytarsus luctuosus* Freeman. Both have a peculiarly shaped median volsella which separates them from all other African *Tanytarsus* species. *T. usambarae* does not fit well into Palaearctic species group concepts. The genus diagnosis of *Tanytarsus* is emended to include species with a scutal tubercle. *Tanytarsus saetosus* Lehmann is transferred to the genus *Neostempellina* Reiss.

Elisabeth Stur & Torbjørn Ekrem, Museum of Zoology, University of Bergen, Muséplass 3, N-5007 Bergen, Norway.

Introduction

Tanytarsus v. d. Wulp, 1874, is well represented in the Afrotropical region. 24 species are listed in the Catalogue of the Diptera of the Afrotropical region (Freeman & Cranston 1980), five more were described by Lehmann (1979, 1981), an additional six by Ekrem (1999b), while Ekrem & Harrison (1999) added one more from East Africa. On the other hand, Ekrem (1999a) transferred *Tanytarsus bukavus* Lehmann, 1979, and *Tanytarsus congolensis* Lehmann, 1979, to *Cladotanytarsus* Kieffer, and Ekrem & Reiss (1999) transferred *Tanytarsus abnormis* Lehmann, 1981, to the genus *Neostempellina* Reiss. Examination of the type material of *Tanytarsus saetosus* Lehmann, 1981, deposited in the Zoologische Staatssammlung München (ZSM), revealed that this species also belongs to the genus *Neostempellina*. Thus, excluding *Neostempellina saetosa*, comb. nov. and including *T. usambarae*, spec. nov., a total of 33 *Tanytarsus* species are so far known from the Afrotropical region.

Material, methods and morphology

The terminology follows Sæther (1980) with the additions and corrections given by Sæther (1990). The term “shoulders” is used for the posteromarginal lobes of the male anal tergite. The mounting procedure is according to Sæther (1969). Measurements are taken according to Schlee (1966) with the additions given by Ekrem (1999b), and presented as ranges followed by the mean.

During the work of reviewing Afrotropical *Tanytarsus*, the junior author examined the type material of *Tanytarsus luctuosus* Freeman, deposited at the Natural History Museum (London). Neither Freeman (1958) nor Lehmann (1979) described *T. luctuosus* to have a scutal tubercle, but examination of the types and of Lehmann’s material from Zaïre (now the Democratic Republic of Congo) revealed

that *T. luctuosus* possesses a well developed scutal tubercle. Since absence of this structure has been regarded as diagnostic for the genus *Tanytarsus* (Cranston et al. 1989) we choose to emend the genus diagnosis to also include species with a scutal tubercle.

Tanytarsus usambarae, spec. nov.

Figs 1A-E

Types. Holotype: ♂, Tanzania, W. Usambara Mts, Mazumbai, Kaputu stream, 1420 m a.s.l., Malaise trap, 14.XI-3.XII.1990, leg. ZMBN's Tanzania expedition (ZMBN Type No. 352). – Paratypes: 4♂♂, as holotype. All slide-mounted in Canada balsam, deposited at Museum of Zoology, University of Bergen, Bergen, Norway. (ZMBN)

Etymology. The species is named after its type locality, the Usambara Mts.

Diagnostic characters. The new species differs from other *Tanytarsus* species by the following combination of characters in the adult male: broad, transverse, brown pigmented bands on tergites II, III, VI and VII; tergites I, IV and V pale yellowish; tergites VIII and IX brown; wing hairy, carrying setae on all veins except M, and more than 25 setae in cell m; frontal tubercles minute, only visible as two white dots; hypopygium with 4-6 median setae on anal tergite, anal point with 8-10 spines between well developed anal crests; superior volsella with posterior tubercle, more than 8 dorsal setae and numerous dorsolateral microtrichia, digitus absent; median volsella large with numerous long, slender, simple lamellae reaching beyond apex of inferior volsella.

Description

Male imago (n = 5 if not otherwise stated).

Total length: 2.7-3.0 mm, 2.9 mm. Wing length 1.55-1.66 mm, 1.59 mm. Total length/wing length 1.7-1.9, 1.8.

Colouration. Cleared specimens with light brown head, darker brown antennae and eyes; thorax (Fig. 1E) with dark brown patches anteriorly on scutum, laterally under parapsidal suture, on median anepisternum II, on posterior anepisternum II, on epimeron II, on postnotum, and on preepisternum; anteprepronotum and scutellum light brownish; legs and halteres light brown; abdominal tergites II, III, VI and VII with broad, transverse, dark band; tergites I, IV and V pale yellowish; tergites VIII and IX brown.

Head. Antennae normally developed, AR 0.92-1.03, 0.95. Ultimate flagellomere 454-522 μm , 476 μm long. Longest antennal seta about 480 μm long. Distance between eyes about 150 μm . Minute frontal tubercles only visible as two white dots. Temporal bristles 10-11; including 3 inner verticals, 3-4 outer verticals and 3-4 postorbitals. Clypeus semicircular, 72-86 μm , 79 μm long with 9-14 setae. Tentorium 104-126 μm , 113 μm long, 29-32 μm , 31 μm wide at sieve plate. Stipes 120-126 μm , 124 μm long, 18-22 μm , 19 μm wide. Cibarial pump 47-48 μm wide. Lengths of palpomeres I-IV (in μm): 36, 36-47, 148 (n = 1), 122 (n = 1), fifth palpomere missing.

Thorax (Fig. 1E). Scutal tubercle absent. Dorsocentrals 8-11, acrostichals 22-25, prealars 1-2, scutellars 6-8. Halteres with 6-7 setae.

Wing. VR 1.16-1.20, 1.18. Sc bare, R with 36-42, 32 setae, R₁ with 28-37, 32; R₄₊₅ with 52-61, 55 setae; M bare; M₁₊₂ with 70-80, 76 setae; M₃₊₄ with 34-40, 38; Cu with 19-27, 22; Cu₁ with 23-25, 24; PCu with 56-71, 63 and An with 31-38, 35 setae. Cell m with 28-45 setae, r₄₊₅ with more than 200, m₁₊₂ with more than 250 including on false vein, m₃₊₄ with about 150 and cu + an together with 250-290 setae.

Legs. Spur on front tibia 32-36 μm , 35 μm long including scale. Spurs of middle tibia 32 μm long including 14-18 μm long comb and 22-29 μm long including 12-18 μm long comb; of hind tibia 36-43 μm long including 18 μm long comb and 32-36 μm long including 18 μm long comb. Length of fore femur 720-763 μm , 743 μm ; fore tibia 317-364 μm , 344 μm ; mid femur 720-760 μm , 737 μm ; mid tibia 590-637 μm , 615 μm ; hind femur 738-810 μm , 774 μm and hind tibia 774-814 μm , 795 μm . All tarsal segments missing.

Hypopygium (Figs 1A-D). Tergite IX 95-107 μm , 99 μm long with 4-6 median setae, 16-20 apical setae; apical margin with small lateral shoulders; lateral teeth not visible; microtrichia free area between anal crests. Anal point 45-61 μm , 57 μm long, 14-16 μm wide at base and 8-9 μm wide at apex. Anal point with 8-10 spines between well developed anal crests. Anal tergite bands caudally curved, almost




Fig. 1. *Tanytarsus usambarae*, spec. nov. A. Hypopygium ventral view. B. Hypopygium dorsal view. C. Superior volsella. D. Median volsella. E. Thorax.

reaching anal point, with median elongations almost connecting the tergite bands. Transverse sternapodeme 50-75 μm , 59 μm long; phallapodeme forked, 86-109 μm , 96 μm long, anterolateral branch being shorter than anteromedian branch. Gonocoxite 100-143 μm , 118 μm long, gonostylus sausage-shaped, 89-114 μm , 104 μm long. Superior volsella more or less oval with small tubercle on posterior margin (Fig. 1C); bearing 9-12 weak dorsal setae and 3 median setae, the apical 2 stronger than the basal one; dorsolateral microtrichia present; digitus absent. Median volsella (Fig. 1D) 82-95 μm , 88 μm long including about 20, 20-45 μm long, simple lamellae reaching beyond apex of inferior volsella. Inferior volsella, club-shaped, comparatively straight, 70-89 μm , 78 μm long, with strong apical setae. HR 1.02-1.25, 1.13; HV 2.54-2.84, 2.68 ($n = 3$).

Systematics

African *Tanytarsus* species generally do not conform well to European species group concepts (Ekrem 1999b), and although *T. usambarae*, spec. nov. shares some characteristics with the species of the *T. lugens* and *T. gregarius* groups, it does not fit the group diagnoses. Based on the adult male morphology, *T. usambarae* keys to the *T. lugens* group when the few microtrichia between the anal crests are ignored (Cranston et al. 1989). When regarded as possessing a field of microtrichia between the anal crests, *T. usambarae* keys to the *T. gregarius* species group. However, although the microtrichia do extend to in between the anal crests, they do not constitute a field as seen in *T. gregarius* Kieffer and *T. inaequalis* Goetghebuer. We therefore regard a field of microtrichia not to be present between the anal crests.

The new species differs from the group diagnosis of the *T. lugens* group (Reiss & Fittkau 1971) in having $AR < 1.4$, a smooth anal point apex, and a particularly large median volsella with numerous simple lamellae. *T. usambarae*, spec. nov. shares these characters with its probable sister species, *T. luctuosus*, from which it differs only in having a posterior tubercle on the superior volsella, pale yellowish abdominal tergites I, IV and V, and in lacking a scutal tubercle. *T. usambarae* and *T. luctuosus* co-exist in the West Usambara Mts., and both were found in the material collected by the same Malaise trap.

Habitat

The type locality of *T. usambarae* is situated close to Mazumbai Forest Reserve on Kwagoroto Hill in the West Usambara Mts. Kaputu stream originates at about 1860 m a.s.l. and runs through almost undisturbed rain forest before it empties into a marshy area at about 1400 m (Andersen & Johanson 1992). The type sample was taken at an altitude of about 1420 m. Here the stream varies from 0.5-2 m in width, and is 10-20 cm deep. The current is regarded as moderate (Andersen & Johanson 1992). The bottom substrate is varied with stones, fine sand and mud and contains some plant litter. At the time of collection, the water temperature varied between 14.6 °C and 17.6 °C, and the pH was measured to be 5.9.

Acknowledgments

The authors wish to thank Ole A. Sæther and Trond Andersen (Bergen) for critically reading the manuscript, and for providing the first author research facilities in Bergen. Thanks also to John Chainey (London) for lending us the type material of *T. luctuosus*, to Gerhard Haszprunar, director of ZSM, for permission to work with the chironomid collection in Munich, and to Martin Baehr and Martin Spies (Munich) for comments on the final version of the manuscript. The material of *T. usambarae* was collected as part of a joint project between the Faculty of Forestry, Sokoine University of Agriculture (SUA), Morogoro, Tanzania and the Museum of Zoology, University of Bergen, Norway. Financial support for the expedition in Tanzania was given by the Norwegian Research Council (NFR) and the Norwegian Agency for International Development (NORAD). Gladys Ramirez made the slide preparations.

References

- Andersen, T. & K. A. Johanson 1992. Caddis flies (Trichoptera) from a mountain rain forest in NE Tanzania. – In: Otto, C. (ed.): Proc. 7th Int. Symp. Trichoptera: 59-64. – Leiden, The Netherlands
- Cranston, P. S., M. E. Dillon, L. C. V. Pinder & F. Reiss 1989. The adult males of Chironominae (Diptera, Chironomidae) of the Holarctic region. Keys and diagnoses. – In: Wiederholm, T. (ed.): Chironomidae of the Holarctic region. Keys and diagnoses. Part 3. Adult males. – Ent. scand. Suppl. 34: 353-502
- Ekrem, T. 1999a. *Cladotanytarsus bukavus* (Lehmann, 1979) comb. n. and *C. congolensis* (Lehmann, 1979) comb. n. from Central Africa (Diptera: Chironomidae). – Ann. Limnol. 35: 185-191
- 1999b. Six new *Tanytarsus* species from Ghana, West Africa (Insecta, Diptera, Chironomidae). – Spixiana 22: 53-68
- & A. D. Harrison 1999. *Tanytarsus minutipalpus*, spec. nov. from the saline lakes in the Rift Valley, East Africa (Insecta, Diptera, Chironomidae). – Spixiana 22: 199-208
- & F. Reiss 1999. Two new *Tanytarsus* species (Diptera: Chironomidae) from Brazil, with reduced median volsella. – Aquatic Insects 21: 205-213
- Freeman, P. 1958. A study of the Chironomidae (Diptera) of Africa south of the Sahara. Part IV. – Bull. Brit. Mus. nat. Hist. Ent. 7: 331-352
- & P. S. Cranston 1980. II. Family Chironomidae. In: Crosskey, R. W. (ed.): Catalogue of the Diptera of the Afrotropical Region. – British Museum (Natural History): 175-202. – London, England
- Lehmann, J. 1979. Chironomidae (Diptera) aus Fließgewässern Zentralafrikas (Systematik, Ökologie, Verbreitung und Produktionsbiologie), Teil I. Kivu-Gebiet, Ostzaire. – Spixiana Suppl. 3: 1-144
- 1981. Chironomidae (Diptera) aus Fließgewässern Zentralafrikas Teil II: Die Region um Kisangani, Zentralzaire. – Spixiana Suppl. 5: 1-85
- Reiss, F. & E. J. Fittkau 1971. Taxonomie und Ökologie europäisch verbreiteter *Tanytarsus*-Arten (Chironomidae, Diptera). – Arch. Hydrobiol. Suppl. 40: 75-200
- Schlee, D. 1966. Präparation und Ermittlung von Meßwerten an Chironomidae (Diptera). – Gewäss. Abwäss. 41/42: 169-193
- Sæther, O. A. 1969. Some Nearctic Podonominae, Diamesinae and Orthocladinae (Diptera: Chironomidae). – Bull. Fish. Res. Bd. Canada 170: 1-154
- 1980. Glossary of chironomid morphology terminology (Diptera: Chironomidae). – Ent. scand. Suppl. 14: 1-51
- 1990. A review of the genus *Limmophyes* Eaton from the Holarctic and Afrotropical regions (Diptera: Chironomidae, Orthocladinae). – Ent. scand. Suppl. 35: 139 pp.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 2000

Band/Volume: [023](#)

Autor(en)/Author(s): Stur Elisabeth, Ekrem Torbjorn

Artikel/Article: [Tanytarsus usambarae, spec.nov. from West Usambara Mts., Tanzania, East Africa \(Insecta, Diptera, Chironomidae\) 219-223](#)