SPIXIANA 32 1 45–110 München, August 2009	SPIXIANA	32 1	45–110	München, August 2009	ISSN 0341-8391
--	----------	-------------	--------	----------------------	----------------

Revision of the western Palearctic species of *Dusona* Cameron

(Hymenoptera, Ichneumonidae, Campopleginae)

Klaus Horstmann

Horstmann, K. (2009): Revision of the western Palearctic species of *Dusona* Cameron (Insecta, Hymenoptera, Ichneumonidae, Campopleginae). – Spixiana **32/1**: 45-110

The western Palearctic species of *Dusona* Cameron are revised. Fifteen new synonymies are indicated, and lectotypes are designated for five taxa. In a checklist, 109 valid names of identified species, 23 nomina dubia and 110 junior synonyms (including five infrasubspecific names) are listed. A key is provided for 109 species. Detailed descriptions of 46 species are given, among these ten new species (63 trans-palearctic species having been described by Hinz & Horstmann 2004).

Address of author: Dr. Klaus Horstmann, Lehrstuhl Zoologie III, Biozentrum, Am Hubland, 97074 Würzburg, Germany.

Introduction

This paper is an extension of an earlier revision of the eastern Palearctic species of *Dusona* Cameron (Hinz & Horstmann 2004). It is based on type revisions by Hinz (1957, 1961, 1962, 1963a, 1963b, 1972, 1973, 1975, 1976, 1977, 1979, 1990) and on his collection that is deposited in the ZSM. In addition, large series of *Dusona* specimens from the M. Schwarz collection and from museums in Edinburgh and Linz, smaller series from the H. Schnee collection and from museums in Leiden, London and Madrid, and specimens of rare or doubtful species from the K. Schmidt and the K. Zwakhals collections and from museums in Helsinki, Paris, Santa Cruz de Tenerife, Stockholm, Vienna and Wageningen were studied. Several types were re-examind.

Ten species are described as new, none from the British Isles (despite the large amount of material studied), three from central Europe (a low number, in view of the very large amount of material studied), three from Spain, one from Italy, two from eastern Europe and one from Turkey. Apparently the fauna of northern, western and central Europe is well known now, and also the fauna of southern and eastern Europe, and of the Near East is rather well known. Few specimens from northwestern Af-

rica were examined; all belonged to species already described from Europe. No specimen was seen from Egypt, and also Shaumar (1966) reported no *Dusona* species from this country.

Several old authors (Audouin, Brischke, Christ, Gmelin, Pollich, Schrank, Sulzer) described species of Ichneumonidae from Europe, which possibly belong to *Dusona* Cameron, but the descriptions are insufficient and the types are lost. These names remain nomina dubia.

The recent studies were supported by many colleagues: E. Diller and S. Schmidt (Zoologische Staatssammlung, Munich, Germany = ZSM) allowed me to remove temporarily the Dusona material of the Hinz collection from Munich to my home in Würzburg. G. Broad (Natural History Museum, London, U.K.), S. Csösz (Magyar Természettudományi Múzeum, Budapest, Hungary), R. Danielsson (Zoologiska Institutionen, Lund, Sweden), A. Freitag (Musée Zoologique, Lausanne, Switzerland), F. Gusenleitner (Biologiezentrum, Landesmuseum, Linz, Austria), A. G. Irwin (Norfolk Museums Service, Norwich, U.K.), I. Izquierdo and M. Paris (Museo Nacional de Ciencias Naturales, Madrid, Spain), Y. Jongema (Laboratory of Entomology, Wageningen, Netherlands), D. R. Kasparyan (Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia), J.-P. Kopelke (Naturmuseum Senckenberg, Frankfurt, Germany), P. Malinen (Zoological Museum, Helsinki, Finland), G. Ortega (Museo de Ciencias Naturales, Santa Cruz de Tenerife, Spain), J. Sawoniewicz (Białystok, Poland), K. Schmidt (Heidelberg, Germany), H. Schnee (Markkleeberg, Germany), M. Schwarz (Kirchschlag, Austria), M. R. Shaw (National Museums of Scotland, Edinburgh, U.K.), A. Taeger (Deutsches Entomologisches Institut, Müncheberg, Germany), E. Tremblay (Museo Zoologia, Napoli, Italy), H. Vårdal (Naturhistoriska Riksmuseet, Stockholm, Sweden), C. Villemant (Muséum National d'Histoire Naturelle, Paris, France), M. Vizek (Naturhistorisches Museum, Vienna, Austria), R. de Vries (Nationaal Natuurhistorisch Museum, Leiden, Netherlands), and C. J. Zwakhals (Arkel, Netherlands) sent types and other material for comparison. C. W. Moeliker (Natuurmuseum, Rotterdam, Netherlands) informed me about two types in the Rotterdam museum. M. R. Shaw assisted me with the English language. I am indebted to them all.

Explanations of terms

Special terms used in the descriptions of Ichneumonidae were defined and figured by Townes (1969: 36 ff.). Some important characters used for the differentiation of *Dusona* species were described and figured by Hinz & Horstmann (2004: 5 ff.). Short definitions of some terms are given here.

- Antennal carina: Carina surrounding the antennal socket. The descriptions refer to the dorsal part only.
- Black lateral stripe: A structure on either side of the 3rd gastral tergite in most of those species in which the epipleurum is not separated from its tergite. It is a horizontal carina on the inside of the tergite, well separated from the ventrolateral edge, which is marked with black, the black colour showing through the tergite.
- Epipleurum (= laterotergite): Lateral part of the 3rd gastral tergite, which is completely or partly or not at all separated from the remaining part of that tergite by a horizontal crease or bulge. If the epipleurum is separated from its tergite, it is usually folded in and concealed on the inside of the latter, the crease being marked with black. If the epipleurum is separated only on the anterior 0.3 or less of the tergite, it is sometimes folded back secondarily and pendent as a triangular or semicircular lobe. If the epipleurum is not separated from its tergite, the former horizontal crease is often replaced by a black lateral stripe.
- Genal carina: Carina separating the temple and the gena from the occiput.

- Glymma: A large or small groove or pit on either side of the petiole.
- Lateral area of the petiole: A narrow area on either side of the petiole, being bordered by fine carinae or defined by a change in the sculpture of the surface (coriaceous instead of smooth).
- Median longitudinal carinae: Carinae of the propodeum bordering the area superomedia and the area petiolaris laterally.
- Mesopleurum: The descriptions usually refer to the central part only. The speculum and the depression in front of the speculum are mentioned separately.
- Oral carina: Carina separating the occiput from the mouth opening.
- Petiolar suture: Suture separating the 1st gastral sternite from its tergite.
- Pleural carina: Carina bordering the metapleurum dorsally and posteriorly, thus separating it from the propodeum.
- Prepectal carina (= epicnemial carina): Carina separating the mesopleurum and mesosternum from the prepectus (= epicnemium). The ventral part lies between its junctions with the transverse carinae; the pleural parts extend between their junctions with the transverse carinae and the anterior edges of the mesopleura.
- Punctation: In a dense punctation the punctures are for the greater part wider than their interspaces. In a dispersed punctation the punctures are for the greater part narrower than their interspaces. In a rather dense punctation many punctures are as wide as their interspaces.
- Radial wrinkles: Wrinkles on the ventral part of the frons, running upwards and sidewards from the dorsal part of the antennal carina.
- Red colour: The colour of the median gastral tergites of most species is decribed as red, but it varies between light reddish brown and dark reddish brown both within and between several species.
- Spiracular carina: Carina connecting the propodeal spiracle with the pleural carina.
- Tangential lines on eyes and temples: Imaginary lines defined to describe the slope of the temples.
- Tarsal claws: The hind tarsal claws are described as short (about as long as the empodium) or elongated (distinctly longer than the empodium).
- Thyridium: The thyridium is described as circular (1.0-1.5 times as long as wide), oval (about 2.0 times as long as wide) or elongated (more than 2.5 times as long as wide).
- Transverse carina: Carina separating the ventral from the pleural part of the prepectus.
- Yellow colour: The yellow colour of the median gastral tergites in some *Dusona* species is rather often partly replaced by dark brown or black marks (with a different pattern in different specimens). Possibly this is a post-mortem effect which is not taken into account in the key.

Definitions of indices

- Epipleural index: Ratio of the length of the crease or bulge separating the epipleurum from the 3rd gastral tergite to the lateral length of that tergite.
- Genal index: Ratio of the length of the oral carina between its junction with the genal carina and the base of mandible to the basal width of a mandible.
- Malar space index: Ratio of the shortest distance between the rim bordering the mandibular grouve and the eye (= malar space) to the basal width of a mandible.
- Nervellar index: Ratio of the length of the upper (anterior) branch of the nervellus (between its junctions with the mediella and the discoidella) to the total length of the nervellus.
- Ocellar index: Ratio of the shortest distance between the eye and the hind ocellus to the longest diameter of the hind ocellus.
- Ovipositor index: Ratio of the visible length of the ovipositor sheaths (seen from the side, in normal position) to the length of a hind basitarsus.
- Spiracular index: Ratio of the length of the spiracular carina (or, if that carina is obliterated, the distance between the spiracle and the pleural carina) to the length of the spiracle.
- Thyridial index: Ratio of the shortest distance between the thyridium and the anterior edge of the 2nd gastral tergite to the length of the thyridium.
- Tibial spur index: Ratio of the length of the shorter to the length of the longer spur of the mid tibia.

Taxonomic notes

Campoplex adjunctus Förster forma discrepans Habermehl

Lectotype (\mathfrak{P}) here designated: "Würzburg 1.6.20 Dr. Enslin", "Karlstadt", "Campoplex adjunctus Först. \mathfrak{P} " (Mus. Frankfurt). A paralectotype (\mathfrak{F}) from Worms is also present in Frankfurt. The taxon is a junior homonym of *C. discrepans* Förster and a junior synonym of *Dusona genalis* (Thomson) (syn. nov.).

Campoplex adriaansei Teunissen

This taxon is a junior synonym of *Dusona carpathica* (Szépligeti) (syn. nov.), according to some specimens in the Hinz collection (ZSM) determined as *C. adriaansei* by Teunissen. The holotype of *C. adriaansei* is preserved in the Rotterdam museum; a postal loan was refused.

Campoplex anceps Holmgren

Lectotype (\$\times\$) here designated (labelled by Hinz): "Hlm" (= Stockholm), "Bhn" (= Boheman) (Mus. Stockholm).

Campoplex angustifrons Förster

Dusona angustifrons (Förster) is variable and poorly differentiated from related species, in particular from D. minor (Provancher) (see key, couplet 49). Usually the flagellum is filiform, with the shortest segment being 0.9-1.0 times as long as wide, but in some specimens the flagellum is slightly fusiform and the shortest segment is 0.8-0.9 times as long as wide, and some specimens are intermediate in this character. In most specimens the genal carina is clearly bent forwards subventrally, but in the holotype ($\mathfrak P$) of Campoplex obreptans Förster, which is tentatively synonymised with D. angustifrons (Hinz 1963: 2), the genal carina is almost straight subventrally. Possibly D. angustifrons as defined here represents a species group.

Campoplex annexus Förster

The lectotype (\mathfrak{P}) of this taxon (Bachmeier 1979: 73) was misinterpreted as conspecific with *Dusona limnobia* (Thomson) by Hinz (1962: 2) and subsequent authors. In fact it belongs to *Dusona americana* (Ashmead) (syn. nov.). Therefore *D. annexa* (Förster) is the valid name for *D. americana*, and it is not a senior synonym of *D. limnobia* (as erroneously published by Hinz & Horstmann 2004: 46). *D. oyamadai* Hinz is another synonym of *D. annexa* (syn. nov.), according to a female and a male from Sakhalin (Russia) in the Hinz collection (ZSM).

Campoplex blandus Förster and C. remotus Förster

Both taxa were described in the same paper. Thomson (1887: 1085) synonymised them and used *C. blandus* as the valid name of the species, thus acting as first reviser (article 24.2.2 of the Code). Although Dalla Torre (1901: 151) and most subsequent authors used *C. remotus* as the valid name, the decision by Thomson should be followed. *C. blandus* cannot be considered as nomen oblitum under article 23.9 of the Code, because the name was used after 1899 by several authors as the valid name of a taxon (see Yu & Horstmann 1997: 152).

Campoplex brevicornis Brischke

The Brischke collection was destroyed, and the types preserved in it are lost. Two syntypes $(1^{\circ}, 1^{\circ})$ are preserved in the Strobl collection in Admont (Strobl 1904: 65), and I studied them many years ago. Both are damaged. The female might belong to *Dusona carpathica* (Szépligeti), but I noted a few differences. I could not place the male. The types should be reexamined, but at present they are unavailable.

Campoplex bucculentus Holmgren

In this species, the prepectal carina is very variable. In some specimens the ventral part merges with the pleural part (and both are low), in others the ventral part merges with the transverse carina (and both are raised). In the latter case, the pleural part of the prepectal carina is present medially and ventrally, or it is present only medially, or it is completely obliterated.

Delopia canariensis Hinz

This taxon was described as a subspecies of *Dusona semiflava* (Costa). But it differs from this species and from the closely related *D. dubitor* Hinz by the characters noted in the key (couplet 77), and should be accorded full species status (stat. nov.). Prior to its description (Hinz 1990: 267), material of this taxon from the Canary Isles was mentioned by Horstmann (1980: 133) under the name *D. semiflava* and by Rey & Izquierdo (1985: 318) under the name *D. dubitor*.

Campoplex carinifer Teunissen

This taxon is a junior synonym of *Dusona montana* (Roman) (syn. nov.), according to three paratypes $(\mathfrak{P}\mathfrak{P})$ of *C. carinifer* in the Teunissen collection (Mus. Leiden) and some specimens in the Hinz collection (ZSM) determined as *C. carinifer* by Teunissen. In his description of the species, Teunissen (1947: 269) forgot to mention the locality label and the depository of the holotype (\mathfrak{P}) . It could not be found.

Campoplex contumax Förster and C. prominulus Förster

Both taxa were described in the same paper. Thomson (1887: 1068) synonymised them and used *C. prominulus* as the valid name of the species, thus acting as first reviser (article 24.2.2 of the Code). Although Dalla Torre (1901: 140) and most subsequent authors used *C. contumax* as the valid name, the decision by Thomson should be followed. *Campoplex prominulus* cannot be considered as nomen oblitum under article 23.9 of the Code, because the name was used after 1899 by several authors as the valid name of a taxon (see Yu & Horstmann 1997: 145).

Campoplex cornellus Teunissen

This taxon is tentatively placed here as a junior synonym of *Dusona angustifrons* (Förster) (? syn. nov.), according to a label added by Teunissen to the holotype (\mathfrak{P}) of his species preserved in the Rotterdam museum (Moeliker in lit.). The type should be examined, but a postal loan was refused.

Campoplex crassipes Thomson

The holotype (\mathfrak{P}) of this taxon (Hinz 1963b: 339) is a rather large specimen of *Dusona aversa* (Förster) (syn. nov.).

Campoplex debilis Förster

The holotype (\mathfrak{P}) of this taxon (Hinz 1962: 10) is a faded specimen of *Dusona flagellator* (Fabricius) (syn. nov.).

Campoplex facialis Holmgren

Apparently the lectotype of this taxon (Hinz 1963b: 343), the gaster of which is lacking, is a female (sculpture of propodeum, colour pattern of mandible). It is a junior synonym of *Dusona annexa* (Förster) (Thomson 1887: 1088) (see above). *Campoplex facialis* Holmgren is a junior homonym of *C. facialis* Boie and was renamed *C. mariae* by Dalla Torre (1901: 146).

Dusona flagellator (Fabricius)

A variety of this species from Kazakhstan was described by Hinz & Horstmann (2004: 72). Apparently the same variety occurs in Europe (see key, couplets 61 and 64). Its status is still unknown.

Campoplex incompletus Bridgman

The holotype (3) of this taxon (Hinz 1972: 47) is very similar to *Dusona tenuis* (Förster) and differs only by colour characters (see key, couplet 101). Several males of *D. incompleta* (Bridgman) from Scotland, England and France were studied (Mus. Edinburgh, London, ZSM). One female from Spain (ZSM) is tentatively associated with these males. Possibly *D. incompleta* is a dark morph of *D. tenuis. Campoplex incompletus* sensu Morley (1915: 87) belongs to *D. leptogaster* (Holmgren), and *D. incompleta* sensu Aubert (1981: 21) belongs to *D. tenuis*.

Campoplex kriechbaumeri Costa

The holotype (\$\partial\$) of this taxon (Scaramozzino 1994: 18) belongs to *Casinaria* Holmgren (comb. nov.).

Delopia lautareti Hinz

The types of this taxon (Coll. Hinz/ZSM) belong to *Dusona genalis* (Thomson) (syn. nov.).

Campoplex libauensis Strand

The holotype (\$) of this taxon (Oehlke 1963: 407) from Libau (= Liepaja/Latvia) belongs to *Dusona anceps* (Holmgren) (syn. nov.).

Campoplex limnobius Thomson

Previous interpretations and descriptions of *Dusona limnobia* (Thomson), including the description by Hinz & Horstmann (2004: 46 f.) under the name *D. annexa* (Förster), also contain the hitherto undescribed species *D. rubidatae* spec. nov. Both species and the closely related species *D. annexa* (syn. *Casinaria americana* Ashmead; see above) are differentiated in the key (couplets 120-121). Specimens previously determined as *D. americana, annexa* or *limnobia* should be re-examined.

Campoplex mactator Förster

In this species, the frons is usually slightly depressed above the antennal sockets, and the depressions are covered with distinct radial wrinkles. In two rather small females (58 flagellar segments; body length 14 mm) from Finland (Mus. Helsinki, ZSM), the depressions are somewhat deeper and coriaceous, whereas the radial wrinkles are short and inconspicuous. The significance of this variation is unknown.

Campoplex nobilitatus Holmgren

Lectotype (a) here designated: "Fg" (= Fogelsång/Scåne), "85." (characteristic number used in the Thomson collection for material determined by Holmgren), Thomson collection (Mus. Lund). The taxon is a junior synonym of *Dusona myrtilla* (Desvignes) (Morley 1915: 76).

Campoplex notabilis Förster

Dusona notabilis (Förster) has not been well understood. Its interpretation is here restricted to material from Europe placed to D. scolator Hinz by Hinz & Horstmann (2004: 151) (see below). D. notabilis differs from the latter by: body length 7-9 mm; mesopleurum distinctly coriaceous and rather dull between the punctures; propodeum without distinct transverse wrinkles or with a few transverse wrinkles in the sulcus; maxillary palpus dark brown. Specimens previously determined as D. notabilis should be re-examined.

Campoplex pineticola Holmgren

In populations of *Dusona pineticola* (Holmgren) in central Europe (Germany, Austria, Switzerland, France; ZSM) the hind tibia is yellowish red or yellowish brown medially, narrowly black basally and narrowly brown apically, and the 5th gastral tergite is yellowish red. The holotype of *Campoplex litigiosus* Habermehl from Black Forest, Germany (Hinz 1963a: 117) belongs to this morph. Hinz (1985: 307)

described a subspecies *sibirica* from Siberia (Chita, Irkutsk), in which both the hind tibia and the 5th gastral tergite are black. But short series of *D. pineticola* from England and Scotland (Mus. Edinburgh) and from central Sweden (Mus. Stockholm) agree in colour with the subspecies *sibirica*, which therefore appears to occur in northern Europe too. Unfortunately, the lectotype of *C. pineticola* which does not bear a locality label but might originate from southern Sweden (Hinz 1963b: 342) is intermediate in colour pattern. Therefore, in this paper the subdivision of *D. pineticola* into subspecies is not accepted.

Campoplex praecox Teunissen

This taxon is a junior synonym of *Dusona pulchripes* (Holmgren) (syn. nov.), according to two paratypes (\mathfrak{P}) of *C. praecox* in the Wageningen museum and one paratype (\mathfrak{P}) in the Hinz collection (ZSM). The holotype (\mathfrak{P}) could not be found in the Wageningen museum.

Heterocharops pugillatorioides Constantineanu

Constantineanu (1932: 77) proposed this name as a junior synonym of *Campoplex pugillator* (Linnaeus). Therefore both the genus and the species names are unavailable (article 11.6 of the Code).

Dusona pulmentariae Hinz

Contrary to the information given in the original description (Hinz 1963a: 119), the holotype (\mathfrak{P}) of this taxon (from Bisamberg near Vienna) is preserved in the Vienna museum.

Campoplex rufipes Holmgren

The lectotype (♂) of this taxon (Hinz 1963b: 343) from "Itl." (= Jemtland/Sweden) belongs to *Dusona alticola* (Gravenhorst) (syn. nov.). *Campoplex rufipes* Holmgren is a junior homonym of *C. rufipes* Gravenhorst and was renamed *C. infelix* by Dalla Torre (1901: 144).

Dusona scolator Hinz

The material from Europe mentioned by Hinz & Horstmann (2004: 151) under this name belongs to *Dusona notabilis* (Förster) (see above). The material of *D. scolator* from Siberia (including the holotype) differs from *D. notabilis* by: body length 6-7 mm; mesopleurum slightly coriaceous and distinctly shining between the punctures; sulcus of propodeum with transverse wrinkles posteriorly (\mathfrak{P}) or medially and posteriorly (\mathfrak{F}); maxillary palpus yellowish red.

Campoplex sericeus Brischke

Lectotype (♀) here designated: "Zoppot 7/8 1879" (= Sopot/Poland), Strobl collection (Mus. Admont) (see Strobl 1904: 65). The taxon is a junior homonym of Campoplex sericeus Holmgren and a junior synonym of Dusona vidua (Gravenhorst) (Schmiedeknecht 1908: 1590).

Campoplex stenocarus Thomson

This taxon, the holotype of which is lost (Fitton 1982: 21), is a junior synonym of *Dusona minor* (Provancher) (syn. nov.), according to a non-type (♀) of *C. steno*carus in the Thomson collection (Mus. Lund) (Hinz 1963b: 339) and material of D. minor from the Nearctic region in the Hinz collection (ZSM).

Campoplex stygius Förster

This taxon is characterized by the shape of the antennal carina and by the position of the propodeal spiracle and the absence of the spiracular carina (see key, couplet 93), but it is variable in two important characters used in the key. The ventral part of the prepectal carina merges with the pleural part, or with the transverse carina. And the anterior part of the ventrolateral edge of the 3rd gastral tergite is sometimes bent inwards, thus forming an epipleurum (epipleural index 0.1-0.2), although a black lateral stripe is situated clearly distant from the ventrolateral edge.

Campoplex subaequalis Förster

The lectotype (δ) of this taxon (Hinz 1962: 7) belongs to Dusona rugifer (Förster) (syn. nov.), both taxa being described in the same paper. D. rugifer is considered here as the valid name of the species.

Campoplex subsulcatus Holmgren

Apparently the holotype of this taxon (Hinz 1963b: 343), the genitalia of which are lacking, is a female (proportions of basal flagellar segments, sculpture of propodeum). Possibly the apex of the gaster was already lacking when Holmgren (1872: 66) described the species. The taxon is a junior synonym of Dusona circumspectans (Förster) (Yu & Horstmann 1997: 145).

Dusona tenerifae Hinz

This species was described from the Canary Isles and is rather often collected on Tenerife. One δ from Bigastro (near Murcia/Spain) (Mus. Madrid) differs from Dusona tenerifae Hinz by the characters noted in the key (couplet 46). It is treated here as a variety of D. tenerifae, but it might represent a separate species. The specimen is damaged (parts of the antennae and the hind legs broken off) and is unsuited for a description. In contrast, the specimen mentioned by Hinz (1977: 50) supposedly from Austria (ZSM) agrees well with D. tenerifae. So far, it is the only specimen of the typical morph known from a locality outside of the Canary Isles. In my opinion, an incorrect locality label cannot be ruled out.

Campoplex tschekii Holmgren

Lectotype (♀) here designated: "Dgb." (= Degeberga/Scåne), "59" (characteristic number used in the Thomson collection for material determined by Holmgren), Thomson collection (Mus. Lund). Holmgren (1872: 54) described the species from a pair collected in copula. Both specimens are preserved in the Thomson collection, the δ being labelled as paralectotype. The taxon is a junior synonym of Dusona aversa (Förster) (Thomson 1887: 1079).

Checklist of western Palearctic species

Those taxa the types of which were recently revised by the present author are marked with an asterisk.

Dusona abdominator Hinz, 1985*

Dusona admontina (Speiser, 1908) (nom. nov. for C. rufiventris Strobl)

Campoplex rufiventris Strobl, 1904 (praeocc.)

Dusona aemula (Förster, 1868)*

Campoplex disseptus Förster, 1868

Campoplex parvulus Förster, 1868

Campoplex discrepans Förster, 1868

Campoplex filicornis Holmgren, 1872

Dusona affinis (Brischke, 1880) (nomen dubium)

Dusona albobasalis spec. nov.

Dusona alpigena Hinz, 1972*

Dusona alpina (Strobl, 1904)

Campoplex lateralis Kriechbaumer, 1883 (praeocc.)

Dusona alticola (Gravenhorst, 1829)

Campoplex floricola Gravenhorst, 1829

Campoplex rufipes Holmgren, 1872* (praeocc.) (syn. nov.)

Campoplex infelix Dalla Torre, 1901 (nom. nov.

for *C. rufipes* Holmgren) (syn. nov.)

Dusona latiseta Hinz, 1961

Dusona anceps (Holmgren, 1860)*

Campoplex disparilis Förster, 1868

Campoplex auriculatus Förster, 1868

Campoplex costulatus Bridgman et Fitch, 1885

Campoplex costulatus Bridgman, 1886 (praeocc.)

Campoplex libauensis Strand, 1918* (syn. nov.)

Dusona andalusica spec. nov.

Dusona angustata (Thomson, 1887)

Dusona angustifrons (Förster, 1868)*

Campoplex obreptans Förster, 1868* (? svn.)

Campoplex zonellus Förster, 1868*

Campoplex cornellus Teunissen, 1947 (? syn. nov.)

Dusona annexa (Förster, 1868)*

Casinaria americana Ashmead, 1890* (syn. nov.) Campoplex facialis Holmgren, 1872* (praeocc.)

Campoplex mariae Dalla Torre, 1901 (nom. nov.

for C. facialis Holmgren)

Dusona oyamadai Hinz, 1994 (syn. nov.)

Dusona areolator (Holmgren, 1856) (nomen dubium)

Dusona auriculator Aubert, 1864*

Dusona aurita (Kriechbaumer, 1883)

Dusona aversa (Förster, 1868)

Campoplex dubiosus Förster, 1868*

Campoplex tschekii Holmgren, 1872*

Campoplex crassipes Thomson, 1887* (syn. nov.)

Dusona baueri Hinz, 1973*

Dusona bellipes (Holmgren, 1872)

Dusona vernalis Hinz, 1957*

Dusona bicolor (Brischke, 1880) (nomen dubium)

Dusona bicoloripes (Ashmead, 1906)*

Campoplex foersteri Roman, 1942*

Dusona blanda (Förster, 1868)

Campoplex remotus Förster, 1868

Campoplex forsselli Holmgren, 1872

Casinaria punctiventris Woldstedt, 1877

Dusona breviceps spec. nov.

Dusona brevicornis (Brischke, 1880) (nomen dubium)

Dusona brischkei (Dalla Torre, 1901) (nom. nov. for

C. lateralis Brischke) (nomen dubium)

Campoplex lateralis Brischke, 1892 (praeocc.)

Dusona bucculenta (Holmgren, 1860)

Campoplex melampus Förster, 1868

Dusona calceata (Brauns, 1895)

Dusona canariensis (Hinz, 1990)* Dusona carinifrons (Holmgren, 1860)

Campoplex minax Förster, 1868

Campoplex geometrae Rudow, 1883

Dusona carpathica (Szépligeti, 1916)*

Campoplex adriaansei Teunissen, 1947 (syn. nov.)

Dusona carpinellae (Schrank, 1802) (nomen dubium)

Dusona castanipes (Thomson, 1887)*

Dusona circumcinctus (Förster, 1868)

Campoplex subcinctus Förster, 1868 (praeocc.) Campoplex subcinctus var. lateralis Ulbricht, 1916

(infrasubspecific)

Dusona circumspectans (Förster, 1868)

Campoplex vagulus Förster, 1868

Campoplex subsulcatus Holmgren, 1872*

Dusona compressa (Sulzer, 1776) (nomen dubium)

Dusona compressa (Christ, 1791) (praeocc.) (nomen dubium)

Dusona compressa (Schrank, 1802) (praeocc.) (nomen dubium)

Dusona confusa (Förster, 1868)

Campoplex lacunosus Kriechbaumer, 1883

Campoplex consimilis Schmiedeknecht, 1908

Dusona consobrina (Holmgren, 1872) (nomen dubium)

Dusona constantineanui Hinz, 1977

Dusona cultrator (Gravenhorst, 1829)

Campoplex cultrator var. nigriventris Ulbricht, 1916 (infrasubspecific)

Dusona disclusa (Förster, 1868)

Dusona dositheae (Audouin, 1834) (nomen dubium)

Dusona dubitor Hinz, 1977*

Dusona einbecki Hinz, 1977*

Dusona erythrogaster (Förster, 1868)

Campoplex indefessus Förster, 1868

Dusona exsculpta (Brischke, 1880) (nomen dubium)

Dusona falcator (Fabricius, 1775)*

Dusona fatigator (Förster, 1868)*

Dusona filator Hinz, 1985*

Dusona flagellator (Fabricius, 1793)*

Campoplex debilis Förster, 1868* (syn. nov.)

Campoplex heterocerus Förster, 1868

Dusona fundator (Hinz, 1990)*

Dusona genalis (Thomson, 1887)*

Campoplex adjunctus f. discrepans Habermehl,

1922* (praeocc.) (syn. nov.)

Delopia lautareti Hinz, 1990* (syn. nov.)

Dusona gracillima (Dalla Torre, 1901) (nom. nov. for C. gracilis Brischke) (nomen dubium)

Campoplex gracilis Brischke, 1888 (praeocc.)

Dusona habermehli (Kriechbaumer, 1898)

Dusona hastulatae spec. nov.

Dusona hispanica spec. nov.

Dusona holmgrenii (Dalla Torre, 1901) (nom. nov. for

C. unicinctus Holmgren)

Campoplex unicinctus Holmgren, 1872* (praeocc.)

Dusona humilis (Förster, 1868)

Campoplex eurynotus Holmgren, 1872

Dusona incompleta (Bridgman, 1889)*

Dusona inermis (Förster, 1868)

Dusona infesta (Förster, 1868)

Campoplex terrificus Förster, 1868

Campoplex infestus var. moldavicus Constantineanu, 1932 (infrasubspecific)

Dusona insignita (Förster, 1868)

Campoplex bistrigosus Holmgren, 1872

Dusona intelligator Aubert, 1966*

Dusona irregularis spec. nov.

Dusona juvenilis (Förster, 1868)

Ichneumon victor Thunberg, 1824 (praeocc.)

Campoplex monozonus Förster, 1868

Campoplex monozonus var. rufifemur Ulbricht, 1926 (infrasubspecific)

Campoplex victor var. rufifemur Ozols, 1959

Dusona lamellator Aubert, 1960*

Dusona leptogaster (Holmgren, 1860)

Campoplex macrostylus Förster, 1868 Campoples silvicola Habermehl, 1922

Dusona libertatis (Teunissen, 1947)*

Dusona limnobia (Thomson, 1887)*

Dusona lividariae Hinz, 1963

Dusona longiseta Hinz, 1961

Dusona luteipes (Thomson, 1887)*

Dusona mactator (Förster, 1868)

Campoplex similatorius Holmgren, 1872

Dusona mercator (Fabricius, 1793)*

Ichneumon venditor Thunberg, 1824 (emendation)

Campoplex oxyacanthae Boie, 1855

Campoplex mesoxanthus Förster, 1868*

Dusona meridionator Aubert, 1960*

Dusona minuta (Holmgren, 1856) (nomen dubium)

Dusona minor (Provancher, 1879)

Campoplex stenocarus Thomson, 1887* (syn. nov.)

Dusona mixta (Pollich, 1781) (nomen dubium)

Dusona montana (Roman, 1929)

Campoplex carinifer Teunissen, 1947* (syn. nov.)

Dusona myrtilla (Desvignes, 1856)

Campoplex tenthredinum Tschek, 1871

Campoplex nobilitatus Holmgren, 1872*

Dusona nebulosa Horstmann, 2004*

Dusona nidulator (Fabricius, 1804)*

Campoplex nitidulator Holmgren, 1856

Campoplex vindex Förster, 1868

Campoplex circumscriptus Förster, 1868

Campoplex martialis Förster, 1868

Campoplex bifidus Thomson, 1887

Campoplex nitidulator var. obscurus Kiss, 1926*

Dusona notabilis (Förster, 1868)*

Dusona obliterata (Holmgren, 1872)

Campoplex limiventris Kriechbaumer, 1883

Dusona occisor (Schrank, 1776) (nomen dubium)

Dusona opaca (Thomson, 1887)

Dusona perditor (Förster, 1868)

Campoplex transversostriatus Schmiedeknecht, 1908

Dusona peregrina (Wollaston, 1858)*

Dusona petiolaris (Brischke, 1880) (nomen dubium)

Dusona petiolator (Fabricius, 1804)*

Campoplex lapponicus Holmgren, 1860

Campoplex callizonus Förster, 1868

Campoplex punctatus Bridgman et Fitch, 1885

Campoplex punctatus Bridgman, 1886 (praeocc.)

Campoplex terebrator var. niger Kiss, 1924* (prae-occ.)

Dusona pineticola (Holmgren, 1872)*

Campoplex litigiosus Habermehl, 1922*

Dusona pineticola sibirica Hinz, 1985* (syn. nov.)

Dusona polita (Förster, 1868)

Campoplex flavipalpus Förster, 1868

Campoplex spoliator Förster, 1868

Campoplex medianus Förster, 1868

Campoplex trisculptus Holmgren, 1872

Campoplex femorator Bridgman et Fitch, 1885

Campoplex femorator Bridgman, 1886 (praeocc.)

Campoplex splendens Thomson, 1887

Campoplex latungulus Thomson, 1887

Dusona prominula (Förster, 1868)

Campoplex contumax Förster, 1868

Campoplex foveolatus Förster, 1868

Dusona pugillator (Linnaeus, 1758)*

Campoplex canaliculatus Förster, 1868 (praeocc.)

Dusona pulchripes (Holmgren, 1872)

Campoplex praecox Teunissen, 1947 (syn. nov.)

Dusona pulmentariae Hinz, 1963*

Dusona recta (Thomson, 1887)

Dusona rossica Hinz, 1979

Dusona rubidatae spec. nov.

Dusona rufonigra (Brischke, 1880) (nomen dubium)

Dusona rugifer (Förster, 1868)*

Campoplex subaequalis Förster, 1868* (syn. nov.)

Campoplex punctus Kriechbaumer, 1883*

Dusona rugulosa (Förster, 1868)

Dusona sagittatae spec. nov.

Dusona semiflava (Ĉosta, 1883)

Campoplex flaviscapus Thomson, 1887

Dusona signator (Brauns, 1895)

Dusona sobolicida (Förster, 1868)

Campoplex ulceratus Holmgren, 1872

Dusona spinipes (Thomson, 1887)

Dusona spinulosa (Brischke, 1880) (nomen dubium)

Dusona splenditor Hinz, 1985

Dusona stenogaster (Förster, 1868)

Campoplex monticola Habermehl, 1922

Dusona stragifex (Förster, 1868)*

Campoplex adjunctus Förster, 1868*

Campoplex areolatus Brauns, 1895*

Dusona stygia (Förster, 1868)

Dusona subfalcata (Gmelin, 1790) (nomen dubium)

Dusona subimpressa (Förster, 1868)

Campoplex transitorius Kiss, 1924*

Dusona surrata (Schrank, 1781) (nomen dubium)

Dusona temnator Hinz & Horstmann, 2004*

Dusona tenerifae Hinz, 1977

Dusona tenuis (Förster, 1868)*

Campoplex proximus Förster, 1868*

Campoplex anxius Förster, 1868*

Campoplex peraffinis Förster, 1868*

Campoplex agnatus Förster, 1868*

Dusona terebrator (Förster, 1868)

Campoplegidea terebrator f. postpetiolata Aubert, 1959 (infrasubspecific)

Dusona thomsoni Hinz, 1963*

Dusona tibialis (Brischke, 1880) (praeocc.) (nomen

Dusona tricolorator Aubert, 1970*

Dusona turcator spec. nov.

Dusona tyrolensis spec. nov.

Dusona ucrainica Hinz, 1972*

Dusona validicornis (Holmgren, 1856) (nomen dubium)

Dusona variator (Hinz, 1990)*
Dusona vidua (Gravenhorst, 1869)
Campoplex sericeus Brischke, 1880* (praeocc.)
Dusona vigilator (Förster, 1868)
Dusona xenocampta (Förster, 1868)*
Campoplex polyxanthus Strobl, 1904
Campoplex balticus Habermehl, 1926
Dusona yezoensis (Uchida, 1928)

Key

The key is not strictly dichotomous. In several cases, variable or intermediate species can be traced through both halves of a couplet.

- Epipleurum of the 3rd gastral tergite completely fused with the tergite, the ventrolateral edge thin and more or less rounded, the epipleurum rarely discernible as a small triangular or semicircular pendent lobe anteriorly (e.g. in *D. intelligator*). Black lateral stripe, if present, distinctly distant from the ventrolateral edge and divergent from it. (Some species which are intermediate or variable in these characters can be traced through both halves of the couplet.)
- At least the 3rd gastral tergite marked with red. (In some aberrant specimens of other species the gaster is black, but usually the colour pattern of the hind tibia is different.)
- 3. 45-46 flagellar segments. Mesopleurum with rather dense or dense punctation on a finely coriaceous and shining background. Speculum for the greater part smooth and shining. Sulcus of the propodeum irregularly wrinkled, with a longitudinal keel posteriorly. Glymma present. Epipleural index 0.8. Body length 11-13 mm... myrtilla (Desvignes, 1856) ♀♂

- 30-34 flagellar segments. Mesopleurum with fine dispersed punctation on a distinctly coriaceous and dull background. Speculum coriaceous and rather dull. Sulcus of the propodeum transversely striate medially and posteriorly. Glymma absent. Epipleural index 0.4-0.5. Body length 6-8 mm.... vidua (Gravenhorst, 1829) ♀♂
- Antennal carina strongly raised and conspicuously modified dorsally. Frons slightly or distinctly depressed above the antennal carinae. Epipleural index 0.2-0.5. 52-63 flagellar segments. Body length 14-18 mm.
- Antennal carina at most slightly raised and not conspicuously modified dorsally. Frons at most slightly depressed above the antennal carinae.
 7.

- Rim of the antennal carina slightly bent upwards (a variable character). Frons with radial wrinkles reaching 0.3-0.5 the distance to the median ocellus. Hind femur conspicuously marked with yellowish red distally. Epipleural index 0.2-0.4. 52-58 flagellar segments. Body length 14-17 mm. (See also couplet 23.)
 -pugillator (Linnaeus, 1758) ♀♂

- Genal carina not raised ventrally. Oral carina not distinctly raised......9.
- 8. Genal carina distinctly raised also medially behind the eye. Glymma distinct. Epipleural index 0.2-0.6. Hind tibia light yellow. Median gastral tergites yellow. 58-62 flagellar segments. Body length 14-16 mm.

...... *libertatis* (Teunissen, 1947) ♀♂

- Ventral part of the prepectal carina merging with the pleural part, both of equal height at the junction, the transverse carina at most as high, often lower or obliterated at the junction, rarely raised some distance from the junction.
- 10. 32-39 flagellar segments. Body length 5.5-8 mm......11.

- 47-62 flagellar segments. Body length 8-17 mm......12.
- 11. Genal carina joining the oral carina at the base of the mandible. Hind tibia black, sometimes tinged with dark brown. Gaster from the posterior 0.2-0.3 of the 2nd gastral tergite onwards reddish brown. Posterior tergites tinged with black dorsally in the ♀, broadly black dorsally or completely black in the ♂. 35-39 flagellar segments. Body length 7-8 mm. (See also couplet 57.)erythrogaster (Förster, 1868) ♀♂
- Genal carina joining the oral carina close to the base of the mandible, genal index 0.1. Hind tibia marked with white or yellowish brown medially on the dorsal side. Gaster black posteriorly (from the 4th or the 5th tergite onwards).
 32-35 flagellar segments. Body length 5.5-7 mm.habermehli (Kriechbaumer, 1898) ♀♂

- 14. Only the 2nd and 3rd gastral tergites marked with red. Hind femur red or marked with red. Hind tibia yellowish brown, sometimes narrowly marked with brown basally. Propodeum deeply depressed, the sulcus distinctly transversely striate. Glymma present. 52-59 flagellar segments. Body length 12-16 mm.

.....cultrator (Gravenhorst, 1829) ♀♂

- Gaster more extensively marked with red, at least the 4th tergite marked with red, and/or hind femur black, and/or distinctly smaller with less flagellar segments......15.
- Epipleural index 0.7-0.8. Epipleurum of the 2nd gastral tergite folded in, the crease usually marked with black. (Species with intermediate specimens can be traced through both halves of the couplet.)

- 16. Petiole on either side with a long row of small transverse grooves or short transverse wrinkles in front of the glymma. Epipleural index 0.2. Epipleurum usually folded in, the crease marked with black, in addition with a black lateral stripe almost parallel with the crease and close to it. 46-48 flagellar segments. Body length 10-11 mm.insignita (Förster, 1868) ♀♂
- Petiole smooth or with few wrinkles laterally in front of the glymmae......17.
- 17. Genal index 0.1-0.2. 41-45 flagellar segments, Body length 8-10 mm.18.
- 18. Speculum finely coriaceous or partly smooth, shining. Impression in front of the speculum with fine longitudinal striae medially and ventrally as well as dorsally. Hind tibia yellowish medially (in some specimens dark yellowish brown), tinged with brown apically. Epipleural index 0.5. 41-44 flagellar segments. Body length 8-9 mm....... prominula (Förster, 1868) ♀♂
- Speculum coriaceous, rather dull. Impression in front of the speculum with fine longitudinal striae dorsally, the ventral 0.6 coriaceous and without striae. Hind tibia yellow medially, broadly black apically, the apical ring extending over 0.2-0.3 of the tibia. Epipleural index 0.2-0.3. 42-45 flagellar segments. Body length 8-10 mm.......recta (Thomson, 1887) ♀♂
- 2nd gastral tergite red posteriorly. (In aberrant specimens of some species the 2nd gastral tergite is dorsally black or dark brown posteriorly, but in these the hind femur is marked with red and/ or the epipleural index is at least 0.4.).........21.

......alpina (Strobl, 1904) ♀♂

- 22. 49-53 flagellar segments. Genal index 0.3-0.4. Hind femur black. Hind tibia yellowish brown medially, distinctly black or dark brown basally and apically, the apical ring extending over 0.2-0.3 of the tibia. Body length 9-11 mm. (See also couplet 13.).....genalis (Thomson, 1887) ♀♂
- 23. Frons slightly depressed above the antennal carinae. Antennal carina usually slightly raised and its rim bent upwards, with fine transverse striae (a rather variable character). Epipleural index 0.2-0.4. Hind femur black, conspicuously marked with yellowish red distally. Propodeum broadly and deeply depressed, with distinct transverse wrinkles in the sulcus. 52-58 flagellar segments. Body length 14-17 mm. (See also couplet 6.)............ pugillator (Linnaeus, 1758) ♀♂
- Frons not depressed above the antennal carinae
 Antennal carina low and narrow. Epipleural index 0.4-0.7 (exception: *D. luteipes*; see couplet 25).

- 24. Median and posterior gastral tergites to a variable extent black dorsally and red ventrolaterally. (If the 3rd and 4th tergites are completely red, then the posterior tergites are broadly marked with red laterally. If the posterior tergites are completely black, then the 3rd and 4th tergites are black dorsally and red laterally.) Flagellum stout, 2nd segment 1.3-1.5 times as long as wide. Hind femur red or marked with red. 2nd gastral tergite at least twice as long as wide. 44-48 flagellar segments. Body length 11-12 mm........ *castanipes* (Thomson, 1887) ♀♂
- Gaster with the anterior and posterior tergites black, the median tergites red, at least the 3rd tergite for the greater part or completely red (not black dorsally and red laterally).............25.

......luteipes (Thomson, 1887) ♀

- Hind femur conspicuously marked with black basally. Epipleural index 0.4-0.7. Body length 9-14 mm......26.
- Propodeum slightly or distinctly depressed, with transverse wrinkles in the sulcus at least posteriorly (sculpture rather variable).......28.

- 28. Frons on either side with a small raised point between eye and antennal socket (these points sometimes inconspicuous). Hind femur red on

- the distal 0.2. Hind tibia yellowish red, scarcely darkened basally and apically, at most marked with light brown. 5th gastral tergite red or marked with red. 54-55 flagellar segments. Body length 11-13 mm. *bellipes* (Holmgren, 1872) ♀♂
- Frons without raised points between eye and antennal socket. Hind femur often completely black. Hind tibia yellow to dark brown medially, marked with dark brown or black basally.
 5th gastral tergite black......29.
- 29. 42-47 flagellar segments. Speculum smooth or almost smooth, strongly shining. Mesopleurum slightly coriaceous and distinctly shining between the punctures. Metapleurum with dispersed or rather dense punctation, slightly coriaceous and distinctly shining between the punctures. Gaster distinctly shining posteriorly. (Some specimens are less shining, but are characterized by the lower number of flagellar segments.) Hind tarsal claws with about eight pectinate teeth situated basally, medially and subapically. Hind tibia yellowish brown medially, narrowly dark brown basally and apically. Body length 10-12 mm.

......polita (Förster, 1868) ♀♂

- 30. Hind tibia broadly yellow medially, narrowly brown basally and apically. ♂ with the apices of the front and mid coxae and the mid trochanter yellow, and with a rather narrow area petiolaris (a rather variable character, the difference discernible only by direct comparison of specimens). Epipleural index 0.4-0.5 in the ♀, 0.7 in the ♂. 51-59 flagellar segments. Body length 12-14 mm.......rugifer (Förster, 1868) ♀♂
- Hind tibia yellowish red to reddish brown medially, narrowly black basally, brown to black apically, the apical mark sometimes absent. ♂ with the front and mid coxae and the mid trochanter black, and with a rather broad area petiolaris (a rather variable character, the difference discernible only by direct comparison of specimens). Epipleural index about 0.7 in both sexes. 48-53 flagellar segments. Body length 10-13 mm......stragifex (Förster, 1868) ♀♂

- 31. Mesopleurum with wrinkles and without punctation centrally, densely rugose-punctate dorsally and ventrally. (In some small specimens the mesopleurum is rugose-punctate also centrally. These differ from *D. admontina* by having the posterior gastral tergites black.) Median longitudinal carinae of propodeum complete anteriorly. Petiolar suture complete. 1st gastral sternite coriaceous medially. Posterior gastral tergites black. 38-43 flagellar segments. Body length 7-8 mm.......

- Mesopleurum with distinctly separated punctures at least dorsally and ventrally. Posterior gastral tergites black. (In *D. castanipes* and *montana* the posterior gastral tergites are marked with red. These species differ from *D. admontina* by having the flagellum with 43-46 segments.)
- Petiole smooth or with fine sculpture laterally in front of the glymmae, at most with some transverse wrinkles anteriorly. 1st gastral sternite often smooth medially. Number of flagellar segments often lower......34.
- 34. 2nd flagellar segment 1.8-2.5 times as long as wide in the ♀, 2.2 times in the ♂. Glymma absent, at most a small and shallow furrow present. Hind tibia rather dark yellowish brown, black

- basally, not distinctly darkened apically. Propodeum not depressed or slightly depressed, in some parts with fine transverse wrinkles, in particular posteriorly. 43-45 flagellar segments. Body length 11-12 mm.
-montana (Roman, 1929) ♀♂ 2nd flagellar segment usually stouter. Glymma

- Genal index 0.4-0.6. Hind femur sometimes marked with red. Hind tibia usually marked with yellowish red to brown medially.......36.
- 36. Temple 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Palpi yellowish. Mandible broadly yellow medially. Hind femur black. Propodeum hardly impressed, for the greater part irregularly wrinkled, with a short median longitudinal keel und some longitudinal wrinkles posteriorly. 4th gastral tergite for the greater part black, with a narrow reddish longitudinal band dorsally (in the only known specimen). 50 flagellar segments. Body length 12 mm. ♀ unknown..... holmgrenii (Dalla Torre, 1901) ♂
- 37. Maxillary palpus blackish in both sexes. Mandible blackish in the ♀, sometimes marked with yellowish red medially in the ♂. Median longitudinal carinae of the propodeum complete or almost complete, joining the costulae anteriorly. Ovipositor index 1.0. Mesopleurum coriaceous and rather dull between the punctures. Hind tibia distinctly marked with black basally and apically. 2nd gastral tergite red on the posterior 0.4-0.5. 39-44 (rarely up to 49) flagellar segments. Body length 9-11 mm. (Small specimens of *D. nidulator*, in which the petiole is almost smooth in front of the glymmae, run here, but they differ from *D. terebrator* in other characters. See couplet 33.)..... *terebrator* (Förster, 1868) ♀♂

- Antennal carina not raised dorsally and/or its rim not distinctly modified. In some species the carina is slightly raised, but its rim is not widened to a dorsal crescent-shaped plate or to a nose-like projection (*D. alticola, auriculator, rubidatae, sobolicida*). In other species the carina is low, but its rim is widened to a smooth crescent-shaped plate (*D. angustifrons, temnator*) or to a small transversely striate projection (*D. fundator, stygia*).
- Genal carina joining the oral carina distant from the base of the mandible......40.
- 40. Genal carina distinctly raised ventrally, joining the distinctly raised oral carina distant from the base of the mandible, both forming a lopsided triangular pyramid, the tip of the latter shifted downwards and situated behind the base of the mandible. Frons distinctly depressed above the antennal sockets, with distinct radial wrinkles. Antennal carina distinctly widened dorsally to a nose-like projection. Ventral part of the prepectal carina merging with the pleural part, the latter joining the anterior edge of the mesopleurum. 50-54 flagellar segments. Body length 10-13 mm.......aurita (Kriechbaumer, 1883) ♀♂
- Genal carina not distinctly raised ventrally...... 41.
- 41. Metapleurum with distinct wrinkles centrally and dorsally. Pleural carina partly indistinct

and covered with wrinkles. Glymma absent. 2nd gastral tergite red on the posterior 0.1-0.2. Antennal carina widened to a nose-like projection dorsally. Spiracular carina indistinct and covered with wrinkles. 50-56 flagellar segments. Body length 14-16 mm.

......pineticola (Holmgren, 1872) ♀♂

- $44. \ \ Hind\ tibia\ completely\ light\ yellow.....45.$
- Hind tibia whitish yellow to black medially, darkened basally and apically......47.
- 45. 2nd gastral tergite broadly black anteriorly, yellowish red on the posterior 0.3. Ventral part of the prepectal carina and transverse carina both

- rather low. 4th tergite for the greater part or completely black. 35-38 flagellar segments. Body length 7-8 mm......
-pulchripes (Holmgren, 1872) ♀♂
- 2nd gastral tergite yellow, at most with a small dorsal black mark anteriorly......46.
- 46. Ventral part of the prepectal carina and transverse carina both strongly raised. Mid trochanter yellow. 4th tergite completely yellow or the posterior half black. 30-32 flagellar segments. Body length 7-8 mm. Probably only Canary Isles......tenerifae Hinz, 1977 ♀♂
- Ventral part of the prepectal carina and transverse carina both rather low. Mid trochanter black in the ♂. 4th gastral tergite black. Both flagellar tips broken off in the only known specimen. Body length 7.5 mm. ♀ of the variety unknown. Spain. tenerifae Hinz, 1977 var. ♂

- 39-50 flagellar segments. Genal carina joining the oral carina close to the base of the mandible, genal index 0.1-0.2 (sometimes difficult to ascertain).......51.
- 49. Hind corner of pronotum produced to a flange, which projects about as far as the tegula (best seen from above) and which is rounded dorsally and excavated ventrally (sometimes less produced, but 5th gastral tergite always marked with red). 5th gastral tergite red or marked with red. Hind tibia usually blackish, sometimes broadly yellowish brown medially. 32-35 flagellar segments in the \$\varphi\$, 34-37 segments in the \$\varphi\$. Body length 6-8 mm.
 - *minor* (Provancher, 1879) ♀♂

- Hind corner of pronotum not much produced, distinctly shorter than the tegula (moderately produced in some specimens of *D. angustifrons*).
 5th gastral tergite black.......50.
- 50. Antennal carina low, its rim more or less bent upwards and widened to a smooth narrow crescent-shaped plate. Flagellum filiform, shortest flagellar segment 0.9-1.0 times, penultimate segment 1.0-1.1 times as long as wide. (Variety: Flagellum slightly fusiform, shortest segment 0.8-0.9 times, penultimate segment 0.9-1.0 times as long as wide.) Ventral part of the prepectal carina low or slightly raised, usually merging with the pleural part (rarely with the transverse carina). Petiole with glymmae at the anterior edge of lateral areas. 2nd gastral tergite dorsally red on the posterior 0.1 (sometimes the red band more extended laterally). 4th gastral tergite usually for the greater part or completely red (sometimes marked with black dorsally). 33-37 flagellar segments in the 9, 36-38 segments in the ♂. Body length 7-9 mm. (Rarely ♀ with 31 flagellar segments; body length 5.5-6 mm.).....angustifrons (Förster, 1868) ♀♂

- Antennal carina slightly raised, somewhat thickened, finely transversely striate. Genal

brown. Gaster from the posterior 0.2-0.3 of the

2nd tergite onwards reddish brown. Posterior

carina slightly raised ventrally, joining the

slightly raised oral carina behind the base of the

	mandible. Hind femur black. 44-50 flagellar segments. Body length 8-10 mm. (See also couplet 68.)sobolicida (Förster, 1868) ♀♂		tergites tinged with black dorsally in the \mathcal{L} , broadly black dorsally or completely black in the \mathcal{L} . Glymma present, sometimes small. 35-39 flagellar segments. Body length 7-8 mm. (See
53.	24-32 flagellar segments		also couplet 11.)
-	33-47 flagellar segments57.		erythrogaster (Förster, 1868) ♀♂
54.	Maxillary palpus and mandible blackish. Pleural part of the prepectal carina complete or almost complete. Hind femur reddish brown or brown. Glymma large. Ovipositor index 1.3, ovipositor sheaths distinctly widened subapically (in lateral view). 26-27 flagellar segments in the $\[\varphi \]$, 29-30 segments in the $\[\varphi \]$. Body length 6 mm. (See also couplet 84.)	_	Pleural part of the prepectal carina usually present at least medially. Hind tibia whitish yellow to yellowish brown medially at least in the ♂ (exception: <i>D. hastulatae</i> , which differs from <i>D. erythrogaster</i> by: 40-42 flagellar segments; glymma absent; posterior gastral tergites black). Posterior gastral tergites black in both sexes
-	Maxillary palpus yellow or yellowish red. Mandible marked with yellow55.	58. –	40-47 flagellar segments
55.	2^{nd} gastral tergite red on the posterior 0.5-0.6. Hind femur black. Glymma distinct. Dorsal 0.3 of the pleural part of the prepectal carina obliterated. Ovipositor index 0.8. 29-30 flagellar segments in the $^{\circ}$, 30-31 segments in the $^{\circ}$. Body length 8-9 mm	59.	Hind tibia broadly light yellow medially, narrowly black basally and apically, the apical black ring extending over 0.1-0.2 of the tibia. 2^{nd} gastral tergite red on the posterior 0.4-0.6 in the 9 , on the posterior 0.3-0.4 in the 3 . 42-46 flagellar segments. Body length 8-12 mm
_	2 nd gastral tergite red on the posterior 0.1-0.3. Hind femur red or black. Glymma small or absent. (Specimens of <i>D. carpathica</i> with 30-32 flagellar segments run here; see couplet 65. They differ from <i>D. aemula</i> and <i>juvenilis</i> by: ventral part of the prepectal carina and transverse ca-	60.	Hind tibia black or tinged with white medially on the dorsal side or narrowly marked with yellow medially, the apical black ring extending over 0.3-0.4 of the tibia
	rina slightly raised; pleural part of the prepectal carina present medially and ventrally, the dorsal 0.4 obliterated.)56.		medially, obliterated dorsally and ventrally. Petiole smooth and rounded laterally, without glymmae or lateral areas. Hind tibia black,
56.	Pleural part of the prepectal carina distinct medially and dorsally, sometimes obliterated ventrally. 4 th gastral tergite often marked with		sometimes tinged with brown medially on the dorsal side. 40-42 flagellar segments,. Body length 8-9 mm hastulatae spec. nov. ♀♂
	red (rarely complete black). Ovipositor index 0.6. 26-30 flagellar segments in the ♀, 28-32 segments in the ♂. Body length 7-8 mm. (Rarely 24	-	Pleural part of the prepectal carina present medially and ventrally. Petiole with small glymmae or narrow lateral areas
	flagellar segments in the $\c ?$, 28 segments in the $\c ?$; body length 5 mm.)	61.	Temple 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the petiole. Genal carina slightly bent side-
_	Pleural part of the prepectal carina for the greater part obliterated, often some wrinkles present medially and/or subdorsally. 4^{th} gastral tergite usually black, sometimes marked with red laterally. Ovipositor index 1.3. 26-28 flagellar segments in the \Im , 28-31 segments in the \Im . Body length 7-9 mm.	_	wards and not bent forwards subventrally. Hind tibia black. Both flagellar tips broken off in the only known $^{\circ}$. 46 flagellar segments in a $^{\circ}$ tentatively placed here. Body length 9 mm
57.	juvenilis (Förster, 1868) ♀♂ Pleural part of the prepectal carina obliterated. Hind tibia black, sometimes tinged with dark		tangential lines on eyes and temples intersect on the scutellum or postscutellum. Genal carina slightly raised and distinctly bent sidewards and forwards subventrally in the typical morph,

slightly bent sidewards and hardly bent forwards in a variety with less flagellar segments. Hind tibia black in the \mathfrak{P} , sometimes tinged with white subbasally on the dorsal side. Hind tibia rather narrowly marked with whitish yellow in the \mathfrak{F} , distinctly black basally and apically, the apical ring extending over 0.3-0.4 of the tibia. 41-47 flagellar segments in the typical morph, 38-41 segments in a variety. Body length 7-9 mm. (See also couplet 64.)

......flagellator (Fabricius, 1793) ♀♂

- 62. Hind tibia broadly yellowish medially, narrowly marked with black basally and apically, the apical ring extending over 0.15-0.2 of the tibia. Propodeum usually with a distinct narrow longitudinal furrow and with short transverse striae in the furrow. 2nd gastral tergite marked with red on the posterior 0.2-0.3, the red mark often tinged with brown or black, the tergite sometimes almost completely black. 3rd gastral tergite red, often marked with black dorsally or posteriorly. 4th tergite black. Ovipositor stout basally, about 2.5 times as high as the apical width of a hind basitarsus. Ovipositor index 0.9. Glymma absent. 33-34 flagellar segments. Body length 9-11 mm......

- 64. Frons with almost smooth areas ventrally above the antennal sockets, without radial wrinkles or with some short radial wrinkles. 32-36 flagellar segments. Flagellum slightly fusiform, shortest segment 0.8-0.9 times as long as wide in the \mathcal{P} , 0.9-1.0 times in the \mathcal{E} . Hind tibia broadly vellowish red medially in both sexes, rather narrowly marked with black basally and apically, the apical ring extending over 0.2-0.3 of the tibia, rarely the hind tibia almost completely black and tinged with brown only medially. Propodeum rather coarsely wrinkled beside the longitudinal furrow, the median longitudinal carinae indicated. Petiole with small glymmae (rarely the glymmae obliterated). Body length 8-9 mm......disclusa (Förster, 1868) ♀♂
 - Frons with distinct and rather long radial wrinkles ventrally above the antennal sockets. 38-41 flagellar segments. Flagellum filiform, shortest segment 1.0-1.1 times as long as wide in the $\[Pi]$, 1.1-1.2 times in the $\[Pi]$. Hind tibia black in the $\[Pi]$, sometimes tinged with white subbasally on the dorsal side. Hind tibia for the greater part black in the $\[Pi]$, marked with whitish yellow medially (often only on the dorsal side), the apical black ring extending over 0.3-0.4 of the tibia. Propodeum with rather fine irregular wrinkles beside the longitudinal furrow, the longitudinal carinae usually obliterated. Petiole without glymmae, with narrow lateral areas. Body length 7.5-8.5 mm. (See also couplet 61.).....

.....flagellator (Fabricius, 1793) var. ♀♂

- Frons with almost smooth areas ventrally above the antennal sockets, with or without distinct radial wrinkles. 33-39 flagellar segments. Flagellum filiform, shortest segment 0.9 times as long as wide in the ♀, 1.0 times in the ♂. Ventral part of the prepectal carina and transverse carina strongly raised, the former distinctly emar-

- 66. Genal carina joining the oral carina absolutely at the base of the mandible. Mandible black in the ♀, narrowly marked with brown in the ♂. Mesopleurum coriaceous between the punctures. Longitudinal carinae of the propodeum obliterated. Sculpture of the propodeum rather variable (with fine or rather coarse irregular wrinkles, sometimes with a few transverse wrinkles, usually with some longitudinal wrinkles posteriorly). 33-39 flagellar segments. Body length 7-9 mm....... notabilis (Förster, 1868) ♀♂
- 67. Genal carina raised ventrally close to the junction with the also raised oral carina, forming a lopsided triangular pyramid, the tip of the latter shifted downwards and situated more or less behind the base of the mandible..............68.
- Genal carina not distinctly raised ventrally, its junction with the oral carina situated above the base of the mandible......79.
- 68. Genal carina slightly raised behind the lower edge of the eye, not raised and sometimes almost obliterated subventrally, slightly raised ventrally, joining the slightly raised oral carina behind the base of the mandible (an inconspicuous and rather variable character). Mesopleurum distinctly coriaceous (rarely almost smooth) centrally between the punctures. Hind tibia broadly yellow or yellowish brown medially, distinctly marked with dark brown or black basally and apically. 2nd gastral tergite red on the posterior 0.2-0.3. 3rd and 4th tergites red, the 4th tergite sometimes marked with black posteriorly. Antennal carina slightly raised, not bent upwards, the rim with fine transverse striae. 44-50 flagellar segments. Body length 8-10 mm. (See also couplet 52.)

..... sobolicida (Förster, 1868) ♀♂

- Ventral part of the prepectal carina merging with the transverse carina, both carinae often distinctly raised. (In some specimens of *D. splenditor* the ventral part of the prepectal carina merges with the pleural part, but in this species the flagellum is distinctly light brown distally.)......70.
- 70. 2nd gastral tergite black, only the posterior lateral corners with small yellow marks, sometimes the posterior edge with a very narrow yellow band.......71.
- 2nd gastral tergite dorsally yellow at least on the posterior 0.1-0.2, laterally yellow on the posterior 0.5-0.7.......73.
- Genal carina very strongly raised ventrally, genal index 0.7-0.8. Junction of the genal carina with the oral carina more or less pointed. Hind tibia yellow, at most narrowly marked with red or brown basally and apically. 4th gastral tergite broadly yellow laterally in both sexes.........72.

72. Pleural part of the prepectal carina complete. Mesopleurum with dispersed or rather dense punctation centrally. Speculum for the greater part smooth. Flagellum and hind tarsus black in both sexes. Mid leg yellow from the apex of the coxa onwards in both sexes. Hind tarsal claws short, with about four short pectinate teeth situated basally. 43-46 flagellar segments. Body length 8-9 mm.

......andalusica spec. nov. ♀♂

......lividariae Hinz, 1963 ♀♂

- Flagellum black distally. Hind tibia yellow, at most very indistinctly marked with light brown basally and apically......74.
- 74. Temple 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. 51-55 flagellar segments. 2nd gastral tergite usually with brown or black lateral lines slightly above the ventrolateral edges of the tergite. Body length 11-13 mm. Genal carina not raised medially (behind the centre of the eye). Mid leg yellow from the apex of the femur onwards in the ♀, from the apex of the coxa onwards in the ♂, tarsus marked with brown. 2nd gastral tergite dorsally yellow on the posterior 0.2-0.3.......

 Temple 0.7-0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the mesoscutum. 44-50 flagellar segments. 2nd gastral tergite with the ventrolateral

..... *mercator* (Fabricius, 1793) ♀♂

- edges partly marked with black. Body length 9-11 mm.......75.

- 76. Frons scarcely depressed, without a low bulge medially. Junction of the genal and the oral carinae more or less strongly shifted downwards (in some ♀♀ as strongly shifted downwards as in D. lamellator). Genal index 0.6-0.8. Mid leg yellow from the distal 0.4-0.8 of the femur onwards in the ♀ (rarely the femur completely yellow), from the apex of the coxa onwards in the ♂, apical tarsal segments brown. Hind tarsal claws of the ? with about five rather long pectinate teeth situated subapically as well as basally and medially (pectinate teeth of the hind claws of the δ short and inconspicuous). 2^{nd} gastral tergite dorsally yellow on the posterior 0.3. 46-50 flagellar segments. Body length 9-12 mm.....angustata (Thomson, 1887) ♀♂
- Frons slightly depressed, with a low bulge between these depressions. Junction of the genal and the oral carinae strongly shifted downwards, its tip almost reaching the ventral edge of the mandibles (in lateral view and with the mandibles closed). Genal index 0.8. Mid leg yellow from the base or the apex of the trochanter onwards in the \$\frac{1}{2}\$, from the apex of the coxa onwards in the \$\frac{1}{2}\$, apical tarsal segments brown. Hind tarsal claws in both sexes with 4-5 short pectinate teeth situated basally and medially. 2nd gastral tergite dorsally yellow on the posterior 0.3-0.4. 46-49 flagellar segments. Body length 11-13 mm.......

.....lamellator Aubert, 1960 ♀♂

......canariensis (Hinz, 1990) ♀♂

- Clypeus with well separated punctures on the ventral 0.3-0.5.....78.
- - semiflava (Costa, 1883) ♀♂
- At most 58 flagellar segments. Body length at most 14 mm......81.
- 80. Depression in front of the speculum for the greater part with longitudinal striae. Hind tibia yellow apically. 62-67 flagellar segments. Body length 17-20 mm.....
 - *falcator* (Fabricius, 1775) ♀♂
- Depression in front of the speculum granulatestrigose in the ventral half. Hind tibia marked with light red apically. 59-66 flagellar segments. Body length 14-17 mm.
- Ventral part of the prepectal carina merging with the pleural part, both of equal height at the junction, the transverse carina at most as

- 83. Genal index 0.1-0.2. Gaster black anteriorly and posteriorly, red medially.84.
- Genal index 0.2-0.3. Gaster reddish brown from the 2nd or 3rd tergite onwards, sometimes marked with brown or black dorsally. Ovipositor index 0.5-0.6.
 85.
- 84. 26-27 flagellar segments in the ♀, 29-30 segments in the ♂. Body length 6 mm. Maxillary palpus and mandible blackish. Glymma large. Ovipositor index 1.3. (See also couplet 54.)......
- Maxillary palpus and mandible marked with yellow. Glymma very small or absent. Ovipositor index 0.6.....120.
- 85. Antennal carina low and narrow. Speculum coriaceous and granulate-strigose, dull. Depression in front of the speculum with longitudinal and irregular wrinkles. Propodeum not depressed, for the greater part irregularly wrinkled. 47-49 flagellar segments. Body length 9-11 mm. Canary Isles.......
 -abdominator Hinz, 1985 ♀♂

- Propodeum more or less depressed, with distinct transverse wrinkles at least posteriorly......90.

- 44-58 flagellar segments. Body length 8-14 mm.
 Genal index 0.2-0.5. Hind tibia yellowish or yellowish brown medially, brown or black basally, brown apically, the apical mark often short or absent.
- 91. Genal index 0.2. Mesopleurum coriaceous and rather dull between the punctures. Propodeum distinctly depressed. 2nd gastral tergite black, with small reddish marks on the lower posterior corners. 44-45 flagellar segments. Body length 9-10 mm......einbecki Hinz, 1977 ♀♂
- Genal index 0.4-0.5. Mesopleurum finely coriaceous and somewhat shining between the punctures, sometimes smooth. Propodeum scarcely depressed. 2nd gastral tergite black dorsally, broadly red laterally.......92.
- 49-58 flagellar segments. Body length 10-14 mm. Speculum shining, usually smooth at least ventrally. Metapleurum usually with distinct punctures (punctate or rugose-punctate)................................signator (Brauns, 1895) ♀♂
- Antennal carina not widened dorsally to a noselike projection......94.
- 94. 2nd gastral tergite red on the posterior 0.1-0.2. Mesopleurum shining, slightly or not at all coriaceous between the punctures. Hind tibia black, rarely tinged with white or brown medially........95.

- 2nd gastral tergite red on the posterior 0.2-0.7. Mesopleurum often distinctly coriaceous and rather dull between the punctures. (Mesopleurum slightly coriaceous and shining in *D. aversa, incompleta* and *tenuis,* which are characterized by having the 2nd gastral tergite red on the posterior 0.3-0.7.) Hind tibia completely black or marked with whitish yellow or yellowish brown medially.......98.
- 95. Ovipositor index about 3.0. Hind claws strongly elongated in the δ , without pectinate teeth in the apical half (not strongly elongated and with pectinate teeth in the apical half in the \Im). Temples slightly constricted behind the eyes, tangential lines on eyes and temples intersect on the 1st gastral segment. Temple 0.9-1.0 times as long as the width of an eye. Pleural part of the prepectal carina present medially or medially and ventrally, obliterated dorsally. Mesopleurum with well separated punctures centrally, sometimes a few short wrinkles present. Pleural carina and spiracular carina present. 4th gastral tergite for the greater part or completely red. 40-46 flagellar segments. Body length 10-12 mm.....longiseta Hinz, 1961 ♀♂
- Ovipositor index at most 1.4. Hind tarsal claws not strongly elongated in either sex......96.

- 97. Temples slightly constricted behind the eyes, tangential lines on eyes and temples intersect on the propodeum or the 1st gastral segment. Temple 0.9-1.0 times as long as the width of an

- eye. Rim of the antennal carina low and narrow dorsally, almost not bent upwards, but distinctly bent inwards. Pleural part of the prepectal carina usually present ventrally and medially, sometimes present only medially or completely obliterated. Pleural carina present. Spiracular carina usually present, sometimes obliterated or covered with wrinkles. 4th gastral tergite for the greater part or completely red. 40-47 flagellar segments. Body length 9-11 mm.......bucculenta (Holmgren, 1860) ♀♂
- Temples distinctly constricted behind the eyes, tangential lines on eyes and temples intersect on the scutellum. Temple 0.7 times as long as the width of an eye. Antennal carina low, its rim moderately bent upwards and distinctly bent inwards, forming a smooth narrow crescent-shaped plate. Pleural part of the prepectal carina obliterated, the pleural part of the prepectus evenly rounded, not separated from the mesopleurum, without wrinkles. Pleural carina fine, usually partly covered with wrinkles. Spiracular carina indistinct and covered with wrinkles. 4th gastral tergite usually for the greater part or completely black, rarely red. 47-54 flagellar segments. Body length 9-10 mm.
 -temnator Hinz & Horstmann, 2004 ♀♂
- Ventral part of the prepectal carina slightly or not at all raised......102.
- -aversa (Förster, 1868) ♀♂
- 100. Genal carina joining the oral carina close to the base of the mandible, genal index 0.1. Genal carina distinct ventrally. Propodeum slightly depressed, with distinct and almost complete

- median longitudinal carinae and with irregular wrinkles in the shallow sulcus between the longitudinal carinae. 37 flagellar segments. Body length 9 mm. ♀ unknown. (See also couplet 66.)......tyrolensis spec. nov. ♂
- Genal carina joining the oral carina distant from the base of the mandible, genal index about 0.3. Genal carina more or less obliterated close to the junction with the somewhat raised oral carina, the gena somewhat depressed (a slightly variable character). Propodeum with a distinct narrow longitudinal furrow and with distinct transverse striae in the furrow. Longitudinal carinae of the propodeum obliterated.......101.
- 101. Front femur black, its distal 0.2 dark yellowish brown, front tibia dark yellowish brown, mid femur and mid tibia black in the ♀. Mandible for the greater part, tegula, front and mid coxae and hind tibia black in the ♂ (sometimes hind tibia tinged with yellowish brown medially). 41-46 flagellar segments. Body length 8-10 mm. ♀ tentatively placed here.......

.....incompleta (Bridgman, 1889) ♂?♀

- Front femur and front tibia yellowish red, mid femur yellowish red on its distal 0.5 and mid tibia yellowish red in the ♀. Mandible, tegula, front and mid coxae and hind tibia distinctly marked with yellow in the ♂. 44-48 flagellar segments. Body length 8-9 mm. (Some specimens are intermediate between *D. incompleta* and tenuis.) tenuis (Förster, 1868) ♀♂

- Striae in front of the speculum merging with

- 104. Genal index 0.2-0.3. Temple with fine and rather dense punctation medially. Hind tibia black in the $\mathfrak P$ (rarely tinged with white subbasally), dorsally marked with white medially or basally and medially in the $\mathfrak F$. Petiole without lateral areas or these areas weakly indicated. Mesopleurum with dense punctation. 43-46 flagellar segments. Body length 8-9 mm. (See also couplets 125 and 127.).....

.....meridionator Aubert, 1960 ♀♂

- Genal index 0.1-0.2. Temple with very fine and dispersed punctation medially. Hind tibia yellowish brown medially, narrowly black basally, broadly black apically. Petiole with lateral areas. 42-47 flagellar segments. Body length 8-9 mm.
- At least 34 flagellar segments. Body usually longer. Hind leg and gaster not as extensively marked with red.......106.

- 108. Antennal carina slightly raised, slightly thickened, its rim smooth. Hind femur red or reddish brown. Propodeum with a narrow longitudinal furrow and with short transverse striae in the furrow. 43-46 flagellar segments. Body length 8-9 mm. (See also couplet 85.).......
 -auriculator Aubert, 1964 ♀♂
- Antennal carina low and narrow. Hind femur black......109.
- 109. Mesopleurum slightly coriaceous or almost smooth and shining between the punctures. Propodeum slightly depressed, sometimes with a shallow and rather narrow longitudinal furrow, with distinct transverse striae at least posteriorly. (In aberrant specimens the transverse striae are almost completely obliterated.) 44-48 flagellar segments. Body length 9-11 mm. (See also couplet 92.)
 -circumcinctus (Förster, 1868) ♀♂
- Mesopleurum distinctly coriaceous and rather dull between the punctures. Propodeum not depressed or scarcely depressed, for the greater part irregularly wrinkled, at most with some short transverse wrinkles posteriorly, usually with a longitudinal keel posteriorly. 39-46 flagellar segments. Body length 7-9 mm. (See also couplet 89.)...... subimpressa (Förster, 1868) ♀♂
- 110. Genal index 0.1-0.2. Hind tibia broadly whitish or light yellow medially, narrowly marked with black basally and apically, the basal mark sometimes lacking, the apical ring 1.0-1.5 times as long as the width of the hind tibia. 2nd gastral tergite dorsally red on the posterior 0.4-0.7, laterally red on the posterior 0.7-1.0.............111.

- 112. Depression in front of the speculum granulatestrigose or with fine longitudinal striae, which merge with an area without punctation anteriorly. Mesopleurum distinctly coriaceous between the punctures. Hind tibia yellow or yellowish brown medially, marked with brown or black basally and apically.......113.
- Depression in front of the speculum with fine or strong longitudinal striae or wrinkles, which merge with the punctation anteriorly.......115.

- 114. 34-40 flagellar segments. Nervellar index 0.8-0.9. Propodeum rather broadly and not deeply depressed, the sulcus coriaceous anteriorly, with irregular wrinkles and a short longitudinal keel posteriorly. Gaster rather stout, 2nd tergite less than twice as long as wide. Body length 6-8 mm.......humilis (Förster, 1868) ♀♂
- 45-53 flagellar segments. Nervellar index usually 0.4-0.6. Propodeum rather broadly and deeply depressed, with transverse wrinkles in the sulcus. Gaster slender, 2nd tergite at least twice as long as wide. Body length 8-12 mm. (See also couplet 102.)
 -xenocampta (Förster, 1868) ♀♂
- 115. Hind tibia dark yellowish red, narrowly marked with black basally, hardly darkened apically. Propodeum rather broadly and not deeply depressed, with short transverse wrinkles in the midline and with long transverse wrinkles posteriorly. Spiracular carina indistinct or obliterated. 43-47 flagellar segments. Body length 9-10 mm.......opaca (Thomson, 1887) ♀♂

- Hind tibia completely black, or black and white (or light yellow), or yellow to yellowish brown medially and distinctly marked with brown or black basally and apically......116.
- 116. Hind tibia yellowish brown medially (also on the ventral side), marked with brown or black basally and apically......117.
- Hind tibia black, or black and tinged with dark brown medially, or black and marked with white on the dorsal side, or whitish basally and black apically......122.
- Temple 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the mesoscutum or the scutellum........118.
- Genal index 0.1-0.25. Petiole without glymmae, with narrow lateral areas......119.
- Mandible marked with yellow medially in the ♀, for the greater part yellow in the ♂. Hind femur red or black......120.
- 120. Antennal carina slightly raised, its rim slightly thickened, with fine transverse striae. 44-49 flagellar segments in the ♀, 45-51 segments in the ♂. Hind femur often black. Genal carina not bent forwards subventrally. Transverse carina present. Mesopleurum coriaceous, dull, with dispersed or rather dense punctation. Body length 7.5-10 mm. (Specimens of *D. sobolicida* in which the genal carina is concealed might run

- Antennal carina low and narrow. 38-44 flagellar segments. Hind femur often red.121.

......limnobia (Thomson, 1887) ♀♂

- Hind tibia black, sometimes tinged with dark brown medially. Propodeum often rather broadly and shallowly depressed (exception: *D. meridionator*). (*D. meridionator*, in which the hind tibia is black in the ♀ and marked with white in the ♂, can be traced through both halves of the couplet.)......126.
- Flagellum black (exception: some specimens of D. albobasalis). Metapleurum and propodeum not coarsely wrinkled. Hind tibia often with a

- Temple with dense or rather dense punctation medially. 40-46 flagellar segments. Glymma absent......125.

- 126. Temple 0.7-0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum (rarely on the propodeum).......127.
- Temple 0.9-1.0 times as long as the width of an eye, tangential lines on eyes and temples intersect on the propodeum or the 1st gastral segment.
- Temple with dense or rather dense punctation medially. Propodeum distinctly and rather nar-

- Temple with dispersed punctation medially.
 Propodeum rather broadly and shallowly depressed. Petiole with narrow lateral areas and/or with glymmae......128.
- 128. Genal index 0.4. Shortest flagellar segment 1.3 times as long as wide. Areola stalked, receiving the 2nd recurrent vein at the middle. Glymma small or absent. 2nd gastral tergite 2.5-3.0 times as long as wide. Maxillary palpus yellowish. 40-43 flagellar segments. Body length 8.5-9.5 mm.breviceps spec. nov. ♀♂
- Genal index 0.6. Shortest flagellar segment 0.9 times as long as wide. Areola pointed or narrowly sessile, receiving the 2nd recurrent vein distinctly distad of the middle. Glymma large. 2nd gastral tergite about 1.5 times as long as wide. Maxillary palpus blackish in the ♀, dark brown in the ♂. 42-44 flagellar segments. Body length 7-9 mm...........variator (Hinz, 1990) ♀♂
- Antennal carina low and narrow dorsally, its rim unmodified. Pleural carina distinct. Spiracular carina present.

- 131. Median longitudinal carina of the frons very fine or obliterated. 2nd gastral tergite red on the posterior 0.1-0.2. Nervellar index 0.7-0.8......95.

- 45-49 flagellar segments. Nervellar index 0.5-0.7.
 Longitudinal furrow of the propodeum with distinct transverse wrinkles medially and posteriorly. Body length 11 mm.

.....intelligator Aubert, 1966 ♀♂

Descriptions

Descriptions of 63 trans-palearctic species were already published by Hinz & Horstmann (2004). *D. annexa* was described under the name *D. americana*.

Dusona abdominator Hinz, 1985

9: Body length 9-11 mm. Temples strongly constricted behind the eyes, 0.65 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.9 of the mesoscutum. Ocellar index 0.6-0.7. Frons not depressed, coriaceous and with fine punctation, partly with wrinkles, with a fine median longitudinal carina which sometimes is partly obliterated, and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.3 of the clypeus coriaceous and densely rugosepunctate and rugose. Clypeus slightly concave, finely coriaceous and with rather dense or dense punctation ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low and straight ventrally, joining the low oral carina distant from the base of the mandible at an acute angle. Genal index 0.3. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 47-49 segments, slender, filiform, slightly attenuated distally, 2nd segment 2.1 times, shortest segment 1.0-1.1 times, penultimate segment 1.1-1.2 times as long as wide. Pronotum for the greater part rugose-punctate and with longitudinal wrinkles, finely coriaceous and densely punctate in the dorsal posterior corner. Mesoscutum coriaceous, dull, with dense or very dense punctation, partly rugose-punctate. Scutellum with lateral carinae on the anterior 0.4-0.5, finely coriaceous and with dispersed punctation. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, distinctly raised, narrowly emarginate medially. Transverse carina raised posteriorly, obliterated anteriorly and medi-

ally. Pleural part of the prepectal carina low, present ventrally and medially, the dorsal 0.3 obliterated. Mesopleurum finely coriaceous, slightly shining, with distinct and dense punctation. Speculum for the greater part coriaceous and granulate-strigose, dull, a small ventral part finely coriaceous and shining. Depression in front of the speculum coriaceous and with fine longitudinal and irregular wrinkles, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose. Pleural carina fine. Areolet large, stalked, receiving the 2nd recurrent vein at or near the middle. Nervellus slightly inclivous, weakly or distinctly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 5.8 times as long as high. Hind tarsal claws short, with a few short pectinate teeth situated basally. Propodeum extending to 0.3 of the hind coxae, not depressed, areolated, but the carinae partly covered with wrinkles. Area superomedia, area petiolaris and lateral areas with transverse or irregular wrinkles, the area superomedia sometimes coriaceous and without wrinkles anteriorly. Propodeal spiracle large, 2.6 times as long as wide. Spiracular carina narrow. Spiracular index 0.5. 1st gastral tergite 3.8 times, 2nd tergite 1.5 times as long as wide. Petiole smooth laterally, without glymmae or lateral areas. Petiolar suture fine, partly obliterated. 1st gastral sternite for the greater part smooth, with few transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter with black lateral stripes well above the ventrolateral edge (the stripes sometimes indistinct). Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown. Mandible tinged with reddish brown medially. Wings hyaline. Stigma dark brown medially. Front and mid legs reddish brown from the femur onwards, the tarsi tinged with dark brown. Hind femur and hind tibia dark reddish brown, the femur sometimes partly tinged with black. Gaster dark reddish brown from the 2nd tergite onwards, the 2nd tergite black on the anterior 0.2, the posterior tergites sometimes tinged with dark brown. Pilosity whitish.

δ: Pleural carina sometimes partly obliterated. Propodeum rounded, in most parts with irregular wrinkles, the carinae almost obliterated. Petiole sometimes with fine longitudinal striae laterally. 1st gastral sternite sometimes finely coriaceous. Front trochanter and apex of the front coxa marked with yellowish red. Genital clasper dark reddish brown. In other characters similar to the ♀.

Dusona albobasalis spec. nov.

Holotype (\$): "SQ-S-Chotin, 26.7.1993, leg. H. Halada" (near Komárno/Slovakia) (Mus. Linz). – Paratypes: 2\$\$, 2\$\$\footnote{\delta}\$ with the same dates (1\$, 1\$\footnote{\delta}\$ Mus. Linz, 1\$\footnote{\delta}\$, 1\$\footnote{\delta}\$ ZSM); 1\$\$\footnote{\delta}\$ "Moldavia, Kišinėv env., June 1995, I. Pavličok lgt." (Mus. Linz); 1\$\footnote{\delta}\$, "Khoper. zap., Varvarino, sosn. ostepn. polyany, Kasparyan, 2 VII 1977" (Voronezh province/ Russia) (Mus. St. Petersburg); 1\$\footnote{\delta}\$ same locality, 8.7.1977 (Mus. St. Petersburg); 1\$\footnote{\delta}\$ "Białystok P, 7.8.70, ..., leg. J. Sawoniewicz" (Poland) (Coll. Sawoniewicz).

9: Body length 7-8 mm. Temples slightly constricted behind the eyes, 0.75 times as long as the width of an eye, tangential lines on eyes and temples intersect on the propodeum. Ocellar index 1.1. Frons not depressed, coriaceous and with very dense punctation, with a fine median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, with dense punctation on a slightly coriaceous background ventrally, its apex slightly rounded medially. Malar space index 0.5. Genal carina low, slightly bent sidewards, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.15. Temple with fine and dense punctation on a very finely coriaceous background medially. Flagellum with 41-42 segments, slender, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.9 times, shortest segment 1.1 times, penultimate segment 1.3 times as long as wide. Pronotum with strong and dense punctaton on a smooth background laterally, with indistinct wrinkles ventrally. Mesoscutum coriaceous, dull, with dense or very dense punctation. Scutellum with lateral carinae on the anterior 0.5, with dense punctation. Prescutellar groove smooth medially, with longitudinal striae laterally. Prepectal carina low, ventral part merging with the pleural part, the latter complete or almost complete dorsally. Transverse carina obliterated. Mesopleurum with distinct dense punctation on a very finely coriaceous or smooth background. Speculum with fine striae dorsally, granulate-strigose and shining ventrally, punctate posteriorly. Depression in front of the speculum with distinct striae, which merge with the punctation anteriorly. Metapleurum with dense or very dense punctation on a very finely coriaceous background. Pleural carina complete. Areolet rather small, short-stalked, receiving the 2nd recurrent vein close to the middle. Nervellus distinctly reclivous, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.4. Hind femur 4.4 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally. Propodeum extending to 0.6 of the hind coxae, with a narrow and rather deep longitudinal furrow, for the greater part irregularly wrinkled, with distinct short transverse wrinkles in the furrow, carinae obliterated. Propodeal spiracle rather small, 3.2 times as long as wide. Spiracular carina slightly widened. Spiracular index 0.4. 1st gastral tergite 4.7 times, 2nd tergite 1.7 times as long as wide. Petiole without glymmae, with weakly indicated narrow lateral areas, smooth in front of these areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster slightly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish red. Mandible yellowish red dorsally and medially. Flagellum dark brown or light brown distally. Tegula yellow. Wings hyaline. Stigma light yellowish brown medially. Front leg from the trochanter onwards yellowish red, trochanter brownish ventrally, tarsus brownish distally. Mid femur yellowish red on the distal 0.4-0.6. Mid tibia for the greater part whitish yellow, marked with light red ventrally and apically. Mid tarsus yellowish proximally, dark brown distally. Hind tibia on the proximal 0.7 broadly whitish yellow dorsally, narrowly blackish ventrally, completely black on the distal 0.3. Hind tarsus black, the segments narrowly yellow basally. 2nd gastral tergite red on the posterior 0.1-0.2. 3rd and 4th tergites red. Rarely 2nd tergite completely black and 3rd and 4th tergites tinged with black dorsally. Pilosity whitish.

 δ : The δ distinctly differs from the \S in the shape of the temples, but agrees well in most other characters (except some sexual differences in colour pattern). Temples almost not constricted behind the eyes, 0.9-1.0 times as long as the width of an eye, tangential lines on eyes and temples intersect behind the apex of the gaster. Genal index 0.25. Flagellum with 40-45 segments. Pronotum with longitudinal wrinkles ventrally. Tibial spur index 0.5-0.6. Petiole without lateral areas. Petiolar suture partly indicated. Maxillary palpus yellow. Mandible for the greater part yellow. Flagellum dark brown distally. Front leg from the apex of the coxa onwards yellow. Mid leg from the apex of the coxa onwards yellow, trochanter and femur marked with brown ventrally, tarsus brown distally. Sometimes mid leg as dark as in the ♀ (coxa, trochanter, trochantellus and proximal 0.5 of femur black). Hind tibia whitish yellow basally and medially (also on the ventral side), black on the distal 0.3 (the black mark shorter dorsally than ventrally). Sometimes hind tibia brownish on the ventral

side basally and medially (but broadly white on the dorsal side). Hind tarsus dark brown. Sometimes the 4^{th} gastral tergite extensively marked with black or the 5^{th} tergite marked with red. In other characters similar to the 9.

Dusona alpigena Hinz, 1972

9: Body length 10-11 mm. Temples slightly constricted behind the eyes, 0.8-0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum or the propodeum. Ocellar index 1.0. Frons scarcely depressed, coriaceous and finely rugose-punctate, partly rugose, with a distinct median longitudinal carina and with short radial wrinkles, which extend to 0.25 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus coriaceous and densely rugose. Clypeus slightly concave, finely coriaceous and with rather dense punctation ventrally, its apex truncate medially. Malar space index 0.5. Genal carina low and rather straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.2-0.3. Temple finely coriaceous and with distinct fine dispersed punctation medially. Flagellum with 40-44 segments, slender, filiform, the distal 0.3 slightly attenuated, 2nd segment 2.0 times, shortest segment 1.0-1.1 times, penultimate segment 1.2 times as long as wide. Pronotum densely rugose-punctate laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum distinctly coriaceous, dull, with fine and dense punctation, partly with fine wrinkles. Scutellum with lateral carinae on the anterior 0.8, coriaceous and with distinct and dense punctation. Prescutellar groove almost smooth, with few fine longitudinal striae. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.2 of the latter obliterated or covered with wrinkles. Transverse carina present or obliterated. Mesopleurum finely coriaceous and with distinct rather dense or dense punctation, somewhat shining. Speculum finely granulate-strigose and with some fine longitudinal striae, somewhat shining. Depression in front of the speculum with distinct rather long striae, which merge with the punctation anteriorly. Metapleurum finely and densely rugose or rugose-punctate. Pleural carina distinct. Areolet rather large, stalked, receiving the 2nd recurrent vein distinctly distad of the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.6. Hind femur 5.0

times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, distinctly and rather narrowly depressed, for the greater part irregularly wrinkled, with short transverse wrinkles in the midline, with a short longitudinal keel posteriorly, longitudinal carinae obliterated or indistinct and covered with wrinkles. Propodeal spiracle large, 2.6 times as long as wide. Spiracular carina distinct and for the greater part narrow. Spiracular index 0.7. 1st gastral tergite 3.5 times, 2nd tergite 1.5 times as long as wide. Petiole with distinct lateral areas and small glymmae, the glymmae sometimes obliterated, smooth in front of the glymmae. Petiolar suture fine, obliterated anteriorly. 1st gastral sternite for the greater part with fine sculpture, shining, with fine transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster slightly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor

Colour: Black. Maxillary palpus dark brown basally, yellowish brown medially and apically. Mandible yellowish brown medially. Wings hyaline. Stigma dark brown medially. Front leg yellowish brown from the femur onwards, the femur basally marked with brown ventrally, the apical tarsal segment brown. Mid leg yellowish brown from the distal 0.4-0.6 of the femur onwards, the tarsus marked with brown. Hind tibia broadly yellowish brown medially, distinctly marked with black basally and apically, the apical ring extending over 0.1-0.2 (rarely 0.3) of the tibia. Hind tarsus for the greater part blackish, the proximal 0.3-0.5 of the basitarsus yellowish brown. 2nd gastral tergite red on the posterior 0.15-0.3. 3rd and 4th tergites red, the 4th tergite rarely darkened posteriorly. Pilosity whitish.

 σ : Flagellum with 42-46 segments, the shortest segment 0.9 times as long as wide. Propodeum with a narrower depression, the longitudinal carinae indicated or almost complete. Mandible yellow dorsally and medally. In other characters similar to the φ .

Variety: Some specimens are distinctly smaller: body length 8-9 mm; flagellum with 39-41 segments. In addition, the propodeum is less deeply depressed, and the red colour of the legs is somewhat more extended: mid femur for the greater part reddish brown, hind femur marked with reddish brown distally. The status of this morph is unknown.

Dusona alpina (Strobl, 1904)

9: Body length 11-12 mm. Temples strongly constricted behind the eyes, 0.65 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.8 of the mesoscutum. Ocellar index 0.9. From slightly depressed above the antennal sockets, finely coriaceous and with fine dispersed punctation, with a fine and partly obliterated median longitudinal carina and with indistinct short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.3 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat or sightly concave, distinctly coriaceous and with distinct rather dense punctation ventrally, its apex truncate medially. Malar space index 0.3. Genal carina low and almost straight ventrally, joining the scarcely raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.4. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 49-52 segments, filiform, the distal 0.2 attenuated, 2nd segment 2.1 times, shortest segment 0.9 times, penultimate segment 1.15 times as long as wide. Pronotum finely and densely rugose and rugose-punctate laterally on the dorsal 0.2, with longitudinal wrinkles or striae ventrally. Mesoscutum coriaceous, dull, with fine and dense punctation. Scutellum with lateral carinae on the anterior 0.8, coriaceous and densely rugose-punctate. Prescutellar groove smooth. Prepectal carina low, the ventral part merging with the pleural part, the latter complete, joining the anterior edge of the mesopleurum. Transverse carina obliterated. Mesopleurum coriaceous, dull, with distinct fine rather dense or dense punctation. Speculum coriaceous, rather dull, partly granulate-strigose. Depression in front of the speculum granulate-strigose, partly with fine striae, which merge with a narrow area without punctation anteriorly. Metapleurum finely coriaceous and with fine and dense punctation, somewhat shining. Pleural carina complete. Areolet rather large, stalked, receiving the 2nd recurrent vein basad of the middle. Nervellus vertical or slightly reclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 4.8 times as long as high. Hind tarsal claws short, with about four rather long pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, the longitudinal carinae fine, almost complete. Area superomedia almost not depressed, coriaceous, partly with fine irregular wrinkles. Area petiolaris distinctly and broadly depressed, for the greater part irregularly wrinkled, with some distinct transverse or longitudinal wrinkles and/or with a short longitudinal keel posteriorly. Propodeal spiracle long and narrow, 4.5 times as long as wide. Spiracular carina

narrow. Spiracular index 0.6. 1st gastral tergite 4.0 times, 2nd tergite 2.0 times as long as wide. Petiole with distinct small or rather large glymmae, smooth in front of the glymmae. Petiolar suture very fine or obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum indistinctly separated from the 3rd gastral tergite anteriorly, folded in or pendent, epipleural index 0.1-0.2. Black lateral stripe distinct, almost joining the epipleurum anteriorly, divergent from the ventrolateral edge posteriorly. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown or black. Mandible yellowish brown medially. Wings hyaline. Stigma dark yellowish brown medially. Front leg yellowish red from the trochantellus onwards, trochantellus and femur tinged with dark brown ventrally, sometimes also the trochanter marked with yellowish red dorsally. Distal 0.2 of the mid femur and mid tibia yellowish brown. Mid tarsus dark brown. Hind tibia dark yellowish brown or brown medially on the dorsal side, distinctly black basally and apically, the apical ring extending over 0.2-0.3 of the tibia. Hind tarsus blackish. 2nd gastral tergite black dorsally, laterally red on the posterior 0.2-0.3. The following tergites black dorsally, red laterally, rarely the 3rd and 4th tergites partly red also dorsally. Gastral sternites red. Pilosity whitish.

 δ : Propodeum less distinctly depressed, with distinct longitudinal wrinkles and/or with a distinct longitudinal keel posteriorly. Maxillary palpus yellow medially and apically. Mandible yellow medially. Front leg from the trochanter onwards yellow or yellowish red, only the femur tinged with brown ventrally. Mid femur marked with yellow distally, mid tibia and mid tarsus yellow, the apical tarsal segment brown. Hind tibia yellowish or yellowish brown medially. Gastral sternites brown. Genital clasper yellowish red. In other characters similar to the $\mathfrak P$.

Dusona andalusica spec. nov.

Holotype (\mathfrak{P}): "21/6", "3522.", "Andalusia. Staudinger." (southern Spain) (Mus. St. Petersburg). – Paratype (\mathfrak{d}): "25/4", "3521.", "Andalusia. Staudinger." (ZSM).

Q: Body length 8 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior half of the mesoscutum. Ocellar index 0.5. Frons hardly depressed above the antennal sockets, coriaceous and densely rugose-punctate, with a fine median longitudinal carina and with distinct

radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and very densely rugose-punctate and rugose. Clypeus for the greater part flat, slightly concave ventrally, the ventral 0.5 with strong and dense punctation on a slightly coriaceous background, its apex slightly rounded medially. Malar space index 0.25. Genal carina slightly raised medially (behind the eye), very strongly raised ventrally close to the junction with the very strongly raised oral carina, both forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the mandible. Gena deeply depressed close to the junction, the depression as deep as the width of the 2nd flagellar segment. Genal index 0.7. Temple slightly coriaceous and with fine dispersed punctation medially. Flagellum with 43 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.5 times, shortest segment 0.9 times, penultimate segment 1.0 times as long as wide. Pronotum slightly coriaceous and densely rugose-punctate laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with distinct and dense punctation, partly rugose-punctate. Scutellum with lateral carinae on the anterior 0.2, with distinct and very dense punctation. Prescutellar groove smooth medially, with fine longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae rather strongly raised, the former rather narrowly and not deeply emarginate medially. Pleural part of the prepectal carina distinct and complete, joining the anterior edge of the mesopleurum. Mesopleurum with dispersed and rather fine punctation on a smooth background. Speculum for the greater part smooth, with some punctures and wrinkles dorsally and posteriorly. Depression in front of the speculum with longitudinal striae dorsally and medially, which merge with the punctation anteriorly, smooth ventrally. Metapleurum distinctly rugose and with longitudinal wrinkles. Pleural carina complete. Areolet large, narrowly sessile, receiving the 2nd recurrent vein at the middle. Nervellus almost vertical, weakly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 5.5 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally. Propodeum extending to 0.6 of the hind coxae, with a distinct and rather narrow longitudinal furrow, with distinct transverse wrinkles in and beside the furrow, the lateral areas with transverse wrinkles too, without longitudinal carinae, but with the posterior transverse carina present laterally. Propodeal spiracle large, 2.4 times as long as wide. Spiracular carina slightly widened. Spiracular index 0.3. 1st gastral tergite 5.9 times, 2nd tergite 3.6 times as long as wide. Petiole smooth laterally, without glymmae or lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths slender subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma medium brown medially. Front and mid legs from the apex of the coxa onwards yellow. Mid tarsus brownish distally. Hind tibia yellow. 2nd gastral tergite dorsally with a very narrow yellow band on the posterior edge, with yellow marks on the posterior lateral corners. 3rd tergite yellow. 4th tergite yellow anteriorly, black posteriorly, the black mark extending to half of the tergite dorsally. Pilosity whitish.

 σ : Body length 9 mm. Flagellum with 46 segments. Mesopleurum with rather dense punctation on a smooth background centrally. Nervellar index 0.7. Spiracular index 0.6. In other characters similar to the φ (also in the colour of the legs and the gaster).

Dusona angustata (Thomson, 1887)

9: Body length 9-10 mm. Temples strongly constricted behind the eyes, 0.75 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.8 of the mesoscutum. Ocellar index 0.8. Frons scarcely depressed above the antennal sockets, densely rugose and rugosepunctate, with a few transverse wrinkles, with a distinct median longitudinal carina and with distinct radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina narrow, slightly raised (inconspicuous). Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat dorsally, slightly concave subventrally, coriaceous and with rather dense or dense punctation ventrally, its apex truncate or slightly rounded medially. Malar space index 0.2. Genal carina distinctly raised behind the ventral half of the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the base of the mandible. Gena deeply depressed close to the junction, the depression 0.8-1.0 times as deep as the width of the 2nd flagellar segment. Genal index 0.6. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 46-49 segments, slightly fusiform, 2nd segment 1.5 times, shortest segment 0.7-0.8 times, penultimate segment 1.0-1.1 times as long as wide. Pronotum densely rugose-punctate and rugose laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with very dense punctation. Scutellum with lateral carinae on the anterior 0.3-0.4, coriaceous and densely rugose-punctate. Prescutellar groove smooth. Ventral part of the prepectal carina distinctly raised, strongly raised submedially behind the front coxae, scarcely emarginate medially, merging with the slightly raised transverse carina. Pleural part of the prepectal carina low, complete or the dorsal 0.2 covered with wrinkles. Mesopleurum with distinct rather dense or dense punctation on a smooth background, with longitudinal wrinkles posteriorly in front of the metapleurum. Speculum shining, finely granulate-strigose, with some fine longitudinal striae, a ventral part smooth. Depression in front of the speculum with distinct rather long striae, which merge with the punctation anteriorly. Metapleurum with distinct longitudinal wrinkles dorsally, rugose and rugosepunctate ventrally. Pleural carina distinct. Areolet large, pointed, receiving the 2nd recurrent vein at or slightly basad of the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.5 times as long as high. Hind tarsal claws short, with about five rather long pectinate teeth situated subapically as well as basally and medially. Propodeum extending to 0.6 of the hind coxae, broadly and deeply depressed, with strong transverse wrinkles in the sulcus and in the lateral areas, longitudinal carinae and costulae partly present. Propodeal spiracle large, 3.5 times as long as wide. Spiracular carina distinct, broad. Spiracular index 0.2. 1st gastral tergite 6.2 times, 2nd tergite 2.8 times as long as wide. Petiole with distinct lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 2.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths slightly widened medially and subapically (in lateral view). Ovipositor index 0.7. Colour: Black. Maxillary palpus yellow or

Colour: Black. Maxillary palpus yellow or yellowish brown. Mandible yellow dorsally and medially. Tegula yellow, with a brown mark. Wings hyaline. Stigma medium brown medially. Front femur yellow on the distal 0.7-0.8. Front tibia and front tarsus yellow. Mid femur yellow on the distal 0.2 (rarely completely yellow). Mid tibia yellow. Mid tarsus for the greater part dark brown. Hind tibia yellow. Hind tarsus black, the basitarsus yellow basally. 2nd gastral tergite dorsally yellow on the posterior 0.3, laterally yellow on the posterior 0.7,

the ventrolateral edges partly marked with black. 3rd and 4th tergites yellow. Pilosity whitish.

δ: Flagellum filiform, the distal 0.2 attenuated, the shortest segment 0.9 times as long as wide. Hind tarsal claws with about four rather short pectinate teeth situated basally and medially. Maxillary palpus yellow. Mandible broadly yellow medially. Front and mid legs from the apex of the coxa onwards yellow, apical tarsal segments brown. In other characters similar to the \mathfrak{P} .

Dusona aversa (Förster, 1868)

9: Body length 10-12 mm. Temples distinctly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 1.1-1.5. Frons not depressed, coriaceous and densely rugose and rugose-punctate, with a distinct median longitudinal carina, without radial wrinkles. Antennal carina low and narrow. Face and clypeus coriaceous and densely rugose and rugose-punctate, only a narrow apical band of the clypeus less densely punctate. Clypeus slightly rounded basally, flat or slightly concave subapically, its apex slightly rounded medially. Malar space index 0.35. Genal carina low and straight subventrally, joining the low oral carina distant from the base of the mandible at an acute angle. Genal index 0.4. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 38-40 segments, filiform, the distal 0,2 attenuated, 2nd segment 2.0 times, shortest segment 0.9-1.0 times, penultimate segment 1.3 times as long as wide. Pronotum densely punctate on an almost smooth background laterally on the dorsal 0.3, densely rugose and rugose-punctate ventrally, with some longitudinal wrinkles. Mesoscutum coriaceous, with fine and very dense punctation. Scutellum with lateral carinae on the anterior 0.7, with distinct punctation on an almost smooth background, partly rugose-punctate. Prescutellar groove smooth. Ventral part of the prepectal carina strongly raised, narrowly and not deeply emarginate medially, merging with the distinctly raised transverse carina. Pleural part of the prepectal carina low, present medially, often obliterated ventrally and dorsally, sometimes almost complete. Mesopleurum distinctly and rather densely or densely punctate on a finely coriaceous and shining background. Speculum for the greater part with fine longitudinal striae, a small ventral part almost smooth. Depression in front of the speculum with rather long striae, which merge with the punctation anteriorly. Metapleurum or densely rugose-punctate on an almost smooth background. Pleural carina distinct. Areolet rather narrow, narrowly sessile, pointed or short-stalked, receiving the 2nd recurrent vein about at the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.4-0.5. Hind femur 4.3 times as long as high. Hind tarsal claws short, with about four rather short pectinate teeth situated basally and medially. Propodeum extending to 0.5 of the hind coxae, broadly and rather deeply depressed, for the greater part with fine irregular wrinkles, with short transverse wrinkles in the midline and long transverse wrinkles posteriorly, carinae almost completely obliterated. Propodeal spiracle rather large, 3.3 times as long as wide. Spiracular carina narrow. Spiracular index 0.5. 1st gastral tergite 4.3 times, 2nd tergite 2.1 times as long as wide. Petiole without glymmae, narrow lateral areas indicated, smooth in front of these areas. Petiolar suture fine, partly obliterated. 1st gastral sternite smooth, rarely with some fine wrinkles. Thyridium oval. Thyridial index 2.0. Epipleurum not separated from the 3rd gastral tergite, the latter with fine or distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus dark brown. Mandible sometimes tinged with brown medially. Wings hyaline. Stigma medium brown medially. Front femur dorsally and apically and front tibia reddish brown or dark brown. Front basitarsus yellowish basally. Mid and hind tibiae rarely tinged with dark brown medially on the dorsal side. 2nd gastral tergite red on the posterior 0.4-0.6. 3rd and 4th tergites red, sometimes with dark dorsal marks. Pilosity whitish.

 δ : Ocellar index 1.0-1.4. Flagellum with 41-44 segments. Tibial spur index 0.5-0.6. Rarely front femur apically and front and mid tibiae light reddish brown. In other characters similar to the \mathfrak{P} .

Dusona baueri Hinz, 1973

♀: Body length 6 mm. Temples slightly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the 2nd gastral segment. Ocellar index 1.0. Frons

not depressed, coriaceous and finely rugose and rugose-punctate, with a fine median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow. Face coriaceous and with fine dense punctation, partly with fine wrinkles. Clypeus flat basally and medially, bent inwards subapically, for the greater part coriaceous and with fine rather dense punctation, smooth subapically, its apex broadly and shallowly emarginate. Malar space index 0.4. Genal carina low and almost straight subventrally, joining the low oral carina distant from the base of the mandible at an acute angle. Genal index 0.25. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 21 segments, slightly clavate, 2nd segment 2.0 times, penultimate segment (which is the shortest segment) 0.9 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely and finely rugose-punctate. Scutellum with lateral carinae on the anterior 0.4, coriaceous and with fine rather dense punctation. Prescutellar groove smooth. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.3 of the latter obliterated. Transverse carina present, low. Mesopleurum coriaceous, rather dull, with fine rather dense or dense punctation. Speculum finely granulate-strigose, slightly shining. Depression in front of the speculum with fine rather long striae, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose-punctate and rugose. Pleural carina present. Areolet rather large, narrowly sessile, receiving the 2nd recurrent vein distinctly distad of the middle. Nervellus inclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.6 times as long as high. Hind tarsal claws somewhat elongated, with about two small pectinate teeth situated basally. Propodeum extending to 0.5 of the hind coxae, broadly and shallowly depressed, coriaceous and for the greater part with irregular wrinkles, with fine and short transverse wrinkles in the midline, with some longitudinal wrinkles posteriorly, carinae almost completely obliterated. Propodeal spiracle small, oval, 2.0 times as long as wide. Spiracular carina narrow. Spiracular index 0.8. 1st gastral tergite 4.3 times, 2nd tergite 1.2 times as long as wide. Petiole with very small glymmae, smooth in front of these glymmae. Petiolar suture fine, for the greater part obliterated. 1st gastral sternite for the greater part smooth, with some fine transverse striae posteriorly. Thyridium small, oval. Thyridial index 2.0. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster slightly compressed posteriorly. Ovipositor sheaths rather narrow subapically (in lateral view). Ovipositor index 0.85.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma medium brown medially. Front and mid legs light red from the apex of the coxa onwards. Hind leg red from the trochanter onwards. Gaster dark reddish brown from the 2nd tergite onwards, anterior 0.2 of the 2nd tergite black, posterior tergites and sternites partly tinged with black. Pilosity whitish.

δ: Apex of the clypeus truncate. Flagellum with 30 segments, almost filiform, the distal 0.2 slightly attenuated, 2nd segment 2.0 times, shortest segment 1.1 times, penultimate segment 1.2 times as long as wide. Propodeum without short transverse wrinkles in the sulcus. Petiole with small lateral areas laterally (instead of glymmae). Posterior gastral tergites and sternites black. Genital clasper dark brown. In other characters similar to the ♀.

Dusona breviceps spec. nov.

Holotype ($\$): "San Martin de Luiña – Asturias, Trampa Malaise, 17-23/VIII-92, C. Rey leg." (Spain) (Mus. Madrid). – Paratypes: $2\$ $\$ 11 $\$ $\$ $\$ from the same locality, dates 17.-29.8.1992 (Mus. Madrid, $1\$ $\$ $\$ 2 $\$ $\$ ZSM); $1\$ Llueva, Voto, Cantabria (Spain), 3.-10.8.2001 (Mus. Madrid); $4\$ $\$ $\$ same locality, dates 5.-30.8.2000 (Mus. Madrid).

9: Body length 8.5-9 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, coriaceous and densely rugose-punctate, with a narrow median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, coriaceous and with dispersed or rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina low and almost straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.4. Temple finely coriaceous and with very fine and very dispersed punctation medially. Flagellum with 40-43 segments, filiform, slender, the distal 0.2 attenuated, 2nd segment 2.1 times, shortest segment 1.3 times, penultimate segment 1.5 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.2, with distinct longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with very dense punctation. Scutellum with lateral carinae on the anterior 0.6, coriaceous and with rather dense punctation. Prescutellar groove almost smooth medially, with some fine longitudinal striae laterally. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.2-0.3 of the latter obliterated. Transverse carina obliterated. Mesopleurum coriaceous and with distinct dense punctation, slightly shining. Speculum granulatestrigose and with fine striae. Depression in front of the speculum with distinct striae, which merge with the punctation anteriorly. Metapleurum coriaceous and finely and densely rugose. Pleural carina fine, complete. Areolet rather small, stalked, receiving the 2nd recurrent vein at the middle. Nervellus slightly reclivous, rather weakly intercepted. Nervellar index 0.6-0.8. Tibial spur index 0.6. Hind femur 5.6 times as long as high. Hind tarsal claws short, with about three distinct pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, with a rather broad and shallow sulcus, for the greater part irregularly wrinkled, with short transverse wrinkles in the midline medially and posteriorly (rarely without transverse wrinkles), carinae for the greater part obliterated, median longitudinal carinae partly indicated. Propodeal spiracle rather small, 2.6 times as long as wide. Spiracular carina narrow. Spiracular index 0.5. 1st gastral tergite 4.7 times, 2nd tergite 2.8 times as long as wide. Petiole with distinct narrow lateral areas, sometimes with small glymmae at the anterior edge of these areas, smooth in front of the lateral areas. Petiolar suture partly present. 1st gastral sternite for the greater part smooth, with some fine transverse striae posteriorly. Thyridium elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish brown. Mandible yellowish brown dorsally and medially. Tegula yellowish brown or dark brown. Wings hyaline. Stigma brownish black. Front leg yellowish brown from the trochanter onwards, tibia yellow dorsally. Mid femur reddish brown, tinged with black basally. Mid tibia yellow dorsally, reddish brown ventrally. Mid tarsus yellowish proximally, brown distally. Hind leg black, tibia tinged with dark brown medially, basitarsus narrowly yellowish basally. 2nd gastral tergite dark red on the posterior 0.3. 3rd and 4th tergites red. 5th tergite red, dorsally marked with black posteriorly (to a variable extent). Pilosity whitish.

&: Mandible broadly yellowish medially. Tegula yellow or yellowish brown. Front and mid trochanters

and trochantelli yellow. In other characters similar to the \mathfrak{L} .

Dusona calceata (Brauns, 1895)

9: Body length 9-10 mm. Temples strongly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. Ocellar index 1.1. Frons slightly depressed above the antennal sockets, coriaceous and distinctly and densely rugose, partly with transverse wrinkles, with a distinct median longitudinal carina and with indistinct radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus densely rugose-punctate and rugose. Clypeus flat, distinctly and densely punctate on a smooth background ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina low medially behind the eye, strongly raised subventrally and ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter rounded, shifted downwards and situated behind the base of the mandible. Gena deeply depressed close to the junction, the depression 0.6 times as deep as the width of the 2nd flagellar segment. Genal index 0.4. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 42-43 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.5 times, shortest segment 0.65 times, penultimate segment 1.1 times as long as wide. Pronotum densely rugose-punctate on a smooth background laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum densely rugose-punctate on a smooth background. Scutellum with lateral carinae on the anterior 0.3, densely punctate on a smooth background. Prescutellar groove smooth medially, with some longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae distinctly raised, the former scarcely emarginate medially. Pleural part of the prepectal carina present medially and ventrally or only medially, or completely obliterated and replaced by some wrinkles. Mesopleurum distinctly and densely punctate on a smooth background. Speculum with distinct longitudinal striae dorsally and medially, almost smooth ventrally. Depression in front of the speculum with distinct striae, which merge with the punctation anteriorly. Metapleurum densely rugose-punctate and rugose on a smooth background. Pleural carina distinct. Areolet large,

short-stalked, receiving the 2nd recurrent vein at or slightly distad of the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.8. Tibial spur index 0.7. Hind femur 4.2 times as long as high. Hind tarsal claws slightly elongated, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.4 of the hind coxae, broadly and not deeply depressed, the sulcus for the greater part with irregular wrinkles, with short transverse wrinkles in the midline (these sometimes obliterated) and long transverse wrinkles posteriorly, the lateral areas with strong transverse wrinkles, longitudinal carinae obliterated. Propodeal spiracle large, 2.4 times as long as wide. Spiracular carina narrow, sometimes indistinct. Spiracular index 0.4. 1st gastral tergite 3.9 times, 2nd tergite 1.7 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 0.7. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus blackish basally, yellowish brown apically. Mandible narrowly tinged with brown medially. Wings hyaline. Stigma yellowish brown medially. Front femur black basally and ventrally, yellow dorsally and apically. Front tibia and front tarsus yellow. Mid femur narrowly yellow apically. Mid tibia yellow. Mid tarsus blackish. Hind tibia yellow basally and medially, sometimes with a small blackish mark basally, distinctly black apically, the apical ring narrow dorsally, extending over 0.1-0.2 of the tibia, rather broad ventrally, extending over 0.3-0.4 of the tibia. 2nd gastral tergite for the greater part black, the posterior edge with a narrow yellow band dorsally and the posterior lateral corners with small yellow marks. 3rd tergite yellow. 4th tergite narrowly yellow anteriorly. Pilosity whitish.

 $\ensuremath{\mathcal{G}}$: Flagellum slightly fusiform, shortest segment 0.85 times, penultimate segment 1.0 times as long as wide. Longitudinal carinae of the propodeum indicated posteriorly. Mandible narrowly yellow medially. Front trochanter and front trochantellus marked with yellow. Front femur for the greater part yellow, marked with black ventrally. Mid tarsus light brown. Hind tibia for the greater part yellow, the distal 0.1-0.2 marked with light brown. 4^{th} gastral tergite sometimes broadly yellow anteriorly. In other characters similar to the $\ensuremath{\mathcal{Q}}$.

Dusona canariensis (Hinz, 1990)

9: Body length 10-12 mm. Temples strongly constricted behind the eyes, 0.6-0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.6-0.7 of the mesoscutum. Ocellar index 0.5. Frons scarcely depressed above the antennal sockets, coriaceous and densely rugosepunctate, with a distinct median longitudinal carina and with radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.2-0.3 of the clypeus densely rugose and rugose-punctate. Clypeus for the greater part flat, slightly concave ventrally, densely punctate (with well separated punctures) on a smooth background medially and ventrally, its apex truncate medially. Malar space index 0.3. Genal carina low medially behind the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the base of the mandible. Gena deeply depressed close to the junction, the depression 0.7 times as deep as the width of the 2nd flagellar segment. Genal index 0.5. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 43-48 segments, filiform, the distal 0.3 attenuated, 2nd segment 2.0 times, shortest segment 0.8 times, penultimate segment 1.0 times as long as wide. Pronotum densely rugose laterally on the dorsal 0.1, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous, partly with a smooth background, densely punctate or rugose-punctate. Scutellum with lateral carinae on the anterior 0.4, finely coriaceous and very densely punctate. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former almost not emarginate medially. Pleural part of the prepectal carina distinct medially, indistinct ventrally, covered with wrinkles on the dorsal 0.2. Mesopleurum distinctly and densely punctate on a smooth background. Speculum with fine longitudinal striae, sometimes granulate-strigose or almost smooth ventrally, shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum strongly and densely punctate or rugose-punctate on a smooth background ventrally, with longitudinal striae dorsally. Pleural carina distinct. Areolet large, narrowly sessile, receiving the 2nd recurrent vein basad of the middle. Nervellus vertical, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.5-6.0 times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally and medially. Propodeum extending to 0.6 of the hind coxae, rather narrowly depressed, almost completely covered with strong transverse wrinkles, the longitudinal carinae lacking. Propodeal spiracle rather large, 2.7 times as long as wide. Spiracular carina narrow, sometimes widened. Spiracular index 0.4. 1st gastral tergite 6.2 times, 2nd tergite 2.4 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 2.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths slightly widened medially and subapically (in lateral view). Ovipositor index 0.8.

Colour: Black. Maxillary palpus yellow. Mandible yellow, teeth brown. Tegula yellow. Wings hyaline. Stigma blackish medially. Front and mid legs yellow from the apex of the coxa onwards, mid coxa sometimes completely black, mid tarsus brown distally. Hind tibia yellow. Hind basitarsus narrowly marked with yellow basally. 2nd gastral tergite dorsally yellow on the posterior 0.3-0.4, laterally yellow on the posterior 0.7, the ventrolateral edges partly marked with black. 3rd and 4th tergites yellow. Pilosity whitish.

 δ : Penultimate segment of the flagellum 1.1-1.2 times as long as wide. Front and mid coxae for the greater part yellow. 5th gastral tergite narrowly yellow anteriorly. In other characters similar to the \mathfrak{P} .

Dusona castanipes (Thomson, 1887)

9: Body length 11-12 mm. Temples distinctly constricted behind the eyes, 0.8-0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the propodeum. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, coriaceous and finely rugose-punctate, with a distinct median longitudinal carina and with short radial wrinkles (which may be indistinct). Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous, with distinct and dense punctation, partly with wrinkles. Clypeus flat, coriaceous and with rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.3. Genal carina low and straight subventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.4. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 44-48 segments, filiform, the distal 0.2 attenuated, 2nd segment 1,3-1.5 times, shortest segment 0.7 times, penultimate segment 1.0 times as long as wide. Pronotum densely rugose-punctate and rugose on a smooth background laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with fine dense or very dense punctation. Scutellum with lateral carinae on the anterior 0.4, finely coriaceous and with rather dense punctation. Prescutellar groove smooth. Prepectal carina low, ventral part merging with the pleural part, the latter complete or the dorsal 0.15 obliterated. Transverse carina indicated anteriorly, obliterated posteriorly. Mesopleurum rather densely or densely punctate on a finely coriaceous or smooth background, a central part often rugose-punctate. Speculum finely granulate-strigose anteriorly, sometimes smooth, with fine punctation posteriorly, shining. Depression in front of the speculum with rather fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum densely rugose-punctate and rugose on a finely coriaceous or smooth background. Pleural carina distinct. Areolet large, short-stalked, receiving the 2nd recurrent vein slightly basad of the middle. Nervellus vertical, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 4.3 times as long as high. Hind tarsal claws slightly elongated, with about five distinct pectinate teeth situated basally, medially and subapically. Propodeum extending to 0.3 of the hind coxae, with more or less complete carinae. Area superomedia flat, coriaceous or with fine irregular wrinkles, with some short transverse wrinkles posteriorly. Area petiolaris broadly and rather shallowly depressed, in most parts with transverse wrinkles, in addition with a short longitudinal keel posteriorly. Propodeal spiracle rather large, 3.1 times as long as wide. Spiracular carina obliterated or covered with wrinkles. Spiracular index 0.6-0.8. 1st gastral tergite 3.3 times, 2nd tergite 2.0 times as long as wide. Petiole with large glymmae, smooth in front of the glymmae. Petiolar suture obliterated or partly present. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.0. Epipleurum separated from the 3rd gastral tergite and folded in, the crease marked with black. Epipleural index 0.7. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus dark brown or black. Mandible yellow dorsally and medially. Wings hyaline. Stigma dark brown medially. Front and mid legs reddish brown from the femur onwards, the femora darkened basally, the mid tarsus light brown. Hind femur black proximally, dark reddish brown distally (to a variable extend, sometimes completely reddish brown). Hind tibia dark reddish brown, indistinctly tinged with black basally and apically. Hind tarsus dark brown or black, the basitarsus

narrowly marked with yellowish brown basally. 2^{nd} gastral tergite red on the posterior 0.3-0.5. Median and posterior tergites red, the posterior tergites often narrowly marked with black dorsally, sometimes the 3^{rd} and 4^{th} tergites black dorsally and/or the 6^{th} and 7^{th} tergites almost completely black and only a narrow ventrolateral band red. Gastral sternites red. Pilosity whitish.

ở: Shortest flagellar segment 0.8-1.0 times as long as wide. Maxillary palpus yellow. Mandible broadly yellow medially. Front leg yellowish red from the trochanter onwards. Mid leg light reddish brown from the femur onwards. 2nd gastral tergite usually red on the posterior 0.3, rarely black dorsally and red laterally. Median and posterior tergites black dorsally and red laterally (to a variable extend), sometimes the 3rd tergite completely red or the posterior tergites completely black. Gastral sternites red or black. Genital clasper red or brown or black. In other characters similar to the ♀.

Dusona circumspectans (Förster, 1868)

9: Body length 7-8 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.7 of the mesoscutum. Ocellar index 1.0. Frons not depressed, coriaceous and densely rugose-punctate, dull, with a fine median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.3 of the clypeus coriaceous and densely rugosepunctate and rugose. Clypeus flat, coriaceous and with fine rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4-0.5. Genal carina low, straight or slightly bent sidewards subventrally, joining the low oral carina distant of the base of the mandible at an acute angle. Genal index 0.2-0.3. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 37-38 segments, slender, filiform, the distal 0.2 attenuated, 2nd segment 2.1 times, shortest segment 1.1 times, penultimate segment 1.3 times as long as wide. Pronotum densely rugose-punctate and rugose laterally on the dorsal 0.5, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, finely and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.7, coriaceous and densely rugose-punctate. Prescutellar groove with fine longitudinal striae. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.2 of the latter covered with wrinkles. Transverse carina obliterated or present only anteriorly. Mesopleurum coriaceous, dull, with rather dense or dense punctation. Speculum finely granulate-strigose or with fine longitudinal striae, shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose-punctate and rugose. Pleural carina distinct. Areolet rather large, shortstalked, receiving the 2nd recurrent vein distad of the middle. Nervellus vertical or slightly inclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.65. Hind femur 5.6 times as long as high. Hind tarsal claws short, with about three short pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, carinae almost completely obliterated. Region of the area superomedia rather flat, coriaceous and with fine irregular wrinkles. Region of the area petiolaris with a narrow longitudinal furrow, for the greater part irregularly wrinkled, with short transverse wrinkles in the furrow, with long transverse wrinkles posteriorly. Propodeal spiracle rather large, 4.0 times as long as wide. Spiracular carina narrow medially. Spiracular index 0.5. 1st gastral tergite 3.7 times, 2nd tergite 2.0 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the areas. Petiolar suture indicated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 2.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown basally, yellowish brown medially and apically. Mandible sometimes tinged with brown medially. Tegula yellowish brown. Wings hyaline. Stigma medium brown medially. Front leg yellowish red from the trochantellus or the femur onwards. Mid femur yellowish red on the distal 0.4. Mid tibia and mid tarsus yellowish red, the tarsus darkened apically. Hind tibia broadly yellowish brown medially, narrowly brown basally, rather broadly dark brown or black apically, the apical ring extending over 0.3 of the tibia. Hind tarsus dark brown, the basitarsus yellowish brown on the basal 0.5-0.7. $2^{\rm nd}$ gastral tergite red on the posterior 0.3. $3^{\rm rd}$ and $4^{\rm th}$ tergites red. Pilosity whitish.

δ: Flagellum stouter, 2nd segment 1.6-1.8 times, shortest segment 1.0 times, penultimate segment 1.2 times as long as wide. Propodeum distinctly and rather broadly depressed, with distinct transverse wrinkles in the sulcus and with almost complete longitudinal carinae. Hind tibia yellowish brown medially, distinctly black basally and apically. Hind tarsus black, the basitarsus yellowish on the basal 0.2. In other characters similar to the ♀.

Dusona confusa (Förster, 1868)

9: Body length 11-14 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the centre of the mesoscutum. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, coriaceous and with dense punctation, with a fine median longitudinal carina and with distinct radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus coriaceous and densely rugose. Clypeus flat, coriaceous and with rather fine dispersed or rather dense punctation ventrally, its apex truncate medially. Malar space index 0.3. Genal carina low, straight or slightly bent sidewards subventrally, joining the low oral carina close to the base of the mandible at an acute angle. Genal index 0.1-0.15. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 47-54 segments, filiform, the distal 0.4 attenuated, 2nd segment 2.0 times, shortest segment 0.9-1.0 times, penultimate segment 1.2 times as long as wide. Pronotum coriaceous and finely rugose laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with fine and dense punctation, partly rugose-punctate. Scutellum with lateral carinae on the anterior 0.7, coriaceous and with distinct dense punctation. Prescutellar groove smooth medially, with fine longitudinal striae laterally. Prepectal carina low, ventral part merging with the pleural part, the latter complete or the dorsal 0.2 covered with wrinkles. Transverse carina distinct anteriorly, fine or obliterated posteriorly. Mesopleurum coriaceous, dull, with distinct dense punctation. Speculum coriaceous, dull, partly granulate-strigose. Depression in front of the speculum with distinct short longitudinal striae, which usually merge with a narrow area without punctation anteriorly (rarely this area obliterated, and the striae merge with the punctation). Metapleurum coriaceous, densely rugose-punctate and rugose. Pleural carina distinct. Areolet rather small, pointed or short-stalked, receiving the 2nd recurrent vein at or distad of the middle. Nervellus vertical or slightly reclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 5.9 times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, rather broadly and deeply depressed, the sulcus with distinct transverse wrinkles, the longitudinal carinae present posteriorly, often also medially. Propodeal spiracle rather large, 2.9 times as long as wide. Spiracular carina narrow, distinct. Spiracular index 0.4. 1st gastral tergite 5.2 times, 2nd tergite 2.2 times as long as wide. Petiole with distinct narrow lateral areas and small glymmae, smooth in front of the glymmae. Petiolar suture present or partly obliterated. 1st gastral sternite usually completely smooth, sometimes with a few transverse striae posteriorly. Thyridium oval. Thyridial index 2.0. Epipleurum of the 3rd gastral tergite present as a small semicircular lobe anteriorly, which usually is pendent, but rarely is folded in. Epipleural index 0.1. 3rd tergite usually with fine black lateral stripes well above the ventrolateral edge, the stripes sometimes obliterated. Gaster strongly compressed posteriorly. Ovipositor sheaths slightly widened medially and subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus dark brown. Mandible black, marked with yellow dorsally. Tegula dark brown. Wings hyaline. Stigma light yellowish brown medially. Front leg yellowish red from the femur onwards, the femur marked with dark brown basally, the tarsus marked with dark brown apically. Mid femur yellowish red on the distal 0.2. Mid tibia yellowish red. Hind femur sometimes marked with red distally. Hind tibia for the greater part light yellow, narrowly reddish or brown basally, narrowly black apically, the apical ring extending over 0.1 of the tibia. Postpetiole sometimes marked with red laterally. 2nd gastral tergite dorsally yellowish red on the posterior 0.5-0.6, laterally yellowish red on the posterior 0.7-0.8. 3rd and 4th tergites yellowish red. 5th tergite often yellowish red anteriorly (rarely median gastral tergites yellow instead of yellowish red). Pilosity whitish.

δ: Propodeum very deeply depressed, with strong transverse wrinkles and with the longitudinal carinae clearly present medially and posteriorly. Maxillary palpus yellowish brown or brown. Mandible broadly yellow medially. Tegula often marked with yellowish brown. Front coxa yellow apically. Front trochanter and trochantellus yellow. Mid femur often for the greater part yellowish red. Mid tarsus yellowish brown basally. Hind tibia whitish yellow medially, narrowly black basally and apically. In other characters similar to the ♀.

Dusona disclusa (Förster, 1868)

2: Body length 8-9 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum or postscutellum. Ocellar index 0.8. Frons scarcely depressed above the antennal sockets, for the greater part coriaceous and densely rugosepunctate and rugose, small ventral areas above the antennal sockets almost smooth, with a distinct me-

dian longitudinal carina, without radial wrinkles or with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus densely rugose-punctate. Clypeus flat or slightly concave, with dense punctation on a smooth background ventrally, a subventral area often without punctation or with fine dispersed punctation, the apex truncate medially. Malar space index 0.4. Genal carina low, distinctly bent sidewards and forwards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 32-36 segments, slightly fusiform, 2nd segment 1.7 times, shortest segment 0.8-0.9 times, penultimate segment 1.0 times as long as wide. Pronotum with distinct and very dense punctation on a smooth background laterally on the dorsal 0.5, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, rather finely and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.6, densely rugose-punctate. Prescutellar groove smooth or with very fine sculpture. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former very strongly raised submedially (behind the front coxae), distinctly and broadly emarginate medially. Pleural part of the prepectal carina low, present ventrally and medially or only medially, the dorsal 0.4 obliterated. Mesopleurum coriaceous, slightly shining, with distinct and dense punctation. Speculum finely granulate-strigose, shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum densely rugosepunctate and rugose. Pleural carina distinct. Areolet rather large, short-stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus vertical, weakly or distinctly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.6. Hind femur 5.0 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.5 of the hind coxae, distinctly and rather narrowly depressed, with distinct transverse wrinkles medially in the longitudinal furrow and rather coarse irregular wrinkles beside the furrow, with long transverse wrinkles posteriorly, the longitudinal carinae usually present medially and posteriorly. Propodeal spiracle rather large, 3.0 times as long as wide. Spiracular carina narrow medially. Spiracular index 0.4. 1st gastral tergite 5.1 times, 2nd tergite 2.2 times as long as wide. Petiole with distinct narrow lateral areas and with small glymmae (rarely the glymmae obliterated), smooth in front of the glymmae. Petiolar suture for the greater part or completely obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd

gastral tergite, the latter with fine or distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus brown or dark brown. Mandible tinged with brown medially. Tegula yellowish brown. Wings hyaline. Stigma yellowish brown medially. Front and mid legs from the femur onwards yellowish red, the tarsi marked with brown distally, the front trochanter and trochantellus sometimes marked with yellowish red. Hind femur often completely black, sometimes reddish brown distally or for the greater part reddish brown. Hind tibia usually broadly yellowish brown medially, narrowly blackish basally, broadly blackish apically, the apical ring extending over 0.2-0.3 of the tibia, rarely almost completely black and tinged with brown medially. Hind basitarsus narrowly marked with yellowish brown basally. 2nd gastral tergite red on the posterior 0.3-0.4 (rarely 0.6). 3rd tergite red. 4th tergite red anteriorly and black posteriorly (to a variable extent), often for the greater part black and only with red lateral marks anteriorly. Pilosity whitish.

ở: Flagellum filiform, the distal 0.3 attenuated, 2nd segment 1.6 times, shortest segment 0.9-1.0 times, penultimate segment 1.0 times as long as wide. Propodeum strongly depressed, with strong transverse wrinkles medially and posteriorly, often also anteriorly, usually with distinct longitudinal carinae. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Front and mid coxae yellow apically. Front and mid trochanters and trochantelli yellow, the mid trochantellus marked with brown. Hind basitarsus broadly yellowish basally. In other characters similar to the ♀.

Dusona dubitor Hinz, 1977

9: Body length 9-11 mm. Temples strongly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.6-0.8 of the mesoscutum. Ocellar index 0.6. Frons scarcely depressed above the antennal sockets, coriaceous and densely rugose-punctate and rugose, with a distinct median longitudinal carina and with radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.4-0.6 of the clypeus densely rugose-punctate and rugose. Clypeus flat with rather dense punctation on a very finely coriaceous or smooth background ventrally, its apex truncate medially. Malar space

index 0.3. Genal carina low medially behind the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the base of the mandible. Gena deeply depressed close to the junction, the depression 0.7 times as deep as the width of the 2nd flagellar segment. Genal index 0.5. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 44-49 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.8 times, shortest segment 0.9 times, penultimate segment 1.1 times as long as wide. Pronotum densely rugose-punctate on a smooth background laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous, rather finely and densely punctate or rugose-punctate. Scutellum with lateral carinae on the anterior 0.3, densely rugose-punctate. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former almost not emarginate medially. Pleural part of the prepectal carina distinct medially, indistinct ventrally, covered with wrinkles on the dorsal 0.3 (rarely almost complete). Mesopleurum distinctly and rather densely or densely punctate on a smooth background. Speculum coriaceous or granulate-strigose dorsally, smooth medially and ventrally, shining. Depression in front of the speculum with strong longitudinal striae, which merge with the punctation anteriorly. Metapleurum strongly and densely punctate or rugose-punctate on a smooth background ventrally, with longitudinal striae dorsally. Pleural carina distinct. Areolet rather large, pointed or short-stalked, receiving the 2nd recurrent vein about at the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 6.0-6.5 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.6 of the hind coxae, rather broadly depressed, almost completely covered with strong transverse wrinkles, the longitudinal carinae lacking, the transverse carinae partly indicated. Propodeal spiracle rather large, 3.6 times as long as wide. Spiracular carina narrow, sometimes widened. Spiracular index 0.5. 1st gastral tergite 5.9 times, 2^{nd} tergite 2.0 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture fine, partly obliterated. 1st gastral sternite smooth, sometimes with fine transverse striae posteriorly. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths slightly

widened medially and subapically (in lateral view). Ovipositor index 0.8.

Colour: Black. Maxillary palpus yellowish or yellowish brown. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma yellowish brown medially. Front leg yellow from the apex of the coxa onwards, sometimes the trochanter and the femur marked with black ventrally. Mid femur yellow on the distal 0.3-0.4. Mid tibia yellow. Mid tarsus yellow basally, yellowish brown apically. Hind tibia yellow. Hind basitarsus narrowly marked with yellow basally. 2nd gastral tergite dorsally yellow on the posterior 0.7-0.9, the ventrolateral edges partly marked with black. 3rd and 4th tergites yellow. Pilosity whitish.

 δ : Maxillary palpus yellow. Front and mid legs for the greater part yellow, only the base of the coxae black. In other characters similar to the \circ .

Dusona einbecki Hinz, 1977

9: Body length 9-10 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the centre of the mesosutum. Ocellar index 0.8. Frons not depressed, coriaceous and finely and densely rugose-punctate, with a distinct median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow dorsally, distinctly bent inwards, the antennal carinae almost touching medially. Face and dorsal 0.6 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, coriaceous and with dense punctation ventrally, its apex truncate or slightly emarginate medially. Malar space index 0.4. Genal carina low, slightly bent sidewards subventrally, joining the low oral carina close to the base of the mandible at an acute angle. Genal index 0.2. Temple with very fine sculpture and with fine very dispersed punctation medially. Flagellum with 44-45 segments, filiform, the distal 0.2 attenuated, 2nd segment 1.9 times, shortest segment 1.0 times, penultimate segment 1.1 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.5, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, finely and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.4, coriaceous and densely rugose-punctate. Prescutellar groove smooth or with fine longitudinal striae. Prepectal carina low, the ventral part merging with the slightly raised transverse carina. Pleural part of the prepectal carina distinct medially, almost obliterated ventrally, the dorsal 0.3 obliterated or covered

with wrinkles. Mesopleurum coriaceous, dull, with distinct rather dense or dense punctation. Speculum granulate-strigose and with fine striae, dull. Depression in front of the speculum with fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous, dull, finely and densely rugose-punctate. Pleural carina complete. Areolet rather small, stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus distinctly inclivous, weakly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 5.5 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally and medially. Propodeum extending to 0.5 of the hind coxae, broadly and rather deeply depressed, the sulcus coriaceous and finely rugose anteriorly (region of the area superomedia), with short transverse wrinkles medially and long transverse wrinkles posteriorly, irregularly wrinkled sublaterally, longitudinal carinae present posteriorly, covered with wrinkles medially. Propodeal spiracle rather large, 2.6 times as long as wide. Spiracular carina narrow medially. Spiracular index 0.4. 1st gastral tergite 5.9 times, 2nd tergite 2.2 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture fine or obliterated. 1st gastral sternite smooth, sometimes with fine transverse striae posteriorly. Thyridium elongated. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths scarcely widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish brown. Mandible black, sometimes tinged with dark brown medially. Tegula black, tinged with brown apically. Wings hyaline. Stigma dark brown medially. Front femur yellowish red, marked with dark brown basally and ventrally. Front tibia and tarsus yellow. Mid femur yellow on the distal 0.2. Mid tibia yellow. Mid tarsus yellowish brown. Hind tibia yellowish brown, distinctly marked with black basally, almost not darkened apically. Hind tarsus yellowish brown or dark brown. 2nd gastral tergite for the greater part black, the posterior lateral corners red. 3rd tergite narrowly black dorsally, broadly red laterally, 4th tergite for the greater part red, narrowly marked with black dorsally and posteriorly. Pilosity whitish.

 δ : Longitudinal carinae of the propodeum present anteriorly and medially as well as posteriorly. Mandible marked with yellow dorsally. Apex of front coxa yellow. Front trochanter and trochantellus marked with yellowish red. In other characters similar to the \circ .

Dusona erythrogaster (Förster, 1868)

9: Body length 7-8 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. Ocellar index 0.7. Frons slightly depressed above the antennal sockets, finely coriaceous and finely and densely punctate, rather dull dorsally, shining ventrally, with a low longitudinal bulge medially and a longitudinal carina on top of this bulge, and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugosepunctate. Clypeus flat dorsally, slightly concave ventrally, finely coriaceous and densely punctate ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low, distinctly bent sidewards and forwards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 35-37 segments, filiform, the distal 0.2 attenuated, 2nd segment 2.0 times, shortest segment 0.9 times, penultimate segment 1.2 times as long as wide. Pronotum densely rugosepunctate laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely punctate. Scutellum with lateral carinae on the anterior 0.7, coriaceous and densely punctate. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae moderately raised, the former distinctly and broadly emarginate medially. Pleural part of the prepectal carina obliterated, a few wrinkles present medially, the lateral part of the prepectus densely punctate. Mesopleurum coriaceous, slightly shining, densely punctate. Speculum coriaceous, slightly shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous, finely and densely rugose-punctate. Pleural carina distinct. Areolet rather small, short-stalked, receiving the 2^{nd} recurrent vein at or distad of the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 4.5 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally. Propodeum extending to 0.3 of the hind coxae, deeply and rather narrowly depressed, the sulcus with distinct transverse wrinkles, longitudinal carinae obliterated. Propodeal spiracle rather large, 2.9 times as long as wide. Spiracular carina narrow. Spiracular index 0.5. 1st gastral tergite 4.2 times, 2nd tergite 1.9 times as long as wide. Petiole with distinct small or rather large glymmae, smooth in front of the glymmae. Petiolar suture for the greater part obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.0. Epipleurum separated from the 3rd gastral tergite and folded in, sometimes folded back and pendent as an indistinct narrow stripe, epipleural index 0.2, the tergite with black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths distinctly widened medially and subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown. Mandible narrowly marked with brown dorsally. Wings hyaline. Stigma medium brown medially. Front femur reddish brown on the distal 0.7. Front tibia and tarsus reddish brown. Mid femur reddish brown on the distal 0.3. Mid tibia and tarsus reddish brown or dark brown. Hind legs black, the femur rarely marked with reddish brown distally, the tibia sometimes tinged with dark brown medially. 2nd gastral tergite dark redish brown on the posterior 0.3. The following tergites dark reddish brown, the posterior tergites tinged with black dorsally. Pilosity whitish.

 σ : Flagellum with 38-39 segments. Posterior gastral tergites marked with black dorsally or completely black. Genital clasper black. In other characters similar to the φ .

Dusona fatigator (Förster, 1868)

9 (only the holotype known): Body length 9 mm. Temples slightly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the petiole. Ocellar index 1.0. Frons not depressed, coriaceous and densely rugose-punctate and rugose, with a fine median longitudinal carina and with distinct radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, finely coriaceous and with dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina slightly raised subventrally, slightly bent sidewards and not bent forwards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple finely coriaceous and with fine dispersed or rather dense punctation medially. Both flagellar tips broken off in the only known ♀, 2nd flagellar segment 2.0 times as long as wide. Pronotum laterally for the greater part with longitudinal wrinkles, only the dorsal posterior corners densely rugose-punctate and without longitudinal wrinkles. Mesoscutum coriaceous and with rather fine very dense punctation. Scutellum with lateral carinae on the anterior 0.7, coriaceous and with rather dense punctation. Prescutellar groove smooth medially, with longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former distinctly and broadly emarginate medially. Pleural part of the prepectal carina low ventrally and medially, somewhat indistinct, the dorsal 0.3 obliterated. Mesopleurum fine coriaceous and with rather dense punctation, slightly shining. Speculum finely granulate-strigose anteriorly, punctate posteriorly, slightly shining. Depression in front of the speculum with wrinkles dorsally, with rather fine longitudinal striae medially and ventrally, the latter merging with the punctation anteriorly. Metapleurum finely coriaceous and densely punctate and rugose-punctate. Pleural carina distinct. Areolet large, narrowly sessile, receiving the 2nd recurrent vein distad of the middle. Nervellus reclivous, rather weakly intercepted. Nervellar index 0.6. Tibial spur index 0.6. Hind femur 4.5 times as long as high. Hind tarsal claws short, with about five distinct pectinate teeth situated basally and medially. Propodeum extending to 0.4 of the hind coxae, distinctly and rather narrowly depressed, with distinct transverse wrinkles in the longitudinal furrow, irregularly wrinkled beside the furrow, with some transverse wrinkles on the lateral parts of the propodeum, carinae almost completely obliterated. Propodeal spiracle large, 2.3 times as long as wide. Spiracular carina broad. Spiracular index 0.4. 1st gastral tergite 3.8 times, 2nd tergite 1.6 times as long as wide. Petiole with indistinct narrow lateral areas and with very small glymmae, smooth in front of the glymmae. Petiolar suture partly present. 1st gastral sternite for the greater part smooth, with some transverse striae posteriorly. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown basally, yellowish brown medially and apically. Mandible broadly yellowish red dorsally and medially. Tegula dark brown. Wings hyaline. Stigma medium brown medially. Front leg yellowish red from the femur onwards, the femur marked with dark brown basally and ventrally, the tarsus darkened distally. Mid leg blackish, the tibia and tarsus tinged with brown. 2nd gastral tergite red on the posterior 0.4. 3rd and 4th tergites red. Pilosity whitish.

?ð: Clypeus for the greater part densely rugosepunctate and rugose, only a small subapical area densely punctate. Flagellum with 46 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1,5 times, shortest segment 1.0 times, penultimate segment 1.0 times as long as wide. Metapleurum densely rugose-punctate and rugose. Nervellus vertical. Nervellar index 0.8. $3^{\rm rd}$ gastral tergite with black lateral stripes. Maxillary palpus brown basally, yellow medially and apically. Mandible broadly yellow medially. Mid femur yellowish red on the distal 0.2. Mid tibia yellowish red. In other characters similar to the $\mathfrak P$. This $\mathfrak F$ from Escorial/Spain (ZSM) may belong to D. fatigator, but it differs from the holotype (which originates from southern France) in the proportion of the basal flagellar segments and in the nervellus (disregarding the differences in colour pattern, which may be sexual differences).

Dusona filator Hinz, 1985

9: Body length 8-9 mm. Temples distinctly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, finely coriaceous and finely rugose and punctate, with a fine median longitudinal carina and with few short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus densely rugose. Clypeus flat, finely coriaceous and with rather dense punctation ventrally, its apex truncate medially. Malar space index 0.3. Genal carina slightly raised subventrally, distinctly bent sidewards and forwards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple finely coriaceous and with fine very dispersed punctation medially. Flagellum with 29-30 segments, stout, slightly fusiform, almost not attenuated distally, 2nd segment 1.9 times, shortest segment 0.7 times, penultimate segment 0.8 times as long as wide. Pronotum densely rugose-punctate laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, densely punctate. Scutellum with lateral carinae on the anterior 0.4, coriaceous and with dense punctation. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former distinctly and rather broadly emarginate medially. Pleural part of the prepectal carina present ventrally and medially, low, the dorsal 0.3 obliterated. Mesopleurum finely coriaceous, densely punctate, slightly shining. Speculum narrow, finely coriaceous anteriorly, punctate posteriorly, slightly shining. Depression in front of the speculum with distinct long longitudinal striae, which merge with the punctation anteriorly. Metapleurum finely coriaceous and finely punctate and rugose, slightly shining. Pleural carina distinct. Areolet rather narrow, pointed, receiving the 2nd recurrent vein distinctly distad of the middle. Nervellus sightly inclivous, distinctly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 4.4 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, deeply and rather narrowly depressed, with long transverse wrinkles in and beside the longitudinal furrow, median longitudinal carinae obliterated. Propodeal spiracle rather small, 2.3 times as long as wide. Spiracular carina distinct, narrow. Spiracular index 0.8. 1st gastral tergite 3.3 times, 2nd tergite 1.5 times as long as wide. Petiole with rather large glymmae, smooth in front of the glymmae. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.8.

Colour: Black. Maxillary palpus yellowish red. Mandible yellowish brown dorsally. Tegula yellowish brown. Wings hyaline. Stigma yellowish brown medially. Front leg yellowish red from the apex of the coxa onwards. Mid leg yellowish red from the femur onwards, the femur tinged with black ventrally, the tarsus darkened apically. Hind tibia dark yellowish brown medially, narrowly marked with black basally, broadly marked with black apically, the apical ring extending over 0.3 of the tibia. Hind basitarsus marked with yellowish brown basally. 2nd gastral tergite red on the posterior 0.5. 3rd tergite red. 4th tergite red, marked with black dorsally and posteriorly. Pilosity whitish.

δ: Flagellum with 30-31 segments, rather slender, filiform, the distal 0.2 attenuated, 2nd segment 1.9 times, shortest segment 1.0 times, penultimate segment 1.3 times as long as wide. Pleural part of the prepectal carina sometimes obliterated ventrally. Glymma sometimes small. 2nd gastral tergite red on the posterior 0.6. In other characters similar to the ♀.

Dusona genalis (Thomson, 1887)

9: Body length 9-11 mm. Temples moderately constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the propodeum or the 1st gastral segment. Ocellar index 1.1-1.4. Frons slightly depressed above the antennal sockets, for the greater part coriaceous and densely rugose-punctate, narrow areas above the antennal sockets shining, with a distinct median

longitudinal carina and with fine or distinct radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.6 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, coriaceous and with rather dense punctation ventrally, its apex smooth, rounded medially. Malar space index 0.4. Genal carina low and straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.3-0.4. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 49-53 segments, filiform, the distal 0.3 distinctly attenuated, 2nd segment 1.6 times, shortest segment 0.9 times, penultimate segment 1.3 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.4, punctate and with longitudinal wrinkles ventrally. Mesoscutum coriaceous and very densely punctate. Scutellum with lateral carinae on the anterior 0.4, finely coriaceous and densely punctate. Prescutellar groove with fine longitudinal striae. Prepectal carina low, ventral part merging with the pleural part or with the transverse carina. Transverse carina complete. Pleural part of the prepectal carina complete or the dorsal 0.1 covered with wrinkles. Mesopleurum finely coriaceous and with distinct rather dense or dense punctation, slightly shining. Speculum finely coriaceous and granulate-strigose, shining. Depression in front of the speculum dorsally with wrinkles, medially and ventrally with fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum finely coriaceous and densely punctate (with well separated punctures), slightly shining. Pleural carina distinct. Areolet rather large, narrow, pointed or narrowly sessile, receiving the 2nd recurrent vein at the middle. Nervellus vertical or slightly inclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.5-0.6. Hind femur 4.4 times as long as high. Hind tarsal claws slightly elongated, with about five distinct pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, rather narrowly and not deeply depressed, with rather short transverse wrinkles in the longitudinal furrow, with irregular wrinkles beside the furrow, with long transvserse wrinkles posteriorly, the longitudinal carinae almost completely obliterated. Propodeal spiracle large, 3.1 times as long as wide. Spiracular carina narrow. Spiracular index 0.4. 1st gastral tergite 3.3 times, 2nd tergite 1.4 times as long as wide. Petiole with distinct rather large glymmae, smooth in front of the glymmae. Petiolar suture fine or almost obliterated. 1st gastral sternite smooth, sometimes with some fine transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum separated from the 3rd gastral tergite anteriorly, folded in, the crease somewhat rounded and sometimes not marked with black. Epipleural index 0.4-0.6. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus dark brown. Mandible marked with redish brown dorsally and medially. Wings hyaline. Stigma yellowish brown medially. Front leg yellowish red from the femur onwards, the femur tinged with brown ventrally. Mid femur yellowish red on the distal 0.2. Mid tibia yellowish red. Mid tarsus brown. Hind tibia yellowish brown medially, distinctly marked with black basally and apically, the apical ring extending over 0.2-0.3 of the tibia. Hind tarsus blackish, the basitarsus narrowly marked with brown basally. 2nd gastral tergite dorsally red on the posterior 0.3, laterally red on the posterior 0.7. 3rd tergite red, sometimes tinged with black dorsally. 4th tergite for the greater part dark red, marked with black dorsally and posteriorly (to a variable extent). Pilosity whitish.

 δ : Propodeum with the longitudinal carinae partly present. Sometimes the 4^{th} gastral tergite completely black. In other characters similar to the \mathfrak{P} .

Dusona habermehli (Kriechbaumer, 1898)

9: Body length 5.5-7 mm. Temples distinctly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 1.1. Frons not depressed, coriaceous and finely rugose, with some transverse wrinkles, the median longitudinal carina short or obliterated, without radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.3 of the clypeus coriaceous and densely rugose, dull. Clypeus flat, coriaceous and with fine rather dense punctation ventrally, its apex rounded medially. Malar space index 0.3. Genal carina low, slightly bent sidewards subventrally, joining the low oral carina close to the base of the mandible at an acute angle. Genal index 0.1. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 32-35 segments, slender, filiform, the distal 0.2 slightly attenuated, 2nd segment 2.5 times, shortest segment 1.2 times, penultimate segment 1.2 times as long as wide. Pronotum coriaceous and finely punctate and rugose laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with fine and dense punctation, partly rugose. Scutellum with lateral carinae on the anterior 0.3, coriaceous, dull, rather densely punctate. Prescutellar groove smooth or with a few longitudinal striae. Prepectal carina low, ventral part merging

with the low transverse carina. Pleural part of the prepectal carina distinct medially, fine or obliterated ventrally, the dorsal 0.3 obliterated. Mesopleurum coriaceous, dull, with fine dispersed punctation. Speculum granulate-strigose and with fine striae, rather dull. Depression in front of the speculum with fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous, dull, finely rugose-punctate and rugose. Pleural carina complete. Areolet rather small, short-stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus almost vertical, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.4. Hind femur 4.8 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, shallowly and rather narrowly depressed, coriaceous and without transverse wrinkles anteriorly in the region of the area superomedia, with short transverse wrinkles medially in the longitudinal furrow, irregularly wrinkled beside the furrow, with long transverse wrinkles posteriorly, the longitudinal carinae indicated only posteriorly. Propodeal spiracle rather small, 3.2 times as long as wide. Spiracular carina narrow. Spiracular index 0.3. 1st gastral tergite 3.9 times, 2nd tergite 1.4 times as long as wide. Petiole rounded laterally, with fine sculpture, without lateral areas or glymmae. Petiolar suture obliterated. 1st gastral sternite with fine sculpture. Thyridium oval. Thyridial index 1.5. Epipleurum separated from the 3rd gastral tergite anteriorly, the crease marked with black. Epipleural index 0.3. Gaster distinctly compressed posteriorly. Ovipositor sheaths slender (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma dark brown medially. Front leg yellowish red from the apex of the coxa onwards, the tarsus darkened apically. Mid femur yellowish red on the distal 0.5-0.7. Mid tibia yellowish red. Mid tarsus light brown basally, dark brown apically. Hind femur red on the distal 0.2. Hind tibia for the greater part black, marked with white or yellowish brown medially on the dorsal side, the apical black ring extending over 0.3-0.4 of the tibia. Hind tarsus blackish, the three basal segments narrowly marked with white basally. 2nd gastral tergite red on the posterior 0.3. 3rd and 4th tergites red, the 4th tergite often marked with black posteriorly (to a variable extent). Pilosity whitish.

 δ : Very similar to the \mathfrak{P} . Sometimes the hind femur black and/or the 2^{nd} gastral tergite almost completely black.

Dusona hastulatae spec. nov.

Holotype (♀): "28.VI.1958 Einbeck" (Germany) (ZSM). – Paratypes: 4♀♀, 1♂ from the same locality, dates 17.6.-1.7.1958; 1♂ "L. 28.7.1958 Einbeck", "1958/XXVII, 43", "ex: *Larentia hastulata* Hbn. (Lep.)"; 1♀ "Ober-Bayern, Garmisch, Kreuzeckweg, bei ca. 900 m, 22.VII.1927, E. Bauer" (all ZSM).

9: Body length 8,5-9 mm. Temples distinctly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the prescutellar groove. Ocellar index 0.85. Frons not depressed, coriaceous, densely rugosepunctate, with a fine median longitudinal carina and with short radial wrinkles, the latter sometimes lacking. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus coriaceous, densely rugosepunctate and rugose. Clypeus flat, coriaceous and with rather dense or dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.5. Genal carina low, bent sidewards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple very slightly coriaceous and with fine dispersed punctation medially. Flagellum with 40-42 segments, filiform, the distal 0.3 slightly attenuated, 2nd segment 2.2 times, shortest segment 1.1 times, penultimate segment 1.2 times as long as wide. Pronotum coriaceous and densely rugosepunctate laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, finely and densely rugose-punctate. Scutellum with strong lateral carinae on the anterior 0.8, coriaceous and with dense punctation. Prescutellar groove for the greater part smooth, with some longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae distinctly raised, the prepectal carina very strongly raised submedially (behind the front coxae), deeply emarginate medially. Pleural part of the prepectal carina present medially, obliterated ventrally and dorsally. Mesopleurum coriaceous, dull, with distinct dense punctation. Speculum with fine striae, dull. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous, finely and densely rugose-punctate and rugose. Pleural carina present. Areolet rather large, distinctly stalked, receiving the 2nd recurrent vein slightly distad of the middle. Nervellus distinctly reclivous, not or almost not intercepted. In the latter case nervellar index 0.6. Tibial spur index 0.6. Hind femur 4.3 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.4 of the hind coxae, with a distinct rather narrow longitudinal furrow, for the greater part with irregular wrinkles, with short transverse wrinkles

in the furrow, with some long transverse wrinkles posteriorly, carinae obliterated. Propodeal spiracle small, 2.6 times as long as wide. Spiracular carina narrow or slightly widened. Spiracular index 0.3. 1st gastral tergite 3.7 times, 2nd tergite 1.7 times as long as wide. Petiole smooth laterally, without glymmae or lateral areas. Petiolar suture fine anteriorly, obliterated posteriorly. 1st gastral sternite smooth, sometimes with fine transverse striae posteriorly. Thyridium small, elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths distinctly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus dark brown. Mandible black, sometimes tinged with brown medially. Wings hyaline. Stigma yellowish brown medially. Front femur black basally and ventrally, yellowish red dorsally and apically. Front tibia and front tarsus yellow, the tarsus light brown apically. Apical 0.2 of mid femur yellowish red. Mid tibia yellowish dorsally, brownish ventrally. Mid tarsus brown, the basitarsus broadly yellowish basally. Hind leg black, the tibia sometimes tinged with brown medially on the dorsal side. 2nd gastral tergite red on the posterior 0.1-0.2, rarely almost completely black. 3rd tergite red. 4th tergite completely black or marked with red to a variable extent anteriorly, often also the posterior edge red. Pilosity whitish.

 δ : Flagellum with 40-42 segments, 2nd segment 1.9 times, shortest segment 0.9 times, penultimate segment 1.1 times as long as wide. In other characters similar to the φ .

Host: *Epirrhoe hastulata* (Hübner, 1790) (Geometridae). The host larva was collected 28.7.1958, the cocoon was formed 1.8.1958, the parasitoid hatched 20.5.1959.

Dusona hispanica spec. nov.

Holotype (\$): "El Ventorillo, 1480 m, Madrid, Trampa Malaise, (20-24).V.1988" (Spain) (Mus. Madrid). – Paratypes: 1\$\partial \text{with the same dates (ZSM); 1\$\partial \text{:"Peña Real, C. Viejo, Madrid, 6-VI-85", "V. Llorente" (Mus. Madrid).

Q: Body length 9-10 mm. Temples slightly constricted behind the eyes, 1.0 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postpetiole. Ocellar index 1.8. Frons hardly depressed above the antennal sockets, coriaceous and densely rugose-punctate and rugose dorsally, shining and with distinct radial wrinkles ventrally, the

wrinkles extending to 0.3 the distance to the median ocellus, with a distinct median longitudinal carina. Antennal carina low and narrow. Face and dorsal 0.6 of the clypeus coriaceous and densely rugosepunctate and rugose. Clypeus flat dorsally and medially, slightly concave ventrally, slightly coriaceous and with distinct dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.6. Genal carina low, slightly bent sidewards subventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.3. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 39-42 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.8 times, shortest segment 0.9 times, penultimate segment 1.1 times as long as wide. Pronotum densely rugose-punctate and rugose on a smooth background laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with distinct dense punctation. Scutellum with lateral carinae on the anterior 0.8, slightly coriaceous and with distinct dispersed or rather dense punctation. Prescutellar groove smooth medially, with fine longitudinal striae laterally. Prepectal carina low, ventral part merging with the pleural part, the latter obliterated and covered with wrinkles on the dorsal 0.3. Transverse carina distinct, about as high as the prepectal carina. Mesopleurum with distinct and dense punctation on a very finely coriaceous or smooth background. Speculum with fine striae anteriorly, with punctation posteriorly. Depression in front of the speculum with distinct rather fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum densely rugosepunctate on a smooth background, partly rugose. Pleural carina complete. Areolet rather narrow, pointed, receiving the 2nd recurrent vein distad of the middle. Nervellus vertical, distinctly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 4.8 times as long as high. Hind tarsal claws slightly elongated, with three distinct pectinate teeth situated basally. Propodeum extending to 0.3 of the hind coxae, with a rather broad and rather shallow sulcus, for the greater part irregularly wrinkled, with few short transverse wrinkles posteriorly, carinae for the greater part obliterated, median longitudinal carinae indicated posteriorly. Propodeal spiracle large, 2.6 times as long as wide. Spiracular carina narrow. Spiracular index 0.35. 1st gastral tergite 3.5 times, 2nd tergite 1.4 times as long as wide. Petiole without glymmae, with slightly indicated narrow lateral areas, smooth in front of these areas. Petiolar suture indicated. 1st gastral sternite for the greater part smooth, with fine transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter

with black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths narrow subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Mandible tinged with dark brown medially. Wings hyaline. Stigma medium brown medially. Front femur yellowish red apically. Front tibia yellowish red. Front tarsus brown. 2nd gastral tergite red on the posterior 0.5-0.6. 3rd and 4th tergites red. Pilosity whitish.

?♂: Mesopleurum with rather dense and rather fine punctation on a distinctly coriaceous background. Metapleurum with distinct and dense punctation on a coriaceous background, partly with fine rugosity. 2nd gastral tergite 2.0 times as long as wide. Maxillary palpus yellowish brown. Mandible yellowish brown dorsally and medially. Front leg yellowish brown from the femur onwards, tarsus brown apically. Mid femur yellowish red on the distal 0.3. Mid tibia yellowish red. 2nd gastral tergite red on the posterior 0.3. In other characters similar to the ♀. These ♂♂ may belong to *D. hispanica* (2♂♂ from Bendery and Kotovskoye, Moldavia; ZSM).

Dusona holmgrenii (Dalla Torre, 1901)

♂ (only the holotype known): Body length 12 mm. Temples moderately constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 1.2. From scarcely depressed above the antennal sockets, for the greater part coriaceous and with fine and dense punctation, with narrow almost smooth and shining areas above the antennal sockets, with a fine median longitudinal carina, without radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, coriaceous and with fine dispersed or rather dense punctation ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low and straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.5. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 50 segments, filiform, the distal 0.4 attenuated, 2nd segment 1.4 times, shortest segment 0.9 times, penultimate segment 1.2 times as long as wide. Pronotum finely coriaceous and densely rugose-punctate laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with fine dense or very dense punctation. Scutellum with lateral carinae on the anterior 0.8, coriaceous, dull, densely rugosepunctate. Prescutellar groove smooth medially, with a few longitudinal striae laterally. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.1 of the latter indistinct and covered with wrinkles. Transverse carina low. Mesopleurum coriaceous, dull, with distinct rather dense or dense punctation. Speculum finely granulate-strigose, shining. Depression in front of the speculum for the greater part with fine longitudinal wrinkles, which merge with the punctation anteriorly, finely granulate-strigose and shining ventrally. Metapleurum coriaceous, dull, densely rugose-punctate. Pleural carina distinct. Areolet rather large, stalked, receiving the 2nd recurrent vein basad of the middle. Nervellus vertical, weakly intercepted. Nervellar index 0.6. Tibial spur index 0.6. Hind femur 5.1 times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, hardly depressed, for the greater part with irregular wrinkles, with a short median longitudinal keel and some short longitudinal wrinkles posteriorly, median longitudinal carinae present posteriorly. Propodeal spiracle large, 2.5 times as long as wide. Spiracular carina narrow. Spiracular index 0.35. 1st gastral tergite 3.5 times, 2nd tergite 2.3 times as long as wide. Petiole with large glymmae, with some fine longitudinal striae in front of the glymmae. Petiolar suture fine. 1st gastral sternite finely coriaceous and with fine wrinkles, with a few transverse striae posteriorly. Thyridium elongated. Thyridial index 1.5. Epipleurum separated from the 3rd gastral tergite and folded in, the crease marked with black. Epipleural index 0.8. Gaster distinctly compressed posteriorly.

Colour: Black. Maxillary palpus yellowish brown. Mandible broadly yellow medially. Tegula dark brown. Wings hyaline. Stigma yellowish brown medially. Front leg yellowish red from the apex of the coxa onwards. Mid leg yellowish red from the femur onwards, the femur marked with brown basally, the tarsus marked with brown distally. Hind tibia broadly yellowish brown medially, narrowly marked with light brown basally and apically. Hind tarsus light brown. 2nd gastral tergite dorsally yellowish brown on the posterior 0.1, laterally yellowish brown on the posterior 0.2, the posterior edge narrowly marked with brown. 3rd tergite yellowish brown, the posterior edge marked with black. 4th tergite for the greater part black, with a narrow longitudinal reddish band dorsally. Genital clasper black. Pilosity whitish.

[♀] unknown.

Dusona incompleta (Bridgman, 1889)

3: Body length 8-10 mm. Temples moderately constricted behind the eyes, 0.85 times as long as the width of an eye, tangential lines on eyes and temples intersect on the 1st gastral segment. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, for the greater part coriaceous and densely rugosepunctate, with narrow smooth areas above the antennal sockets, with a fine and indistinct median longitudinal carina, without radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, finely coriaceous and with fine rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina low, slightly bent forwards subventrally, joining the distinctly raised oral carina distant from the base of the mandible at a right angle. Genal carina fine and indistinct and gena slightly excavated beside the junction. Genal index 0.3. Temple finely coriaceous and with fine dispersed or rather dense punctation medially. Flagellum with 41-46 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.4 times, shortest segment 1.0 times, penultimate segment 1.2 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with fine dense punctation. Scutellum with lateral carinae on the anterior 0.6, finely coriaceous and densely punctate. Prescutellar groove smooth medially, with longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former distinctly and broadly emarginate medially. Pleural part of the prepectal carina low, the dorsal 0.2 covered with wrinkles. Mesopleurum finely coriaceous and with distinct dense or very dense punctation, slightly shining. Speculum finely granulate-strigose, shining. Depression in front of the speculum with distinct longitudinal wrinkles, which merge with the punctation anteriorly. Metapleurum coriaceous, distinctly and densely punctate, partly rugose-punctate. Pleural carina complete. Areolet rather large, stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.8. Tibial spur index 0.65. Hind femur 4.8 times as long as high. Hind tarsal claws slightly elongated, with about two distinct pectinate teeth situated basally. Propodeum extending to 0.5 of the hind coxae, deeply and rather narrowly depressed, with transverse wrinkles in the longitudinal furrow, irregularly wrinkled beside the furrow, the longitudinal carinae obliterated. Propodeal spiracle rather small, 3.0 times as long as wide. Spiracular carina narrow. Spiracular index 0.6. 1st gastral tergite 3.9 times, 2nd tergite 2.3 times as long as wide. Petiole without glymmae, lateral areas weakly indicated, smooth in front of the lateral areas. Petiolar suture fine. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly.

Colour: Black. Maxillary palpus medium brown. Mandible for the greater part black, narrowly marked with yellow or red dorsally. Wings hyaline. Stigma dark yellowish brown medially. Front leg from the femur onwards yellowish red, the femur sometimes marked with black ventrally, the tarsus darkened distally. Front trochanter and front trochantellus sometimes marked with yellowish red distally. Mid femur sometimes brown on the distal 0.2. Mid tibia yellowish brown or brown. Mid tarsus dark brown, the basitarsus marked with yellowish basally. Hind tibia sometimes tinged with yellowish brown medially. Hind basitarsus marked with yellowish basally. 2nd gastral tergite red on the posterior 0.3-0.4. 3rd and 4th tergites red. Pilosity whitish.

?9: Flagellum with 46 segments. Hind tarsal claws short, with about four pectinate teeth situated basally and medially. 1st gastral tergite 4.4 times, 2nd tergite 1.7 times as long as wide. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5. Maxillary palpus and mandible black. Legs almost completely black, distal 0.2 of the front femur and the front tibia dark yellowish brown. In other characters similar to the 9.

Dusona intelligator Aubert, 1966

9: Body length 11 mm. Temples slightly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the 1st gastral segment. Ocellar index 1.5. Frons slightly depressed above the antennal sockets, finely rugose-punctate and rugose, with a distinct median longitudinal carina and with radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus densely rugose-punctate. Clypeus slightly concave, coriaceous and densely punctate ventrally, its apex slightly rounded medially. Malar space index 0.3. Genal carina low and straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.3. Temple finely coriaceous and with fine rather dense punctation medially. Flagellum with 45-49 segments, filiform, the distal 0.2 attenuated, 2nd segment 1.6 times, shortest seg-

ment 1.0 times, penultimate segment 1.4 times as long as wide. Pronotum densely rugose-punctate laterally on the dorsal 0.6, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous and with dense punctation, slightly shining. Scutellum with lateral carinae on the anterior 0.7, finely coriaceous and densely punctate. Prescutellar groove smooth. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.4 of the latter obliterated. Transverse carina present anteriorly, obliterated posteriorly. Mesopleurum with rather dense or dense punctation on a smooth background. Speculum smooth or finely granulate-strigose and shining anteriorly, punctate posteriorly. Depression in front of the speculum with distinct longitudinal or irregular wrinkles, which merge with the punctation anteriorly. Metapleurum with very dense punctation on a smooth background. Pleural carina present. Areolet rather narrow, pointed or short-stalked, receiving the 2nd recurrent vein at or distad of the middle. Nervellus slightly reclivous, distinctly intercepted. Nervellar index 0.5-0.6 (rarely 0.7). Tibial spur index 0.6. Hind femur 4.3 times as long as high. Hind tarsal claws slightly elongated, with about five distinct pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, rather shallowly and rather narrowly depressed, with short transverse wrinkles in the longitudinal furrow, irregularly wrinkled beside the furrow, with long transverse wrinkles posteriorly, longitudinal carinae obliterated. Propodeal spiracle large, 2.8 times as long as wide. Spiracular carina usually indistinct. Spiracular index 0.3. 1st gastral tergite 4.0 times, 2nd tergite 1.6 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture fine, partly obliterated. 1st gastral sternite smooth, sometimes with some transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum present as a small semicircular pendent lobe anteriorly, not separated from the 3rd gastral tergite by a crease or a black stripe, the tergite with distinct black lateral stripes well above the ventrolateral edge. Gaster moderately compressed posteriorly. Ovipositor sheaths slightly widened medially and subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus dark brown basally, yellowish red medially and apically. Mandible narrowly reddish brown or dark brown medially. Wings hyaline. Stigma yellowish brown medially. Front leg reddish brown from the femur onwards, the tarsus marked with dark brown distally. Mid femur reddish brown, marked with black basally. Mid tibia reddish brown. Mid tarsus dark brown. Hind tibia often tinged with dark brown medially on the dorsal side. 2nd gastral tergite reddish brown or dark brown

on the posterior 0.6-0.7. 3rd and 4th tergites reddish brown or dark brown. Pilosity whitish.

\delta: Front trochanter and front trochantellus reddish brown. In other characters similar to the \mathcal{P} .

Dusona irregularis spec. nov.

Holotype (♀): "Jura-Krakowska-Czestochowska, wies Bolechowice, 30.4.61, J.P." (Poland) (Coll. Sawoniewicz). – Paratype: 1♂ "Vitebskaya obl., Berezinskiy zap., 6.VI.985, ... A. Tereshkin leg." (Belorussiya) (ZSM).

9: Body length 14 mm. Temples distinctly constricted behind the eyes, 0.85 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 0.6. Frons hardly depressed ventrally, coriaceous and with fine dense punctation dorsally and medially, almost smooth ventrally above the antennal sockets, with a fine median longitudinal carina and without radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, coriaceous and with fine rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.2. Genal carina low and straight ventrally, joining the low oral carina distant from the base of the mandible at an acute angle. Genal index 0.45. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 46 segments, stout, filiform, the distal 0.2 attenuated, 2nd segment 1.4 times, shortest segment 0.8 times, penultimate segment 1.1 times as long as wide. Pronotum with fine punctation and fine longitudinal wrinkles laterally. Mesoscutum coriaceous and with distinct dense or very dense punctation. Scutellum with lateral carinae on the anterior 0.3, coriaceous and densely rugose-punctate and rugose. Prescutellar groove smooth. Prepectal carina low, ventral part merging with the pleural part, the latter almost complete, covered with wrinkles dorsally. Transverse carina present, low. Mesopleurum finely coriaceous and with distinct dense or very dense punctation, with longitudinal wrinkles below the speculum. Speculum large, smooth. Depression in front of the speculum almost smooth ventrally, in other parts with fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum finely and densely rugose-punctate and rugose. Pleural carina distinct. Areolet large, stalked, receiving the 2nd recurrent vein at the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.75. Tibial spur index 0.6. Hind femur 5.0 times as long as high. Hind tarsal claws slightly elongated, with 7-8 short pectinate teeth situated basally, medially and subapically. Propodeum extending to 0.3 of the hind coxae, broadly and rather shallowly depressed, for the greater part with irregular wrinkles, with few distinct longitudinal wrinkles posteriorly, carinae almost completely obliterated. Propodeal spiracle rather large, 2.3 times as long as wide. Spiracular carina fine and narrow. Spiracular index 0.7. 1st gastral tergite 3.4 times, 2nd tergite 2.4 times as long as wide. Petiole with large and deep glymmae, smooth or with very fine longitudinal striae in front of the glymmae. Petiolar suture partly indicated. 1st gastral sternite with fine irregular sculpture anteriorly and medially, with distinct transverse striae posteriorly. Thyridium elongated. Thyridial index 1.5. Epipleurum separated from the 3rd gastral tergite and folded in, the crease marked with black. Epipleural index 0.6-0.7. Gastral tergites strongly shining. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus yellowish brown. Mandible yellow dorsally and medially. Tegula dark brown. Wings hyaline. Stigma light yellowish brown medially. Front leg yellowish red from the femur onwards. Mid femur yellowish red, slightly darkened basally. Mid tibia yellowish red. Mid tarsus dark brown, marked with light brown basally. Hind femur red on the distal 0.4. Hind tibia for the greater part yellowish red, narrowly dark brown basally, narrowly light brown apically. Hind tarsus dark brown, marked with yellowish red basally on the ventral side. 2nd gastral tergite red on the posterior 0.3, the posterior edge dark brown dorsally. 3rd tergite red, narrowly marked with black posteriorly. Pilosity whitish.

3: Flagellum slender, median segments 1.1 times as long as wide (both flagellar tips broken off). Metapleurum with fine dispersed punctation on a smooth background centrally. Hind tarsal claws distinctly elongated, with 7-8 short pectinate teeth situated basally and medially. Median longitudinal carinae of the propodeum indicated medially and posteriorly. 1st gastral sternite with very fine sculpture, partly smooth. Gastral tergites not strongly shining. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula light brown. Front and mid legs yellow from the apex of the coxa onwards, the femora yellowish red. Hind tibia broadly yellow medially. Hind tarsus light brown. 2nd gastral tergite dorsally red on the posterior 0.4, the posterior edge narrowly dark brown, almost completely red laterally. 3rd tergite red. 4th tergite red, the posterior 0.3 black. In other characters similar to the ♀.

Dusona lamellator Aubert, 1960

9: Body length 11-13 mm. Temples strongly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the centre of the mesoscutum. Ocellar index 0.6. Frons slightly depressed above the antennal sockets, coriaceous and densely rugose-punctate dorsally, somewhat shining ventrally, with a low median longitudinal bulge and a fine longitudinal carina on top of this bulge, and with distinct radial wrinkles, which extend to 0.6 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.4 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, finely coriaceous and densely punctate ventrally, its apex truncate medially. Malar space index 0.3. Genal carina distinctly raised behind the ventral half of the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, strongly shifted downwards, almost reaching the ventral edge of the mandibles (in lateral view and with the mandibles closed). Gena deeply depressed close to the junction, the depression 1.5 times as deep as the width of the 2nd flagellar segment. Genal index 0.8. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 46-49 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.3 times, shortest segment 0.8 times, penultimate segment 1.0 times as long as wide. Pronotum densely rugosepunctate and rugose laterally on the dorsal 0.6, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.4, distinctly and densely punctate on a smooth background. Prescutellar groove smooth medially, with a few longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae distinctly raised, the former slightly and rather narrowly emarginate medially. Pleural part of the prepectal carina low, almost complete, the dorsal 0.1-0.2 covered with wrinkles. Mesopleurum with distinct and dense punctation on a smooth background, with some longitudinal striae posteriorly in front of the metapleurum. Speculum finely coriaceous and granulate-strigose, almost smooth ventrally, shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum with distinct longitudinal wrinkles dorsally, irregularly wrinkled or rugosepunctate ventrally. Pleural carina distinct. Areolet rather large, pointed or short-stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus vertical or slightly reclivous, weakly intercepted.

Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 5.5 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally and medially. Propodeum extending to 0.7 of the hind coxae, broadly and rather deeply depressed, with long transverse wrinkles in the sulcus and in the lateral areas, median longitudinal carinae indicated medially and posteriorly. Propodeal spiracle large, 3.0 times as long as wide. Spiracular carina narrow, sometimes indistinct. Spiracular index 0.4. 1st gastral tergite 5.5 times, 2nd tergite 2.7 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture fine or partly obliterated. 1st gastral sternite smooth, sometimes with fine transverse striae posteriorly. Thyridium oval. Thyridial index 2.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma medium brown medially. Front leg from the apex of the coxa onwards yellow. Mid leg from the trochanter onwards yellow, the trochanter black basally, the tarsus brownish distally. Hind trochantellus marked with yellow apically and hind femur narrowly marked with yellow basally. Hind tibia yellow. Hind tarsus for the greater part dark brown, the basitarsus broadly yellow basally. 2nd gastral tergite dorsally yellow on the posterior 0.3-0.4, laterally yellow on the posterior 0.5-0.6. 3rd and 4th tergites yellow. Pilosity whitish.

 ${\mathfrak S}$: Mid leg from the apex of the coxa onwards yellow, the tarsus brownish distally. In other characters similar to the ${\mathfrak P}$.

Dusona libertatis (Teunissen, 1947)

Q: Body length 14-16 mm. Temples strongly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.7 of the mesoscutum. Ocellar index 1.0. Frons slightly depressed above the antennal sockets, finely coriaceous and with fine rather dense punctation dorsally, shining ventrally, with a distinct median longitudinal bulge and a fine median longitudinal carina on top of this bulge, with distinct radial wrinkles, which extend to 0.4 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat dorsally, slightly concave ventrally,

finely coriaceous and with rather dense punctation ventrally, its apex truncate medially. Malar space index 0.3. Genal carina distinctly raised behind the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible at an acute angle, the junction not shifted downwards but situated above the base of the mandible. Genal index 0.6. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 58-62 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.4 times, shortest segment 0.8 times, penultimate segment 1.1 times as long as wide. Pronotum finely coriaceous and with fine dense punctation laterally on the dorsal 0.5, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with fine dense punctation. Scutellum with lateral carinae on the anterior 0.3, coriaceous and with dense punctation. Prescutellar groove with fine longitudinal striae. Ventral part of the prepectal carina low, merging with the slightly raised transverse carina (rarely merging with the pleural part). Pleural part of the prepectal carina complete, joining the anterior edge of the mesopleurum. Mesopleurum finely coriaceous, shining, with rather fine dispersed punctation centrally. Speculum finely coriaceous and granulatestrigose, slightly shining. Depression in front of the speculum dorsally with fine longitudinal striae, which merge with the punctation anteriorly, ventrally granulate-strigose. Metapleurum coriaceous and densely rugose. Pleural carina distinct. Areolet large, short-stalked, receiving the 2nd recurrent vein basad of the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.5. Hind femur 5.5 times as long as high. Hind tarsal claws rather short, with about eight long pectinate teeth situated basally, medially and subapically. Propodeum extending to 0.3 of the hind coxae, broadly and rather shallowly depressed, the sulcus with transverse wrinkles, the median longitudinal carinae present medially and posteriorly, covered with wrinkles anteriorly. Propodeal spiracle large, 3.3 times as long as wide. Spiracular carina narrow. Spiracular index 0.4. 1st gastral tergite 4.7 times, 2nd tergite 1.8 times as long as wide. Petiole with large glymmae, smooth in front of the glymmae. Petiolar suture indicated anteriorly, obliterated posteriorly. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.0. Epipleurum separated from the 3rd gastral tergite anteriorly by a crease or bulge, folded in or pendent as a narrow stripe, the crease or bulge partly marked with black. Epipleural index 0.2-0.6. Gaster distinctly compressed posteriorly. Ovipositor sheaths scarcely widened medially and subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish red. Mandible yellowish red medially. Tegula yellow. Wings slightly tinged with light brown. Stigma yellowish brown medially. Front leg yellow or yellowish red from the trochanter onwards, the tarsus darkened distally. Mid femur yellow on the distal 0.2. Mid tibia yellow. Mid tarsus medium brown. Hind femur narrowly marked with yellowish red apically on the dorsal side. Hind tibia yellow. Hind tarsus dark brown. 2nd gastral tergite dorsally yellow on the posterior 0.4, laterally yellow over its total length, the ventrolateral edge marked with black. 3rd and 4th tergites yellow. 5th tergite narrowly yellow anteriorly. Anterior sternites yellow, posterior sternites black. Pilosity whitish.

 δ : Apex of the front coxa yellow. Mid femur black basally and ventrally, yellowish red dorsally and apically. In other characters similar to the \circ .

Dusona limnobia (Thomson, 1887)

9: Body length 7-9 mm. Temples strongly constricted behind the eyes, 0.6 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.7 of the mesoscutum. Ocellar index 0.6. From scarcely depressed above the antennal sockets, for the greater part coriaceous and densely rugose-punctate, slightly shining ventrally, with a fine and sometimes indistinct median longitudinal carina and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugosepunctate and rugose. Clypeus flat, coriaceous and with rather dense or dispersed punctation ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low ventrally, slightly bent sidewards and distinctly (but not strongly) bent forwards subventrally, joining the low oral carina close to the base of the mandible at an acute angle. Genal index 0.1-0.2. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 39-43 segments, slender, filiform, the distal 0.2 slightly attenuated, 2nd segment 2.1 times, shortest segment 1.0 times, penultimate segment 1.4 times as long as wide. Pronotum densely rugose-punctate and rugose laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.8, coriaceous, dull, very densely punctate. Prescutellar groove smooth medially, with longitudinal striae laterally. Ventral part of the prepectal carina low, slightly emarginate medially, usually merging with the transverse carina, rarely merging with the pleural part, the transversa carina distinct and complete, the dorsal 0.4 of the pleural part of the prepectal carina obliterated. Mesopleurum coriaceous, often dull, rarely slightly shining, with rather dense or dense punctation (the punctation often dense in the centre and rather dense below the speculum). Speculum with fine longitudinal striae, slightly shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum densely rugose. Pleural carina complete. Areolet rather large, short-stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus vertical or slightly reclivous, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.6. Hind femur 4.6 times as long as high. Hind tarsal claws short, with about four pectinate teeth situated basally. Propodeum extending to 0.5 of the hind coxae, distinctly and rather narrowly depressed, with short transverse wrinkles in the longitudinal furrow, irregularly wrinkled beside the furrow, median longitudinal carinae indicated posteriorly. Propodeal spiracle rather small, 2.5 times as long as wide. Spiracular carina narrow. Spiracular index 0.7. 1st gastral tergite 5.0 times, 2nd tergite 2.3 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture fine, partly obliterated. 1st gastral sternite for the greater part smooth, with some transverse striae posteriorly. Thyridium elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths distinctly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus yellow or yellowish brown. Mandible marked with yellow dorsally and medially. Tegula light brown. Wings hyaline. Stigma dark brown medially. Front leg yellowish red from the trochanter onwards. Mid leg yellowish red from the femur onwards. Hind femur often reddish brown, sometimes black or marked with black. Hind tibia broadly yellowish brown medially, narrowly marked with blackish basally, rather broadly marked with blackish apically, the apical ring extending over 0.15-0.3 of the tibia. Hind tarsus dark brown, the basal 0.5 of the basitarsus yellowish brown. 2^{nd} gastral tergite red on the posterior 0.2-0.3. 3^{rd} and 4^{th} tergites red, the 4th tergite sometimes marked with black dorsally. 5th tergite rarely marked with red anteriorly. Pilosity whitish.

6: Flagellum with 41-44 segments. Mesopleurum coriaceous, dull, densely punctate. Propodeum often somewhat more coarsely wrinkled, the median longitudinal carinae present medially and posteriorly. Mandible broadly yellow medially. Tegula yellow or yellowish brown. Apex of the front and mid coxae yellow. Median gastral tergites sometimes tinged with black dorsally. In other characters similar to the $\mathfrak{P}.$

Dusona longiseta Hinz, 1961

9: Body length 10-12 mm. Temples slightly constricted behind the eyes, 0.9-1.0 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postpetiole. Ocellar index 0.9. Frons not depressed, coriaceous and densely punctate, sometimes partly rugose, a median longitudinal carina absent or weakly indicated, the radial wrinkles short or obliterated. Antennal carina low and narrow. Face finely coriaceous and densely punctate, partly rugose-punctate. Clypeus for the greater part flat, slightly concave ventrally, densely punctate on an almost smooth background, its apex slightly rounded medially. Malar space index 0.4. Genal carina low, slightly bent sidewards subventrally, joining the low oral carina at an almost right angle. Genal index 0.2-0.35. Temple finely coriaceous and with fine rather dense or dense punctation medially. Flagellum with 40-42 segments, filiform, the distal 0.2 attenuated, 2nd segment 1.9 times, shortest segment 0.8 times, penultimate segment 1.0 times as long as wide. Pronotum very densely punctate on a smooth background laterally on the dorsal 0.3, with longitudinal wrinkles ventrally. Mesoscutum rather densely or densely punctate on a slightly coriaceous or smooth background. Scutellum with lateral carinae on the anterior 0.8, rather densely or densely punctate on a smooth background. Prescutellar groove with fine longitudinal striae. Usually the ventral part of the prepectal carina merging with the transverse carina, both carinae low, the pleural part of the prepectal carina present medially, fine and sometimes obliterated ventrally. Rarely the ventral part of the prepectal carina merging with the pleural part, and the transverse carina obliterated posteriorly. Dorsal 0.4 of the pleural part of the prepectal carina always obliterated. Pleural part of the prepectus with dense punctation on a smooth background. Mesopleurum distinctly and rather densely or densely punctate on a smooth (rarely finely coriaceous) background. Speculum small, with fine sculpture and strongly shining anteriorly, punctate posteriorly. Depression in front of the speculum with short longitudinal wrinkles, which merge with the punctation anteriorly. Metapleurum densely rugose-punctate and rugose. Pleural carina distinct. Areolet large, pointed, receiving the 2nd recurrent vein distad of the middle. Nervellus almost vertical, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.6. Hind femur 4.0 times as long as high. Hind tarsal claws slightly elongated, with five distinct pectinate teeth situated basally, medially and subapically. Propodeum extending to 0.3 of the hind coxae, broadly and rather shallowly depressed, the sulcus with transverse wrinkles, the median longitudinal carinae present posteriorly, obliterated medially. Propodeal spiracle rather small, 2.4 times as long as wide. Spiracular carina narrow. Spiracular index 0.7. 1st gastral tergite 3.0 times, 2nd tergite 1.4 times as long as wide. Petiole with narrow lateral areas (these sometimes almost obliterated), without glymmae, smooth in front of the lateral areas. Petiolar suture fine. 1st gastral sternite smooth. Thyridium circular. Thyridial index 2.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths not widened subapically (in lateral view). Ovipositor slightly bent upwards. Ovipositor index about 3.0.

Colour: Black. Mandible yellowish red or brown medially. Wings hyaline. Stigma dark brown medially. Front femur yellowish red anteriorly. Front tibia and tarsus yellowish red. Mid tibia and tarsus dark brown, the tibia with a yellowish stripe anteriorly. Hind tibia black, sometimes tinged with brown medially. 2nd gastral tergite red on the posterior 0.1-0.2. 3rd and 4th tergites red, the 4th tergite sometimes marked with black dorsally. Pilosity whitish.

δ: Flagellum with 43-46 segments, 2nd segment 1.6 times, shortest segment 1.0 times, penultimate segment 1.2 times as long as wide. Hind claws distinctly elongated, with a long apical tooth and about four rather short pectinate teeth in the basal half. Front leg for the greater part black, with anterior yellow marks on the coxa and the trochanter and anterior yellowish stripes on the femur and the tibia. Mid and hind legs black, the mid femur sometimes with a small yellow mark apically. 2nd gastral tergite sometimes completely black. In other characters similar to the ♀.

Dusona luteipes (Thomon, 1887)

Q (only the holotype known): Body length 8 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 0.9. Frons scarcely depressed above the antennal sockets, coriaceous and densely rugose-punctate dorsally, shining ventrally, with a distinct median longitudinal carina and with radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.2 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat,

finely coriaceous and with fine rather dense punctation ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low and straight ventrally, joining the hardly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.3. Temple finely coriaceous and with very fine dispersed punctation medially. Both flagellar tips broken off in the only known specimen, flagellum filiform, 2nd segment 1.7 times, shortest segment 1.0 times as long as wide. Pronotum coriaceous and densely punctate laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.8, coriaceous and with dispersed or rather dense punctation. Prescutellar groove smooth. Prepectal carina low, the ventral part merging with the pleural part, the dorsal 0.1 of the latter covered with wrinkles. Transverse carina present anteriorly, obliterated posteriorly. Mesopleurum coriaceous, dull, with rather fine and rather dense or dense punctation. Speculum granulatestrigose and with fine longitudinal striae, somewhat shining. Depression in front of the speculum with distinct fine longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous and finely rugose. Pleural carina distinct. Areolet rather small, stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.3 times as long as high. Hind tarsal claws short, with about three short pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, rather shallowly and narrowly depressed, with short transverse wrinkles in the longitudinal furrow, irregularly wrinkled beside the furrow, median longitudinal carinae indicated posteriorly. Propodeal spiracle rather small, 2.8 times as long as wide. Spiracular carina narrow. Spiracular index 0.6. 1st gastral tergite 3.6 times, 2nd tergite 1.6 times as long as wide. Petiole with narrow lateral areas and with small glymmae, smooth in front of the glymmae. Petiolar suture fine, obliterated posteriorly. 1st gastral sternite for the greater part smooth, with fine transverse striae posteriorly. Thyridium circular. Thyridial index 1.5. Epipleurum separated from the 3rd gastral tergite anteriorly and folded in, the crease marked with black. Epipleural index 0.3. Gaster distinctly compressed posteriorly. Ovipositor sheaths scarcely widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish brown. Mandible narrowly marked with reddish brown subapically. Wings hyaline. Stigma dark brown medially. Front, mid and hind legs yellowish red from the femur onwards. Hind tarsus brown. 2nd gastral tergite dorsally red on the posterior 0.6,

laterally completely red. 3rd and 4th tergites red. 5th tergite with small reddish marks laterally. Pilosity whitish.

ð unknown.

Dusona notabilis (Förster, 1868)

9: Body length 7-9 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum. Ocellar index 0.9. Frons not depressed, coriaceous and with fine rather dense punctation dorsally and medially, with narrow almost smooth areas ventrally above the antennal sockets, with a fine median longitudinal carina and with distinct radial wrinkles, which extend to 0.4 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.2 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, finely coriaceous and with fine rather dense or dense punctation ventrally, shining, its apex slightly rounded medially. Malar space index 0.6. Genal carina low ventrally, slightly bent forwards and distinctly bent sidewards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 33-38 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.8 times, shortest segment 0.9 times, penultimate segment 1.0-1.1 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and with fine and very dense punctation, partly rugose-punctate. Scutellum with lateral carinae on the anterior 0.7, coriaceous and with rather dense punctation. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae distinctly raised, the former distinctly emarginate medially. Pleural part of the prepectal carina for the greater part low, slightly raised ventrally (at the junction with the transverse carina), the dorsal 0.1-0.2 covered with wrinkles. Mesopleurum coriaceous, dull or slightly shining, with distinct fine and dense punctation. Speculum granulate-strigose, slightly shining. Depression in front of the speculum with wrinkles dorsally, with fine longitudinal striae and granulate-strigose medially and ventrally, the striae merging with the punctation anteriorly. Metapleurum coriaceous, dull, densely rugose-punctate and rugose. Pleural carina distinct. Areolet rather large, pointed or short-stalked, receiving the 2nd recurrent vein at or slightly distad of the middle. Nervellus vertical, weakly intercepted. Nervellar

index 0.7-0.8. Tibial spur index 0.65. Hind femur 4.8 times as long as high. Hind tarsal claws short, with about four short pectinate teeth situated basally and medially. Propodeum extending to 0.4 of the hind coxae, rather broadly and rather deeply depressed, the sulcus for the greater part with fine or rather coarse irregular wrinkles, sometimes with some fine and short transverse wrinkles in the midline, or with some longitudinal wrinkles posteriorly, the median longitudinal carinae usually obliterated, sometimes scarcely indicated medially and posteriorly. Propodeal spiracle rather small, 2.6 times as long as wide. Spiracular carina narrow. Spiracular index 0.7. 1st gastral tergite 4.5 times, 2nd tergite 1.9 times as long as wide. Petiole with rather small glymmae, or the glymmae obliterated, smooth in front of the glymmae. Petiolar suture almost completely obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster moderately compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus dark brown. Wings hyaline. Stigma light yellowish brown medially. Front leg from the femur onwards yellowish red, the femur darkened basally, the tarsus darkened apically. Mid femur yellowish red on the distal 0.3. Mid tibia yellowish red. Mid tarsus brown. Hind tibia yellowish brown or dark yellowish brown medially, narrowly black basally, broadly black apically, the apical ring extending over 0.3-0.4 of the tibia. Hind tarsus dark brown, the basitarsus narrowly marked with yellowish basally. 2nd gastral tergite red on the posterior 0.2-0.3. 3rd tergite red. 4th tergite red on the anterior 0.5 or completely red. Pilosity whitish.

 σ : Flagellum with 35-39 segments. Propodeum slightly depressed, irregularly wrinkled, usually with a short longitudinal keel posteriorly. Maxillary palpus yellowish brown. Mandible marked with brown dorsally and medially. In other characters similar to the φ .

Dusona peregrina (Wollaston, 1858)

2: Body length 8-9 mm. Temples moderately constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postscutellum. Ocellar index 0.6. Frons not depressed, coriaceous and densely rugose-punctate and rugose, with a fine median longitudinal carina and with a few radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina

low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugose. Clypeus flat, coriaceous and with rather dense punctation ventrally, its apex truncate medially. Malar space index 0.3. Genal carina low medially behind the eye, strongly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the base of the mandible. Gena moderately depressed close to the junction, the depression 0.5 times as deep as the width of the 2nd flagellar segment. Genal index 0.3. Temple finely coriaceous and with very fine dispersed punctation medially. Flagellum with 36-39 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.8 times, shortest segment 0.9 times, penultimate segment 1.0 times as long as wide. Pronotum coriaceous and densely rugose laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.3, coriaceous and densely punctate. Prescutellar groove smooth or with indistinct longitudinal striae. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.1-0.2 covered with wrinkles. Transverse carina present, sometimes obliterated posteriorly. Mesopleurum finely coriaceous, slightly shining, with dispersed or rather dense punctation. Speculum granulatestrigose dorsally, finely coriaceous and shining ventrally. Depression in front of the speculum with fine longitudinal striae dorsally and medially, which merge with the punctation anteriorly, almost smooth ventrally. Metapleurum finely coriaceous and with rather dense or dense punctation, with some fine longitudinal striae dorsally. Pleural carina distinct. Areolet large, narrowly sessile, receiving the 2nd recurrent vein at or slightly basad of the middle. Nervellus vertical or slightly reclivous, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 5.1 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.5 of the hind coxae, broadly and not deeply depressed, the sulcus with short or long transverse wrinkles, the median longitudinal carinae almost completely obliterated. Propodeal spiracle rather large, 2.7 times as long as wide. Spiracular carina narrow. Spiracular index 0.5. 1st gastral tergite 5.1 times, 2nd tergite 2.3 times as long as wide. Petiole with indistinct narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths almost not widened subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma dark brown medially. Front leg from the apex of the coxa onwards yellow. Mid leg from the trochanter onwards yellow, the tarsus brown apically. Hind tibia yellow. Hind tarsus dark brown, the basitarsus narrowly marked with yellowish basally. 2nd gastral tergite yellow on the posterior 0.2. 3rd and 4th tergites yellow. Pilosity whitish.

 σ : Flagellum with 40-43 segments, 2nd segment 1.9 times, shortest segment 1.0 times, penultimate segment 1.2 times as long as wide. Depression in front of the speculum often with longitudinal striae also ventrally. Median longitudinal carinae of the propodeum sometimes indicated medially and posteriorly. Mid coxa yellow apically. 2nd gastral tergite yellow on the posterior 0.3. In other characters similar to the σ .

Dusona pulchripes (Holmgren, 1872)

9: Body length 7-8 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum. Ocellar index 0.65. Frons not depressed, coriaceous and densely rugose-punctate, with a median longitudinal carina and with rather short radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate and rugose, the clypeus with some longitudinal wrinkles. Clypeus almost flat, finely coriaceous and with fine rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina slightly raised ventrally, almost straight, joining the oral carina at the base of the mandible at an acute angle. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 35-38 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.6 times, shortest segment 0.85 times, penultimate segment 1.0 times as long as wide. Pronotum coriaceous and densely rugose-punctate and rugose laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and finely and densely rugose-punctate and rugose. Scutellum with lateral carinae on the anterior 0.4, coriaceous and densely rugose-punctate. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae low. Pleural part of the prepectal carina distinct medially, very fine and sometimes indistinct ventrally, the

dorsal 0.3 obliterated. Mesopleurum finely coriaceous, shining, with distinct and dense punctation. Speculum with fine striae and granulate-strigose dorsally, almost smooth and shining ventrally. Depression in front of the speculum with fine long longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose. Pleural carina complete. Areolet rather large, pointed or short-stalked, receiving the 2nd recurrent vein at or distad of the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 5.7 times as long as high. Hind tarsal claws short, with a few short pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, rather narrowly and rather deeply depressed, with short transverse wrinkles in the longitudinal furrow anteriorly and medially, with irregular wrinkles beside the furrow, with long wrinkles posteriorly, the median longitudinal carinae obliterated. Propodeal spiracle small, 2.2 times as long as wide. Spiracular carina narrow. Spiracular index 1.0. 1st gastral tergite 4.2 times, 2nd tergite 1.6 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5-2.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma dark brown medially. Front leg from the apex of the coxa onwards yellow. Mid leg from the femur onwards yellow, the femur sometimes marked with dark brown on the basal 0.2-0.4, the tarsus brown. Hind tibia yellow. Hind tarsus dark brown, the basitarsus narrowly marked with yellowish basally. 2nd gastral tergite yellow on the posterior 0.3. 3rd tergite yellow, sometimes narrowly tinged with brown posteriorly. 4th tergite sometimes narrowly marked with yellow anteriorly. Pilosity whitish.

 δ : Median longitudinal carinae of the propodeum indicated posteriorly. Mid leg from the trochanter onwards yellow. In other characters similar to the \circ .

Dusona rossica Hinz, 1979

2: Body length 14-15 mm. Temples slightly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior part of the propodeum.

Ocellar index 1.0. Frons slightly depressed above the antennal sockets, coriaceous and densely rugosepunctate dorsally, with narrow almost smooth areas above the antennal sockets, with a fine median longitudinal carina and with some short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugosepunctate and rugose. Clypeus slightly concave, coriaceous and with dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.3. Genal carina low and straight ventrally, joining the low oral carina close to the base of the mandible at an acute angle. Genal index 0.2. Temple finely coriaceous and with fine rather dense punctation medially. Flagellum with 56 segments, filiform, the distal 0.3 distinctly attenuated, 2nd segment 1.6 times, shortest segment 1.0 times, penultimate segment 1.3 times as long as wide. Pronotum densely rugose-punctate on an almost smooth background laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous, shining, with fine dense punctation. Scutellum with lateral carinae on the anterior 0.6, densely punctate on a smooth background. Prescutellar groove smooth medially, with fine longitudinal striae laterally. Prepectal carina low, the ventral part merging with the pleural part, the latter almost complete, the dorsal 0.1 covered with wrinkles. Transverse carina present, low. Mesopleurum finely coriaceous, shining, with fine rather dense or dense punctation. Speculum finely granulate-strigose, shining. Depression in front of the speculum with distinct rather long longitudinal striae, which merge with the punctation anteriorly. Metapleurum finely coriaceous, slightly shining, densely punctate. Pleural carina distinct. Areolet narrow, pointed, receiving the 2nd recurrent vein at the middle. Nervellus slightly inclivous, weakly intercepted. Nervellar index 0.7-0.8. Tibial spur index 0.7. Hind femur 4.3 times as long as high. Hind tarsal claws somewhat elongated, with about six short pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, broadly and rather shallowly depressed, with short transverse wrinkles anteriorly and medially in the midline, with irregular wrinkles laterally, with long transverse wrinkles posteriorly, the median longitudinal carinae indicated medially and posteriorly. Propodeal spiracle large, 3.2 times as long as wide. Spiracular carina narrow. Spiracular index 0.3. 1st gastral tergite 2.8 times, 2nd tergite 1.2 times as long as wide. Petiole with large glymmae, smooth in front of the glymmae. Petiolar suture fine. 1st gastral sternite for the greater part smooth, with some fine transverse striae posteriorly. Thyridium oval. Thyridial index 1.0. Epipleurum separated from the 3rd gastral tergite anteriorly and folded in, the crease marked with black or the tergite with a black stripe close to the crease. Epipleural index 0.4. Gaster distinctly compressed posteriorly. Ovipositor sheaths scarcely widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish brown. Mandible marked with yellowish red dorsally and medially. Tegula dark brown. Wings hyaline. Stigma light yellowish brown medially. Front and mid legs from the femur onwards light reddish brown, the tarsi tinged with dark brown apically. Hind femur reddish brown or marked with reddish brown distally. Hind tibia yellowish red, narrowly marked with dark brown basally, narrowly and indistinctly marked with reddish brown apically. Hind tarsus yellowish red, tinged with dark brown distally. 2nd gastral tergite red on the posterior 0.5. 3rd and 4th tergites red. Pilosity whitish.

♂ not available for study.

Dusona rubidatae spec. nov.

Holotype (♀): "L. 7.IX.1974 Einbeck" (Germany), "1974/XXIII 25", "E094930 N514100", "ex: *Cidaria rubidata* F. (Lep.)" (ZSM). – Paratypes: 6♀♀, 4♂♂ with the same dates (ZSM); 3♀♀, 2♂♂ "L. 7.IX.1974 Einbeck", "1974/XVIII", "ex: R618", "ex: *Epirrhoe molluginata* Hb. (Lep.)" (ZSM).

9: Body length 7.5-10 mm. Temples strongly constricted behind the eyes, 0.6 times as long as the width of an eye, tangential lines on eyes and temples intersect on the centre of the mesoscutum. Ocellar index 0.7. Frons scarcely depressed, coriaceous and with fine dispersed punctation dorsally and medially, almost smooth and with short radial wrinkles ventrally above the antennal sockets, with a fine median longitudinal carina. Antennal carina slightly raised, its rim slightly thickened and with fine transverse striae. Face and dorsal half of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, finely coriaceous and with fine dispersed punctation ventrally, its apex almost straight medially. Malar space index 0.2. Genal carina low, slightly bent sideward and hardly bent forwards subventrally, joining the low oral carina distant from the base of the mandible at an acute angle. Genal index 0.2. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 44-49 segments, slender, filiform, the distal 0.2 attenuated, 2nd segment 2.3 times, shortest segment 1.0-1.1 times, penultimate segment 1.4 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.2,

with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely punctate. Scutellum with lateral carinae on the anterior 0.8, finely coriaceous and with distinct dense punctation. Prescutellar groove smooth medially, with few longitudinal striae laterally. Prepectal carina low, ventral part usually merging with the pleural part, rarely with the transverse carina, dorsal 0.1-0.2 of the pleural part obliterated. Transverse carina present, raised anteriorly, rather low posteriorly. Mesopleurum coriaceous, rather dull, with dispersed (rarely rather dense) punctation. Speculum granulate-strigose, rather dull. Depression in front of the speculum with fine longitudinal striae, which merge with the punctation or with a narrow area without punctation anteriorly. Metapleurum coriaceous, dull, rugose anteriorly, with additional very fine punctures posteriorly. Pleural carina distinct. Areolet rather small, stalked, receiving the 2nd recurrent vein at the middle. Nervellus reclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 5.2 times as long as high. Hind tarsal claws short, with about four rather long pectinate teeth situated basally and medially. Propodeum extending to 0.5 of the hind coxae, rather broadly and rather deeply depressed, the longitudinal furrow with irregular wrinkles anteriorly, with short transverse wrinkles medially and posteriorly, the longitudinal carinae obliterated. Propodeal spiracle 4.0 times as long as wide. Spiracular carina narrow. Spiracular index 0.6. 1st gastral tergite 4.3 times, 2nd tergite 2.2 times as long as wide. Petiole without glymmae, with narrow lateral areas, smooth in front of these areas. Petiolar suture fine, partly obliterated. 1st gastral sternite for the greater part smooth, with fine transverse wrinkles posteriorly. Thyridium elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellow. Mandible narrowly yellowish brown medially. Tegula dark brown. Wings hyaline. Stigma yellowish brown medially. Front and mid legs yellowish red from the trochanter onwards. Mid femur sometimes darkened proximally. Mid tarsus darkened distally. Hind femur black, dark brown or red (red in about 20 % of the specimens). Hind tibia yellowish brown medially, narrowly dark brown basally, rather broadly dark brown apically, the apical ring extending over 0.15-0.3 of the tibia. Hind tarsus dark brown. 2nd gastral tergite red on the posterior 0.3. 3rd and 4th tergites red. Pilosity whitish.

ở: Flagellum with 45-51 segments, 2nd segment 2.0 times, shortest segment 1.1 times, penultimate segment 1.3 times as long as wide. Median longitudinal carinae of the propodeum partly present. Mandible yellow (teeth dark brown). Tegula yellow or yellowish brown. Apex of the front coxa, front trochanter and front trochantellus yellow. 2nd gastral tergite red on the posterior 0.1-0.3. 4th tergite sometimes marked with black dorsally or posteriorly. In other characters similar to the ♀.

Hosts: *Catarhoe rubidata* (Denis & Schiffermüller), *Epirrhoe molluginata* (Hübner) (Geometridae). The host larvae were collected 7.9.1974, the cocoons were formed in September 1974, the parasitoids hatched in June 1975 (1 already in October 1974).

Dusona rugulosa (Förster, 1868)

9: Body length 17-18 mm. Temples strongly constricted behind the eyes, 0.85 times as long as the width of an eye, tangential lines on eyes and temples intersect on the centre of the mesoscutum. Ocellar index 0.8. Frons slightly depressed above the antennal sockets, coriaceous and with fine dense punctation dorsally and medially, almost smooth ventrally above the antennal sockets, with a distinct median longitudinal carina and with radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus densely rugose-punctate and rugose. Clypeus almost flat, densely punctate on a smooth background ventrally, its apex truncate medially. Malar space index 0.4. Genal carina low medially behind the eye, straight and distinctly raised ventrally, joining the strongly raised oral carina distant from the base of the mandible at an acute angle, the junction of both carinae not shifted downwards. Genal index 0.6. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 63-66 segments, slender, filiform, the distal 0.4 distinctly attenuated, 2nd segment 1.3 times, shortest segment 0.9 times, penultimate segment 1.5 times as long as wide. Pronotum for the greater part with punctures and longitudinal wrinkles laterally, the dorsal posterior corner with fine dense punctation on a smooth background. Mesoscutum coriaceous and with fine dense or very dense punctation. Scutellum with lateral carinae on the anterior 0.6, coriaceous and densely punctate. Prescutellar groove smooth medially, with a few longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, the prepectal carina low, the transverse carina slightly raised. Pleural part of the prepectal carina for the greater part obliter-

ated, replaced by some wrinkles medially. Pleural part of the prepectus rugose-punctate on a smooth background. Mesopleurum with dispersed rather fine punctation on a smooth background centrally, with rather dense or dense punctation anteriorly and dorsally. Speculum finely coriaceous and granulate strigose or almost smooth anteriorly, distinctly shining, with fine punctation posteriorly. Depression in front of the speculum with distinct longitudinal striae dorsally, finely granulate-strigose or almost smooth ventrally, the striae merging with the punctation anteriorly. Metapleurum with longitudinal striae dorsally, with fine punctation or fine wrinkles on a smooth background ventrally. Pleural carina complete. Areolet large, pointed, receiving the 2nd recurrent vein basad of the middle. Nervellus almost vertical, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 4.3 times as long as high. Hind tarsal claws slightly elongated, with about eight rather long pectinate teeth situated basally and medially. Propodeum extending to 0.4 of the hind coxae, broadly and shallowly depressed, with long transverse wrinkles in the sulcus and in the lateral areas, the median longitudinal carinae obliterated, the transverse carinae present and only the posterior transverse carina obliterated medially. Propodeal spiracle large, 3.7 times as long as wide. Spiracular carina narrow, partly obliterated. Spiracular index 0.2. 1st gastral tergite 3.6 times, 2nd tergite 1.6 times as long as wide. Petiole smooth laterally, glymmae very small or obliterated. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.3. Epipleurum separated from the 3rd gastral tergite anteriorly, folded in or pendent as a triangular lobe, the crease or bulge separating the epipleurum from its tergite indistinctly marked with black. Epipleural index 0.2-0.3. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.8.

Colour: Black. Maxillary palpus yellowish brown. Mandible yellow, the teeth brown. Tegula yellowish brown. Wings hyaline. Stigma light yellowish brown medially. Front leg from the apex of the coxa onwards yellowish red. Mid leg from the femur onwards yellowish red, the femur marked with black ventrally, the tarsus tinged with brown distally. Hind femur yellowish red on the distal 0.1-0.2. Hind tibia reddish brown, narrowly darkened basally. Hind tarsus light brown. 2nd gastral tergite marked with red dorsally in front of the posterior edge, almost completely red laterally, the posterior edge black, sometimes the 2nd tergite completely black dorsally and marked with dark reddish brown laterally. 3rd and 4th tergite red, with small blackish marks, or for the greater part black and with reddish brown marks laterally. Posterior tergites tinged with reddish brown laterally. Gastral sternites red or black. Pilosity whitish.

ð unknown.

Dusona sagittatae spec. nov.

Holotype (♀): "Linz, Schörgenhub, ObÖst., el. 15.7.33, Dr. L. Müller" (Austria) (Mus. Linz). – Paratypes: 2♂♂ with the same dates (1♂ with an additional label "R. v. Larentia sagittata auf Thalictrum") (1♂ Mus. Linz, 1♂ ZSM); 2♀♀, 1♂ "Parasit in L. sagittata Rp. ... em 29.5.66, leg. K. Mazzucco, Salzb." (= Salzburg/Austria) (or with similar dates) (1♀, 1♂ Coll. Schwarz, 1♀ ZSM).

9: Body length 8 mm. Temples distinctly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum. Ocellar index 0.7. Frons hardly depressed above the antennal sockets, coriaceous and densely rugose-punctate and rugose, with a fine median longitudinal carina and with distinct radial wrinkles, which extend to 0.3 the distance to the median ocellus. Antennal carina low, narrow. Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat, finely coriaceous and with fine rather dense or dense punctation ventrally, its apex straight medially. Malar space index 0.4. Genal carina low, straight subventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.35. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 38-40 segments, filiform, the distal 0.2 attenuated, 2nd segment 1.7 times, shortest segment 0.85 times, penultimate segment 1.0 times as long as wide. Pronotum coriaceous and finely and densely rugose-punctate and rugose laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, with fine and dense punctation. Scutellum with lateral carinae on the anterior 0.7, coriaceous and with dense punctation. Prescutellar groove with fine longitudinal striae. Ventral part of the prepectal carina merging with the transverse carina, both carinae slightly raised, the former not emarginate medially. Pleural part of the prepectal carina low, its dorsal 0.3 obliterated. Mesopleurum coriaceous, dull, with distinct and dense punctation. Speculum coriaceous and with fine striae. Depression in front of the speculum with distinct longitudinal striae, which merge with an area without punctation anteriorly. Metapleurum densely rugose, partly with dorsoventral wrinkles (similar to the sculpture of the lateral areas of the propodeum). Pleural carina complete. Areolet large, stalked, receiving the 2nd recurrent vein at the middle. Nervellus slightly inclivous, distinctly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 4.4 times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, distinctly and rather broadly depressed, for the greater part with irregular wrinkles, in the midline with some short transverse wrinkles medially and posteriorly, costulae distinct, median longitudinal carinae indicated posteriorly. Propodeal spiracle rather small, 3.4 times as long as wide. Spiracular carina narrow. Spiracular index 0.7. 1st gastral tergite 3.5 times, 2nd tergite 1.4 times as long as wide. Petiole without glymmae, with indistinct narrow lateral areas, smooth in front of these areas. Petiolar suture almost completely obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellowish brown. Mandible yellowish brown dorsally and medially. Tegula yellowish brown. Wings hyaline. Stigma yellowish brown medially. Front leg from the femur onwards yellowish red, the femur slightly darkened ventrally (sometimes the apex of the front coxa and the front trochanter and trochantellus vellowish too). Mid femur black proximally, light red distally. Mid tibia whitish yellow. Mid tarsus yellowish brown. Hind tibia broadly whitish yellow medially, narrowly black basally, rather broadly black apically, the apical ring extending over 0.2 of the tibia. Hind tarsus blackish, the basitarsus narrowly yellowish basally. 2nd gastral tergite red on the posterior 0.2. 3rd tergite light red. 4th tergite light red on the anterior half. Pilosity whitish.

 δ : Flagellum with 37-41 segments. Mandible broadly yellowish red medially. Front leg yellowish red from the apex of the coxa onwards. Mid femur black basally and ventrally, yellowish red dorsally and apically. In other characters similar to the \circ .

Host: Perizoma sagittatum (Fabricius) (Geometridae).

Dusona semiflava (Costa, 1883)

2: Body length 9-10 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect

on the centre of the mesoscutum. Ocellar index 0.7. Frons scarcely depressed above the antennal sockets, coriaceous and finely and densely rugose-punctate, with a distinct median longitudinal carina and with distinct radial wrinkles, which extend to 0.4 the distance to the median ocellus. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus densely rugose-punctate and rugose. Clypeus flat dorsally and medially, slightly concave ventrally, very finely coriaceous and with rather dense punctation ventrally, its apex truncate or slightly rounded medially. Malar space index 0.35. Genal carina low medially behind the eye, strongly raised subventrally and ventrally, joining the strongly raised oral carina distant from the base of the mandible, both carinae forming a lopsided triangular pyramid, the tip of the latter pointed, shifted downwards and situated behind the base of the mandible. Gena deeply depressed close to the junction, the depression 0.8-1.0 times as deep as the width of the 2nd flagellar segment. Genal index 0.5. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 44-48 segments, filiform, the distal 0.3 attenuated, 2nd segment 1.5 times, shortest segment 0.8-0.9 times, penultimate segment 1.1-1.2 times as long as wide. Pronotum densely rugose-punctate and rugose on a smooth background laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous, rather finely and very densely punctate or rugose-punctate. Scutellum with lateral carinae on the anterior 0.3, densely rugose-punctate. Prescutellar groove smooth medially, with some longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former almost not emarginate medially. Pleural part of the prepectal carina complete, rarely covered with wrinkles on the dorsal 0.1. Mesopleurum distinctly and densely punctate on a smooth background. Speculum coriaceous or granulate-strigose dorsally, smooth medially and ventrally, shining. Depression in front of the speculum with strong longitudinal striae, which merge with the punctation anteriorly. Metapleurum strongly and densely punctate on a smooth background ventrally, with longitudinal striae dorsally. Pleural carina distinct. Areolet large, pointed, receiving the 2nd recurrent vein at or slightly distad of the middle. Nervellus slightly inclivous, weakly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 5.7 times as long as high. Hind tarsal claws short, with about five distinct pectinate teeth situated basally and medially. Propodeum extending to 0.6 of the hind coxae, rather narrowly depressed, almost completely covered with strong transverse wrinkles, the longitudinal carinae lacking, the transverse carinae partly indicated. Propodeal spiracle large, 3.4 times as long as wide. Spiracular carina widened. Spiracular index 0.3. 1st gastral tergite 5.1 times, 2nd tergite 2.2 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5-2.0. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster strongly compressed posteriorly. Ovipositor sheaths slightly widened medially and subapically (in lateral view). Ovipositor index 0.6.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Scape usually broadly marked with yellow ventrally, sometimes black. Tegula yellow. Wings hyaline. Stigma medium brown medially. Front leg yellow from the apex of the coxa onwards. Mid leg yellow from the trochanter or the femur onwards, the tarsus light brown distally. Hind tibia yellow. Hind basitarsus narrowly marked with yellow basally. 2nd gastral tergite dorsally yellow on the posterior 0.1-0.2, laterally yellow on the posterior 0.7, the ventrolateral edges partly marked with black. 3rd tergite yellow. 4th tergite yellow anteriorly, black posteriorly. Pilosity whitish.

 δ : Scape usually broadly marked with yellow ventrally, sometimes only with a small yellow spot. Front and mid legs for the greater part yellow, only the base of the coxae black. In other characters similar to the \mathfrak{P} .

Dusona stenogaster (Förster, 1868)

2: Body length 9-11 mm. Temples strongly constricted behind the eyes, 0.7 times as long as the width of an eye, tangential lines on eyes and temples intersect on the posterior 0.8 of the mesoscutum. Ocellar index 0.8. Frons scarcely depressed above the antennal sockets, with fine dispersed punctation, distinctly coriaceous and dull dorsally, finely coriaceous and shining ventrally, with a distinct median longitudinal carina, without radial wrinkles or with a few radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.5 of the clypeus coriaceous and densely rugose-punctate and rugose. Clypeus flat dorsally, slightly concave ventrally, finely coriaceous and with fine rather dense or dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.35. Genal carina low ventrally, distinctly bent forwards and sidewards subventrally, joining the oral carina at the base of the mandible at a right angle. Temple coriaceous and with very fine dispersed punctation medially. Flagellum with 33-34 segments, slightly clavate, 2nd segment 2.0 times, shortest segment 0.7 times, penultimate segment 0.8 times as long punctate laterally on the dorsal 0.4, with longitudinal wrinkles ventrally. Mesoscutum coriaceous, dull, densely rugose-punctate and rugose. Scutellum with lateral carinae on the anterior 0.4, finely coriaceous and with fine dispersed or rather dense punctation. Prescutellar groove smooth medially, with fine longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former deeply emarginate medially. Pleural part of the prepectal carina present medially, obliterated ventrally and dorsally. Mesopleurum finely coriaceous, somewhat shining, with rather fine and rather dense punctation. Speculum finely coriaceous, shining. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum rugose anteriorly, finely coriaceous and with fine dense punctation posteriorly. Pleural carina distinct. Areolet rather small, short-stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus inclivous, weakly intercepted. Nervellar index 0.8. Tibial spur index 0.6. Hind femur 4.3 times as long as high. Hind tarsal claws short, with about four distinct pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, rather narrowly and rather deeply depressed, the longitudinal furrow coriaceous and with irregular wrinkles anteriorly, with short transverse wrinkles medially, with long transverse wrinkles posteriorly, the median longitudinal carinae obliterated. Propodeal spiracle small, 2.7 times as long as wide. Spiracular carina widened. Spiracular index 0.3. 1st gastral tergite 3.5 times, 2nd tergite 2.4 times as long as wide. Petiole with narrow lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture partly indicated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 2.5. Epipleurum not separated from the 3rd gastral tergite, the latter with indistinct black lateral stripes well above the ventrolateral edge (the stripes sometimes obliterated). Gaster strongly compressed posteriorly. Ovipositor sheaths strongly widened medially and subapically (in lateral view). Ovipositor remarkably stout basally, about 2.5 times as high as the apical width of a hind basitarsus. Ovipositor index 0.9. Colour: Black. Maxillary palpus yellowish

as wide. Pronotum coriaceous and densely rugose-

Colour: Black. Maxillary palpus yellowish brown. Mandible marked with yellow dorsally and medially. Tegula yellow. Wings hyaline. Stigma yellowish brown medially. Front leg from the trochanter onwards yellowish red, the tarsus dark brown apically, sometimes the coxa marked with yellowish red apically. Mid leg from the femur onwards yellowish red, the tarsus dark brown apically. Hind tibia broadly marked with yellowish medially on the dorsal side, narrowly black basally, marked with

black ventrally, narrowly black apically, the apical ring extending over 0.2 of the tibia. Hind tarsus blackish, the basitarsus narrowly marked with yellowish brown basally. 2nd gastral tergite dorsally red on the posterior 0.2, laterally red on the posterior 0.4, sometimes almost completely black and marked with red only laterally. 3rd tergite red, marked with black dorsally (to a variable extent). Pilosity whitish.

 δ : Flagellum filiform, 2^{nd} segment 2.0 times, shortest segment 1.0 times, penultimate segment 1.0 times as long as wide. Propodeum narrowly and deeply depressed, rather coarsely wrinkled, the median longitudinal carinae often indicated medially and posteriorly. Front and mid legs from the apex of the coxa onwards yellowish red, the tarsi brown apically. Hind tibia broadly yellowish medially (also on the ventral side), narrowly marked with black basally and apically, the apical ring extending over 0.15-0.2 of the tibia. In other characters similar to the φ .

Dusona tenerifae Hinz, 1977

9: Body length 7-8 mm. Temples distinctly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the scutellum. Ocellar index 0.5-0.6. Frons not depressed, coriaceous and finely punctate and rugose-punctate, with a fine median longitudinal carina (sometimes indistinct) and with short radial wrinkles. Antennal carina low and narrow. Face and dorsal 0.7 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, finely coriaceous and with dispersed or rather dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina low ventrally, slightly bent forwards and sidewards subventrally, joining the oral carina at the base of the mandible at an acute angle. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 30 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 2.0 times, shortest segment 1.0 times, penultimate segment 1.0 times as long as wide. Pronotum coriaceous and densely rugose-punctate laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaecous and densely rugose-punctate. Scutellum with lateral carinae on the anterior 0.3, coriaceous and with dense punctation. Prescutellar groove smooth. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former not emarginate medially. Pleural part of the prepectal carina replaced by several wrinkles. Pleural part of the prepectus rugose. Mesopleurum coriaceous and with rather dense or dense punctation. Speculum and depression in front of the speculum with long longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose. Pleural carina complete. Areolet rather large, short-stalked, receiving the 2nd recurrent vein at the middle. Nervellus inclivous, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.4 times as long as high. Hind tarsal claws short, with about five short pectinate teeth situated basally. Propodeum extending to 0.5 of the hind coxae, narrowly and shallowly depressed, for the greater part with irregular wrinkles, with short transverse wrinkles medially and posteriorly in the shallow longitudinal furrow, median longitudinal carinae indistinct or distinct. Propodeal spiracle small, 1.8 times as long as wide. Spiracular carina narrow. Spiracular index 0.9. 1st gastral tergite 5.8 times, 2nd tergite 2.2 times as long as wide. Petiole with indistinct lateral areas, without glymmae, smooth in front of the lateral areas. Petiolar suture obliterated. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter without black lateral stripes. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.7.

Colour: Black. Maxillary palpus yellowish brown. Mandible narrowly yellowish red medially. Tegula yellow. Wings hyaline. Stigma dark brown medially. Front leg from the apex of the coxa or from the trochanter onwards yellow. Mid leg from the trochanter onwards yellow, the tarsus dark brown apically. Hind tibia yellow. Hind tarsus dark brown, the basitarsus narrowly marked with yellow basally. Postpetiole sometimes marked with yellow posteriorly. 2nd to 4th gastral tergites yellow, the 2nd tergite with or without a small black dorsal mark anteriorly, sometimes the 4th tergite marked with black posteriorly. Pilosity whitish.

δ: Flagellum with 31-32 segments. Longitudinal furrow of the propodeum somewhat deeper, with the transverse wrinkles somewhat longer. In other characters similar to the ♀.

Variety: see taxonomic notes.

Dusona tricolorator Aubert, 1970

9: Body length 7-9 mm. Temples slightly constricted behind the eyes, 0.8 times as long as the width of an eye, tangential lines on eyes and temples intersect on the petiole. Ocellar index 1.2. Frons not depressed, coriaceous and densely rugose punctate dorsally and medially, almost smooth ventrally above the antennal sockets, with a fine median longitudinal carina and with short radial wrinkles. Antennal

carina low and narrow. Face coriaceous and densely rugose-punctate. Clypeus flat, densely punctate on an almost smooth background, its apex slightly rounded medially. Malar space index 0.4. Genal carina low and almost straight ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.2. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 36-37 segments, slightly fusiform, 2nd segment 1.7 times, shortest segment 0.9 times, penultimate segment 1.0 times as long as wide. Pronotum distinctly and densely rugose-punctate laterally on the dorsal 0.6, with longitudinal wrinkles ventrally. Mesoscutum finely coriaceous, shining, distinctly and very densely punctate. Scutellum with lateral carinae on the anterior 0.8, densely punctate on a smooth background. Prescutellar groove smooth medially, with longitudinal striae laterally. Prepectal carina low, the ventral part merging with the pleural part, the dorsal 0.3 of the latter covered with wrinkles. Transverse carina indistinct. Mesopleurum very densely punctate on a smooth background. Speculum with fine striae anteriorly, rather dull, punctate posteriorly. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum with very dense punctation on a smooth background. Pleural carina distinct. Areolet rather small, stalked, receiving the 2nd recurrent vein distad of the middle. Nervellus almost vertical, distinctly intercepted. Nervellar index 0.7. Tibial spur index 0.6. Hind femur 5.0 times as long as high. Hind tarsal claws short, with some short pectinate teeth situated basally. Propodeum extending to 0.6 of the hind coxae, narrowly and deeply depressed, with short transverse wrinkles in the longitudinal furrow, with irregular wrinkles beside the furrow, median longitudinal carinae obliterated. Propodeal spiracle rather small, 2.8 times as long as wide. Spiracular carina narrow medially. Spiracular index 0.7. 1st gastral tergite 4.4 times, 2nd tergite 2.2 times as long as wide. Petiole with distinct rather small glymmae, smooth in front of the glymmae. Petiolar suture indicated anteriorly, obliterated posteriorly. 1st gastral sternite smooth. Thyridium oval. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly. Ovipositor sheaths slightly widened subapically (in lateral view). Ovipositor index 0.5.

Colour: Black. Maxillary palpus yellow. Mandible narrowly marked with yellowish brown or reddish brown dorsally and medially. Wings hyaline. Stigma yellowish brown medially. Front femur yellowish red, sometimes marked with black basally and ventrally. Front tibia and tarsus yellow. Mid femur

yellow on the distal 0.2. Mid tibia and tarsus yellow, the tarsus marked with brown distally. Hind tibia marked with whitish yellow basally and medially on the dorsal side, black ventrally and apically, the black apical ring extending over 0.2-0.3 of the tibia. Hind tarsus brown, the basitarsus yellowish brown on the proximal half. 2nd gastral tergite red on the posterior 0.1-0.2. 3rd and 4th tergites red, sometimes with dorsal black marks. Pilosity whitish.

 δ : Hind tibia whitish yellow basally and medially (also on the ventral side), black apically. 3^{rd} and 4^{th} gastral tergites black dorsally. In other characters similar to the \circ .

Dusona turcator spec. nov.

Holotype (♂): "TK – 40 km. E. Urfa. 14-V-1975. leg. Kl. Warncke" (Turkey) (ZSM).

♂: Body length 9 mm. Temples slightly constricted behind the eyes, 0.85 times as long as the width of an eye, tangential lines on eyes and temples intersect on the postpetiole. Ocellar index 0.85. From slightly depressed above the antennal sockets, densely rugosepunctate, with a distinct median longitudinal carina which is slightly raised medially, and with distinct radial wrinkles, which extend to 0.4 the distance to the median ocellus. Antennal carina rather low and narrow. Face and dorsal 0.7 of the clypeus densely rugose-punctate. Clypeus flat, densely punctate on a finely coriaceous background ventrally, its apex slightly rounded medially. Malar space index 0.4. Genal carina low and slightly sinuous ventrally, joining the slightly raised oral carina distant from the base of the mandible at an acute angle. Genal index 0.2. Temple with fine dispersed punctation on a finely coriaceous background medially. Flagellum with 43 segments, slender, filiform, the distal 0.2 attenuated, 2nd segment 1.7 times, shortest segment 1.0 times, penultimate segment 1.3 times as long as wide. Pronotum densely punctate on a smooth background laterally on the dorsal 0.7, rugose-punctate ventrally. Mesoscutum with dense or very dense punctation on a slightly coriaceous and shining background. Scutellum with lateral carinae on the anterior 0.6, with distinct and dense punctation on a smooth background. Prescutellar groove smooth. Prepectal carina low, ventral part merging with the pleural part, the dorsal 0.3 of the latter obliterated. Transverse carina obliterated. Mesopleurum with dense or very dense punctation on a smooth background. Speculum with fine striae anteriorly, with dense punctation posteriorly. Depression in front of the speculum with distinct longitudinal striae, which merge with the punctation anteriorly. Metapleurum with distinct and dense punctation on a smooth background. Pleural carina distinct. Areolet rather large, pointed, receiving the 2nd recurrent vein slightly basad of the middle. Nervellus slightly reclivous, weakly intercepted. Nervellar index 0.6. Tibial spur index 0.6. Hind femur 3.7 times as long as high. Hind tarsal claws rather short, with about four distinct pectinate teeth situated basally and medially. Propodeum extending to 0.3 of the hind coxae, with a distinct narrow and rather shallow longitudinal furrow, with short transverse striae in the furrow anteriorly and medially, with irregular wrinkles beside the furrow, with long transverse striae posteriorly, the longitudinal carinae obliterated. Propodeal spiracle large, 2.9 times as long as wide. Spiracular carina obliterated. Spiracular index 0.4. 1st gastral tergite 3.7 times, 2nd tergite 1.9 times as long as wide. Petiole smooth laterally, without glymmae, with weakly indicated narrow lateral areas. Petiolar suture present anteriorly, obliterated medially and posteriorly. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.5. Epipleurum not separated from the 3rd gastral tergite, the latter with black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow. Wings hyaline. Stigma medium brown medially. Front leg from the trochanter onwards yellow. Mid femur yellowish red. Mid tibia and mid tarsus yellow, the tarsal segments marked with light brown apically. Hind tibia broadly whitish medially, very narrowly black basally, rather narrowly black apically, the apical ring extending over 2.0 of the tibia. Hind tarsal segments whitish basally, light brown apically. 2nd gastral tergite dorsally red on the posterior 0.7, completely red laterally. 3rd tergite red. 4th tergite red on the anterior 0.4. Pilosity whitish.

♀ unknown.

Dusona tyrolensis spec. nov.

Holotype (♂): "Italia (Bolzano), Brixen 2000, 23.VI. 1976, C. J. Zwakhals" (ZSM).

♂: Body length 9 mm. Temples slightly constricted behind the eyes, 0.9 times as long as the width of an eye, tangential lines on eyes and temples intersect on the propodeum. Ocellar index 1.0. Frons hardly depressed above the antennal sockets, coriaceous and densely rugose dorsally, smooth ventrally, with long radial wrinkles subventrally, which extend to half the distance to the median ocellus, with a fine median longitudinal carina. Antennal carina low and narrow. Face and dorsal 0.3 of the clypeus coriaceous and densely rugose-punctate. Clypeus flat, finely coriaceous and with distinct dense punctation ventrally, its apex slightly rounded medially. Malar space index 0.5. Genal carina low, slightly bent sidewards and forwards subventrally, joining the oral carina close to the base of the mandible at a right angle. Genal index 0.1. Temple finely coriaceous and with fine dispersed punctation medially. Flagellum with 37 segments, filiform, the distal 0.2 slightly attenuated, 2nd segment 1.9 times, shortest segment 1.0 times, penultimate segment 1.15 times as long as wide. Pronotum densely rugose-punctate and rugose laterally on the dorsal 0.2, with longitudinal wrinkles ventrally. Mesoscutum coriaceous and densely punctate. Scutellum with lateral carinae on the anterior 0.7, coriaceous and densely rugosepunctate. Prescutellar groove smooth medially, with longitudinal striae laterally. Ventral part of the prepectal carina merging with the transverse carina, both carinae strongly raised, the former deeply emarginate medially. Pleural part of the prepectal carina low ventrally and medially, the dorsal 0.4 covered with wrinkles and obliterated. Mesopleurum with distinct dense punctation on a finely coriaceous and shining background. Speculum for the greater part finely granulate-strigose, with a small smooth area ventrally. Depression in front of the speculum with rather fine long longitudinal striae, which merge with the punctation anteriorly. Metapleurum coriaceous and densely rugose-punctate and rugose. Pleural carina distinct. Areolet rather large, pointed, receiving the 2nd recurrent vein distad of the middle. Nervellus slightly reclivous, very weakly intercepted. Nervellar index 0.7. Tibial spur index 0.7. Hind femur 5.0 times as long as high. Hind tarsal claws short, with about three short pectinate teeth situated basally. Propodeum extending to 0.4 of the hind coxae, slightly depressed, for the greater part with irregular wrinkles, with few transverse wrinkles medially, with short longitudinal wrinkles posteriorly, almost completely areolated, the area superomedia indicated, but the median longitudinal carinae partly indistinct and covered with wrinkles. Propodeal spiracle rather small, 2.9 times as long as wide. Spiracular carina fine, narrow. Spiracular index 0.7. 1st gastral tergite 4.2 times, 2nd tergite 2.3 times as long as wide. Petiole smooth laterally, without glymmae or lateral areas. Petiolar suture indicated. 1st gastral sternite smooth. Thyridium elongated. Thyridial index 1.0. Epipleurum not separated from the 3rd gastral tergite, the latter with distinct black lateral stripes well above the ventrolateral edge. Gaster distinctly compressed posteriorly.

Colour: Black. Maxillary palpus yellow. Mandible broadly yellow medially. Tegula yellow, with a brown mark apically. Wings hyaline. Stigma yel-

lowish brown medially. Front trochanter and front trochantellus yellowish, with brown marks. Front leg yellowish red from the femur onwards. Mid femur black basally and ventrally, reddish brown dorsally and apically. Mid tibia and mid tarsus yellowish red, the tarsus marked with brown distally. Hind tibia yellowish medially, narrowly black basally, broadly black apically, the apical ring extending over 0.3 of the tibia. Hind tarsus brownish black, the basitarsus narrowly yellowish basally. $2^{\rm nd}$ gastral tergite dark red on the posterior 0.3. $3^{\rm rd}$ and $4^{\rm th}$ tergites dark red. Pilosity whitish.

9 unknown.

References

Aubert, J.-F. 1981. Syllogismes, illogismes et innovations chez les Ichneumonides. Bulletin de la Société Entomologique de Mulhouse 1981: 17-22.

Bachmaier, F. 1979. Das in der Zoologischen Staatssammlung München aufbewahrte Typenmaterial der Gattung *Dusona* Cameron 1900 (= *Campoplex* auct.) und sein Erhaltungszustand (Hymenoptera, Ichneumonidae, Porizontinae). Spixiana 2: 69-93.

Constantineanu, M. I. 1932. Cas d'anomalie chez quelques espèces d'Ichneumonides. Archives de zoologie expérimentale et générale 74: 69-79.

Dalla Torre, C. G. de 1901. Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. III: Trigonalidae, Megalyridae, Stephanidae, Ichneumonidae, Agriotypidae, Evaniidae, Pelecinidae. Lipsiae, pp. 1-544.

Fitton, M. G. 1982. A catalogue and reclassification of the Ichneumonidae (Hymenoptera) described by C. G. Thomson. Bulletin of the British Museum (Natural History) Entomology 45(1): 1-119.

Hinz, R. 1957. Zur Systematik und Ökologie der Ichneumoniden I (Hym.). Deutsche entomologische Zeitschrift, Neue Folge 4: 86-90.

- 1961. Zur Systematik und Ökologie der Ichneumoniden II (Hym.). Deutsche entomologische Zeitschrift, Neue Folge 8: 250-257.
- 1962. Die Försterschen Typen der Gattung *Dusona* Cam. (*Campoplex* auct.) (Hym.) Opuscula zoologica 66: 1-12.
- 1963a. Zur Systematik und Ökologie der Ichneumoniden III (Hym.). Deutsche entomologische Zeitschrift, Neue Folge 10: 116-121.
- 1963b. Über einige Typen der Gattung *Dusona* Cameron (Hymenoptera: Ichneumonidae). Beiträge zur Entomologie 13: 335-344.
- 1972. Zur Systematik und Ökologie der Ichneumoniden IV (Hym.). Deutsche entomologische Zeitschrift, Neue Folge 19: 45-54.
- 1973. Zwei Bemerkungen zur Systematik der Gattung Dusona Cameron (Campoplex auct.) (Hym., Ichneumonidae). Nachrichtenblatt der bayerischen Entomologen 22: 29-30.

- 1975. Vier Bemerkungen zur Systematik der Ichneumonidae (Hym.). Nachrichtenblatt der bayerischen Entomologen 24: 65-68.
- 1976. Zur Systematik und Ökologie der Ichneumoniden V (Hymenoptera). Deutsche entomologische Zeitschrift, Neue Folge 23: 99-105.
- 1977. Über einige Arten der Gattung *Dusona* Cameron (Hymenoptera, Ichneumonidae). Nachrichtenblatt der bayerischen Entomologen 26: 47-54.
- 1979. Beiträge zur Taxonomie der ostpalaearktischen Arten der Gattung *Dusona* Cameron (Hymenoptera, Ichneumonidae). Deutsche entomologische Zeitschrift, Neue Folge 26: 215-227.
- 1985. Neue sibirische *Dusona*-Arten (Hymenoptera, Ichneumonidae). Deutsche entomologische Zeitschrift, Neue Folge 32: 297-317.
- 1990. Beiträge zur Taxonomie der Gattung *Delopia* Cameron, 1903 (*Dusona* auct.) (Hymenoptera, Ichneumonidae). Entomofauna 11: 261-271.
- Hinz, R. & K. Horstmann 2004. Revision of the eastern Palearctic species of *Dusona* Cameron (Insecta, Hymenoptera, Ichneumonidae, Campopleginae). Spixiana, Supplement 29: 1-183.
- Holmgren, A. E. 1872. Om de scandinaviska arterna af ophionidslägtet *Campoplex*. Bihang till Kungliga Svenska Vetenskapsakademiens handlingar 1(2): 89 pp.
- Horstmann, K. 1980. Über die Campopleginae der Makaronesischen Inseln (Hymenoptera, Ichneumonidae). Spixiana 3: 121-136.
- Morley, C. 1915. Ichneumonologia Britannica, V. The Ichneumons of Great Britain. Ophioninae. London, X & 400 pp.

- Oehlke, J. 1963. Revision der im Deutschen Entomologischen Institut aufbewahrten Typen paläarktischer Ichneumoniden (Hymenoptera: Ichneumonidae). Beiträge zur Entomologie 13: 403-410.
- Rey, G. & I. Izquierdo. 1985. Sobre las especias canarias de *Dusona* Cameron, 1900 (Hym., Ichn., Porizontinae). Boletin Asociacion Espanola Entomologia 9: 317-320.
- Scaramozzino, P. L. 1994. Nota preliminare sui tipi delle specie di Ichneumonidae, Evaniidae e Gasteruptionidae descritte da Achille Costa. Notiziario Imenotterologico Italiano 5: 17-19.
- Schmiedeknecht, O. 1908. Opuscula Ichneumonologica. IV. Band. Ophioninae. Fasc. XX. Blankenburg in Thüringen, pp. 1521-1600.
- Shaumar, N. 1966. Les Ichneumonides d'Égypte. Entomophaga 11: 441-469.
- Strobl, G. 1904. Ichneumoniden Steiermarks (und der Nachbarländer). V. Fam. Ophionidae. Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark 40 (1903): 43-111.
- Teunissen, H. G. M. 1947. Het genus *Campoplex* (Subfam. Ophioninae. Fam. Ichneumonidae). Tijdschrift voor Entomologie 88 (1945): 249-270.
- Thomson, C. G. 1887. Försök till uppställning och beskrifning af arterna inom slägtet *Campoplex* (Grav.). Opuscula entomologica (Lund), Fasc. XI: pp. 1043-1182.
- Townes, H. 1969. The genera of Ichneumonidae, part. 1. Memoirs of the American Entomological Institute 11: II & 307 pp.
- Yu, D. & K. Horstmann. 1997. A catalogue of World Ichneumonidae (Hymenoptera). Memoirs of the American Entomological Institute 58 (1-2): VI & 1558 pp.