SPIXIANA	40	1	7-12	München, August 2017	ISSN 0341-8391
----------	----	---	------	----------------------	----------------

# New records of the scaled squid, *Lepidoteuthis grimaldii* Joubin, 1895 in the Canary Islands, Eastern Atlantic Ocean

(Cephalopoda, Oegopsida, Lepidoteuthidae)

## Alejandro Escánez, Ángel Guerra, Francisco Rocha & Fernando Lozano-Soldevilla

Escánez, A., Guerra, Á., Rocha, F. & Lozano-Soldevilla, F. 2017. New records of the scaled squid, *Lepidoteuthis grimaldii* Joubin, 1895 in the Canary Islands, Eastern Atlantic Ocean (Cephalopoda, Oegopsida, Lepidoteuthidae). Spixiana 40(1): 7–12.

Two well conserved female specimens of the uncommon soft-scaled squid *Lepidoteuthis grimaldii* Joubin, 1895 (Lepidoteuthidae, Oegopsida) were caught in Canary Island waters: a maturing female of 580 mm dorsal mantle length found dead on the surface at 27°38'N, 18°01'W (near El Hierro Island); an immature female of 350 mm dorsal mantle length captured by commercial open midwater trawl at 28°48'N, 16°00'W (north of Tenerife Island), between 342 and 487 m depth. This is the first report of an adult of this species from the Canary Islands. Morphological data are presented and the bathymetric distribution of this species in Canary waters is discussed.

Alejandro Escánez & Francisco Rocha, BioCephALab Cephalopod Biology and Aquaculture Laboratory, Departamento de Ecoloxía e Bioloxía Animal, Edificio de Ciencias Experimentais, Campus As Lagoas-Marcosende, Universidade de Vigo, 36310 Vigo, Spain; e-mail: aescanez@msn.com

Ángel Guerra, ECOBIOMAR, Grupo de Ecología y Biodiversidad Marina, Instituto de Investigaciones Marinas de Vigo-CSIC, C/Eduardo Cabello, 6, 36208 Vigo, Spain

Fernando Lozano-Soldevilla, Departamento de Biología Animal y Edafología y Geología, Facultad de Biología, Universidad de La Laguna, Avenida Astrofísico Francisco Sánchez S/N, Campus Universitario de Anchieta, 38206 San Cristobal de La Laguna, Spain

#### Introduction

Our knowledge on the biodiversity, biology and distribution of oceanic cephalopod species is unequal. Pelagic species with high fishery potential, such as Ommastrephidae, have been intensively studied (Hoving 2008, Jereb & Roper 2010) in contrast to those with low commercial value. For different reasons (small mouth areas of midwater nets, trawl speeds, evasion behaviour, etc.), large oceanic and deep-sea cephalopods are rarely caught using direct sampling techniques at sea (Clarke 1977, 2006). However, there are several effective techniques for paralarvae or juvenile stages of large species and adults of small species (e.g. Moreno et al. 2009).

At present, one of these poorly known species is the scaled squid, *Lepidoteuthis grimaldii* Joubin, 1895. This large squid reaches sizes of up to 100 cm dorsal mantle length (DML) and is characterised by its distinctive hexagonal shaped dermal cushion, formerly called "scaled", that covers the mantle except for the posteroventral part under the posterior half of the fins. Adults are also characterised by the loss of tentacles (Nesis & Nikitina 1990, Jereb & Roper 2010). *Lepidoteuthis grimaldii* is a sexually dimorphic species in which mature females attain twice the size



**Fig. 1.** The two *Lepidoteuthis grimaldii* collected in the Canary Islands. **A.** Ventral view of the female caught in El Hierro Island. **B.** Dorsal view of the female captured on the north side of Tenerife Island. Scale bars = 20 cm.

of mature males. Mature males lack a hectocotylus and have a relatively long terminal organ (penis), which projects freely for a considerable portion of the organ's length (Jackson & O'Shea 2009).

Lepidoteuthis grimaldii inhabits tropical and subtropical oceans with disjunct records in the Pacific Ocean (Hawaii, Japan, New Caledonia, New Zealand, Australia, and Tasmanian Sea) and a broad distribution in the Atlantic Ocean (Clarke 1962, Clarke 2006, Jereb & Roper 2010). The vertical distribution of this species is unclear. Paralarvae have been fished as shallow as 100 m at night while young and subadult stages have been collected at 700 m day and night. Until now, only three large specimens have been fished, all of them in bottom trawls between 812–1100 m depth suggesting that *L. grimaldii* is a demersal species (Jackson & O'Shea 2009, Jereb & Roper 2010, Young & Vecchione 2016). In contrast, scaled squid has been commonly reported in predator stomach contents, such as sperm whales, *Physeter macrocephalus* Linnaeus, 1758, and as young stages in plankton hauls (see Table 2).



**Fig. 2. A.** *L. grimaldii* female in stage III of maturation, collected in El Hierro (580 mm DML). **B.** Female in stage II of maturation captured on the north side of Tenerife Island (330 mm DML). 1 = ovary; 2 = nidamental gland.



**Fig. 3.** *Lepidoteuthis grimaldii* beak aspect. **A.** Upper and lower beak of maturing female (580 mm DML). **B.** Upper and lower beak of immature female (330 mm DML). Scale bars = 0.5 cm.

Clarke (1964) reported for the first time the capture of two young individuals in the "Canary Islands region". Other records of this species in the archipelago are beaks found in stomach contents of stranded cetaceans (Fernández et al. 2009, Hernández-García & Martín 1994) and slender alfonsino, *Beryx splendens* Lowe, 1834 (Dürr & González 2002), captured in the area.

The captures of two large scaled squid in the Canary Islands, which we present in this communication, represent the first record of adult specimen in the region. These individuals allow us to report additional morphometric and biological data from this uncommon deep-sea squid species.

#### Methods

Two specimens of the scaled squid were captured in Canary Archipelago waters. Specimen 1 was found dead in El Hierro Island (Mar de Las Calmas) surface waters at 27°38'N, 18°01'W, during cetacean photoidentification surveys on 25 May of 2013. Specimen 2 was caught on 20 April 2012 by RV Cornide de Saavedra during the "CETOBAPH" mesopelagic survey, at 28°48' N, 16°00' W (north of Tenerife) at a fishing depth between 342 to 487 m over a seafloor of 2000 m. The fishing gear employed was a commercial open midwater trawl with a modified cod-end (10 mm mesh size). Both specimens were frozen on board. Identification was based on the diagnostic characters given by Nesis & Nikitina (1990) and Jereb & Roper (2010). After defrosting specimens at room temperature, sex was identified and measurements were taken following Roper & Voss (1983). Radula and beaks from each specimen were removed and measured following Clarke (1962). Beak pigmentation stage was determined using the scale of Fang et al. (2016). Maturity was assessed according to Lipinski (1979).

#### **Results and discussion**

Morphometric characters of the body and beak, weight, sex and maturity stage of both specimens are presented in Table I.

Specimen 1 (Fig. 1A) was a 580 mm DML, stage III female (Fig. 2A). Its enlarged nidamental glands (180 mm length) covered some internal organs; no mature oocytes were observed in the oviducts, which were still not very swollen. Immature oocytes were observed in the ovary (Fig. 2A). Specimen 2 was a 350 mm DML stage II (immature) female (Fig. 2B). Internal organs of both specimens were in very good condition.

The radula is formed by seven rows of teeth, with a tricuspid rachidian tooth; small lateral cusps.

First lateral tooth externally bicuspid; small lateral cusp. Second lateral and marginal row with conical teeth, slightly curved towards the middle row, in agreement with Clarke & Maul (1962).

Beaks of specimen 1 and 2 were at pigmentation stages 5 and 3 respectively (Fang et al. 2016). In comparison, beaks of female *Ommastrephes bartramii* (Lesueur, 1821) between 330–430 mm DML were at pigmentation stages 3 and 4, while adults over 400 mm DML were at pigmentation stages 6 and 7 (Fang et al. 2016). The dissimilarity in pigmentation pattern may be due to the different maximum sizes

**Table 1.** Measurements of *Lepidoteuthis grimaldii* specimens caught in the Canary Islands between 2011 and 2013. All measurements in mm.

Measurements	Specimen 1 El Hierro 25/05/2013	Specimen 2 Tenerife 20/04/2012
Dorsal mantle length	580	330
Ventral mantle length	545	310
Mantle width	183	110
Fin length	225	158
Fin width	215	150
Head length	100	95
Head width	60	45
Funnel length	90	45
Funnel width	80	35
Funnel cartilage length	50	27.58
Right Arms		
Arms length I	355	sectioned
Arms length II	320	270
Arms length III	300	130
Arms length IV	sectioned	sectioned
Left Arms		
Arms length I	315	sectioned
Arms length II	355	sectioned
Arms length III	335	sectioned
Arms length IV	280	293
Weight (g)	2700	545.89
Upper beak		
Upper hood length	32.45	19.96
Upper rostrum length	15.75	9.35
Upper crest length	58.80	30.96
Lower beak		
Lower hood length	7.50	5.92
Lower rostrum length	15.20	8.48
Lower crest length	19.40	13.10
Lower wing length	12.00	8.91
Sex	Female	Female
Maturity stage	III	II

Method	Location	n	Sex	DML	Depth (m)	Reference
				(mm)		
Bottom trawl	Tasmania, Cascade Rise	1	male	442	950-1100	Jackson & O'Shea 2009
Bottom trawl	New Zealand, Chatham Rise		female	617	812	Young & Vecchione 2016
Bottom trawl	New Zealand	1	immature	270	-	Jackson & O'Shea 2009
Bottom trawl	New Zealand		-	> 500	-	Jackson & O'Shea 2009
Plankton trawl	Gulf of Guinea	1	young stages	29.4-39.1	50-23	Lu & Clarke 1975
Plankton trawl	Azores Island	1	young stages	-	-	Clarke 2006
Plankton trawl	Spain, Galician Bank	4	young stages	-	-	Clarke 2006
Plankton trawl	Spain, Canary Region	2	young stages	60-85	-	Clarke 1964
Plankton trawl	Madeira island	1	young stages	-	-	Clarke 2006
Plankton trawl	Abyssal plain near Madeira	1	young stages	-	-	Clarke 2006
Stomach content	Azores Island	3	2 females	720-860	-	Joubin 1895, 1900
			+ 1 ind.			
Stomach content	Madeira Island	9	4 females	155-970	-	Clarke 1960,
			+ 5 ind.			Clarke & Maul 1962
Stomach content	South Atlantic	1	female	510	-	Merret 1963

**Table 2.** Records of *Lepidoteuthis grimaldii* in the literature. n = number of specimens; DML = dorsal mantle length; ind. = indeterminate; <math>- = no data.

of the two species, which is 600 mm in *O. bartramii* and 1000 mm in *L. grimaldii*, but may also be due to distinct growth rates and feeding habits (Fang et al. 2016).

The cause of death of the specimen 1 is unknown. It was found in very good condition and floating in the surface waters of El Hierro Island (Mar de Las Calmas). This is an area of the Canary Islands frequently used by a resident beaked whale population, and where mesopelagic and benthopelagic fauna remains floating at the surface; for example, hatchet fish *Argyropelecus* spp. Cocco, 1829 and histioteuthid squid are frequently found (Escánez et al. 2011). The depth where specimen 2 was caught is consistent with the known bathymetric distribution of *L. grimaldii*, which is reported as a mesopelagic to bathypelagic species (Jereb & Roper 2010).

### Acknowledgements

This work has been supported by the project "Cetáceos, Oceanografía y Biodiversidad de las Aguas Profundas de La Palma y El Hierro" funded by "Ministerio de Ciencia e Innovación" of the Spanish Government, grant number CETOBAPH-CGL2009-1311218. We wish to thank the Instituto Español de Oceanografía for the use of the RV Cornide de Saavedra and its crew. We are grateful to Dr. Natacha Aguilar for collecting specimens. We would like to thank Dr. Louise Allcock for her valuable comments on the paper.

#### References

- Clarke, M. R. 1960. *Lepidoteuthis grimaldii* a squid with scales. Nature 4754: 955–956.
- -- 1962. The identification of the cephalopods beaks and the relationships between beak size and total body weight. Bulletin of the British Museum (Natural History), Zoology Series 8(10): 419–480.
- 1964. Young stages of *Lepidoteuthis grimaldi* (Cephalopoda, Decapoda). Proceedings of the Malacological Society of London 36: 69–78.
- 1977. A brief review of sampling techniques and tools of marine biology. Pp. 439–469 in: Angel, M. (ed.). A voyage of discovery. Oxford and New York (Pergamon Press).
- 2006. Oceanic cephalopod distribution and species diversity in the eastern North Atlantic. Arquipélago, Life and Marine Sciences 23A: 27–46.
- -- & Maul, G. E. 1962. A description of the "scaled" squid *Lepidoteuthis grimaldi* Joubin 1895. Proceedings of the Zoological Society of London 139: 97-118.
- Dürr, J. & González, J. A. 2002. Feeding habits of *Beryx* splendens and *Beryx decadactylus* (Berycidae) off the Canary Islands. Fisheries Research 54: 363–374. doi:10.1016/S0165-7836(01)00269-7
- Escánez, A., Guerra, A., González, A. Tobeña, M., Landeria, J. M. & Aguilar, N. 2011. Trophic resources of blainville's beaked whales (*Mesoplodon densirostris*) and cuvier's beaked whales (*Ziphius cavirostris*) in El Hierro, Canary Island. P. 239 in: Gauffier, P. & Verborgh, P. (eds). Abstract book. 25th Conference of the European Cetacean Society. Long term datasets on marine mammals: learning from the past to manage the future. 21<sup>st</sup>–23<sup>rd</sup> March 2011, Cádiz, Spain.

- Fang, Z., Liu, B., Chen, X., Jin, Y., Li, J. & Chen, Y. 2016. Sexual asynchrony in the development of beak pigmentation for the neon flying squid *Ommastrephes bartramii* in the North Pacific Ocean. Fisheries Science 82: 737–746. doi:10.1007/s12562-016-1011-y
- Fernández, R., Santos, M. B., Carrillo, M., Tejedor, M. & Pierce, G. J. 2009. Stomach contents of cetaceans stranded in the Canary Islands 1996–2006. Journal of the Marine Biological Association of United Kingdom 89(5): 873–883. doi:10.1017/S0025315409000290
- Hernández-García, V. & Martín, V. 1994. Stomach contents of two short-finned pilot whale (*Globicephala macrorhynchus* Gray, 1846) (Cetacea, Delphinidae) off the Canary Islands: a preliminary note. International Council for the Exploration of the Sea (ICES) CM 1994, n. 16.
- Hoving, H. J. T. 2008. Reproductive biology of oceanic decapodiform cephalopods. 187 pp., Ph.D. Thesis, University of Groningen, The Netherlands.
- Jackson, G. D. & O'Shea, S. 2009. Unique hooks in the male scaled squid *Lepidoteuthis grimaldi*. Journal of the Marine Biological Association of the United Kingdom 83: 1099–1100.
- Jereb, P. & Roper, C. F. E. 2010. Cephalopods of the world. An annotated and illustrated catalogue of cephalopod species known to date. Vol. 2. Myopsid and Oegopsid squids. FAO Species Catalogue for Fishery Purposes 4(2): 1–605.
- Joubin, L. 1895. Contribution à l'Étude des Céphalopods de l'Atlantique Nord. Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, Fascicule IX, 63 pp. [In French].
- 1900. Céphalopods provenant des campagnes de la Princesse-Alice (1891–1897)]. Résultats des Cam-

pagnes Scientifiques Accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, Fascicule XVII, 135 pp. [In French].

- Lipinski, M. R. 1979. Universal maturity scale for the commercially-important squids (Cephalopoda: Teuthoidea). The results of maturity classification of the *Illex illecebrosus* (LeSueur, 1821). Population for the years 1973–1977. ICNAF Research Document 79/II/38, Serial No. 5364, 40 pp.
- Lu, C. C. & Clarke, M. R. 1975. Vertical distribution of cephalopods at 11°N, 20°W in the north Atlantic. Journal of the Marine Biological Association of the United Kingdom 55: 369–389.
- Merret, N. R. 1963. *Lepidoteuthis grimaldi* Joubin (Cephalopoda) from the south Atlantic. Annals and Magazine of Natural History 13(6): 635–636.
- Moreno, A., Dos Santos, A., Piatkowski, U., Miguel P. Santos, A. & Cabral, H. 2009. Distribution of cephalopod paralarvae in relation to the regional oceanography of the western Iberia. Journal of Plankton Research 31 (1): 73–91. doi:10.1093/plankt/fbn103
- Nesis, K. N. & Nikitina, I. V. 1990. Revisión of the squid family Lepidoteuthidae. Zoologichesky Zhurnal 69 (10): 38–49. In: Sweeney, M. J. (comp.) 2003. English translations of selected publications on cephalopods by Kir N. Nesis. Vol. 1 (2): 773–787, Washington, D.C. (Smithsonian Institution Libraries).
- Roper, C. F. E. & Voss, G. L. 1983. Guidelines for taxonomic descriptions of cephalopods species. Memoirs of the National Museum of Victoria, 44: 49–63.
- Young, R. E. & Vecchione, M. 2016. Lepidoteuthidae Pfeffer 1912. Lepidoteuthis grimaldii Joubin, 1895. The scaled squid. Tree of life project. World Wide Web electronic publication http://tolweb.org/Lepidoteuthis\_grimaldii/19833 [accessed 16-Nov-2016].

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Spixiana, Zeitschrift für Zoologie

Jahr/Year: 2017

Band/Volume: 040

Autor(en)/Author(s): Escanez Alejandro, Guerra Angel, Rocha Francisco, Soldevilla Fernando Lozano

Artikel/Article: <u>New records of the scaled squid, Lepidoteuthis grimaldii Joubin, 1895 in</u> the Canary Islands, Eastern Atlantic Ocean (Cephalopoda, Oegopsida, Lepidoteuthidae) 7-12