

Scientific note

The unstoppable march of *Halyomorpha halys*: new first country records

(Hemiptera, Pentatomidae)

Fabio Cianferoni, Francesca Graziani & Filippo Ceccolini

Halyomorpha halys (Stål, 1855) is a stink bug native to East Asia, which has been introduced into several European countries, North America, Chile, and intercepted also in Oceania (see Cianferoni et al. 2018 and Claerebout et al. 2018 for further details). Concerning Europe, in several northern countries it is considered only intercepted (Claerebout et al. 2018); in particular in Great Britain this insect was intercepted twice, one in 2010 in London, in association with passenger luggage flown in from the USA, and the other in 2013 in North Yorkshire, associated with a consignment of stone imported from China (Malumphy 2014, Cianferoni et al. 2018). Eastwards this insect reached European Turkey and Caucasus, but to date no records existed for the Asian part of Turkey.

New first occurrence records updating the distribution of this invasive and pest species (Bergmann et al. 2016) are given.

The examined material, from the platform “iNaturalist” (www.inaturalist.org), was revised by the authors and it is listed below. Geographical coordinates are in decimal degrees (datum WGS84). The uncertainty (in metres) of data was indicated according to the point-radius method (Wieczorek et al. 2004).

Greece. Lesbos Island, Pyrgi Thermis, 39.173592°N 26.503971°E (uncertainty = 335 m), 23.IV.2019, 1 adult (Fig. 1), photo by Savvas Zafeiriou.

Great Britain. England, Farnham, Zoo Birdworld, 51.181439°N 0.841524°E (uncertainty = 4 m), 27.IV.2019, 1 adult, photo by Jaime Rockhill.

Turkey. Istanbul, Kadıköy, 40.98098°N 29.02079°E (uncertainty = 3 m), 23.IX.2018, 1 adult, photo by Ahmet Kutsi Nircan; Istanbul, Üsküdar, 41.0187°N 29.008997°E (uncertainty = 5 m), 27.X.2018, 1 adult, photo by “c_nurdan_ozturk”.

USA, Louisiana. New Orleans, 29.95057°N 90.04649°W (uncertainty = 2 km), 30.IV.2018, 1 adult, photo by Thomas Werts. **Oklahoma.** Tulsa, Broken Arrow, 36.007804°N 95.817144°W (uncertainty = 5 m), 2.XII.2018, 1 adult, photo by “autumn4444”.

In the USA *H. halys* is now recorded for all the states (Cianferoni et al. 2018), completing its unstoppable invasion of the country.

Concerning Europe, a further recent record for England, where the species was known only through multiple introductions (but only intercepted), is given. Now, it will be important to continue monitoring in order to ascertain if the species has established or not in the country. The Turkish records are the first east of

the Bosphorus and, although they are near to the previous published record from Istanbul (Çerçi & Koçak 2017), these are actually the first ones for the Asiatic Turkey. The record from Lesbos Island (Greece), very near to Anatolia, is the easternmost for this country (Cianferoni et al. 2018, Claerebout et al. 2018). It is very likely that Anatolia will be invaded soon, since *H. halys* has now crossed the Bosphorus and landed on the Asian side and is also very close to Asian Turkey in Georgia in the east (see Claerebout et al. 2018).

These data unfortunately confirm again the very quick spreading of this allochthonous true bug in the Western Palaearctic, threatening several new countries northwards (i. e. Great Britain) and from Eastern Europe towards Near East (Cianferoni et al. 2018).

Acknowledgements. We thank all the photographers who uploaded these observations on the platform “iNaturalist”.

References

- Bergmann, E. J., Venugopal, P. D., Martinson, H. M., Raupp, M. J. & Shrewsbury, P. M. 2016. Host plant use by the invasive *Halyomorpha halys* (Stål) on woody ornamental trees and shrubs. *PLoS One* 11 (2): e0149975.
- Çerçi, B. & Koçak, Ö. 2017. Further contribution to the Heteroptera (Hemiptera) fauna of Turkey with a new synonymy. *Acta Biologica Turcica* 30 (4): 121–127.
- Cianferoni, F., Graziani, F., Dioli, P. & Ceccolini, F. 2018. Review of the occurrence of *Halyomorpha halys* (Hemiptera: Heteroptera: Pentatomidae) in Italy, with an update of its European and World distribution. *Biologia* 73 (6): 599–607.
- Claerebout, S., Haye, T., Ólafsson, E., Pannier, E. & Bultot, J. 2018. Premières occurrences de *Halyomorpha halys* (Stål, 1855) pour la Belgique et actualisation de sa répartition en Europe (Hemiptera: Heteroptera: Pentatomidae). *Bulletin de la Société Royale Belge d'Entomologie* 154: 205–227.
- Malumphy, C. 2014. Second interception of *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae) in Britain. *Het News* 21 (3): 4–5.
- Wieczorek, J., Guo, Q. & Hijmans, R. J. 2004. The point-radius method for georeferencing locality descriptions and calculating associated uncertainty. *International Journal of Geographical Information Science* 18 (8): 745–767.



Fig. 1. Adult of *Halyomorpha halys* from Lesbos Island (photo by Savvas Zafeiriou).

Fabio Cianferoni, Zoological Section “La Specola”, Natural History Museum, University of Florence, Via Romana 17, 50125 Florence, Italy; and Research Institute on Terrestrial Ecosystems, CNR – National Research Council of Italy, Via Madonna del Piano 10, 50019 Sesto Fiorentino (Florence), Italy; e-mail: cianferoni.fabio@gmail.com

Francesca Graziani & Filippo Ceccolini, Zoological Section “La Specola”, Natural History Museum, University of Florence, Via Romana 17, 50125 Florence, Italy; e-mails: francescagraziani78@gmail.com, ceccolinif@virgilio.it

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 2019

Band/Volume: [042](#)

Autor(en)/Author(s): Cianferoni Fabio, Graziani Francesca, Ceccolini Filippo

Artikel/Article: [The unstoppable march of Halyomorpha halys: new first country records](#)
[60](#)