

## Scientific note

***Aproaerema cinctelloides* (Nel & Varenne, 2012)  
is not a Mediterranean island endemic**

(Lepidoptera, Gelechiidae, Anacampsinæ)

Andreas H. Segerer &amp; Peter Huemer

The genus *Aproaerema* Durrant, 1897 comprises in Europe a total of 30 species many of which are difficult to identify due to significant overall similarity (Elsner et al. 1999, Huemer & Karsholt 2020). Recently, *Aproaerema cinctelloides* was described as a new species, morphologically very close to, though significantly distinct from, *A. cinctella* (Clerck, 1759) (Nel & Varenne 2012). Originally, *A. cinctelloides* has been supposed to be endemic to Corsica. Here, we report that *A. cinctelloides* does also occur in several different places in mainland Europe.

The first specimens outside of Corsica were collected in Germany between 2010–2019 and identified from DNA barcodes using the holotype of *A. cinctelloides* as a reference, which is present in the private collection of Thierry Varenne (Nice, France). According to genetic analysis using the tools provided in the BOLD database (Ratnasingham & Hebert 2007; <http://boldsystems.org/>), *A. cinctelloides* and *A. cinctella* are separated from

each other by a mean genetic K-2-P distance of 5 %. The maximum intraspecific variation in *A. cinctelloides* is 1.28 % with two separate clusters, one from Germany and Corsica, the other from Lower Austria to the Balkans (see Huemer et al. 2020 for details).

At present, only one German population of *A. cinctelloides* is known, occurring highly locally in the Danube valley east of Regensburg, central Bavaria within steep formations of Jurassic limestone with a markedly xerothermophilic microclimate (Figs 1, 2). In the course of the recent extensive DNA barcoding survey of European Gelechiidae, *A. cinctelloides* was also detected in SE Europe (Greece, northern Macedonia) (Huemer et al. 2020), and in Lower Austria (Stark, unpublished record).

From our recent data, we conclude that *A. cinctelloides* is a local, xerothermophilic species which may have been overlooked in mainland Europe due to the overall similarity to the sibling taxon, *A. cinctella*, which is common and has a much broader ecological ampli-



**Fig. 1.** *Aproaerema cinctelloides*, new for Central Europe (Bavaria: Donauhänge near Regensburg, August 21 2010, BC ZSM Lep 61448, leg. Segerer, coll. ZSM). Photo: A. H. Segerer (scale bar: 2 mm).

Andreas H. Segerer (corresponding author), SNSB – Zoologische Staatssammlung München, Münchhausenstr. 21, 81247 München, Germany; e-mail: [segerer@snsb.de](mailto:segerer@snsb.de)

Peter Huemer, Naturwissenschaftliche Sammlungen, Tiroler Landesmuseen Betriebsges. m.b.H., 6060 Hall, Austria; e-mail: [p.huemer@tiroler-landesmuseen.at](mailto:p.huemer@tiroler-landesmuseen.at)



Fig. 2. Habitat of *A. cinctelloides* in central Bavaria. Photo: A. H. Segerer.

tude. Presumably, *A. cinctelloides* can be found in suitable habitats also elsewhere in Europe. Any *A. cinctella*-like specimens collected at hot, dry locations should be thoroughly checked for identity.

**Acknowledgements.** Funding for barcoding the Bavarian specimens was provided through the project Barcoding Fauna Bavarica (BFB) by the Bavarian Ministry of Science and Art (Bayerisches Staatsministerium für Wissenschaft und Kunst, Munich, Germany). DNA barcoding of specimens outside Germany was supported by the Promotion of Educational Policies, University and Research Department of the Autonomous Province of Bolzano, South Tyrol with funds to the projects “Genetische Artabgrenzung ausgewählter arktalpiner und boreomontaner Tiere Südtirols” and “Erstellung einer DNA-Barcode-Bibliothek der Schmetterlinge des zentralen Alpenraumes (Süd-, Nord- und Osttirol)”.

## References

- Elsner, G., Huemer, P. & Tokár, Z. 1999. Die Palpenmotten (Lepidoptera, Gelechiidae) Mitteleuropas. 208 pp., Bratislava (František Slamka).
- Huemer, P. & Karsholt, O. 2020. Commented checklist of European Gelechiidae (Lepidoptera). ZooKeys 921: 65–140.
- , Karsholt, O., Aarvik, L., Berggren, K., Bidzilya, O., Junnilainen, J., Landry, J.-F., Mutanen, M., Nupponen, K., Segerer, A., Šumpich, J., Wieser, C., Wiesmair, B. & Hebert, P. D. N. 2020. DNA barcode library for European Gelechiidae (Lepidoptera) suggests greatly underestimated species diversity. Zookeys 921: 141–157.
- Nel, J. & Varenne, T. 2012. *Pseudopostega cyrneoalcopepla* n. sp., *Monochroa cyrneoagonella* n. sp., *Syncopacma cinctelloides* n. sp., espèces nouvelles découvertes en Corse (Lep. Opostegidae, Gelechiidae). Oreina 17: 11–13.
- Ratnasingham, S. & Hebert, P. D. N. 2007. BOLD: The Barcode of Life Data System ([www.barcodinglife.org](http://www.barcodinglife.org)). Molecular Ecology Notes 7: 355–364.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 2020

Band/Volume: [043](#)

Autor(en)/Author(s): Segerer Andreas H., Huemer Peter

Artikel/Article: [Aproaerema cinctelloides \(Nel & Varenne, 2012\) is not a Mediterranean island endemic \(Lepidoptera, Gelechiidae, Anacampsinae\) 147-148](#)