


Stuttgarter Beiträge zur Naturkunde

Serie B (Geologie und Paläontologie)

Herausgeber:

Staatliches Museum für Naturkunde, Rosenstein 1, 70509 Stuttgart



Stuttgarter Beitr. Naturk.	Ser. B	Nr. 221	7 pp., 10 figs.	Stuttgart, 20. 3. 1995
----------------------------	--------	---------	-----------------	------------------------

New Upper Jurassic Diptera (Limoniidae, Eoptychopteridae) from the Solnhofen Lithographic Limestone (Bavaria, Germany)

By Wieslaw Krzeminski, Kraków, and Jörg Ansorge, Rostock

With 10 figures

Summary

Two new Diptera are described from the Upper Jurassic Solnhofen Lithographic Limestone (Bavaria, Germany): *Crenoptychoptera bavarica* n. sp. (Eoptychopteridae) and *Tipunia intermedia* n. g. n. sp. (Limoniidae). *Tipunia* exhibits characters that are typical of both the Limoniidae and Tipulidae.

Zusammenfassung

Aus den oberjurassischen Solnhofener Plattenkalken (Bayern, Deutschland) werden zwei neue Dipteren Spezies beschrieben: *Crenoptychoptera bavarica* n. sp. (Eoptychopteridae) und *Tipunia intermedia* (Limoniidae). *Tipunia* n. g. n. sp. besitzt typische Merkmale der Limoniidae und Tipulidae.

1. Introduction

The Upper Jurassic Solnhofen Lithographic Limestone (Bavaria, Germany) is well known not only for *Archaeopteryx* but also for its numerous fossil insects. Till now the representatives of the following orders have been found: Odonata, Ephemeroptera, Saltatoria, Blattodea, Homoptera, Heteroptera, Coleoptera, Neuroptera, Hymenoptera, Trichoptera and Diptera.

Diptera seem to be very rare, as up to now only one species has been described (*Prohirmoneura jurassica* HANDLIRSCH 1906, probably a representative of the Nemestrinidae (Brachycera): BEQUAERT & CARPENTER 1936, USSATCHOV 1968).

PONOMARENKO 1985 recorded a culiciform midge in the collection of the Museum of Natural History, Vienna.

Empidia wulpi WEYENBERGH 1869, *Asilicus lithophilus* WEYENBERGH 1869 and *Tipularia teyleri* WEYENBERGH 1869 are very doubtful dipterans which need revision.




Fig. 1. *Crenoptychoptera bavarica* n. sp. (Eoptychopteridae), holotype (SMNS 62460) from the Lower Tithonian Solnhofen Lithographic Limestone. — Scale 5 mm.

The here described new dipterans from the Solnhofen Lithographic Limestone are comparatively large, and it seems that small dipteran species have simply been overlooked.

Recently a tipulomorph-like fly has been figured by FRICKHINGER 1994 (Fig. 354).

2. Description

Suborder Neoneura

Infraorder Ptychopteromorpha

Family Eoptychopteridae HANDLIRSCH 1906

Genus *Crenoptychoptera* KALUGINA 1985

Crenoptychoptera bavarica n. sp.

Figs. 1–2

Holotype: Complete specimen (SMNS 62460), housed in Staatliches Museum für Naturkunde, Stuttgart, Germany, coll. W. Ludwig 1992.

Locus typicus: Wegscheid, near Eichstätt (Bavaria).

Stratum typicum: Upper Jurassic Lithographic Limestone, Lower Tithonian (Hybodontum Zone).

Derivatio nominis: After the Latin name Bavaria for Bayern.




Fig. 2. *Crenoptychoptera bavarica* n. sp. (Eoptychopteridae), wing venation.

Diagnosis. — Sc ending just before R_{4+5} -fork; crossvein m-m almost straight.

Description. — Complete fly, legs (15 mm) longer than wings. Only the right wing is well preserved, but the anal part. Wing length ca. 11 mm. Sc ending just before fork of R_{4+5} . R_1 long; crossvein r-r (R_2) positioned ca. three times its length before R_1 tip. Rs rather long (ca. $\frac{1}{4}$ of wing length and about $\frac{1}{4}$ longer than R_{4+5}). R_3 2.3 times as long as Rs. R_4 2.3 times as long as R_{4+5} . Crossvein r-m at the fork of Rs. d-cell long and narrow, its shape being characteristic of the genus. Only three medial veins present. M_{1+2} almost twice as long as M_1 . Crossvein m-m almost straight, only very slightly wavy. Terminal section of M_{3+4} invisible. Tip of Cu and anal veins not preserved.

Remarks. — Till now 3 species of the genus *Crenoptychoptera* have been described from the Jurassic and Lower Cretaceous of Asia (KALUGINA 1985, 1989). Only one specimen was found in Europe, in the Lower Jurassic of Dobbertin, Germany (ANSORGE, in preparation).

The species described here is larger than *C. antica* KALUGINA 1985 and *C. defossa* KALUGINA from the Middle Jurassic of Kubekowo (Siberia). *C. bavarica* has a shorter Sc than *C. gronskayae* KALUGINA 1989, which is almost of the same size.

Suborder Polyneura

Infraorder Tipulomorpha

Family Limoniidae HENDEL 1936

Subfamily Architipulinae HANDLIRSCH 1906

Genus *Tipunia* n. g.

Type species: *Tipunia intermedia* n. g. n. sp.

Derivatio nominis: A compound of *Tipula* (Fam. Tipulidae) and *Limonia* (Fam. Limoniidae).


Diagnosis. — Sc terminating in C; distal part of d-cell widened, which is caused by extension of m-m; m-cu long and almost parallel to the wing margin.

Tipunia intermedia n. g. n. sp.

Figs. 3–8

Holotype: Complete female (SMNS 9633), housed in Staatliches Museum für Naturkunde, Stuttgart, Germany.

Locus typicus: Solnhofen.




Figs. 3–4. *Tipunia intermedia* n. g. n. sp. (Limoniidae), holotype (SMNS 9633) from the Lower Tithonian Solnhofen Lithographic Limestone. — 3: General view, female; scale 5 mm. — 4: Left wing; scale 1 mm.

Stratum typicum: Upper Jurassic Lithographic Limestone, Lower Tithonian (Hybodontum Zone).

Derivatio nominis: After the intermediate position between Tipulidae and Limoniidae.

Diagnosis. — Crossvein sc-r at three times its length before the end of Sc. R_4 2.5 times as long as R_3 . Crossvein m-m ca. $\frac{1}{3}$ longer than the section of M_3 within the d-cell.




Figs. 5–7. *Tipunia intermedia* n. g. n. sp. (Limoniidae). — 5: General view. — 6: Wing venation. — 7: Ovipositor.

Description. — Body 18 mm long; head partially preserved, antennae and palpi invisible. Wings 13,2 mm long; Sc terminating in C opposite Rs-fork. Crossvein sc-r at three times its length before the end of Sc. R_1 ending close to R_{3+4} -fork. Crossvein r-r (R_2) short, almost parallel to costal wing margin. R_3 2.5 times as long as R_2 . Distal part of d-cell conspicuously widened, which is caused by extension of the crossvein m-m. m-m ca. $\frac{1}{3}$ longer than the section of M_3 within d-cell. M_4 wavy. Crossvein m-cu extended, connecting M_4 and Cu, positioned almost parallel to the wing margin. cu-cell very broad, short. A_2 rather long, straight. Wing membrane along Cu wrinkled as in some larger Tipulidae and Limoniidae. Female terminalia: ovipositor short, visible in dorsal view.

3. Remarks on the systematic position of *Tipunia*

The new genus possesses features characteristic of both the Limoniidae (Sc terminating in the wing margin; R_3 relatively long — Fig. 9) and the Tipulidae (r-r almost parallel to costal wing margin; cu-cell very broad — Fig. 10).



Figs. 8–10. Venation of the distal part of the wings of Tipulomorpha. — 8: *Tipunia intermedia* n. sp. (Limoniidae). — 9: *Architipula fragmentosa* (BODE) (Limoniidae: Architipulinae). — 10: *Tipula (Lunatipula) lunata* L. (Tipulidae).

Unfortunately, features of the head (tip of the rostrum, palpi and antennae) are unidentifiable. Furthermore the position of the ovipositor does not allow a reconstruction of its shape in lateral view. In this situation the fact that Sc terminates in the costa remains as a character that inclines us to place this genus in the family Limoniidae. The position of m-cu is characteristic of the subfamily Architipulinae.

Very probably, *Architipula austeni*, a species described by JARZEMBOWSKI (1991) from the Lower Cretaceous Weald Clay (Great Britain) belongs to this genus as well. This can be concluded from the photo in JARZEMBOWSKI (1991, Fig. 13a) and the author's remark that „Sc is directed to the anterior margin“. Beyond any doubt this species does not belong to *Architipula* HANDLIRSCH 1906.

Till now the oldest Limoniidae are known from the Upper Triassic (KRZEMINSKI 1992), and the Tipulidae from the beginning of the Upper Cretaceous (KRZEMINSKI 1992 a, b).

Acknowledgement

We thank Dr. G. Dietl, Staatliches Museum für Naturkunde in Stuttgart, for the loan of the specimens, Prof. Dr. R. Willmann, Göttingen, for comments on the manuscript.

4. References

- BEQUAERT, J. C. & CARPENTER, F. M. (1936): The Nemestrinidae of the Miocene of Florissant, Colorado, and their relations to the recent fauna. — *Journal of Paleontology*, **10**: 395–407, 7 figs.; Lawrence/Kansas.
- FRICKHINGER, K. A. (1994): Die Fossilien von Solnhofen. 356 S.; Korb (Goldschneck).
- JARZEMBOWSKI, E. A. (1991): New insects from the Weald Clay of the Weald. — *Proceedings Geological Association*, **102/2**: 93–108, 18 figs.; London.
- KALUGINA, N. S. (1989): Novye psikhodomorphnye dwukrylye nasekomye mezozoya Sibiri (Diptera: Eoptychopteridae, Ptychopteridae). — *Paleontologičeskii Zhurnal* **1989/1**: 65–77, 5 figs.; Moscow.
- KALUGINA, N. S. & KOVALEV, V. G. (1985): Dvukrylye Jury Sibiri. — *Nauka*, 198 pp., 121 figs., 8 pls.; Moscow.
- KRZEMINSKI, W. (1992): Triassic and Lower Jurassic stage of Diptera Evolution. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **65**: 39–59, 17 figs.; Zürich.
- (1992): *Tipula* (s. lato) *evae* n. sp. from Cretaceous (East Asia) — the oldest representative of the family Tipulidae (Diptera, Polyneura). — *Acta zoologica cracoviensis*, **35/1**: 43–44; Krakow. — [1992a]
- (1992): The oldest Polyneura and their importance to the phylogeny of the group. — *Acta zoologica cracoviensis*, **35/1**: 45–52, Krakow. — [1992b]
- PONOMARENKO, A. G. (1985): Fossil insects from the Tithonian „Solnhofener Plattenkalke“ in the Museum of Natural History, Vienna. — *Annalen des Naturhistorischen Museums Wien*, **87 (A)**: 135–144; Wien.
- USSATCHOV, D. A. (1968): Novye jurskie Asilomorpha (Diptera) fauny Karatau. — *Entomologiceskoye Obozreniye*, **47**: 617–628, 8 figs.; Moscow.

Authors adresses:

Dr. W. Krzeminski, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, ul. Sławkowska 17, 31-016 Kraków, Poland.
Jörg Ansorge, Danziger Str. 32, D-18107 Rostock, Germany.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Stuttgarter Beiträge Naturkunde Serie B \[Paläontologie\]](#)

Jahr/Year: 1995

Band/Volume: [221_B](#)

Autor(en)/Author(s): Krzeminski Wieslaw, Ansorge Jörg

Artikel/Article: [New Upper Jurassic Diptera \(Limoniidae, Eoptychopteridae\) from the Solnhofen Lithographie Limestone \(Bavaria, Germany\) 1-7](#)