

Stuttgarter Beiträge zur Naturkunde

Serie A (Biologie)

Herausgeber:

Staatliches Museum für Naturkunde, Rosenstein 1, D-70191 Stuttgart

Stuttgarter Beitr. Naturk.	Ser. A	Nr. 650	11 S., 18 Abb.	Stuttgart, 16. VI. 2003
----------------------------	--------	---------	----------------	-------------------------

New species and records of *Prostomis* Latreille, including the first fossil records from Baltic amber and a checklist of the species (Coleoptera: Prostomidae)

WOLFGANG SCHAWALLER

Abstract

New species and new distributional data of the known species of the genus *Prostomis* Latreille are presented, and diagnostic characters of some species are figured. New species: *Prostomis apoica* n. sp. and *Prostomis mindanaoica* n. sp. from the southern Philippines (Mindanao), *Prostomis weigeli* n. sp. from western New Guinea (Irian Jaya). *Prostomis americanus* Crotch, 1874 from northwestern America is considered as a valid species. The genus *Prostomis* is presented for the first time from Tertiary Baltic amber; the fossils could not be named to species because their aedeagus remains unknown. A checklist of all species with distributional data is compiled and a complete bibliography of taxonomic papers is added.

Key words: Coleoptera, Prostomidae, *Prostomis*, new species, fossil record, Baltic amber, checklist, bibliography.

Zusammenfassung

Neue Arten und neue Verbreitungsangaben schon bekannter Arten der Gattung *Prostomis* Latreille werden präsentiert, diagnostische Merkmale einiger Arten abgebildet. Neue Arten: *Prostomis apoica* n. sp. und *Prostomis mindanaoica* n. sp. von den südlichen Philippinen (Mindanao), *Prostomis weigeli* n. sp. vom westlichen Neu Guinea (Irian Jaya). *Prostomis americanus* Crotch, 1874, aus dem nordwestlichen Amerika wird als valide Art betrachtet. Die Gattung *Prostomis* wird das erste Mal aus tertiärem Baltischen Bernstein vorgestellt; die Fossilien können nicht zur Art bestimmt werden, weil deren Aedoeagus unbekannt bleibt. Eine Checkliste aller Arten mit Verbreitungsangaben ist zusammengestellt und eine vollständige Bibliographie taxonomischer Arbeiten beigefügt.

Contents

1	Introduction	2
2	New species	3
2.1	General remarks	3
2.2	<i>Prostomis apoica</i> n. sp.	3
2.3	<i>Prostomis mindanaoica</i> n. sp.	3
2.4	<i>Prostomis weigeli</i> n. sp.	5
3	New records of known species	5
3.1	<i>Prostomis africana</i> Grouvelle, 1896	5

3.2	<i>Prostomis americanus</i> Crotch, 1874	5
3.3	<i>Prostomis beatae</i> Schawaller, 1991	6
3.4	<i>Prostomis edithae</i> Schawaller, 1991	6
3.5	<i>Prostomis katrinae</i> Schawaller, 1991	7
3.6	<i>Prostomis kinabaluca</i> Schawaller, 1992	7
3.7	<i>Prostomis</i> cf. <i>lawrencei</i> Schawaller, 1993	7
3.8	<i>Prostomis mandibularis</i> (Fabricius, 1801)	7
3.9	<i>Prostomis mordax</i> Reitter, 1887	7
3.10	<i>Prostomis morsitans</i> Pascoe, 1860	8
3.11	<i>Prostomis pacifica</i> Fairmaire, 1881	8
3.12	<i>Prostomis susannae</i> Schawaller, 1991	8
4	The first fossil records from Baltic Amber	8
4.1	<i>Prostomis</i> sp. A	8
4.2	<i>Prostomis</i> sp. B	8
4.3	<i>Prostomis</i> sp. C	8
4.4	Discussion	9
5	Biology	9
6	Checklist of the species of <i>Prostomis</i> with their distribution	9
7	References	10

1 Introduction

The genus *Prostomis* Latreille has a wide range in the Holarctic, Oriental, Papuan-Pacific, Australian and South African regions. 24 valid species have been described till now (see chapter 5 and ARROW 1927, BLACKBURN 1897, 1903, CROTCH 1874, FABRICIUS 1801, FAIRMAIRE 1881, FLEISCHER 1919, GROUVELLE 1896, HETSCHKO 1930, OLLIFF 1884, PASCOE 1860, REITTER 1889, SCHAWALLER 1991, 1992, 1993, 1994, SZALLIES 1994, WATERHOUSE 1877). Besides the structure of the aedeagus, which is unusually small in comparison to the body size, the shape of the jugular processes on the ventral side of the head is considered as an important diagnostic species character, although the biological function of this striking structure which is equal in both sexes is unknown (SCHAWALLER 1993). These processes are surely not just “instruments” for moving forward in red-rotten wood, because in this case they would not have evolved to such modified characters in more or less identical “wood-conditions” worldwide.

Since my previous contributions about this genus (see references) several new specimens from different collections came at hands which are summarised in the present paper. As a result, three new species are described from the southern Philippines (Mindanao) and from western (Indonesian) New Guinea, *Prostomis americanus* Crotch, 1874 is considered as a valid taxon and several new records of known species enlarge our knowledge about their distributional patterns. I take the chance and add a new checklist of the species and a complete bibliography of taxonomic papers (see references). The phylogenetic situation within the genus is still unsatisfactory mainly due to the small number of known diagnostic characters.

The genus *Prostomis* is recorded for the first time from Tertiary Baltic amber, three fossil specimens are presented herein. They could not be named to species because their aedeagus remains unknown.

Acronyms of depositories

- BMNH The Natural History Museum, London/UK (MAX BARCLAY)
 BRIO Biosystematic Research Institute, Ottawa/Canada (Dr. ALEŠ SMETANA)

DEI	Deutsches Entomologisches Institut, Eberswalde/Germany (Dr. LOTHAR ZERCHE)
HNHM	Hungarian Natural History Museum, Budapest/Hungary (Dr. OTTÓ MERKL)
MHNG	Muséum d'Histoire Naturelle, Genève/Switzerland (Dr. IVAN LÖBL)
NHMB	Naturhistorisches Museum, Basel/Switzerland (Dr. DANIEL BURCKHARDT)
NME	Naturkundemuseum, Erfurt/Germany (MATTHIAS HARTMANN)
SMNS	Staatliches Museum für Naturkunde, Stuttgart/Germany
SMTD	Staatliches Museum für Tierkunde, Dresden/Germany (OLAF JÄGER)
TMSA	Transvaal Museum, Pretoria/South Africa (RUTH MÜLLER)

Acknowledgements

I would like to thank all colleagues and friends (see list of depositories) for the loan of material under their care. Dr. WOLFGANG WEITSCHAT (University Hamburg) kindly arranged the loan of the amber fossils from the private collections G. HERRLING (Bramsche/Germany) and F. KERNEGGER (Hamburg/Germany).

2 New species

2.1 General remarks

The characters of species within *Prostomis* have been discussed earlier (see for example SCHAWALLER 1993); therefore I describe and figure here only the diagnostic characters (jugular processes, mandibles, aedeagus). A longer description of all the other external characters seems worthless, because they are identical (apart from some variability) in all species worldwide.

2.2 *Prostomis apoica* n. sp. (Figs. 1–2, 13)

Holotype (♂): Philippines, SE Mindanao, Mt. Apo, Ilomavis, 1400 m, 18.–19.V.1996, leg. L. BOLM, SMNS.

Paratype: Same data as holotype, 1 ex. SMNS.

Etymology: Named after Mt. Apo on Mindanao, where the type series was collected.

Diagnostic characters: Jugular processes (Fig. 2) asymmetrical, left process longer and distinctly swollen shortly before rounded tip, right process with acute tip and without other modifications. Mandibles at the base with earlike dilatation (Fig. 1). Aedeagus see Fig. 13. Body length 7.3–8.0 mm.


Remarks: Differs from *Prostomis luzonica* Schawaller, 1992, described from Luzon, by completely different jugular processes and aedeagus, and from *Prostomis mindanaoica* n. sp., described herein also from Mindanao, also by different jugular processes (the aedeagi cannot be compared because the holotype of *mindanaoica* n. sp. is a female). The type localities of both described species from Mindanao are situated in different, isolated mountain ranges.

2.3 *Prostomis mindanaoica* n. sp. (Figs. 3–4)

Holotype (♀): Philippines, N Mindanao, 30 km W Maramag, 1600 m, 28.–30. XII.1990, leg. L. BOLM, SMNS.

Etymology: Named after the southern Philippine Island Mindanao, where the holotype has been collected.

Diagnostic characters: Jugular processes (Fig. 4) asymmetrical, left process longer and with a hammer-like tip, right process with an acute tip pointing some-



Figs. 1–12. Mandibles from dorsal (1, 3, 5, 11) and jugular processes from ventral (2, 4, 6, 7–8, 9–12). – 1–2. *P. apoica* n. sp., holotype. 3–4. *P. mindanaoica* n. sp., holotype. 5–6. *P. weigeli* n. sp., holotype. 7. *P. edithae* from Vietnam. 8. *P. edithae* from Sichuan. 9. *P. africana* from Saasveld. 10. *P. susannae* from Malaysia. 11–12. *P. cf. lawrencei* from Testega. – Scale line: 1 mm.

what outwards. Mandibles laterally at the base with earlike dilatation (Fig. 3). Body length 7.2 mm.

Remarks: The shape of the jugular processes differs from those in *Prostomis luzonica* Schawaller, 1992, described from the Philippines (Luzon), where the left jugular process bears a tip consisting of 3 teeth. Without having intermediate forms

at hand (possibly from the central Philippine islands), I consider this difference as species-specific and not as a gradual infraspecific variation. Thus, the specimen from Minadanao is described as a different species, although only a single female is available and unfortunately the aedeagi cannot be compared.

2.4 *Prostomis weigeli* n. sp. (Figs. 5–6, 14)

Holotype (♂): Indonesia, Irian Jaya, Fakfak, 2 km E airport, 16.–18.VII.1996, leg. P. SCHÜLE & P. STÜBEN, SMNS.

Paratypes: Indonesia, Irian Jaya, 50 km S Nabire, Pusppenssat, 750 m, 30.–31. XII.1997, leg. A. WEIGEL, 3 ex. NME, 1 ex. SMNS. – Indonesia, Irian Jaya, 50 km S Nabire, Flaga road, Pusppenssat, 18. II.1998, leg. A. WEIGEL, 2 ex. NME.

Etymology: Named after ANDREAS WEIGEL (Wernburg/Germany), one of the collectors of the type series.

Diagnostic characters: Jugular processes (Fig. 6) more or less symmetrical, both tips rounded and pointing forwards, inner side of both processes without teeth but medially with a more or less developed knob-like dilatation, outer side of both processes without teeth and distinctly convex. Mandibles laterally at the base with earlike dilatation (Fig. 5). Aedeagus see Fig. 14. Body length 5.5–7.7 mm.

Remarks: Two other species have been described from New Guinea (Papua), namely *Prostomis lawrencei* Schawaller, 1993 (occurring probably also in Irian Jaya, see new records below, and Queensland) and *Prostomis papuana* Schawaller, 1993, having different jugular processes, mandibles and aedeagi.

3 New records of known species

3.1 *Prostomis africana* Grouvelle, 1896 (Fig. 9)


New records: South Africa, Cape Prov., Tsitsikama Pt., 6.–8.VIII.1987, leg. G. MINET, 2 ex. SMNS. – South Africa, Tsitsikama Forest, Humansdorp D., I.1961, leg. N. LELEUP, 8 ex. TMSA. – South Africa, Knysna Forest, Swart R. Bos (=Forest), I.1961, leg. N. LELEUP, 1 ex. TMSA. – South Africa, N Knysna, Jonkersberg, XI.1941, leg. G. VAN SON, 2 ex. TMSA. – South Africa, S Cape Prov., George-Saasveld, 22.IX.1985, leg. S. ENDRÖDY-YOUNGA, 1 ex. TMSA.

Remarks: The bigger series now at hand shows that the figure of the jugular processes given in SCHAWALLER (1991) is based on a single small and underdeveloped specimen, for the “new” diagnostic shape of this structure see Fig. 9; the somewhat asymmetrical processes are distinctly bent outwards forming a hook.

3.2 *Prostomis americanus* Crotch, 1874

New records: Canada, British Columbia, Langley, 23. XII.1931, leg. K. GRAHAM, 4 ex. BMNH. – USA, California, Mendocino National Park, Potter Valley and Pilsbury Lake, 750 m, 7. VII.1991, leg. D. POLLOCK, 2 ex. SMNS.

Remarks: This species is considered herein as a valid taxon as described by CROTCH (1874) and not as a synonym of the Euro-Caucasian *Prostomis mandibularis* (Fabricius, 1801) as listed in the Coleopterorum Catalogus (HETSCHKO 1930). It is also different from the eastern Siberian species *Prostomis mordax* Reitter, 1887. I could not find any reference, where such a synonymy has been established with arguments, thus a formal revalidation seems not necessary. The American species in-



Figs. 13–15. Aedeagus. – 13. *P. apoica* n. sp., holotype. 14. *P. weigeli* n. sp., holotype. 15. *P.* cf. *lawrencei* from Testega. – Scale line: 0.3 mm.

cluding comparative morphology of the larvae will be treated elsewhere (D. POLLOCK in litt.). Type locality is Vancouver Island.

3.3 *Prostomis beatae* Schawaller, 1991

New record: Nepal, Sindhupalchok Distr., Manegero, 2500 m, 13.VI.1989, leg. M. BRANCUCCI, 1 ex. NHMB.

3.4 *Prostomis edithae* Schawaller, 1991 (Figs. 7–8)

New records: Nepal, Myagdi Distr., upper Myagdi Khola N Dobang, 2800–3100 m, 22.–24.IV.1995, leg. J. MARTENS & W. SCHAWALLER, 4 ex. SMNS. – Nepal, Kaski Distr., Bantanti, 2300 m, 21.IV.2000, leg. F. WOLF, 3 ex. NME. – Nepal, Manang Distr., Pisang, 3200 m, 31.V.1993, leg. J. SCHMIDT, 3 ex. NME. – Nepal, Manaslu Himal, Bara Pokhari, 3000 m, 4.IV.1999, leg. LAU & J. SCHMIDT, 1 ex. NME. – Nepal, SE Annapurna, Krapa Danda near Pilgrimshouse, 3300 m, 4.–5.VI.1997, leg. O. JÄGER, 2 ex. SMTD. – Nepal, Solukhumbu Distr., above Pangum, 2900–3000 m, 16.V.1997, leg. W. SCHAWALLER, 1 ex. SMNS. – China, N Yunnan, Yulong Shan, Ganhaizi Pass, 3000–3500 m, 18.–23.VII.1990, leg. D. KRÁL, 1 ex. NHMB. – China, Sichuan, Gongga Shan, Hailuogou, above camp 3, 3050 m, 8.VII.1996, leg. A. SMETANA, J. FARCAČ & P. KABÁTEK, 1 ex. BRIO. – N Vietnam, Tonkin, Sa Pa, 16.–20.V.1990, leg. O. SAUSA, 1 ex. SMNS.

Remarks: For the jugular processes of the new records in Vietnam and Sichuan see Figs. 7–8.

3.5 *Prostomis katrinae* Schawaller, 1991

New record: NE Thailand, Chiang Mai Prov., Doi Suthep, 19.–22.IV.1991, leg. S. BILY, 1 ex. SMNS.

3.6 *Prostomis kinabaluca* Schawaller, 1992

New records: Malaysia, Borneo, W Sarawak, Mt. Matang, 2000 ft. [= 600 m], 6.–8.II.1914, leg. G. E. BRYANT, 2 ex. BMNH. – Indonesia, W Sumatra, Bukittinggi, Gunung Singgalang, 2100–2600 m, 16.X.1990, leg. A. RIEDEL, 1 ex. SMNS. – Indonesia, Sumatra, Jambi, Gunung Kerinci, 1800–2100 m, 6.–7.III.1991, leg. L. BOČÁK & M. BOČÁKOVÁ, 4 ex. NHMB, 1 ex. SMNS. – Indonesia, W Java, Tangkuban Prahur Volcano, 30 km N Bandung, 1600–1800 m, 6.–9.X.1995, leg. H. J. BREMER, 5 ex. SMNS. – Indonesia, W Java, Gede Volcano, 50 km SE Bogor, 2800–2950 m, 5.XI.1984, leg. AGOSTI, I. LÖBL & D. BURCKHARDT, 1 ex. MHNG.

Remarks: A few specimens of this newly collected material, particularly from Sumatra, might be transitional forms concerning the jugular processes between *P. kinabaluca* Schawaller, 1992 (described from Borneo) and *P. katrinae* Schawaller, 1991 (described from Thailand). In other words: more material from other Oriental localities might show that we face only a single biospecies and not two different taxa.

3.7 *Prostomis* cf. *lawrencei* Schawaller, 1993 (Figs. 11–12, 15)

New records: Indonesia, Irian Jaya, Manokwari Prov., Testega, 1100–1300 m, 30.III.–2.IV.1993, leg. A. RIEDEL, 1 ex. SMNS. – Indonesia, Irian Jaya, Manokwari Prov., Testega-Meydoudga, 1100 m, 4.IV.1993, leg. A. RIEDEL, 1 ex. SMNS. – Indonesia, Irian Jaya, Jayawijaya Prov., between Theila and Habbema Lake, 2800–2950 m, 22.X.1993, leg. A. RIEDEL, 2 ex. SMNS.

Remarks: These specimens from the western (Indonesian) part of New Guinea coincide with type material of *Prostomis lawrencei* Schawaller, 1993, from Papua New Guinea and Queensland concerning the jugular processes (Fig. 12), and the aedeagus (Fig. 15), but differ by distinctly dilated mandibles (Fig. 11) (without lateral dilatation in the type material). At the present state of knowledge I consider these differences not as specific.

3.8 *Prostomis mandibularis* (Fabricius, 1801)

New records: Spain (“Hispania”), without date and collector, 2 ex. HNHM. – Italy, Alpi Apuane, SW Piazza al Serchio, 650 m, 11.IV.1992, leg. W. SCHAWALLER, 1 ex. SMNS. – Slovenia, NE Kočevje, Pragoszd Pečka, 850 m, 15.V.2002, leg. W. SCHAWALLER, 12 ex. SMNS. – Greece, Steréa Elada, Evritania, W Timfristos, 1.VI.1990, leg. V. BRACHAT, 1 ex. SMNS. – Greece, Makedonia, Smolikas, 2000 m, 5.–11.VIII.1980, leg. H. MÜHLE, 1 ex. SMNS. – Turkey, Prov. Artvin, Aralik near Borcka, 21.VIII.1991, leg. F. LANGE, 1 ex. SMNS. – Ukraine, Crimea, Iaila Mts., without date, leg. WINKLER, 4 ex. HNHM, 2 ex. SMNS. – Russia, Caucasus, Dagestan, forest at Samur river at the mouth into the Caspian Sea, 7.–8.VII.1991, leg. W. SCHAWALLER, 1 ex. SMNS. – Azerbaijan, Caucasus, Nabran 30 km W Khachmas, 21.–22.IV.1987, leg. S. GOLOVATCH & Y. ESKOV, 1 ex. SMNS.

3.9 *Prostomis mordax* Reitter, 1887

New records: Russia, Kuril Islands, Kunashir Island, 5.VII.1991, leg. S. KURBATOV, 2 ex. SMNS. – Russia, S Sachalin, Tschechova Mt., 29.VI.1973, leg. KASPARYAN, 1 ex. SMNS. – Rus-

sia, Primorskiy Kray, Przhivalski Mts., 53 km SE Ussuriysk, 250 m, 13.VI.1993, leg. L. ZERCHE, 1 ex. DEI.

3.10 *Prostomis morsitans* Pascoe, 1860

New record: India, Darjeeling, Gopaldhara, 4720 ft. [= 1450 m], XII.1911, leg. H. STEVENS, 1 ex. BMNH.

3.11 *Prostomis pacifica* Fairmaire, 1881

New record: Fiji, Viti Levu Island, Lombau, 26 km W Suva, 12.X.1985, leg. G. BORNE-MISSZA, 1 ex. SMNS.

3.12 *Prostomis susannae* Schawaller, 1991 (Fig. 10)

New records: Nepal, Humla Distr., 20 km W Simikot, 5–6 km SE Chala, 3500–3600 m, 28.VI.2001, leg. A. KOPETZ, 5 ex. NME, 1 ex. SMNS. – Nepal, Humla Distr., 13 km S Simikot, NE Malikaasthan, 3800–3400 m, 8.VII.2001, leg. A. WEIGEL, 1 ex. NME. – Nepal, Humla Distr., 20 km W Simikot, 3.8 km SE Chala, 3500 m, 27.V.2001, leg. A. WEIGEL, 2 ex. NME. – Nepal, Jumla Distr., Gothichaur, 3000 m, 1.V.1995, leg. A. WEIGEL, 7 ex. NME, 2 ex. SMNS. – Nepal, Jumla Distr., Khali Lagna Pass, 3500 m, 16.–17.VI.1998, leg. W. SCHAWALLER, 8 ex. SMNS. – Nepal, Mustang Distr., right bank of Lethe Khola near Lethe, 2400 m, 5.–7.V.1995, leg. J. MARTENS & W. SCHAWALLER, 3 ex. SMNS. – Nepal, Annapurna, Bratang, 2900 m, 22.IX.1992, leg. A. WEIGEL, 1 ex. NME. – Malaysia, Genting Highland, 12.IV.1990, leg. A. RIEDEL, 1 ex. SMNS.

Remarks: The specimen from Malaysia has the same shape of the jugular processes (Fig. 10) as material from Nepal (type locality).

4 The first fossil records from Baltic Amber

4.1 *Prostomis* sp. A (Figs. 16–17)


Material: Specimen from the private collection F. KERNEGGER, Hamburg. Amber piece embedded in plastic, fossil in good condition. Body length with mandibles 4.5 mm. For shape of the jugular processes see Fig. 16, left process somewhat thicker than right one. Mandibles with weak dilatations (Fig. 17).

4.2 *Prostomis* sp. B (Fig. 18)

Material: Specimen from the private collection G. HERRLING, Bramsche, no. 958. Amber piece not embedded, fossil in fair condition, ventral and dorsal side partly with milky cover. Body length with mandibles 5.5 mm. For shape of the jugular processes see Fig. 18, both processes nearly symmetrical. Shape of the mandibles not visible.

4.3 *Prostomis* sp. C

Material: Specimen from the private collection G. HERRLING, Bramsche, no. 963. Amber piece not embedded, fossil in bad condition, ventral side completely and dorsal side mostly with milky cover. Body length with mandibles 5.3 mm. Shape of the jugular processes and of the mandibles not visible.



Figs. 16–18. Mandible from dorsal (17) and jugular processes from ventral (16–18) in the fossil specimens from Tertiary Baltic amber. – 16–17. *Prostomis* sp. A from the KERNEGGER amber collection. 18. *Prostomis* sp. B from the HERRLING amber collection no. 958. – Scale line: 1 mm.

4.4 Discussion

The above listed Tertiary fossils of *Prostomis* can not be named to species, because the structure of the aedeagus remains unknown. It even seems not clear if the fossils represent a single species or different ones. The shapes of the jugular processes of at least 2 specimens are somewhat different. This can be interpreted as either infraspecific variation or as specific difference. Even additional fossil specimens might not solve this problem because it is very unlikely that the aedeagus will be visible without body dissection. The jugular processes of the recent European *Prostomis mandibularis* are similar in shape but not identical (see fig. 4 in SCHAWALLER 1991); but again, without knowing the aedeagus a distinct separation is impossible.

5 Biology

The species of *Prostomis* are characteristic elements of old red-rotten trees, where adults and larvae live syntopically. They probably more depend on the conditions of the rotten wood than being adapted to particular tree species. The European *Prostomis mandibularis* occurs in *Picea*, *Abies*, *Pinus*, *Fagus* and in other trees. Obviously, beetles and their larvae stay for several generations in such a rotten habitat and are not forced to leave those trees for years. Only after natural or artificial changing of the preferred ecological conditions, beetles spread out by nocturnal flight. Then, they might be caught by interception traps of coleopterologists as well as by sticky amber trees.

6 Checklist of the species of *Prostomis* with their distribution

africana Grouvelle, 1896
americanus Crotch, 1874
apoica n. sp.
atkinsoni Waterhouse, 1877
beatae Schawaller, 1991
cameronica Schawaller, 1992

Southern Africa
 Northwestern America
 Philippines: Mindanao
 Australia: Tasmania
 Himalayas: Nepal
 Malaysia

<i>cornuta</i> Waterhouse, 1877	Australia: Victoria, New South Wales, Australian Capital Territory, Queensland
<i>edithae</i> Schawaller, 1991	Himalayas: Nepal; China: Yunnan and Sichuan, Vietnam
<i>gladiator</i> Blackburn, 1903	Australia: New South Wales
<i>intermedia</i> Blackburn, 1897	Australia: Victoria, New South Wales, Australian Capital Territory, Tasmania
<i>katrinae</i> Schawaller, 1991	Thailand
<i>kinabaluca</i> Schawaller, 1992	Borneo, Java, Sumatra
<i>latoris</i> Reitter, 1889	Japan, Taiwan
<i>lawrencei</i> Schawaller, 1993	New Guinea, Australia: Queensland
<i>luzonica</i> Schawaller, 1992	Philippines: Luzon
<i>mandibularis</i> (Fabricius, 1801)	Europe, Crimea, Don, Caucasus, Elburs Mts.
syn. <i>elburica</i> Fleischer, 1919	
<i>mindanaoica</i> n. sp.	Philippines: Mindanao
<i>mordax</i> Reitter, 1887	Eastern Siberia, southern Kuriles, Sachalin, northern Japan
<i>morsitans</i> Pascoe, 1860	Himalayas: Nepal and Darjeeling
<i>novacaledonica</i> Schawaller, 1994	New Caledonia
<i>pacifica</i> Fairmaire, 1881	Fiji: Viti Levu, Vanua Levu, Kadavu
<i>papuana</i> Schawaller, 1993	New Guinea
<i>samoensis</i> Arrow, 1927	Samoa Group
<i>schlegeli</i> Olliff, 1884	Sri Lanka
<i>subtilis</i> Szallies, 1994	Southern Turkey
<i>susannae</i> Schawaller, 1991	Himalayas: Nepal; Malaysia
<i>weigeli</i> n. sp.	New Guinea

7 References

- ARROW, G. J. (1927): Coleoptera. Clavicornia and Lamellicornia. – Insects of Samoa 4, pp. 35–66; London (British Museum).
- BLACKBURN, T. (1897): Further notes on Australian Coleoptera, with descriptions of new genera and species. – Transactions of the Royal Society of South Australia 21: 88–98.
- BLACKBURN, T. (1903): Further notes on Australian Coleoptera, with descriptions of new genera and species. – Transactions of the Royal Society of South Australia 27: 91–182.
- CROTCH, G. R. (1874): Descriptions of new species of Coleoptera from the Pacific coast of the United States. – Transactions of the American entomological Society 5: 73–80.
- FABRICIUS, J. C. (1801): Systema Eleutheratorum secundum ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus, vol. I: XXIV + 506 pp.; Kiliae (Bibliopolii Academici).
- FAIRMAIRE, L. (1881): Essai sur les Coléoptères des îles Viti (Fidgi). – Annales de la Société entomologique de France (6) 1: 243–318.
- FLEISCHER, A. (1919): Eine neue *Prostomis*-Art aus dem Elbursgebirge. – Entomologische Blätter 15: 211.
- GROUVELLE, A. (1896): Descriptions de Clavicornes d’Afrique et de Madagascar. – Annales de la Société entomologique de France 65: 71–94.
- HETSCHKO, A. (1930): Fam. Cucujidae. – Coleopterorum Catalogus 109: 1–122; Berlin (W. Junk).
- OLLIFF, A. S. (1884): Description of a new species of *Prostomis* (Cucujidae) from Ceylon and a short account of its larva. – Notes of the Leyden Museum 6: 100–102.
- PASCOE, F. B. (1860): Notices of new or little known genera and species of Coleoptera. – Journal of Entomology 1: 98–132.
- REITTER, E. (1889): Verzeichnis der Cucujiden Japans mit Beschreibung neuer Arten. – Wiener entomologische Zeitung 8: 313–320.
- SCHAWALLER, W. (1991): Prostomidae (Coleoptera) aus dem Himalaya mit einem Beitrag zur Larvalmorphologie. – Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie) 461: 17 pp.

- SCHAWALLER, W. (1992): Prostomidae (Coleoptera) aus Südost-Asien. – *Revue Suisse de Zoologie* **99**: 255–262.
- SCHAWALLER, W. (1993): The genus *Prostomis* (Coleoptera: Prostomidae) in Australia and adjacent regions. – *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* **489**: 12 pp.
- SCHAWALLER, W. (1994): A new species of *Prostomis* Latreille from New Caledonia (Coleoptera: Prostomidae). – *Doriana* **6**: 1–3.
- SZALLIES, A. (1994): Drei neue Coleopteren-Arten aus der Türkei (Coleoptera: Prostomidae, Cerambycidae). – *Entomologische Zeitschrift* **104**: 259–263.
- WATERHOUSE, O. (1877): Descriptions of new Coleoptera from various localities. – *Entomologist's monthly Magazine* **14**: 23–28.

Author's address:

Dr. WOLFGANG SCHAWALLER, Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany; e-mail: schawaller.smns@naturkundemuseum-bw.de

Manuscript received: 12. III. 2003, accepted: 10. IV. 2003.

ISSN 0341-0145

Autoren-Richtlinien: <http://www.naturkundemuseum-bw.de/stuttgart/schriften>
Schriftleitung: Dr. Hans-Peter Tschornig, Rosenstein 1, 70191 Stuttgart
Gesamtherstellung: Gulde-Druck GmbH, 72072 Tübingen

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Stuttgarter Beiträge Naturkunde Serie A \[Biologie\]](#)

Jahr/Year: 2003

Band/Volume: [650_A](#)

Autor(en)/Author(s): Schawaller Wolfgang

Artikel/Article: [New species and records of *Prostomis* Latreille, including the first fossil records from Baltic amber and a checklist of the species \(Coleoptera: Prostomidae\) 1-11](#)