

Four new species of Rhysodini (Coleoptera: Carabidae) with revised keys to *Grouvellina* Bell & Bell and the *mishmicus* group of *Rhyzodiastes* Fairmaire

ROSS T. BELL & JOYCE R. BELL

Abstract

Yamatosa bacca n. sp. (China), *Shyrodes nakladali* n. sp. (China), *Rhyzodiastes (Temoana) puetzi* n. sp. (China), and *Grouvellina nzwani* n. sp. (Comoro Islands) are described and illustrated. New keys for *Grouvellina* and *mishmicus* group of *Rhyzodiastes (Temoana)* Bell & Bell are provided.

Key words: Carabidae, Rhysodini, new species, keys, China, Comoro Islands.

Zusammenfassung

Yamatosa bacca n. sp. (China), *Shyrodes nakladali* n. sp. (China), *Rhyzodiastes (Temoana) puetzi* n. sp. (China) und *Grouvellina nzwani* n. sp. (Komoren) werden beschrieben und illustriert. Neue Bestimmungsschlüssel für die Gattung *Grouvellina* und die *mishmicus*-Gruppe von *Rhyzodiastes (Temoana)* Bell & Bell werden vorgestellt.

Contents

1	Introduction	129
2	Genus <i>Yamatosa</i> Bell & Bell.....	129
3	Genus <i>Shyrodes</i> Grouvelle.....	131
4	Genus <i>Rhyzodiastes</i> , subgenus <i>Temoana</i> Bell & Bell.....	132
5	Genus <i>Grouvellina</i> Bell & Bell.....	133
6	References	135

1 Introduction

Recorded here are three additional species in the rich rhysodine fauna of western China as well as a second species of *Grouvellina* from the Comoro Islands. *Shyrodes nakladali* n. sp. is only the second species discovered in the isolated genus *Shyrodes*, the other species being *S. dohertyi* Grouvelle, 1903, from the Ruby Mines of northern Burma.

Abbreviations

CAS	California Academy of Sciences, San Francisco, CA., USA
CMNH	Carnegie Museum of Natural History, Pittsburgh, PA., USA
DEI	Deutsches Entomologisches Institut, Müncheberg, Germany
IOZ	Institute of Zoology, Chinese Academy of Sciences, Beijing, P. R. China
L/GW	Length/Greatest Width of pronotum
MSNF	Museo Zoologico “La Specola”, Sezione del Museo di Storia Naturale dell’Università degli Studi di Firenze, Florence, Italy
SMNS	Staatliches Museum für Naturkunde, Stuttgart, Germany

Acknowledgments

We thank the following museum curators for sending loans of specimens: D. KAVANAUGH (CAS), R. DAVIDSON (CMNH), L. BARTOLOZZI (MSNF) and W. SCHAWALLER (SMNS). We are also grateful to the following private collectors for the privilege of studying their rhysodines: O. NAKLÁDAL and R. KMECO (Czech Republic), A. PÜTZ and D. WRASE (Germany), and especially to H. BRUSTEL (France) for sending us many rhysodines from many places in the world over the years and to A. SMETANA (Canada) for finding and sending the specimens from the European private collectors. We also thank our colleagues at the University of Vermont, B. PARKER (Plant & Soil Science) and D. BARRINGTON (Botany), for providing work space.

2 Genus *Yamatosa* Bell & Bell

Yamatoa BELL, 1977.

Yamatosa BELL & BELL, 1979.

Yamatosa bacca n. sp.
(Figs. 1–4)

Holotype ♂: “China: N-Yunnan [c2005-16] Nujian Lisu Aut. Pref., Gongshan Co., Gaoligong Shan, side valley, 3000–3050 m, 27°47.90'N, 98°30.19'E, conif. forest with *Rhododen-*

dron, broad leaved bushes, litter, moss, dead wood, sifted along creek, & snow fields, 21-VI-2005, M. SCHÜLKE". (SMNS).

Paratypes: 1 ♂: Same data as holotype (SMNS); 3 ♂♂, 3 ♀♀: "S. China, Yunnan Prov., Pass 50 km W. of Judian, 11–13.6.2005, OTO NAKLÁDAL leg" (1 ♂, 1 ♀ Wrase Colln.; 1 ♂, 1 ♀ R. KMECO Colln.; 1 ♂, 1 ♀ CMNH); 1 ♂: "China, Yunnan, Gongshan County, Cikai Township, 40 km W of Gongshan on Dulong Valley Road at Dabadi, 2900 m, N27.79619°/E098.51867°, 29 September 2002, Stop #HBL-2002-20, H.-B. LIANG collector, Rhysodine sp. CH-02, det. D. H. KAVANAUGH 2004" (IOZ); 1 ♀: "China, Yunnan, Fugong County, Lishadi Township, 10 km W of Shibali on Shibali Road, 3221 m, N27.20055°/E098.71399°, 16 August 2005, Stop #PP-3905, P. PAQUIN collector" (IOZ). [The last two specimens listed were returned to D. KAVANAUGH at CAS and he will transfer them to IOZ].

Etymology

The word "bacca" in Latin means bead or nut, referring to the eye which is very small and projects from the marginal curve of the temporal lobe.

Diagnosis

A species similar to *Y. smetanorum* Bell & Bell but easily separated by the apparent absence of coarse punctures from the abdominal sterna and the metasternum. The abdominal pits are differently shaped and are not oblique.

Description


Body length 5.7–7.0 mm. Antennal segment XI obtuse, stylet absent; head cordate, sides evenly rounded, frontal grooves narrow, well-defined; eye reduced, slightly more than in *Y. smetanorum*, temporal lobes slightly separated (Fig. 1); mentum without punctures.

Pronotum (Fig. 1) with L/GW=1.32, slightly shorter than in *Y. smetanorum*; lateral margin not sinuate near base, or slightly so; basal impression about 0.5 of length of pronotum; marginal groove nearly complete, effaced only very near anterior margin; precoxal carina absent.

Elytral setae very reduced, one near apex of stria IV and one external to apical tubercle (Fig. 2). Metasternum (Fig. 3) entirely without punctures except for extremely minute ones; same for abdominal sterna; both sexes with deep lateral pits on sternum IV (Fig. 3). Anterior femur not toothed in either sex; middle tibia of male with calcar shape similar to that of *Y. smetanorum* but smaller; calcar of metatibia sinuate (Fig. 4).

The key in BELL & BELL (2002) can be modified as follows:

- 10 Mentum punctate medially, ♂ with tooth on femur of anterior leg 11



Figs. 1–4. *Yamatosa bacca* n. sp. – 1. Head and pronotum, dorsal view. 2. Left elytron, dorsal view. 3. Metasternum, abdomen ♂ or ♀. 4. Metafemur, metatibia ♂, lateral view. – Scale: 1 mm.

- Mentum without punctures, ♂ without tooth on femur of anterior leg. **12**
- 12** Metasternum without coarse punctures along lateral margins; abdominal sterna not coarsely punctate; both sexes with semicircular lateral pits on sternum IV, those of ♀ larger than in ♂. ***bacca* n. sp.**
- Metasternum with some coarse punctures near lateral margins; abdominal sterna coarsely punctate; both sexes with deep lateral pits on sternum IV, those of ♂ simple, those of ♀ divided by carinae into two parts. ***smetanorum***

3 Genus *Shyrodes* Grouvelle

Shyrodes GROUVELLE, 1903.

Shyrodes nakladali n. sp.

(Figs. 5–10)

Holotype ♂: “China, Yunnan [CH07-13], Baoshan Pref., Gaoligong Shan, E pass, 36 km SE Tengchong, 2200 m, 24°49'32"N, 98°46'06"E, decid. forest, litter, wood, fungi sifted, 31-V-2007, leg. A. PÜTZ” (PÜTZ Colln.) [to be deposited in the future at DEI].

Paratypes: 1 ♂: “S. China, Yunnan Prov., Gaoligong Shan Mts., Pass S.W. of Baoshan, 4–8.6.2005, OTO NAKLÁDAL leg.” (KMECO Colln.); 1 ♀: “China, Yunnan Province, Gaoli-

gongshan Mountains, Baoshan Prefecture, Baoshan County, Nankang Yakou, 24°49.9'N/98°46.0'E, 2130 m, 4–7 November 1998, Stop #98-129C, D. H. KAVANAUGH, C. E. GRISWOLD, C.-L. LONG, R. LI & H.-X. HE collectors” (IOZ) [forwarded by D. KAVANAUGH, CAS, to IOZ].

Etymology


Named for the collector OTO NAKLÁDAL, who has brought back this important discovery from western China. It is only the second species assigned to this genus.

Diagnosis

Similar to *S. doherthyi*, but lacking a discal striole, and the ventral tooth on the anterior femur is barely suggested.

Description

Body length 6.0–6.8 mm. Antenna with short, flattened apical stylet, rings of minor setae present on segments V–X, basal setae absent; head (Fig. 5) with frontal grooves deeper than in *S. doherthyi*, not reaching antennal grooves; frontal space small; medial angles of temporal lobes obtuse, narrowly separated; frontal grooves deeply incised, anteriorly connected to postclypeal grooves by much shallower portion; anterior part of head (Fig. 6)



Figs. 5–10. *Shyrodes nakladali* n. sp. – 5. Head and pronotum, dorsal view. 6. Head, left lateral view. 7. Left elytron, dorsal view. 8. Metasternum, abdomen ♂ or ♀. 9. Anterior femur ♂, lateral view. 10. Metatibia ♂, lateral view. – Scale: 1 mm.

more elongate than in *S. dohertyi*, mentum more elongate, length of mentum (from gular tubercle to median tooth) divided by greatest width=1.80, compared to 1.66 for *S. dohertyi*; temporal lobes with obtuse hind angles; eye protruded but even smaller than in *S. dohertyi*, less than one half diameter of first antennal segment.

Pronotum (Fig. 5) more elongate than in *S. dohertyi*, L/GW 1.37 (compared to 1.33 for *S. dohertyi*); marginal groove fine, nearly complete; basal impressions small, deep, open posteriorly, without discal striole; hind angles turned, forming obtuse prominences overhanging basal impressions; no setae on prothorax.

Elytra (Fig. 7) deeply impressed towards scutellum, anterior to oblique line running medially from humerus; stria I impressed, closely, finely punctate; striae II–VII represented by rows of very fine punctures; striae not extended into depressed post-scutellar area, nor to region of humeral angle. Metasternum (Fig. 8) with lateral carina extended 0.6 of its length. Abdominal sterna III–V (Fig. 8) each with one transverse row of coarse punctures; sternum VI with scattered coarse punctures; male and female with lateral pits on sternum IV, those of female deeper. Male with only minute vestige of femoral ventral tooth (Fig. 9) present, none on female; calcar of middle tibia scarcely evident, that of hind tibia (Fig. 10) distinct though small.

4 Genus *Rhizodiastes*, subgenus *Temoana* Bell & Bell

Rhizodiastes FAIRMAIRE, 1895.

Temoana BELL & BELL, 1985.

Rhizodiastes (Temoana) puetzi n. sp.

(Figs. 11–15)


Holotype ♂: “China Yunnan [CHO7-14], Baoshan Pref., Gaoligong Shan, 33 km SE Tengchong, 2100–2200 m, 24°51'22"N, 98°45'36"E, decid. forest, litter, wood, fungi sifted, 31-V-2007, leg. A. PÜTZ” (PÜTZ Colln.). [A. PÜTZ notes that at some future date he would like to deposit his holotype at DEI].

Paratypes: 1 ♀: Same data as holotype (WRASE Colln.); 1 ♂: “China, Yunnan Province, Gaoligongshan Mountains, Baoshan Prefecture, Baoshan County, Nankang Yakou, 24°49.9'N/98°46.0'E, 2130 m, 4–7 November 1998, Stop #98-129, D. H. KAVANAUGH, C. E. GRISWOLD, C-L. LONG, R. LI & H.-X. HE collectors” (IOZ) [transferred by D. KAVANAUGH, CAS].

Non-type material: 1 ♀: “S. China, Yunnan pr., 65 km E. Tengshong, humid forest, 2200–2350 m, 21–22.V.2003. leg. S. MURZIN” (BRUSTEL Colln.). [Note: Presumably Tengshong is the same place as Tengchong and is just a variant transcription.]

Etymology

Named for the collector, ANDREAS PÜTZ, another adventurous collector, who has brought back an additional species of the unexpectedly rich rhysodine fauna of western China.



Figs. 11–15. *Rhizodiastes (Temoana) puetzi* n. sp. – 11. Head and pronotum, dorsal view. 12. Metasternum, abdomen ♂ or ♀. 13. Left elytron, dorsal view. 14. Profemur ♂, lateral view. 15. Metatibia ♂, lateral view. – Scale: 1 mm.

Diagnosis

A member of the *waterhousei* group, most similar to *R. preorbitalis* Bell & Bell. Both lack a median metasternal suture and transverse grooves on abdominal sterna III–V. In *R. puetzi* there is a transverse groove on each side of sternum VI, joined laterally to the marginal groove of that sclerite.

Description

Body length 5.8–7.9 mm. Antennal stylet short, acute; antennal segments VI–XI with basal setae; head (Fig. 11) as broad as long; median lobe of head longer than in *R. preorbitalis*, extended to level of eyes; posterior margin of temporal lobe fringed with pollinosity; orbital groove absent; preorbital pit present; eye narrow, crescentic, longer than in *R. preorbitalis*.

Pronotum (Fig. 11) shorter and broader than in *R. preorbitalis* (L/GW = 1.35), sides curved, widest near middle, apex narrowed, base less narrowed; median groove fine, linear; both median pits displaced towards middle of pronotum, posterior median pit at end of long, narrow depression; inner carina sloped gradually to paramedian groove; basal impressions broad, deep, triangular, tapered anteriorly into paramedian groove; outer carina rather broad at middle, less than half width of inner carina; marginal groove fine, complete, visible in dorsal view.

Elytra (Fig. 13) rather short, sides nearly parallel; sutural stria fine, becoming faint near base, finely punctate; parasutural stria much deeper, faintly punctate; interval III distinctly convex; intratubercular stria complete, impressed; marginal stria interrupted anteriorly. Abdominal sterna III–V (Fig. 12) with lateral pits, those of sternum IV enlarged in both sexes; sterna III–V without transverse sulci; sternum VI with widely separated transverse sulci, these each joined to apices of submarginal sulcus; sternum VI with one pair of apical setae. Male with low ventral carina on proximal half of anterior femur (Fig. 14), ended in an abrupt angle; calcars small (Fig. 15).

The species key of the *mishmicus* group of *Rhyzodistes* (*Temoana*) in BELL & BELL (1985) can be substituted for couplets 7 and 8 as follows:

- | | | |
|-------|---|---------------------|
| 7. | Metasternum with complete median sulcus..... | 8 |
| 7'. | Metasternum without median sulcus..... | 8.1 |
| 8. | Abdominal sterna III–V with transverse sulci..... | |
| | <i>waterhousei</i> | |
| 8'. | Abdominal sterna III–V without transverse sulci..... | |
| | <i>mishmicus</i> | |
| 8.1. | Abdominal sterna III–V with transverse sulci..... | <i>orestes</i> |
| 8.1'. | Abdominal sterna III–V without transverse sulci..... | 8.2 |
| 8.2. | Abdominal sternum VI with transverse sulci usually joined to marginal groove..... | <i>puetzi</i> n.sp. |
| 8.2'. | Abdominal sternum VI without transverse sulci..... | |
| | <i>preorbitalis</i> | |

5 Genus *Grouvellina* Bell & Bell

Grouvellina BELL & BELL, 1978.

Grouvellina nzwani n. sp.

(Figs. 16–20)

Holotype ♂: “Is. Comore, Anjouan, Nioumakele I., 1988, legit R. VIOSSAT” (MSNF).

Paratypes: 4 ♀♀: Same data as holotype (MSNF).

Etymology

“Nzwani” is the local name for the island usually called “Anjouan”.

Diagnosis


A species of *Grouvellina*, tracing to couplet 6 in our key to species (BELL & BELL 1979) and closest to *G. hexadon* (BELL & BELL 1985). The most obvious differences are in the male. In this species, the ventral tooth on the femur of the second leg is low and rounded, while in *G. hexadon*, it is sharp and prominent.

Description

Body length 5.9–8.0 mm. Antennal stylet short, acute; tufts of minor setae present on antennomeres V–X; basal setae V–X present; antennomeres II–X with both basal rings of pollinosity. Head (Fig. 16) approximately as wide as long, labrum with four setae; median head lobe short, broad; antennal lobes with broad borders and small glabrous bosses; antennal and postantennal grooves deep; median margin of temporal lobe broadly curved, lateral and basal margins of temporal lobes broadly margined with pollinosity; 2–3 temporal setae.

Pronotum (Fig. 16) much less elongate than in *G. hexadon*, L/GW = 1.22 (L/GW in *G. hexadon* = 1.35); pronotal carinae narrower than pronotal grooves, inner carina broadest near base, outer carina broadest near apex; marginal groove relatively broad; five marginal setae; prosternum with distinct precoxal carina in extensive precoxal pollinose area.

Elytra (Fig. 17) elongate, parallel-sided; striae deep, as broad as intervals; striae pollinose, very coarsely punctate, giving undulating margins to intervals; base of interval II elevated, forming small tooth; humeral angle elevated, tooth-like; stria II with 8 setae, stria IV with 7 setae, stria VII with about 8 setae. Metasternum (Fig. 20) entirely coarsely punctate, pollinose along anterior lateral margins, pollinosity broadened at anterior lateral angle. Abdominal sterna (Fig. 20) each with continuous pollinose strips; sternum VI coarsely punctate, with anterior lateral pollinose transverse grooves widely separated medially. Male with small ventral tooth on front femur, minute tooth on front tibia; middle femur (Fig. 18) with slight ventral swelling; hind femur (Fig. 19) with distinct ventral



Figs. 16–20. *Grouvellina nzwani* n. sp. – 16. Head and pronotum, dorsal view. 17. Left elytron, dorsal view. 18. Mesofemur, mesotibia ♂. 19. Metafemur, metatibia ♂. 20. Metasternum, abdomen ♂ or ♀. – Scale: 1 mm.

swelling. Female with ventral tooth on front femur but front tibia without tooth; female with slight ventral swellings on middle and hind femora; male with acutely pointed calcars on middle and hind tibiae (Fig. 19) (that of hind tibia of *G. hexadon* is blunt).

Species key

This is a revision of an earlier key found in BELL & BELL (1979).

- | | |
|---|---|
| <p>1 Outer carina of pronotum both complete and glabrous; marginal carina linear in dorsal view..... 2</p> <p>– Outer carina not both complete and glabrous; marginal carina broader, appearing as an additional ridge lateral to outer carina..... 16</p> <p>2 Anterior femur with ventral tooth in both sexes; anterior tibia of male with tooth on inner side..... 3</p> <p>– Anterior femur and tibia without teeth..... 10</p> <p>3 Eye in contact with glabrous area of temporal lobe, orbital groove incomplete. – Madagascar..... <i>montana</i></p> <p>– Eye isolated from temporal lobe by pollinose orbital groove..... 4</p> <p>4 Distinct glabrous parafrontal boss present..... 5</p> <p>– Parafrontal boss absent..... 9</p> <p>5 Pronotum widest at base. – Madagascar..... <i>cuneata</i></p> <p>– Pronotum widest anterior to base..... 6</p> <p>6 Postmentum pollinose; antenna with only antennomere II with pollinose band. – Madagascar..... <i>gigas</i></p> | <p>– Postmentum glabrous; antenna with pollinose bands on antennomeres II–X..... 7</p> <p>7 Anterior end of elytral interval II not projected. – Madagascar..... <i>tubericeps</i></p> <p>– Anterior end of interval II projected anteriorly as small tooth..... 8</p> <p>8 Middle femur of male with acute ventral tooth; pronotum elongate, L/GW = 1.22; hind calcar blunt. – Comores..... <i>hexadon</i></p> <p>– Middle femur of male with rounded ventral swelling; pronotum not elongate, L/GW = 1.35; hind calcar acute. – Comores..... <i>nzwani</i> n. sp.</p> <p>9 Four or five temporal setae; striae II and IV each with about ten setae; mentum short, its lateral margins parallel. – Madagascar..... <i>hova</i></p> <p>– One temporal seta; stria II with two, stria IV with four or five setae; mentum longer, with its lateral margins converging anteriorly. – Madagascar..... <i>janaki</i></p> <p>10 Labrum with four setae..... 11</p> <p>– Labrum with two setae..... 13</p> <p>11 Precoxal carina absent; elytral intervals carinate; base of interval II not projected; body length 9–10 mm. – Madagascar..... <i>ranavalona</i></p> <p>– Precoxal carina present; elytral intervals not carinate; base of interval II projected anteriorly as small tooth; body length 5–7 mm..... 12</p> <p>12 Parafrontal boss present; two temporal setae; four lateral setae on pronotum. – Madagascar..... <i>edentata</i></p> <p>– Parafrontal boss absent; three temporal setae; pronotum with two lateral setae. – Madagascar..... <i>cooperi</i></p> |
|---|---|

- 13** Median lobe of head long, narrow, its sides parallel. **14**
 – Median lobe shorter, broader, its sides not parallel. **15**
- 14** Pronotal carina not wider than pronotal grooves; pronotum more elongate, L/GW=1.35; pronotum widest near middle; five temporal setae; antennomeres I–IV with pilosity. – Madagascar..... *radama*
 – Pronotal carinae distinctly wider than grooves; pronotum shorter, L/GW=1.30; pronotum widest anterior to middle; three temporal setae; antennomeres I–X with pilosity. – Madagascar..... *bulirschi*
- 15** Tufts of minor setae on antennomeres V–X; medial angle of temporal lobe rounded; four lateral setae on pronotum. – Madagascar..... *descarpentriesi*
 – Tufts of minor setae on antennomeres VI–X; medial angle of temporal lobe acute; one lateral seta on pronotum. – Madagascar..... *divergens*
- 16** Outer carina of pronotum entirely pollinose or with very small glabrous area; outer groove of pronotum abbreviated both anteriorly and posteriorly; temporal lobe largely pilose; two or three temporal setae, arising within temporal pilosity; five pronotal setae. – Madagascar. *grouvellei*
 – Outer carina of pronotum entirely glabrous, abbreviated posteriorly; outer groove of pronotum complete, joining inner groove at basal impression; each temporal seta concealed within isolated tuft of pilosity; two pronotal setae..... **17**
- 17** Both sexes with ventral tooth on anterior femur; tufts on temporal lobe arranged in curve following posterolateral margin. – Madagascar..... *dentipes*
 – Anterior femur without ventral tooth; tufts of temporal lobe in transverse rows, one or two in anterior row, two or three in posterior row. – Madagascar..... *cinerea*

6 References

- BELL, R. T. (1977): Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel (Coleoptera: Fam. Rhysodidae). – *Entomologica basiliensis* **2**: 151–158.
- BELL, R. T. & BELL, J. R. (1978): Rhysodini of the world. Part I. A new classification of the tribe, and a synopsis of *Omoglymmius* subgenus *Nitiglymmius* new subgenus (Coleoptera: Carabidae or Rhysodidae). – *Quaestiones entomologicae* **14**: 43–88.
- BELL, R. T. & BELL, J. R. (1979): Rhysodini of the world. Part II. Revisions of the smaller genera (Coleoptera: Carabidae or Rhysodidae). – *Quaestiones entomologicae* **15**: 377–446.
- BELL, R. T. & BELL, J. R. (1985): Rhysodini of the world. Part IV. Revisions of *Rhyzodiastes* Fairmaire and *Clinidium* Kirby, with new species in other genera (Coleoptera: Carabidae or Rhysodidae). – *Quaestiones entomologicae* **21**: 1–172.
- BELL, R. T. & BELL, J. R. (2002): Two new species of Rhysodini (Coleoptera: Carabidae) with revised keys to *Yamatosa* Bell & Bell and *Omoglymmius* (*Pyxiglymmius*) Bell & Bell. – *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* **636**: 7 pp.
- FAIRMAIRE, L. (1895): Descriptions de quelques Coléoptères de Madagascar. – *Annales de la Société entomologique de Belgique* **39**: 8–40.
- GROUVELLE, A. (1903): Synopsis des Rhysodides et descriptions d'espèces nouvelles. – *Revue d'Entomologie* **22**: 85–148.

Authors' address:

Prof. Dr. ROSS T. BELL, JOYCE R. BELL, University of Vermont, Biology Department, Marsh Life Science Building, Burlington, Vermont 05401, USA;
 e-mail: rtbell@uvm.edu

Manuscript received: 7.IX.2010, accepted: 17.XI.2010.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Stuttgarter Beiträge Naturkunde Serie A \[Biologie\]](#)

Jahr/Year: 2011

Band/Volume: [NS_4_A](#)

Autor(en)/Author(s): Bell Ross T., Bell Joyce R.

Artikel/Article: [Four new species of Rhysodini \(Coleoptera: Carabidae\) with revised keys to Grouvellina Bell & Bell and the mishmicus group of Rhyzodiastes Fairmaire 129-135](#)