

A new species of *Promethis* Pascoe from West Papua with unusual head armature (Coleoptera: Tenebrionidae: Cnodalonini)

Author(s): Roland Grimm

Source: Stuttgarter Beiträge zur Naturkunde A, 9(1):181-183.

Published By: Stuttgart State Museum of Natural History

<https://doi.org/10.18476/sbna.v9.a9>

URL: <http://www.bioone.org/doi/full/10.18476/sbna.v9.a9>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

A new species of *Promethis* Pascoe from West Papua with unusual head armature (Coleoptera: Tenebrionidae: Cnodalonini)

ROLAND GRIMM

Abstract

Promethis skalei n. sp. from West Papua is described, illustrated and differentiated from its congeners by unusual head armature and last abdominal ventrite morphology.

Key words: Tenebrionidae, Cnodalonini, *Promethis*, new species, West Papua.

Zusammenfassung

Promethis skalei n. sp. aus West Papua wird beschrieben und abgebildet. Gegenüber den bereits bekannten Arten der Gattung zeichnet sich *P. skalei* durch den apikolateralen Fortsatz der Wangen und den apikalen Zahn des letzten abdominalen Ventrites aus.

Contents

1	Introduction	181
2	Description of the new species.....	181
3	References	183

1 Introduction

The species of the genus *Promethis* Pascoe, 1869 (Tenebrionidae: Stenochiinae: Cnodalonini) are typical forest dwellers of the Oriental, Papuan and Australian Regions, and were revised by KASZAB (1988a, b). Further species were described by BA & REN (2009) from Hainan Island (China), GRIMM (2011, 2013) from Borneo, MASUMOTO et al. (2005) from Taiwan, REN & BAI (2005) from Gansu Province (China), REN & HUA (2006) from Guizhou Province (China), and REN & YANG (2004) from Guangxi Province (China). ANDRÉ SKALE (Hof/Saale, Germany) collected in West Papua a species new for science which is described in the present paper.

Acknowledgements

I am grateful to JOHANNES REIBNITZ (Stuttgart) for taking the photographs and assembling the plate. LIU SHANSHAN (Baoding) and Dr. WOLFGANG SCHAWALLER (Stuttgart) kindly helped in getting copies of papers on Chinese *Promethis*. Dr. OTTÓ MERKL (Budapest) and Dr. WOLFGANG SCHAWALLER (Stuttgart) reviewed the manuscript.

Acronyms of depositories

CRG	Collection Dr. ROLAND GRIMM, Neuenbürg, Germany
SMNS	Staatliches Museum für Naturkunde Stuttgart, Germany

2 Description of the new species

Promethis skalei n. sp.

(Figs. 1–2)

Holotype ♂: W Papua, Manokwari prov., 18 km NW Ransiki, Anggi Gida, Kampung Itkau, 1830 m, cutting area, primary forest, 01°24.23'S/133°35.53'E, 4.III.2007, leg. A. SKALE (CRG).

Paratypes: Same data as holotype, 1 ♀ (CRG), 1 ♀ (SMNS).


Etymology

This species is named in honour of ANDRÉ SKALE (Hof/Saale, Germany), the collector of the type specimens.

Description

Elongate-oval, black, body length 23.6–25.5 mm, body width 9.0–10.0 mm.

Head widest across middle of eyes; anterior border of clypeus straight, laterally arcuate towards clypeo-genal meeting; fronto-clypeal and clypeo-genal sutures well incised, frontal suture insignificant. Genae terminating apico-laterally in a spine-like process (Fig. 1a). Surface densely, but finely punctured. Antennae short, reaching middle of pronotum. Mentum subcordiform, apico-laterally declivous, basal border arcuate; surface in the ♂ densely hirsute; in the ♀♀ densely coarsely to rugosely



Figs. 1–2. *Promethis skalei* n. sp., ♂ holotype. – **1.** Dorsal view. **1a.** Dorsal view of head. **1b.** Ventrites 4–5. **2.** Aedeagus, dorsal (a) and lateral (b) views.

punctured with only scattered long erect hairs (remaining hairs possibly abraded?).

Pronotum widest in the middle, transverse, width/length ratio (measured along transversal and longitudinal midlines) 1.42–1.56. Anterior border shallowly emarginate, lateral borders slightly arcuate towards apical angles, nearly straight and slightly converging towards basal angles, and shallowly emarginate before base. Apical angles obtusely rounded, basal angles subrectangular. All borders distinctly margined, apical margin interrupted in the middle. Surface densely punctured, punctures weakly impressed, with long, adpressed hairs, interspaces weakly microreticulate; disc with basally keeled midlongitudinal sulcus, somewhat flattened along lateral borders. Propleura nearly smooth, basally shallowly rugulose with scattered punctures; apically with fine, scattered punctures or microgranulate. Prosternal process broad, flat, bisulcate, slightly arcuate towards apex.

Elytra striate-punctate; intervals convex, with scattered minute punctures. Lateral margins in dorsal view visible along entire length. Metaventricle rugulose, with interspersed fine, hairy punctures or microgranules. Abdominal ventrite 1 rugulose, with interspersed setigerous punctures; ventrites 2–3 anteriorly rugulose, posteriorly finely punctured; ventrites 4–5 minutely punctured; ventrite 5 not bordered, but apically with distinct tooth (Fig. 1b).

Legs without modifications, but apex of ♂ protibiae widened and flattened ventrally, and fringed with yellow hairs on inner side.

Aedeagus see Fig. 2.

Differential diagnosis

Promethis skalei n. sp. can be easily separated from all described congeners by the apicolateral spine-like process

of genae (Fig. 1a) and by the apical tooth of last abdominal ventrite (Fig. 1, 1b).

3 References

- BA, Y.-B. & REN, G.-D. (2009): Taxonomy of *Promethis* Pascoe (Coleoptera, Tenebrionidae) on Hainan Island, China. – *Zootaxa* **2064**: 27–38.
- GRIMM, R. (2011): New and little known species of Tenebrionidae (Coleoptera) from Borneo (2). – *Stuttgarter Beiträge zur Naturkunde A, Neue Serie* **4**: 249–257.
- GRIMM, R. (2013): New and little known species of Tenebrionidae (Coleoptera) from Borneo (3). – *Stuttgarter Beiträge zur Naturkunde A, Neue Serie* **6**: 175–181.
- KASZAB, Z. (1988a): Katalog und Bestimmungstabelle der Gattung *Promethis* Pascoe, 1869 (Coleoptera, Tenebrionidae). – *Acta Zoologica Academiae Scientiarum Hungaricae* **34**: 67–170.
- KASZAB, Z. (1988b): Faunistische Angaben der Gattung *Promethis* Pascoe, 1869 (Coleoptera, Tenebrionidae). – *Folia Entomologica Hungarica* **49**: 55–116.
- MASUMOTO, K., AKITA, K. & LEE, C.-F. (2005): New tenebrionid beetles from Taiwan (1). – *Entomological Review of Japan* **60**: 247–254.
- REN, G.-D. & BAI, M. (2005): Coleoptera, Tenebrionidae. – In: YANG, X.-K. (ed.): *Insect Fauna of Middle-West Qingling Range and South Mountains of Gansu Province*, pp. 379–389; Beijing (China Science and Technology Press) [In Chinese with English summary].
- REN, G.-D. & HUA, H.-R. (2006): Tenebrionidae. – In: LI, Z.-Z. & JIN, D.-C. (eds.): *Fanjingshan Jingguan Kunchong*, pp. 265–274; Guiyang (Guizhou Science and Technology Publishing House) [In Chinese with English summary].
- REN, G.-D. & YANG, X.-J. (2004): Coleoptera: Tenebrionidae. – In: YANG, X.-K. (ed.): *Insects from Mt. Shiwandashan Area of Guangxi*, pp. 311–319; Beijing (China Forestry Publishing House) [In Chinese with English summary].

Author's address:

Dr. ROLAND GRIMM, Unterer Sägerweg 74, 75305 Neuenbürg, Germany;
e-mail: grimm.tenebrio@t-online.de

Manuscript received: 7.IX.2015, accepted: 10.IX.2015.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Stuttgarter Beiträge Naturkunde Serie A \[Biologie\]](#)

Jahr/Year: 2016

Band/Volume: [NS_9_A](#)

Autor(en)/Author(s): Grimm Roland

Artikel/Article: [A new species of *Promethis Pascoe* from West Papua with unusual head armature \(Coleoptera: Tenebrionidae: Cnodalonini\) 181-183](#)