verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum

A new Species of Mucor from India

Madhava D. Mehrotra. (907, Kalyani Devi, Allahabad, India.)

With 2 fig. in the text.

During the Winter of 1962, soil from different places in Suhagi (M. P.) was collected. Later on, the different soil samples were examined for new forms belonging to the order *Mucorales*. A. number of fungi were isolated and out of them, one appeared to be interesting. Single spore cultures of the fungus were made, and on examination, it was found to belong to the genus *Mucor*, but differed from



Fig. I. — 1—12. Mucor subagiensis, 1. A sporangiophore. Note the presence of septation and circinately borne lateral sporangia, 2—4. Sporangiophores showing terminal columellae. 5—7. Branching of the lateral sporangiophores, 8. Note the circinate tip of the lateral branch of the sporangiophore, 9. Columellae of terminal sporangia, 10. Columellae of lateral sporangia, 11. Sporangiospores. Note the range in size and shape, 12. Some enlarged sporangiospores showing echinulations.

all the described species of the section Sphaerosporus in certain features. The fungus was also sent to Dr. C. W. Hesseltine by Dr. B. S. Mehrotra who identified it as a strain of *Mucor brunneus*. It was, therefore, thought desirable to compare the present isolate with *Mucor brunneus*. A culture of *M. brunneus* was obtained through the courtesy of the Director, Central Bureau Voor Schimmel Cultures, Baarn, Holland. A detailed comparative study was made on various solid media. The author noticed several distinguishing characters between the two, and the differences are enumerated below.



Fig. II. -4-2. Mucor subagiensis. 1. Photomicrograph showing terminal sporangia, \times 150. 2. Photomicrograph showing columellae. Note the presence of circinate lateral sporangiophores, \times 130.

The present isolate comes in the section Sphaerosporus of the key to the genus Mucor, because majority of the spores are spherical. In this section it comes in the group with species growing at room temperature and with sporangiospores larger than 4.0μ . Out of the species included in this group, the present isolate comes close to Mucor brunneus Naumov, mainly because of the presence of echinulate sporangiospores, but differs from it in the absence of cylindrical columella with a spine at its tip; the presence of markedly septate sporangiophores; the presence of sporangia of two size ranges — the terminal sporangia markedly larger than the lateral ones; the lateral sporangia usually circinately borne; mostly spherical sporangio-spores instead of spherical to irregular; and the absence of chlamy-dospores.

Taking all the above facts into consideration, it appears that this isolate is one hitherto unknown. It is described as a new species and named after the place of origin.

Mucor suhagiensis Mehrotra, sp. nov.

Caespituli in PDA¹ et SMA² celeriter crescentes, primum caerulescentes postea brunnei; sporangiophora "Petri" paterae tectum attingentia, sporangio terminata, septata, 14.0—30 µ diam., irregulariter ramulosa; ramuli laterales iterum ramulosi, sporangiis globulosis terminati; sporangia biformia: terminalia majora, globulosa, asperula, fragilia, 56.5—96.0 µ; sporangia lateralia minora, sed terminalibus similia, 16.5—47.5 µ; columella sporangiorum terminalium globosa vel hemisphaerica, 40.0—67.8, sed sporangia lateralia 6.0-40 µ; sporangiosporae globulosae, ovoideae vel oblongae, echinulatae, 4.5—15.5 × 4.5—9.9 µ; zygosporae non visae.

Colonies on PDA¹ and SMA² growing rapidly, at first bluish later becoming brown; sporangiophores touching the lid of the Petri dish, terminating into a sporangium, septate, $14.0 - 30 \mu$ in diam branching profusely; side branches also branching, bearing circinate sporangia at their tip, $40. - 10.0 \mu$; sporangia of two types, the terminal sporangia larger, globular, rough-walled, fragile, $56.5-96.0 \mu$; the lateral sporangia smaller but like the terminal ones, $16.5 - 47.5 \mu$; columellae of the terminal sporangia globular to hemispherical, $40.0 - 67.8 \mu$ in diam, while of the lateral sporangia $6.0 - 40 \mu$; sporangiospores globular, oval to oblong, echinulate, $4.5 - 15.5 \times 4.5 - 9.9 \mu$; zygospores not seen.

- PDA¹: Peeled potatoes, 200gm; glucose, 20gm; agar agar, 20gm; dist. water, 1000ml.
- SMA²: Dextrose, 40gm; asparagine, 2gm; KH₂PO₄, 0.5gm; MgSO₄. 7H₂O, 0.25gm; thiamine chloride, 0.5mg; agar agar, 20gm; dist. water, 1000ml.

Isolated from soil collected from Suhagi, pH 6.4. Type: Culture deposited at NURD, Peoria, Illinois, U.S.A. Also, a culture deposited in the culture collection of the Botany Department (PCCC) as No. MD 10.

Acknowledgements

The author is grateful to the Director, Central Bureau Voor Schimmel Cultures, Baarn, Holland for sending a culture of *Mucor brunneus*, to Dr. F. Petrak for the Latin diagnosis and to Prof. Dr. R. N. Tandon for providing the necessary laboratory facilities. Thanks are also due to the Ministry of Scientific Research and Cultural Affairs, Government of India for the award of Research Training Scholarship during the tenure of this work & to Mr. R. K. Kakkar for his help.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Sydowia

Jahr/Year: 1964

Band/Volume: 17

Autor(en)/Author(s): Mehrotra Madhava Das

Artikel/Article: A new Species of Mucor from India. 282-284