

Coniribution to our Knowledge of Ascomycetes of India. XIII.

By S. T. Tilak

(Botany Department, Marathwada University, Aurangabad (Dn), India.

In earlier papers (1960—1966) 40 ascomycetes have been described. The present paper constitutes thirteenth in the series. In this paper are described *Lophiostoma lecanthii* sp. nov. on *Lecanthus peduncularis* (Wall.) Weld. *Hypoxyylon acaciae* sp. nov. on *Acacia arabica* Willd. and *Dimeriella trichosanthii* sp. nov. on *Trichosanthes* sp.

41. *Lophiostoma lecanthi* sp. nov.

Perithecia immersa, dispersa, globosa vel lageniformia, atra, erumpentia, ostiolo compresso praedita, 240—300 × 250—320 μ ; ascii numerosi, anguste clavati, crasse tunicati, breviter stipitati, 8-spori, 85—114 × 10—15 μ ; sporae irregulariter biseriatae fusiformes, usque 6-septatae, medio parum constrictae, primum hyalinae, postea olivaceae 38—48 × 3—5 μ .




Fig. 1. *Lophiostoma lecanthi*.

Pseudothecia immersed in substrata, scattered, globoid to flask shaped, black, erumpent, ostiolate, ostioles slit like, 240—300 × 250—320 μ . Ascii numerous, hyaline, bitunicate, narrowly clavate, stalk fairly short, 8 spored, 85—114 × 10—15 μ . Ascospores irregularly biseriate, fusiform, 38—48 × 3—5 μ . Transverse septa upto 6, no vertical septa, slightly constricted in the middle septum, hyaline later changing to olivaceous green,

Collected on dead stems of *Lecanthus peduncularis* (Wall.) Weld. in the month of July 1965, at Aurangabad, Leg. S. T. Tilak and deposited in the herbarium of Marathwada University under number MUH. 199 (Tilak type).

Besides being a new species the genus is an addition to the generic list of India.

42. *Hypoxyylon acaciae* sp. nov.

Perithecia globosa vel lageniformia, stromati primum pallide colorato, postea brunneo immersa, ostioli papillatis prominula; pariete a stromate bene distincto praedita, $450-600 \times 240-345 \mu$; asci cylindracei vel clavati, apice iodo ope colorati, 8-spori, tenuiter tunicati, paraphysati, $80-104 \times 8-15 \mu$; spora uniseriatae, postice attenuatae, ellipsoideae, obscure brunneae, $11,2-14,4 \times 6,4-7,2 \mu$.




Fig. 2. *Hypoxyylon acaciae*.

Perithecia globose to flask shaped, embedded in stroma which is light in colour, later changing to brown or dark. Stromatic surface is papillate due to projecting ostioles. Perithecia are distinctly separated from the stroma by a wall of its own and measures from $450-600 \times 240-345 \mu$. Asci cylindrical to clavate, hyaline, with apical ring which stains with iodine, 8-spored, unitunicate, originating from basal region, numerous, paraphysate, $80-104 \times 8-15 \mu$. Ascospores uniseriate, flattened on one side, ellipsoid, dark brown at maturity, $11,2-14,4 \times 6,4-7,2 \mu$.

Collected on dead stems of *Acacia arabica* Willd., in the month of April 1965, at Aurangabad, Leg. S. T. Tilak and deposited in the herbarium of Marathwada University under number MUH. 200 (Tilak type).

With the addition of this species the genus is now represented by species from India.

43. *Dimeriella trichosanthis* sp. nov.

Mycelium hypophyllo, plerumque totam folii superficiem obtegente; Perithecia superficialia, atra, ostiolata, globosa, setis nonnullis praedita, $105-135 \times 90-120 \mu$; pariete pseudoparenchymatico; ascii ellipsoidei vel piriformes, crasse tunicati, antice distincte incrassati, paraphysati, 8-spori, $35-45 \times 9,6-12,8 \mu$; sporae biseriatae, medio septatae, non constrictae, hyalinae, $9-12,8 \times 4-5 \mu$.

Hypophyllous appearing as sooty growth and occupying the laminar portion of leaf, fruit bodies in groups or solitary, superficial, black, ostiolate, with few setae, globose. Walls of the perithecia formed of




Fig. 3. *Dimeriella trichosanthis*.

pseudoparenchymatous cells, $105-135 \times 90-120 \mu$. Ascii bitunicate, hyaline, the outer wall heavily thickened at apex, elliptical to pyriform, paraphysate, numerous, 8-spored, $35-45 \times 9,6-12,8 \mu$. Ascospores two celled, hyaline, biseriate, transversely septate, not constricted at the septum, $9-12,8 \times 4-5 \mu$.

Collected on leaves of *Trichosanthes* sp., in the month of Sept. 1964, at Aurangabad and deposited in the herbarium of Marathwada University under number MUH. 201 (Tilak type).

The genus *Dimeriella* has been extensively studied by Marie Farr (1966) and according to her the valid name should be *Dimeriella* and the older name *Wentiomycetes* Koord should not be accepted as valid in the absence of type species.

Acknowledgements

Thanks are due to Marathwada University for the laboratory facilities and Dr. F. Petrak for the latin diagnosis of the new species.

References

- Farr Marie L. 1965. *Dimeriella*, *Wentiomycetes*, *Episphaerella* and *Epi-polaeum*. *Taxon* 14: 18-21.
— 1966. The Didymosporous dimeriaceous fungi described from leaves of gramineae. *Mycologia* LVIII: 221-248.

- Tilak, S. T. & S. B. K a l e . 1964. Second contribution to our knowledge of Indian Ascomycetes. *Mycopath. & Mycol. appl.* XXVII: 60—64.
- Tilak, S. T. 1966. Contribution to our knowledge of Ascomycetes of India. III. *Mycopath. & Mycol. appl.* XXVIII: 82—85.
- 1966. ibdem IV ibid. XXVIII: 86—89.
- Tilak, S. T. & R a m c h a n d r a R a o , 1966. ibdem V ibid. XXVIII: 90—94.
- Tilak, S. T. 1966. ibdem VI.—XII. l. c. in press.
By S. T. Tilak
- V i s h w a n a t h a n , T. S. & S. T. Tilak. 1960. Additions to our knowledge of Ascomycetes of India. ibdem. XIII: 237—242.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1966/1968

Band/Volume: [20](#)

Autor(en)/Author(s): Tilak S. T.

Artikel/Article: [Coniribution to our Knowledge of Ascomyetes of India. XIII.
264-267](#)