

A new species of *Physalospora* from India.

By A. K. Roy *)

(Division of Mycology, Department of Agriculture, Assam, Jorhat, India)

With one text-figure

Alternantheria sessilis Br. (fam. *Amaranthaceae*) commonly known as 'matikaduri' or 'senchi' is used for culinary and medicinal purposes by the people of Assam. From 1964 onwards during the rainy season the author observed a leaf spot on it, the incitant of which seem to be an undescribed species of *Physalospora* Niessl (Saccardo, 1882). This genus is represented in India by another 15 species (Saccardo, l. c.; Subramanian and Ramakrishnan, 1956; Butler and Bisby, 1960) including one on *Achyranthus*, belonging to the same family *Amaranthaceae* (Ramakrishnan, 1951).

Symptoms appear as minute diffuse purple spots which gradually increase in size but usually do not extend beyond 4 mm. and well demarcated at the margins. Spots may be a few or numerous per leaf blade and in severe case of spotting the tissues become yellow and the leaves fall off. In the middle of the spots black dot like minute perithecia develop which are partially immersed in the host tissues. They arise solitary or in groups. The following is a brief description of the fungus.

Physalospora alternantherae A. K. Roy, sp. nov.

Maculae primum minutae, punctiformes, paulatim accrescentes, orbicularares purpureae, zonula lutescente marginata, usque 4 mm diam., poste in centro canescentes; perithecia amphigena sed plerumque epiphylla, innata, solitaria vel 2—3 plus minusve aggregata, globosa, ostiolata, 78—132 μ diam.; pariete brunneo, pseudoparenchymatico; asci clavati, breviter stipitati, 8-spori, 30—60 \times 6.6—13 μ ; sporae ovoideae vel oblongae, inaequaliter distichae, continuae, 8.5—14 \times 4—7 μ , hyalinae, plasmate granuloso farctae; paraphyses tenerimae, 28—35 \times 2.5—4.2 μ .

Spots first appear as minute dots on the leaves which gradually increase in size, globose, purple, generally with yellowish halo at the begining, upto 4 mm. diameter; centre becoming greyish on which very minute black dot like perithecia develop. Perithecia ampigenous but

*) Present address: School of Agriculture, University of Edinburgh, West Mains Road, Edinburgh-9, Scotland, U. K.

mostly epiphyllous, innate erumpent, solitary or occasionally 2 to 3 in groups, globose, ostiolate, brown, pseudoparenchymatous, 78—132 μ diameter; asci, clavate, short stalked, octosporous, packed closely, 30—60 \times 6.6—13 μ ; ascospores hyaline, oval to oblong, single celled, granular, arranged irregularly biseriate, 8.5—14 \times 4—7 μ ; paraphyses hyaline, delicate, 28—35 \times 2.5—4.2 μ .

On the leaves of *Alternanthera sessilis* Br., Borbheta, Assam, 27. 7. '65, leg. A. K. Roy, deposited in the Herbarium Cryptogamiae Indiae Orientalis, New Delhi, accession No. 29150.

This fungus to some extent is akin to *P. cynodontis* G. Delacroix occurring on *Cynodon dactylon* Pers. (Saccardo, 1891) and *P. mimosae* Rehm. on *Mimosa* sp. (Saccardo, 1902) in the measurement of ascii and ascospores but disagrees in the size of perithecia and some other characters. *Laestadia cephalariae* (Awd.) Sacc. var. *alternantherae* Sacc. recorded on the same host agrees much with the present fungus but on the nature of a paraphysate character of perithecia, this genus is separated from *Physalospora* (Saccardo, 1882). Clements and Shear (1954) in making *Laestadia* Auersw. a synonym of *Phomatospora* Sacc. have also emphasized on this character. Ramakrishnan (l. c.) described *P. achyranthis* Ramakr. on *Achyranthus aspera* L. from Kodaikanal Hill which is differentiated from the present fungus on the following characters.

Fungus	Leaf spot	Perithecia	Asci	Ascospores
<i>P. achyranthis</i>	Indefinite	upto 200 μ	80—95 \times 21—26 μ	17—22 \times 6—10 μ
<i>P. alternantherae</i>	Definite	78—132 μ	30—60 \times 6.6—13 μ	8.5—14 \times 4—6 μ

Acknowledgement

Thanks are due to Dr. T. S. Ramakrishnan Director Institute for Advancement of Science, Manipal, South Kanara, for kindly supplying the literature on *P. achyranthis* and to Dr. F. Petrank for rendering Latin diagnosis of the species.

References

- Butler, E. J. and Bisby, G. R. 1960. The Fungi of India. Revised ed., I. C. A. R., New Delhi: 63—64.
- Clements, E. and Shear, L. 1954. The Genera of Fungi. Hafner Publishing Co., New York: 62 and 260.
- Ramakrishnan, T. S. 1951. Additions to the Fungi of Madras-XI, Proc. Indian Acad. Sci., B, 34: 161.
- Saccardo, P. A. 1882. Sylloge Fungorum, 1: 420 and 446.
— 1891. Ibid., 9: 599.
— 1899. Ibid., 14: 518 and 522.
— 1902. Ibid., 16: 457—458.
- Subramanian, C. V. and Ramakrishnan, K. 1956. List of Indian Fungi, 1952—56, J. Madras Univ., B, 26: 364.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1968/1969

Band/Volume: [22](#)

Autor(en)/Author(s): Roy A. K.

Artikel/Article: [A new species of Physalospora from India. - Addendum. 183-184](#)