/erlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.

## **Revision of Keissleriella podocarpi BUTIN**

H. A. VAN DER AA

Centraalbureau voor Schimmelcultures, P. O. B. 273, 3740 AG Baarn, Netherlands

On a collection of dead needles of *Podacarpus drouyniana* collected by W. GAMS & al. near Pemberton, W. Australia, a unitunicate pyrenomycete with phaeophragmospores was found, which could be identified as a species of *Lepteutypa*. The same needles contained numerous conidiomata of a *Pestalotiopsis* and repeated isolations from single ascospores proved the connection between these states.

Pestalotiopsis STEYAERT (= Pestalotia ss. GUBA, 1961; VON ARX, 1981, pro parte) is well known as anamorph of several species of Lepteutypa PETRAK (= Pestalosphaeria BARR, 1975); the present species could not be identified with any of them, but only with Keissleriella podocarpi BUTIN (1975), described from Podocarpus nubigenus in Chile. In the description of the latter, no attention was given to the structure of the ascus wall, but investigation of the holotype specimen (ZT) confirmed the supposition that the fungus also had a unitunicate ascus with amyloid apical structure, similar to the West Australian collection. BUTIN (l. c.) already hesitated, whether to classify the fungus in Keissleriella and the present observations, especially concerning the ascus wall, demonstrate that this species has to be excluded from Keissleriella, a genus of the Pleosporaceae (Dothideales), characterized by bitunicate asci without amyloid structures in the ascus tip (VON ARX & MULLER, 1975; SIVANESAN, 1984).

For these reasons the fungus is here reclassified and an emended description is provided for the teleomorph, completed with a description of the anamorph, in vivo and in vitro.

Lepteutypa podocarpi (BUTIN) VAN DER AA, comb. nov. – Figs. 1–3.

Basionym: Keissleriella podocarpi BUTIN – Sydowia 27: 273. 1973. Anamorph: Pestalotiopsis sp. – Fig. 3.

Perithecia deeply immersed in the host tissue, subglobose or

conical, usually with a flattened base,  $250-350\,\mu m$  diam,  $200-250\,\mu m$  high, erumpent through the covering layers of the host

1 Sydowia, Vol. XXXIX, 1986

/erlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum



Fig. 1: Lepteutypa podocarpi, holotypus (ZT); ascoma.

tissue, with a cylindrical or conical papilla, which is up to 100  $\mu m$  long, vaulting the epidermis conspicuously and visible as a contrasting white spot, before opening by a central, 10–15(–25) um wide pore. – Perithecial wall 15–25(–35)  $\mu m$  thick, in the apical part of the perithecium of irregular thickness, in the lower parts regularly 15–20  $\mu m$  thick and composed of 4–8(–10) layers with isodiametrical, locally flattened, pale olivaceous or brownish, thin-walled cells, 5–18  $\mu m$  in size. In the apical part of the perithecium and inside the papilla the wall is covered with thin-walled, hyaline or subhyaline, 1.5–2.0  $\mu m$  wide periphyses, with about 10  $\mu m$  long free ends, growing in the ostiolar cavity. In the basal part the wall merges into a tissue of thin-walled, isodiametrical cells, 4–10  $\mu m$  diam, in young ascomata forming several layers, in ripe ascomata strongly reduced and partly disappearing. This tissue bears the numerous asci and paraphyses.

Paraphyses numerous, thin-walled, hyaline, repeatedly septate, occasionally branched in the lower half,  $1.5-5(-7) \ \mu m$  wide. – Asci cylindrical,  $(60-)85-100(-110) \times 8-10 \ \mu m$ , unitunicate, usually

2

Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.



Fig. 2: Lepteutypa podocarpi, holotypus (ZT); a. ascus. - b. ascospores.

with a short, thick stalk, apically rounded and with a thickened wall provided with a distinct,  $3-3.5~\mu m$  wide, amyloid ring, containing 8 ascospores, uni- or indistinctly biseriate. – Ascospores ellipsoid or slightly fusiform, seldom somewhat curved, ends usually rounded, seldom somewhat tapering or even truncate, 2-euseptate, not or only

1\*

3

/erlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum



Fig. 3: Lepteutypa podocarpi, Pestalotiopsis anamorph in vivo (H. A. VAN DER AA 8974, CBS); a. conidioma. – b. developing conidia. – c. conidia.

slightly constricted at the septa, at first pale yellowish brown, later dark brown or with some shade of red-brown, the 3 cells homogeneously coloured or the central cell distinctly darker than the distal ones, the whole spores  $16-18.6(-21) \times 6.5-7.5(-8) \mu m$ .

Conidiomata in vivo at first pycnidial, subconical or irregular in shape, with a flattened base and a subconical apex which is closely connected with the covering host tissue, often very irregular in outline and intermixed with surrounding host cells, without preformed ostiolar structures, soon opening by fissure of the upper part and becoming acervular, up to 550 µm diam and 180 µm high. -Wall of variable thickness due to the close connection with the surrounding irregular host tissue, often about 3–5 cells (10–15 µm) thick and composed of thin-walled, pale olivaceous or hyaline cells, 5-12(-18) µm in size. - Conidiophores lining the basal and lateral parts of the conidiomata, the basal cells usually not differentiated from the inner wall cells. Conidiogenous cells short conical or cylindrical, even thread-like,  $4-22 \times 2-5 \mu m$ , integrated or discrete, holoblastic, annellidic. - Conidia fusiform, smooth, 4-euseptate,  $(30-)32-38(-40) \times 6.8-8.5$  µm, basal cell almost hvaline but strongly cvanophilic, after liberation developing an endogenous, cellular, simple, 2.5-5(-8.5) µm long appendage; the central 3 cells with somewhat thicker walls, second cell from below pale yellowish brown to olivaceous brown, the central cell the darkest, hazel to umber brown, the sub-apical cell olivaceous brown and the apical cell hyaline, not cyanophilic, tapering into an up to 20 µm long, hyaline, cellular, usually bent appendage, bearing at the convex side (3-)4-7(-9) branches which are hyaline, 10-18 µm long and nonspathulate.

Cultural characters. — Isolates from single ascospores and from single conidia similar. — Colonies on cornmeal agar spreading rapidly, attaining a diameter of 7—8 cm in 8 days at 240 C under near UV. Aerial mycelium white, becoming slightly greyish, typically zonate: Well developed and floccose in concentric rings, alternating with zones with scarce aerial mycelium and abundant conidiomata. Reverse of the colony at first slightly yellowish or faintly greenish, soon turning greyish red or brownish orange.

Conidiomata abundant, formed in concentric rings, with blackish exudate of conidia from about the 6th day onward. Conidiomata at first completely closed, subspherical, lens-shaped or in irregular flat crusts, up to  $1200 \,\mu\text{m}$  diam, covered by an upper wall, 1-5, laterally up to  $12 \,\text{cells}$  thick, pseudoparenchymatous or prosenchymatous and composed of thinwalled, pale olivaceous cells,  $5-12 \,\mu\text{m}$  diam. No ostiole formed, but sometimes opening by fissure at maturity in the thinner central part, sometimes with two distinct valves but usually more irregularly. Morphology of the conidia as on the host plant.

Material examined. – Keissleriella podocarpi BUTIN, on needles of Podocarpus nubigenus, Chile, Tres Chiflones, La Unión, 600 m alt., coll. BUTIN, 29 August 1968; holotypus containing teleomorph only, (ZT). – On needles of Podocarpus drouyniana, Beedlup National Park near Pemberton, W. Australia; coll. GAMS, DIXON & MCCREDE, 4 August 1983 (= VAN DER AA 8974, CBS, anamorph and teleomorph, and single-ascospore isolate CBS 113.84).

Discussion. – The teleomorph is a typical species of *Lepteuty*pa PETRAK, according to the somewhat extended concept given by VON ARX (1981), who also listed further synonyms. Recent descriptions were given by SWART (1973) and BARR (1975, sub Pestalosphaeria) and NAG RAJ (1985 b, sub Pestalosphaeria). Most of the ten species described in this genus differ from L. podocarpi in perithecial morphology and the number of septa of the ascospores. L. concentrica (BARR) VON ARX has 3-celled ascospores but with striated exosporium, whereas the conidial state, Pestalotiopsis quepini (DESM.) STEYAERT Var. macrotricha (KLEBAHN) SUTTON, is quite different from the anamorph of L. podocarpi. L. elaeidis (BOOTH & ROBERTSON) VON ARX has also some similarity in the teleomorph, but the Pestalotiopsis anamorph has only 2-3 apical appendages, which are distincly spathulate. According to the description the type species Lepteutypa fuckelii (NITSCHKE) PETRAK has three septate ascospores (Petrak, 1923; Shoemaker & Müller, 1965). Other species also resemble the teleomorph but have quite different anamorphs belonging to the genera Seiridium NEES: FR. or Labridella BRENCLE (sub Labridiella in von Arx, 1981).

Amongst hundreds of *Pestalotiopsis* species, merely described as *Pestalotia* (STEYAERT, 1948 a and b, 1949, 1961; GUBA, 1961; SUTTON, 1961; NAG RAJ, 1985 a and b) there is no species with this combination of measurements and typical apical cell with lateral branches. *Pestalotia heterocornis* GUBA, *P. multiseta* (SPEG.) GUBA and *P. thujae* SAWADA have some similarity but can be differentiated on the base of the number of appendages, their length and branching pattern.

## References

- ARX, J. A. VON (1981). The genera of fungi sporulating in pure culture. J. Cramer, Vaduz, 424 pp.
  - & MULLER, E. (1975). A re-evaluation of the bitunicate ascomycetes with keys to families and genera. - Stud. Mycol. 9: 159 pp.
- BARR, M. (1975). Pestalosphaeria, a new genus in the Amphisphaeriaceae. Mycologia 67: 187–194.

BUTIN, H. (1975). Beitrag zur Ascomyzetenflora von Chile. – Sydowia 27: 267–292 ("1973/74").

GUBA, E. F. (1961). Monograph of Monochaetia and Pestalotia. – Harvard University Press, Cambridge, Massachusetts, 342 pp.

NAG RAJ, T. R. (1985 a). Redisposals and redescriptions in the Monochaetia-Seiridium, Pestalotia – Pestalotiopsis complexes. I. The correct name for the type species of Pestalotiopsis. Mycotaxon 22: 43-51.

 (1985 b). Redisposals and redescriptions in the Monochaetia-Seiridium, Pestalotia-Pestalotiopsis complexes. II. Pestalotiopsis besseyii (GUBA) comb. nov. and Pestalosphaeria varia sp. nov. – Mycotaxon 22: 52–63.

PETRAK, F. (1923): Mykologische Notizen VI. – Annls mycol. 21: 182-335.

SHOEMAKER, R. A. & MULLER, E. (1965). Types of the pyrenomycete genera Hymenopleella and Lepteutypa. - Can. J. Bot. 43: 1457-1460.

SIVANESAN, A. (1984). The bitunicate Ascomycetes and their anamorphs. – J. Cramer, Vaduz, 701 pp.

STEYAERT, R. L. (1948 a). Quelques *Pestalotia* de la flore Belge. – Bull. Jard. bot. Etat Brux. 19: 65–72.

 (1948 b). Contribution à l'étude des *Pestalotia* du Congo Belge. – Bull. Jard. bot. Etat. Brux. 19: 173–186.

 (1949). Contribution à l'étude monographique de Pestalotia de Nor. et Monochaetia SACC. (Truncatella gen. nov. et Pestalotiopsis gen. nov.). – Bull. Jard. bot. Etat Brux. 19: 285–354.

 (1961). Type specimenes of SPEGAZZINI's collections in the "Pestalotiopsis" and related genera (Fungi imperfecti, "Melanconiales"). – Darwiniana 12: 157–190.

SUTTON, B. C. (1961). Coelomycetes I. – Mycol. Pap. 80: 16 pp.

SWART, H. J. (1973). The fungus causing cypress canker. – Trans. Br. mycol. Soc. 61: 71–82.

## ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Sydowia

Jahr/Year: 1986/1987

Band/Volume: 39

Autor(en)/Author(s): Aa H. A., van der

Artikel/Article: Revision of Keissleriella podocarpi BUTIN. 1-7