Wuestneia molokaiensis and its anamorph Harknessia molokaiensis sp. nov. from Eucalyptus

Pedro W. Crous¹ & Jack D. Rogers²

¹ Department of Plant Pathology, University of Stellenbosch, P. Bag X1, Matieland 7602, South Africa ² Department of Plant Pathology, Washington State University, Pullman,

WA 99164-6430, USA

P. W. Crous & J. D. Rogers (2001). Wuestneia molokaiensis and its Harknessia anamorph sp. nov. from Eucalyptus. – Sydowia 53(1): 74–80.

Wuestneia molokaiensis is newly described from Eucalyptus robusta twigs collected in Hawaii. Single ascospore strains produced a Harknessia anamorph and microconidial state in culture. W. molokaiensis and H. molokaiensis are compared with previously described species in these genera and described as new. W. molokaiensis is distinguished based on its smaller ascospores with persistent mucous sheaths, and H. molokaiensis based on its smaller, ellipsoidal conidia.

Keywords: Eucalyptus, Harknessia, leaf spot, stem canker.

The genus *Harknessia* Cooke has been the subject of several recent monographs (Sutton, 1971; Nag Raj & DiCosmo, 1981; Nag Raj, 1993), with the latter treatment recognising 26 species. Since then 9 additional species have been described (Crous & al., 1993; Furlanetto & Dianese, 1998; Swart & al., 1998; Yuan & al., 2000). Teleomorphs have been described in *Wuestneia* Auersw. (Reid & Booth, 1989; Sutton & Pascoe, 1989). The first cultural confirmation of this relationship was made by Crous & al. (1993), and later by Yuan & Mohammed (1997), bringing the total number of accepted taxa in *Wuestneia* up to 12 species and one variety.

Although not many data are available relating to their pathogenicity, species of *Harknessia* are commonly associated with leaf spots, as well as stem and branch cankers (Crous & al., 1989, 1993; Yuan & Mohammed, 1997; Yuan & al., 2000). Preliminary results from stem inoculations on eucalypts with *H. eucalypti* Cooke apud Cooke & Harkn. by Yuan & al. (2000) failed to induce lesions, whereas leaf inoculations with *H. hawaiiensis* F. Stevens & E. Young on 12 different eucalypt species produced prominent lesions on some species, and none on others (Crous, unpubl. data). These observations, as well as the fact that *Harknessia* spp. are most commonly encountered on lesions with other primary pathogens,

suggest that the majority of these species are possibly secondary or weak pathogens.

The present paper results from a collection of an undescribed *Wuestneia* species and its *Harknessia* anamorph from *Eucalyptus*. The aim of this study was thus to describe these taxa, and to compare them with others known in these genera.

Materials and methods

Perithecia of a Wuestneia sp. obtained from Eucalyptus twigs were dissected in sterile water on plates containing 2% malt extract agar (MEA) (Biolab, Midrand, Johannesburg, South Africa). Plates were inverted, and incubated at 25 C in the dark. Single germinating ascospores were transferred to fresh MEA plates, and later to divided plates containing MEA, and carnation leaf agar (CLA) (Fisher & al., 1982; Crous & al., 1992). Plates were incubated at 25 C under continuous near-ultraviolet light to enhance sporulation. Wherever possible, thirty measurements were made of structures mounted in clear lactophenol, and the averages determined. Colony colours (top and bottom) were rated after 14 d on MEA at 25 C in the dark, using the colour charts of Rayner (1970). The holotype specimen was lodged at the Bishop Museum, Honolulu (BISH), and isotypes at the Dept. of Plant Pathology, Washington State University, Pullman (WSP), and the National Collection of Fungi in Pretoria, South Africa (PREM), and ex-type cultures maintained in the culture collection of the Department of Plant Pathology at the University of Stellenbosch, South Africa (STE-U), and at the Centraalbureau voor Schimmelcultures (CBS) in the Netherlands.

Results and discussion

Wuestneia molokaiensis Crous & J. D. Rogers, sp. nov. – Figs. 1, 4–8.

Anamorph: *Harknessia molokaiensis* Crous & J. D. Rogers, sp. nov.

Perithecia solitaria, in ramis inclusa, collis emergentibus, ectostromate cellularum furfuracearum brunnearum vel fulvarum circumcincta, brunnea vel fusca, globosa vel subglobosa, 250–350 µm diam.; peridio 10–15 µm lato, 4–6 stratis cellularum pseudoparenchymatarum sectione texturae angularis praedito, superficie texturae epidermoideae composita; ostiola usque ad 60 µm lata. Asci clavati vel subcylindrici, 65–110(–130) µm longitudine tota × 11–17(–20) µm crassi, unitunicati, brevistipitati, annulo apicali distincto destituti, octo ascosporis uniseriatis vel biseriatis praediti. Ascosporae hyalinae, vagina irregulari 1–3 µm crassa circumcinctae, ellipsoideae, pariete crasso praeditae, intus granulosae, $14-22(-24) \times 5-8$ µm.



Figs. 1–3. – Wuestneia molokaiensis and its Harknessia molokaiensis anamorph (holotype). – 1. Asci and ascospores. – 2. Conidia and conidiophores. – 3. Microconidia and conidiogenous cells. – Bar = 10 μ m.

Harknessia molokaiensis Crous & J. D. Rogers, sp. nov. – Figs. 2, 3, 9–14.

Ab Harknessia eucalyptorum differt in macroconidiis 14–17×8–11 μm et in microconidiis 4–5×2 $\mu m.$

Etymology. – named after the collection locality, Molokai Island, Hawaii.

Ascomata caulicolous, perithecial, solitary, subepidermal with emergent necks, surrounded by an ectostromatic layer of furfuraceous brown to yellow-brown cells. - Perithecia medium to dark brown, globose to subglobose, 250–350 µm diam.; peridium 10–15 µm wide, consisting of 4-6 layers of medium brown pseudoparenchymatous cells of textura angularis (in section), of textura epidermoidea in surface view; ostiole up to 60 µm wide. - Asci clavate to subcylindrical, $65-110(-130) \times 11-17(-20) \mu m$, unitunicate, short-stipitate, lacking a well-defined apical apparatus, containing 8, uni- to biseriate ascospores. – Ascospores hyaline, surrounded by an irregular sheath, 1–3 µm wide, ellipsoid, thick-walled, granular, $14-22(-24) \times 5-8 \ \mu\text{m}$ ($\bar{x} = 18 \times 6 \ \mu\text{m}$). - Conidiomata produced on MEA in culture, separate, scattered, pycnidial, unilocular, globose to subglobose, 300-450 µm diam.; walls of 5-7 layers of medium brown textura angularis, 20-30 µm diam. -Conidiophores lining conidiomatal cavity, reduced to conidiogenous cells. - Conidiogenous cells ampulliform to lageniform, hyaline, smooth, formed from inner layer of the conidiomatal wall, proliferating once or twice percurrently, $7-10 \times 4-6$ µm. – Macroconidia ellipsoid to ovoid with a truncate base, medium brown, aseptate, apex obtuse to bluntly apiculate, smooth, except for longitudinal striations in restricted areas on some conidia, with a lighter central area consisting of small, aggregated droplets, $14-17 \times 8-11$ ($\bar{x} = 15 \times 10 \mu m$); basal appendage hyaline, unbranched, frequently surrounded by a mucous sheath. 5–17×2–3 μm. – Microconidiogenous cells in same or separate conidioma, subcylindrical, hyaline, smooth, with minute apical periclinal thickening, $5-6 \times 1.5-2 \mu m$. – Microconidia hyaline, smooth, aseptate, limoniform to ellipsoid or fusoid with obtuse apex and truncate base, $4-5 \times 2$ µm.

Cultural characteristics. – Colonies white to pale luteous (19f) (surface), and pale luteous to luteous (19f–17b) (bottom), sienna in the centre (13i), with feathery margins and moderate, fluffy, off-white aerial mycelium after 14 d at 25 $^{\circ}$ C in the dark.

Host. – Eucalyptus robusta Sm. (Myrtaceae).



Figs. 4–14. – Wuestneia molokaiensis and its Harknessia molokaiensis anamorph (holotype). – 4, 5. Asci. – 6. Ascospore with mucous sheath (arrow). – 7. Vertical section through pseudothecial wall. – 8. Pseudothecial wall in surface view. – 9, 10. Conidiogenous cells with persistent sheath covering conidium and appendage up to conidiogenous locus (arrows). – 11. Microconidia. – 12. Macroconidium with mucous sheath enclosing basal appendage (arrow). – 13. Macroconidium with striations (arrow). – 14. Macroconidia with basal appendage. – Bars = 10 μm.

Distribution. – U.S.A. (Molokai Island, Hawaii).

Specimen examined. – U.S.A., Hawaii, Molokai Island, Kamacou Forest Reserve, dead branches of *E. robusta*, J. D. Rogers, 9 Jan. 2000, holotypes in BISH, isotypes in WSP and PREM, culture ex-type STE-U 3797.

Wuestneia molokaiensis can be compared with W. karvarrae (B. Sutton & I. Pascoe) Z.Q. Yuan, W. eucalyptorum Crous, M. J. Wingf. & Nag Raj and W. epispora Z.Q. Yuan. Ascospores of W. molokaiensis are surrounded by a mucilaginous sheath, as are those of W. epispora, which was also described from eucalypt branches collected in Tasmania (Yuan & Mohammed, 1997). In W. epispora, however, the ascospores are larger $(20-35 \times 9-15 \ \mu m)$, and the sheath more prominent than in W. molokaiensis. Ascospores of both W. eucalyptorum $(13-28 \times 13-20 \ \mu m;$ Crous & al., 1993) and W. karvarrae $(23-27 \times 9-11 \ \mu m;$ Sutton & Pascoe, 1989) are without any sheath, and are also larger than those of W. molokaiensis $(14-22 \times 5-8 \ \mu m)$.

Several species of Harknessia are comparable with H. molokaiensis based on macroconidium shape and appendage length. Of these species, however, H. molokaiensis has smaller conidia $(14-17 \times 8-11 \ \mu\text{m})$ than *H. eucalypti* Cooke $(19-28 \times 11-1)$ 15 μ m), H. eucalyptorum Crous, M. J. Wingf. & Nag Raj (16–29 \times 9– 14 μ m), *H. karvarrae* B. Sutton & I. Pascoe (17–18 × 12–12.5 μ m) and the *Harknessia* anamorph of *W. epispora* $(18-35 \times 12-16 \mu m)$ (Sutton & Pascoe, 1989; Crous & al., 1993; Nag Raj, 1993; Yuan & Mohammed, 1997). Based on macroconidium dimensions, therefore, H. molokaiensis is easily distinguishable from other species of Harknessia.

References

- Crous, P. W., P. S. Knox-Davies & M. J. Wingfield (1989). A summary of fungal leaf pathogens of *Eucalyptus* and the diseases they cause in South Africa. S. Afr. For. J. 149: 9–16.
- —, A. J. L. Phillips & M. J. Wingfield (1992). Effects of cultural conditions on vesicle and conidium morphology in species of *Cylindrocladium* and *Cylindrocladiella*. – Mycologia 84: 497–504.
- —, M. J. Wingfield & T. R. Nag Raj. (1993). Harknessia species occurring in South Africa. – Mycologia 85: 108–118.
- Fisher, N. L., L. W. Burgess, T. A. Toussoun & P. E. Nelson (1982). Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. – Phytopathology 72: 151–153.
- Furlanetto, C. & J. C. Dianese (1988). Some coelomycetes from central Brazil. Mycol. Res. 102: 19–29.
- Nag Raj, T. R. (1993). Coelomycetous anamorphs with appendage-bearing conidia. Mycologue Publications, Waterloo, Canada.
- —, & F. Di Cosmo (1981). A monograph of *Harknessia* and *Mastigosporella* with notes on associated teleomorphs. – Bibl. Mycol. 80: 1–62.

Rayner, R. W. (1970). A mycological colour chart. – CMI and British Mycological Society. Kew, Surrey, England.

Reid, J. & C. Booth (1989). On *Cryptosporella* and *Wuestneia*. – Can. J. Bot. 67: 879–908.

Sutton, B. C. (1971). Coelomycetes. IV. The genus Harknessia, and similar fungi on Eucalyptus. – Mycol. Pap. 123: 1–46.

- —, & I. G. Pascoe (1989). Addenda to Harknessia (Coelomycetes). Mycol. Res. 92: 431–439.
- Swart, L., P. W. Crous, S. Denman & M. E. Palm (1998). Fungi occurring on Proteaceae. I.– S. Afr. J. Bot. 64: 137–145.
- Yuan, Z. Q. & C. Mohammed (1997). Wuestneia epispora sp. nov. on stems of eucalypts from Australia. – Mycol. Res. 101: 196–200.
- —, T. Wardlaw & C. Mohammed (2000). Harknessia species occurring on eucalypt leaves in Tasmania, Australia. – Mycol. Res. 104: 888–892.

(Manuscript accepted 23rd December 2000)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Sydowia

Jahr/Year: 2001

Band/Volume: 53

Autor(en)/Author(s): Crous Pedro W., Rogers Jack D.

Artikel/Article: <u>Wuestneia molokaiensis and ist anamorph Harknessia</u> molokaiensis sp. nov. from Eucalyptus. 74-80