

Two new species of *Hypoxylon* from China*

H.-X. Ma^{1,2**}, L. N. Vasilyeva³ & Y. Li²

¹ Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, People's Republic of China

² Institute of Mycology, Jilin Agricultural University, Changchun 130118, People's Republic of China

³ Institute of Biology and Soil Science, Far East Branch of the Russian Academy of Sciences, Vladivostok 690022, Russia

Ma H.-X., Vasilyeva L. N. & Li Y. (2011) Two new species of *Hypoxylon* from China. – *Sydowia* 64 (1): 103–108.

Hypoxylon hubeiense and *Hypoxylon yunnanense* (Xylariales, Xylariaceae) are described as new species from China. Morphological descriptions and photographs of stromata and microstructures are provided based on the Chinese material, and the delimitation of these species from similar taxa is discussed.

Keywords: Ascomycota, pyrenomycetous fungi, taxonomy.

Hypoxylon Bull. is a large and cosmopolitan genus, which has its highest diversity in the tropics and subtropics (Martín 1968 a, b; Ju & Tzean 1985; Van der Gucht et al. 1997; Romero & Hladki 2009). The genus *Hypoxylon* has been studied by many mycologists (Miller 1928, 1932; Martín 1967, 1968 a, b; Whalley 1977, 1993; Rogers & Candoussau 1980, 1982; Rogers 1985; Rogers & Samuels 1985; Granmo 1999) and two world monographs were published (Miller 1961; Ju & Rogers 1996). The genus is still poorly known in China, although Tai (1979) listed 35 species most of which (33) were previously reported by Teng (1963). Later, Abe & Liu (1995) found eight species of *Hypoxylon* in Zhejiang province, but the species diversity of *Hypoxylon* in China needs further investigations. This paper reports two new species of *Hypoxylon*, namely, *H. hubeiense* and *H. yunnanense*. Illustrations and descriptions of these species are provided.

Materials and methods

The studied specimens are deposited at the Herbarium of Mycology of Jilin Agricultural University (HMJAU). Microscopic features and measurements were made from slide preparations mounted in water, 10 % KOH and Melzer's iodine reagent. The photographs of asci, ascus apical ring, and ascospores were taken by using a VHX-600E microscope of the Keyence Cor-

* *Hypoxylon* from China 1

** e-mail: mahaixia0429@163.com

poration. The photographs of stromatal surface were taken with a ZSA30w microscope and S70 Canon camera. External stromatal colors were recorded and coded after Rayner (1970). The methods of collecting, preservation, and identification of the examined specimens follow those of Ju and Rogers (1996).

Taxonomy

Hypoxylon hubeiense Hai-Xia Ma, Lar. N.Vassiljeva & Yu Li, sp. nov. – Fig. 1. MycoBank no.: MB 563692

A Hypoxylo rickii in perisporio sine ornamentis spiralibus inconspicuis, in ascosporis rima germinativa stricta praeditis et in peritheciiis globosis vel obovoideis differt.

Holotypus. – CHINA, Hubei Province, Suizhou City, on the bark of fallen branch, 21 Nov 2009, *leg.* Haixia Ma, HMJAU 20819.

Stromata pulvinate to effused-pulvinate, with inconspicuous or slightly conspicuous perithecial mounds, 0.3–1.5 cm long \times 0.2–0.8 cm wide \times 0.5–1.5 mm thick; surface brick (59); orange granules immediately beneath surface and between perithecia, with KOH-extractable pigments orange (7); the tissue below the perithecial layer black brown, inconspicuous to 0.9 mm thick. Perithecia spherical to ovoid, 0.4–0.5 mm in diam. \times 0.5–0.6 mm high. Ostioles lower than the stromatal surface. Asci 41–48 \times 4.5–6 μ m in the spore-bearing portions, the stipes 40–70 μ m long, with apical ring lightly bluing in Melzer's iodine reagent, discoid, 0.5–0.8 μ m high \times 1.2–1.8 μ m wide. Ascospores brown, unicellular, ellipsoid, most inequilateral, infrequently equilateral, with broadly rounded ends, 6.5–8.5 \times 4–5 μ m, with straight germ slit slightly less than to nearly spore-length; perispore dehiscent in 10 % KOH.

Etymology. – Refers to the type locality Hubei.

Habitat. – The fungus grows on the bark of fallen branches in mixed evergreen and deciduous broadleaf forests at about 600 m altitude in a sub-tropical monsoon climate.

Distribution. – Known only from the type locality.

Remarks. – According to the key by Ju & Rogers (199&9, *Hypoxylon hubeiense* belongs to group D and subgroup 1. It is very similar to *Hypoxylon rickii* M. Ju & J. D. Rogers in stromatal morphology and color, but the latter has an inconspicuous coil-like ornamentation on the perispore. In addition, *H. rickii* has a slightly sigmoid germ slit in spore-length, whereas *H. hubeiense* has a straight germ slit slightly less than to nearly spore-length. Furthermore, *H. hubeiense* could be separated from *H. rickii* by the perithecial shape (spherical to ovoid versus tubular). It differs from *H. subgilvum* var. *microsporum* (Y. Abe) Y.M. Ju & J.D. Rogers by larger ascospores and larger perithecia.

Hypoxylon yunnanense Hai-Xia Ma, Lar. N.Vassiljeva & Yu Li, sp. nov. – Fig. 2. MycoBank no.: MB 564008

A Hypoxylo dieckmannii in ascosporis (9,5)10,5–11,5(12,5) \times 4,5–6 μ m differt; *a Hypoxylo hughesii* in superficie stromatica albomaculata et in peritheciiis obovoideis 0,1–0,3 \times 0,2–0,4 mm differt.




Fig.1. – *Hypoxylon hubeiense*, holotype. **a.** Stroma, bar 2.5 mm. **b.** Stromatal surface, bar 0.8 mm. **c.** Ascospores, bar: 6 μ m. **d.** Asci, bar: 42 μ m.

Holotypus. – CHINA, Yunnan Province, Yeya Lake, on the bark of a fallen branch, 1 Sep 2010, leg. Haixia Ma, HMJAU 22605.

Stromata effused-pulvinate, with inconspicuous perithecial mounds, 0.5–2.5 cm long \times 0.4–1.5 cm wide \times 0.5–1 mm thick; surface brown vinaceous (84), dark brown granules immediately beneath surface and between perithecia, without apparent KOH-extractable pigments; the tissue below the perithecial layer black, inconspicuous to up to 0.6 μ m. Perithecia obovoid, 0.1–0.3 mm in diam. \times 0.2–0.4 mm high. Ostioles lower than the stromatal surface. Asci 130–150 μ m total length \times 6–7.5 μ m wide, the spore-bearing parts 68–80 μ m long, with apical ring bluing in Melzer's iodine reagent, discoid, 0.5–0.8 μ m high \times 1.2–1.4 μ m wide. Ascospores light brown to brown, unicellular, ellipsoid-equilateral, with usually narrowly rounded ends, often almost acute, (9.5)10.5–11.5(12.5) \times 4.5–6 μ m, with straight germ slit slightly less than spore-length; perispore indehiscent, or very infrequently dehiscent, in 10 % KOH.

Etymology. – Refers to the type locality Yunnan.

Habitat. – The fungus grows on the bark of fallen branches in mixed evergreen and deciduous broadleaf forests at about 2100 m altitude in a sub-tropical climate.

Distribution. – Known only from the type locality.




Fig. 2. – *Hypoxyylon yunnanense*, holotype. **a.** Stroma, bar: 3 mm. **b.** Stromatal surface, bar: 0.8 mm. **c.** Asci, bar: 48 μ m. **d.** Ascospores, bar: 13 μ m. **e.** Germ slit, bar: 6.5 μ m. **f.** Ascus apical ring, bar: 13.5 μ m.

Remarks. – *Hypoxyylon yunnanense* is somewhat similar to *H. dieckmannii* Theiss in stromatal morphology. Miller (1961) indicated rather small ascospores for *H. dieckmannii* ($6-8 \times 3-3.8 \mu$ m) and considered it as a small-spored variety of *H. rubiginosum* (ascospores $9-12 \times 4-6 \mu$ m), but Martín (1969), Van der Gucht & Whalley (1992) and Ju & Rogers (1996) considered the two species to be distinctly different in ascospore morphology and KOH-extractable pigments of stromata. Ju & Rogers's (1996) indicated the wider range of ascospore size, $6.5-10(11) \times 3.5-4 \mu$ m, for *H. dieckmannii*, but the

Chinese specimen has even larger ascospores, $(9.5)10.5\text{--}11.5(12.5) \times 4.5\text{--}6\text{ }\mu\text{m}$, on average.

Hypoxyylon hughesii Y.-M. Ju & J. D. Rogers (Ju & Rogers 1996) has similar ascospore morphology and stromatal color, but it can be separated from *H. yunnanense* by the surface of the stromata, which has white spots in *H. yunnanense* (Fig. 2 b). In addition, there are differences in the perithecial shape and size. The perithecia of *H. hughesii* are spherical $0.1\text{--}0.2\text{ mm}$ in diam., whereas those in *H. yunnanense* are obovoid $0.1\text{--}0.3 \times 0.2\text{--}0.4\text{ mm}$. Furthermore, the apical ring of *H. hughesii* is larger ($1\text{--}1.5\text{ }\mu\text{m}$ high \times $2.5\text{--}3\text{ }\mu\text{m}$ wide).

Acknowledgements

This study was supported by the National Natural Science Foundation of China (grant no. 30770005) and the Start-up Costs for Doctors in Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences. We are also grateful to the Ministry of Agriculture of China for funding the field study in the project entitled "The Project of System Construction of Modern Agricultural Technology".

References

- Abe Y., Liu Z. (1995) An annotated list of xylariaceous and diatrypaceous fungi collected from Mt. Fengyangshan and Mt. Baishanzu, Zhejiang Prov. in East China. *Bulletin of National Science Museum, Tokyo, Series B*, **21**: 75–86.
- Granmo A. (1999) Morphotaxonomy and chorology of the genus *Hypoxyylon* (Xylariaceae) in Norway. – *Sommerfeltia* **26**: 1–81.
- Ju Y. M., Rogers J. D. (1996) *A revision of the genus Hypoxyylon*. American Phytopathological Society Press, St. Paul, Minnesota.
- Ju Y. M., Tzean S. S. (1985) Investigation of Xylariaceae in Taiwan I. The teleomorph of *Hypoxyylon*. *Transactions of the Mycological Society of the Republic of China* **1**: 13–27.
- Martin P. (1967) Studies in the Xylariaceae: II. *Rosellinia* and the *Primocinerea* Section of *Hypoxyylon*. *Journal of South African Botany* **33**: 315–328.
- Martin P. (1968 a) Studies in the Xylariaceae: III. South African and foreign species of *Hypoxyylon* Section *Entoleuca*. *Journal of South African Botany* **34**: 153–199.
- Martin P. (1968 b) Studies in the Xylariaceae IV. *Hypoxyylon*. Sections *Papillata* and *Anulata*. *Journal of South African Botany* **34**: 303–330.
- Martin P. (1969) Studies in the Xylariaceae: V. *Euhypoxyylon*. *Journal of South African Botany* **35**: 149–206.
- Miller J. H. (1928) Biologic studies in the Sphaeriales. II. *Mycologia* **20**: 205–239.
- Miller J. H. (1932) British Xylariaceae II. *Transactions of the British Mycological Society* **17**: 125–135.
- Miller J. H. (1961) *A monograph of the world species of Hypoxyylon*. University of Georgia Press, Athens, USA.
- Rayner R. W. (1970) *A mycological color chart*. Commonwealth Mycological Institute, Kew.
- Rogers J. D. (1985) *Hypoxyylon duranii* sp. nov. and the anamorphs of *H. caries*, *H. papillatum*, and *Rosellinia subiculata*. *Mycotaxon* **23**: 429–437.
- Rogers J. D., Candoussau F. (1980) A new variety of *Hypoxyylon cohaerens* from France. *Mycologia* **72**: 826–832.
- Rogers J. D., Candoussau F. (1982) *Hypoxyylon gillesii*, a new species with ornamented ascospores from Madagascar. *Mycotaxon* **15**: 507–514.

- Rogers J. D., Samuels G. J. (1985) New taxa of *Hypoxyylon*. *Mycotaxon* **22**: 367–373.
- Romero A. I., Hladki A. I. (2009) Taxonomic and nomenclatural aspects of *Hypoxyylon* taxa from southern South America proposed by Spegazzini. *Mycologia* **101**(5): 733–744.
- Tai F. L. (1979) *Sylloge fungorum Sinicorum*. Science Press, Peking.
- Teng S. Q. (1963) *Fungi of China*. Science Press, Peking.
- Van der Gucht K., Whalley A. J. S. (1992) A new variety and combination for *Hypoxyylon citrifforme*. *Mycological Research* **96**: 895–896.
- Van der Gucht K., Ju Y. M., Rogers J. D. (1997) New *Hypoxyylon* species from Papua New Guinea and notes on some other taxa. *Mycologia* **89**: 503–511.
- Whalley A. J. S. (1977) Key to the British species of *Hypoxyylon*. *Bulletin of the British Mycological Society* **11**: 45–47.
- Whalley A. J. S. (1993) Tropical Xylariaceae: their distribution and ecological characteristics. In: *Aspects of tropical mycology* (eds. Isaac, S., Frankland J. C., Watling R., Whalley A. J. S.). Cambridge University Press, Cambridge, UK: 103–119

(Manuscript accepted 30 Mar 2012; Corresponding Editor: I. Krisai-Greilhuber)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 2012

Band/Volume: [64](#)

Autor(en)/Author(s): Ma Hai-Xia, Vasilyeva Larissa N., Li Yu

Artikel/Article: [Two new species of Hypoxylon from China. 103-108](#)