© Hans Malicky/Austria; download unter www.biologiezentrum.at

T R I C H O P T E R O L O G I C A L L I T E R A T U R E

This list is intended to be as complete as possible. Any paper which contains any information on caddisflies will be included if there is at least one well-defined caddisfly taxon from which the identity of the species may be derived or reconstructed. - Please help to complete this list. If you know any publication which was not listed here, send a copy of it or at least the correct and complete reference to H.Malicky. It will be added in the next number of the Newsletter.

* *

1975 (additions)

Engster,M.S., Invertebrate secretory mechanisms, with special reference to silk secretion in the Trichoptera. - Diss. Abstr.int.(B) 35(11):5247.

Gallepp,G.W.,jr., The behavioural ecology of larval caddisflies, Brachycentrus americanus and Brachycentrus occidentalis. - Diss.Abstr.int.(B) 35(9):4532.

Givens,D.R., Biology and taxonomy of the western Arctopsychinae (Trichoptera;Hydropsychidae). - Proc.Wash.St.ent.Soc. 37:391-393.

Hilsenhoff,W.L., Aquatic insects of Wisconsin. - Tech.Bull. Dep.nat.Resour.Wisc. 39:1-52.

Murgoci,A., Trichoptera, pp.101-107. In: M.Ionescu(ed.), Fauna Ser.Monografica. Acad.Rep.Soc.Rom.,Porţile de Fier.1-316.

Olmi,M., Degradazione biotica e abiotica della risaia italiana e sue conseguenze sulla entomofauna. - Riso 24:373-382.

Shotton,F.W., Animal remains incorporated in caddis fly larval cases at Ufton Fields. - Proc.Coventry Distr.nat.Hist.sci. Soc. 4:283-286.

Sveshnikov,V.A., Faidysh,E.A., Filimanov,P.M., Utilization of the pain threshold to distinguish stages in the instinctive behaviour of insects. - Proc.Acad.Sci.USSR,Biol.Sci. 223(1-6):297-298.

Winner,R.W., Van Dyke,J.S., Caris,N., Farrel,M.P., Response of the macroinvertebrate fauna to a copper gradient in an experimentally-polluted stream. - Verh.IVL 19:2121-2127.

1976 (additions)

Brindle,A., Caddis-flies of Yorkshire. - Naturalist Leeds No. 937:47-50.

Davydova,E.D., Comparative histology of the retrocerebral complex in larvae of caddis-flies(Trichoptera). - Zool. Zh. 55:538-546.

Engster,M.S., Studies on silk secretion in the Trichoptera (F.Limnephilidae). 1.Histology, histochemistry, and ultrastructure of the silk glands. - J.Morph. 150:183-212.

Klausnitzer,B., Braasch,D., Jacob,U., Joost,W., Zimmermann,W., Gegenwärtiger Stand der faunistischen Erforschung der aquatischen Insektenordnungen in der DDR. - Ent.Nachr. (Dresden) 20:133-159.

Litt,R., Notes et observations sur Phryganea striata L. (Trichoptère-stégoptère). - Rev.verviét.Hist.nat. 33(4-6): 28-32, 33(7-9):44-47.

McDaniel,V.R., Smith,K.L., Cave fauna of Arkansas: selected invertebrate taxa. - Proc.Arkansas Acad.Sci. 30:57-60.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1976

Masteller,E.C., Cunningham,H.N.,jr., Leavers,D.R., Tucker,E.,jr. Biological, chemical and geological characteristics during August-September of Lake Erie tributaries of Erie County,P. - Proc.Pa.Acad.Sci. 50:45-59.

Matida,Y.,Kimura,S.,Tanaka,H.,Yokote,M., Effects of some herbicides applied in the forest to the freshwater fisher and other aquatic organisms. 3.Experiments on the assessment of acute toxicity of herbicides on aquatic organisms. - Bull.freshw.Fish.Res.Lab.Tokyo 26:79-84.

Moog,O., Christian,E., Erstnachweis einer cavernicolen Köcher= fliegen-Art für Österreich. - Höhle(Wien) 27:94.

Namuush,G., Pyelzhidmaa,S., On the study of Trichoptera in M.P.R.:52-53. In:Shagdar,D.(ed.): International Conference on research into the natural conditions and resources of the Mongolian People's Republic. Ulan Bator, 1-92 (Russian).

Poinar.G.O.,jr., Presence of Mermithidae (Nematoda) in inver= tebrate paratenic hosts. - J.Parasitol.62:843-844.

Resh,V.H., Life cycles of invertebrate predators of freshwater sponge. 299-314. In:Harrison,F.W.,Cowden,R.R.(eds.): Aspects of sponge biology. Acad.Press NY, 1-354.

Svensson,B.W., Ulfstrand,S., Biotopvalet som anpassningsmekanism hos vattendragens djursamhällen. - Fauna Flora(Stockholm) 1976,6:274-283.

Sykora,J., Walker,W., Vest,D., Weaver,J.S.III, Notes on the seasonal distribution and taxonomy of Pycnopsyche adults (Trichoptera:Limnephilidae). - Proc.Pa.Acad.Sci. 50:133-135.

Sykora,J.L.,Weaver,J.S.III, A new species of Rhyacophila (Trichoptera, Rhyacophilidae) from western Pennsylvania. Ann.Carnegie Mus. 46(3):29-32.

Tani,Kozo, The Caddisflies of Tsushima Islands(Trichoptera). - Biological of Tsushima Islands(BiolSoc.of Nagasaki Pref.), pp.391-396.(Japanese)

Tarter,D.C., Limnology in West Virginia. A lecture and laboratory manual. - Marshall Univ.Book Store,Huntingdon WV 25701, 249 pp., polycopied.

1977 (additions)

Schuster,G.A., Hamilton,S.W., Records of the Trichopteran families Hydroptilidae, Philopotamidae, and Psychomyiidae from Kansas. - New.records of the fauna and flora of Kansas for 1977. Eds. R.E.Brooks, R.L.McGregor. - Tech.Publ.State Biol.Surv.Kansas 6:36-47.

Solem,J.O., Sendstad,E., Bergvik,T., Hegstad,A., New record of Apatania zonella Zett.(Trichoptera,Limnephilidae) from Svalbard. - Norw.J.Ent. 24:85.

Tani,Kozo, Trichoptera. In:Coloured illustrations of the insects of Japan, vol.II.,Hoikusha, pp.184-206.(Japanese)

1978 (additions)

Cepić,V., Marinković-Gospodnetić,M., Zoobentos Rijeke Toplice. God.Biol.Inst.Univ.Sarajevu 31:25-32.

Chandler,C.M., A preliminary checklist of benthic macroinvertebrate families from basket sampling during early summer in the West Fork of Stone's River, Tennessee. - J.Tenn. Acad.Sci. 53:25-28.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1978

Marinković-Gospodnetić,M., The caddis-flies(Trichoptera,Insecta) of Hercegovina (Yugoslavia). - God.Biol.Inst.Univ.Sarajevu 31:115-131.
Moseley,K.A., A preliminary report on Quaternary fossil caddis larvae (Trichoptera). - Quaternary Newsletter 1978:2-12. (published by the Quaternary Research Association, Glasgow).
Schürch,Markus, Trophische Beziehungen invertebrater Räuber in der Lebensgemeinschaft eines Baches. - Diplomarbeit, Zool.Mus.Univ.Zürich, 39pp.
Solem,J.O., Naeringnett hos hvirrellØse dyr i littoralsonen i innsjØer. - Fauna (Oslo) 31:17-22.
Tani,Kozo, The caddisflies of Niigata Prefecture 1.(Trichoptera). - Etsusa Res.Group of Insects, no.49:2-26. (Japanese).

1979 (additions)

Andersen,T., Some caddis flies (Trichoptera) in western Norway, and their arrival pattern in light traps. - Fauna Norwegica, ser.B, 26:12-17.
Hamilton,S.W., Schuster,G.A., Records of Trichoptera from Kansas,II:.The families Glossosomatidae, Helicopsychidae, Hydropsychidae and Rhyacophilidae. - Tech.Publ.State Biol. Surv.Kansas 8:15-22.
Kiss,O., The Trichoptera of the Bükk Mountains. - Acta biol. Debrecina 16:45-55.
Lager,T.M., Johnson,M.D., Williams,S.N., McCulloch,J.L., A preliminary report on the Plecoptera and Trichoptera of northeastern Minnesota. - Great Lakes Ent. 12:109-114.
Leader,J.P., Bedford,J.J., Oviposition by the marine caddisfly Philanisus plebeius (Walk.). - Search 10:275-276.
Marinković-Gospodnetić,M., The species of the genus Hydropsyche of the group pellucidula (Trichoptera) in the Dinarides. - Glas.Zemaljsk.Muz.NS 18, Prir.nauke:165-171, Sarajevo.
Solem,J.O., A comparison of species diversity indices in Trichoptera communities. - In:Grassle,J.F.,Patil,G.P., Smith,W.,Taillie,C.,(eds.):Ecological diversity in theory and practice, pp.255-267. Int.Co-op.Publ.House,Fairland, Maryland.
Späh,H., Ökologische Untersuchungen an organisch belasteten Bächen im Stadtbereich von Bielefeld. - Ber.Naturw.Ver. Bielefeld 24:383-410.
Townsend,C.R., Hildrew,A.G., Foraging strategies and co-existence in a seasonal environment. - Oecologia 38:231-234.
Wiggins,G.B., Trichoptera. In: Merritt,R.W., Cummins,K.W., (eds.): An introduction to the aquatic insects of North America, pp.147-185. Kendall/Hunt Publ.Co.,Dubuque,Iowa.

1980 (additions)

Alstad,D.N., Comparative biology of the common Utah Hydropsychidae (Trichoptera). - Amer.Midl.Nat. 103-167-174.
Belluck,D.A., Pennington,B.A., Unzicker,J.D., Aberrant oviposition by the caddisfly Triaenodes tardus Milne (Trichoptera:Leptoceridae) - Ent.News 91:173-175.
Blickle,R.L., A new Oxyethira (Hydroptilidae,Trichoptera) of theaeola Ross group; with a key to separate the five males of the group. - Pan-Pac.Ent. 56:101-104.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1980

Bloesch,J., Bodenfaunistische Untersuchungen in Aare und Rhein. - Schweiz.Z.Hydrol. 42:285-308.
Botosaneanu,L., Etude de quelques types de Trichoptères ouest-paléarctiques, déposés au muséum de Paris (Insecta, Trichoptera). - Bull.ZoolMus.Univ.Amsterdam 7:189-195.
Bournaud,M., Keck,G., Diversité spécifique et structure des peuplements de macro-invertébrés benthiques au long d'un cours d'eau: le Furans (Ain). - Acta Oecologica/Oecol. Gener. 1:131-150.
Bueno Soria,J., Refugio y hábitos alimenticios de las larvas de Macronema variipenne Flint & Bueno (Trichoptera:Hydropsychidae). - Fol.Ent.Mex. 45:34-35.
Colbo,M.H., Undeen,A.H., Effect of Bacillus thuringiensis var. israelensis on non target insects in stream trials for control of Simuliidae. - Mosquito News 40:368-371.
Cudney,M.D.,Wallace,J.B., Life cycles, microdistribution and production dynamics of six species of net-spinning caddisflies in a large southeastern (USA) river. - Holarctic Ecol. 3:169-182.
Cuppen,H.P.J.J., De macrofauna in een aantal droogvallende - en permanente stillstaande wateren in het ruilverkavelingsgebied Brummen-Voorst. - Regionale Milieuraad Oost-Veluwe, polycopied, 112 pp.
Cuppen,H.P.J.J., Oosterloo,W., Een oriënterend onderzoek naar de makrofauna van de nijmolense beek (gem.epe). - Reg.Milieuraad Oost-Veluwe, 50 pp., polycopied.
Davis,J.R., Species compodition and diversity of benthis macroinvertebrates in the upper Rio Grande, Texas. - Southwest.Nat. 25:137-150.
Davis,J.R., Species composition and diversity of benthic macroinvertebrate population of the Pecos River,Texas. - Southwest.Nat. 25:241-256.
Davis,J.R., Species composition and diversity of benthic macroinvertebrates of lower Devil's River, Texas. - Southw.Nat. 25:379-384.
Denis,C., Action de la photopériode sur le cycle biologique et la diapause d'Anabolia nervosa Curtis (Trichoptera, Limnephilidae). - Annls.Limnol. 16:191-202.
Dratnal,E., Kasprzak,K., The response of the invertebrate fauna to organic pollution in a well oxygenated karst stream exemplified by the Prądnik Stream (South Poland). - Acta Hydrobiol.(Kraków) 22:263-278.
Ferrier,M.D., Wissing,T.E., A scanning electron microscopic study of the mouthparts of Cheumatopsyche analis (Trichoptera:Hydropsychidae) larvae. - Ohio J.Sci. 80:38-40.
Flint,O.S.,jr., VI.Trichoptera. In:Roback,S.W.(ed.): The results of the Catherwood Foundation Bolivian-Peruvian Altiplano Expedition, Part I. - Proc.Acad.Nat.Sci.Philadelphia 132:213-217.
Flint,O.S.,jr., Wallace,J.B., Studies of neotropical caddisflies. XXV: The immature stages of Blepharopus diaphanus and Leptonema columbianum (Trichoptera:Hydropsychidae). - Proc.Biol.Soc.Wash. 93:178-193.
Florin,J., Die Insektenfauna des Hochmoores Balmoos bei Hasle, Kanton Luzern. IV.Trichoptera (Köcherfliegen). - Ent.Ber. Luzern 3:86-93.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

Franz,H.P., Limnologische Untersuchung des Gewässersystems Dhron (Hunsrück). - Decheniana 133:155-179.
Fuller,R.L., Mackay,R.J., Field and laboratory studies of net-spinning activity by Hydropsyche laevae (Trichoptera: Hydropsychidae). - Can.J.Zool. 58:2006-2014.
Fuller,R.L., Mackay,R.J., Feeding ecology of three species of Hydropsyche (Trichoptera:Hydropsychidae) in southern Ontario. - Can.J.Zool. 58:2239-2251.
Geijskes,D.C., Higler,L.W.G., Handleiding voor het projekt kokerjuffers of Schietmotten (Trichoptera). - Instrukties voor medewerkers ETS-Nederland 5:1-24. Rijksmuseum van Natuurlijke Historie, Leiden).
Giudicelli,J., Dakki,M., Les Agapetus du Maroc - description de deux espèces nouvelles (Trichoptera, Glossosomatidae). Bijdr.Dierk. 50:227-234.
Giudicelli,J., Dia,A., Legier,P., Etude hydrobiologique d'une rivière de région méditerranéenne, l'Argens (Var,France). Habitats, hydrochimie, distribution de la faune benthique. Bijdr.Dierk. 50:303-341.
Giudicelli,J., Moubayed,Z., Tourenq,J.N., Un biotope hydrobiologique remarquable: Les Laurons de la Crau (Bouches-du--Rhône, France). La communauté animale et ses relations avec le peuplement des biotopes aquatiques voisins. - Annls.Limnol. 16:271-297.
Givens,D.R., Smith,S.D., A synopsis of the western Arctopsychidae (Trichoptera:Hydropsychidae). - Melanderia 35:1-24.
Grafius,E., Anderson,N.H., Population dynamics and role of two species of Lepidostoma (Trichoptera:Lepidostomatidae) in an Oregon coniferous forest stream. - Ecology 61:808-816.
Hamilton,S.W., Schuster,G.A., Notes on the taxonomy and biology of Phylocentropus (Trichoptera: Polycentropodidae). - J.Kansas Ent.Soc. 53:586.
Hamilton,S.W., Schuster,G.A., Records of Trichoptera from Kansas, III: The families Limnephilidae, Phryganeidae, Polycentropodidae, and Sericostomatidae. - Tech.Publ. State Biol.Surv.Kansas 9:20-29.
Henricson,J., Müller,K., Stream regulation in Sweden with some examples. In:Ward & Stanford (eds.): The ecology of regulated streams. Plenum Publ.Co., pp.183-199.
Ito,T., Morphological and ecological studies on the caddisfly genus Dinarthrodes in Hokkaido, Japan (Trichoptera, Lepidostomatidae). II. Life histories of two coexisting species, D.complicatus and D.satai. - Kontyû 48:311-320.
Jacquemart,S., Un Trichoptère Hydroptilide nouveau du nord du Chili: Metrichia thirysae sp.n. - Brenesia 17:303-318.
Jacquemart,S., A propos de Tinodes cinerea Hagen (Trichoptère Psychomyide) de Madère. - Bull.Inst.r.Sci.nat.Belg.Ent. 52(7):1-6.
Jacquemart,S., Un Trichoptère nouveau de l'Air: Hydroptila airensis sp.n. (Hydroptilide). - Bull.Inst.r.Sci.nat.Belg. Ent. 52(13):1-5.
Jacquemart,S., Description de la larve et de la nymphe d' Australomyia masatierra (Schmid) et considérations sur la larve d'Australomyia masafuera (Schmid)(Trichoptères) provenant de l'Archipel Juan Fernandez(Chili). - Bull. Inst.r.Sci.nat.Belg.Ent. 52(26):1-11.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1980

Kachalova,O.L., Kornoukhova,I.I., Caddisflies (Trichoptera) of Ardon River in the northern Caucasus. - Ent.Obozr. 59:269-275.

Klausnitzer,B., Lehnert,J., Klima,F., Wasserinsekten aus verschiedenen Parkteichen und Springbrunnen von Leipzig. - Wiss.Z.Karl-Marx-Univ.Leipzig, math.-natw.R.29:653-658.

Kumanski,K., To the knowledge of Chaetopterygini from the Caucasus (USSR)(Trichoptera, Limnephilidae). - Reichenbachia 18:153-160.

McLean,D.B., MacLean,B.K., Report of new Trichoptera records for Ohio. - Great Lakes Ent. 13:37-39.

Markarian,R.K., A study of the relationship between aquatic insect growth and water temperature in a small stream. - Hydrobiologia 75:81-95.

Masteller,E.C., Flint,O.S.,jr., Emergence phenology of Trichoptera from Six Mile Creek, Erie County, Pennsylvania, USA. - Aquat.Insects 2:197-210.

Mey,W., Neue Fundorte von Köcherfliegen in der MVR (Insecta, Trichoptera) (Ergebnisse der Mongolisch-Deutschen Biologischen Expeditionen seit 1962, Nr.84). - Mitt.zool.Mus. Berlin 56:103-109.

Moretti,G.P., Trentini,M., Prime osservazioni cromosomiche sul germario Potamophylax cingulatus Steph. femmina e Halesus radiatus Curt. femmina (Trichoptera, Limnephilidae). - Riv.Biol. 73:263-266.

Mori,S., Miura,T., List of plant and animal species living in Lake Biwa. - Mem.Fac.Sci.Kyoto Univ.,Ser.Biol.8-33.

Neveu,A., La derive des invertebres aquatiques et terrestres dans un petit fleuve cotier de l'ouest des Pyrénées, la Nivelle. - Acta Oecol.,Oecol.Appl. 1:317-339.

Roemhild,G., Pheromone glands of microcaddisflies (Trichoptera:Hydroptilidae). - J.Morphol.163:9-12.

Schleich,H.H., Die Golezard-Höhle im Nordiran. - Die Höhle (Wien) 31:97-103.

Schmid,F., Esquisse pour une classification et une phylogénie des Goérides(Trichoptera). - Nat.can. 107:185-194.

Sedlák,E., Rad Chrostici - Trichoptera. In: Rozkosny,R.(ed.), Klic vodnich larev hmyzu, 163-220. Csl.Akad.ved.

Short,R.A., Ward,J.V., Macroinvertebrates of a Colorado high mountain stream. - Southwest.Nat. 25:23-32.

Späh,H., Ökologische Untersuchung des Makroinvertebraten-Fauna der Sennebäche zwischen Stukenbrock und Bad Lippspringe. - Ber.Naturw.Ver.Bielefeld, Sonderh. 2:101-132.

Späh,H., Die Invertebratenfauna der oberen Hunte und einiger ihrer Nebenbäche. - Inf.Natursch.Landschaftspfl.(Wardenburg) 2:347-363.

Szczęsny,B., Caddis-flies (Trichoptera) in the collection of the Institute of Systematic and Experimental Zoology, Polish Academy of Sciences in Cracow. - Acta Zool.Cracov. 24: 449-486.

Tani,K., Larvae of caddisflies from Japan (Trichoptera). - Insects and Nature 15(8):24-30. (Japanese).

Vannote,R.L., Sweeney,B.W., Geographic analysis of thermal equilibria: a conceptual model for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. - Amer.Nat.115:667-695.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1980

Wallace,I.D., Caddis larvae (Trichoptera), with particular reference to those found on the Wirral Peninsular. - Ann.Rep.Proc.Lancash.Chesh.Ent.Soc. 100:79-85.
Wells,A., A review of the Australian genera Orphninotrichia Mosely and Maydenoptila Neboiss (Trichoptera:Hydroptilidae), with descriptions of new species. - Aust.J.Zool.28:627-645.
White,T.R., Weaver,J.S.III, Fox,R.C., Phoretic relationships between Chironomidae (Diptera) and benthic macroinvertebrates. - Ent.News 91:69-74.
Wichard,W., Schmitz,M., Anpassungsmechanismen der osmoregulatorischen Ionenabsorption bei Limnephilidae-Larven (Insecta, Trichoptera). - Gewässer und Abwässer 66/67:102-118.
Williams,D.D., Temporal patterns in recolonization of stream benthos. - Arch.Hydrobiol. 90:56-74.
Winterbourn,M.J., Benthic communities of South Island beech forest streams. - Beech Res.News 7:5-10.
Winterbourn,M.J., The freshwater insects of Australasia and their affinities. - Paleogeogr.,Paleoclimat.,Palaeoecol. 31:235-249.
Winterbourn,M.J., Anderson,N.H., The life history of Philanisus plebeius Walker (Trichoptera: Chathamiidae), a caddisfly whose eggs were found in a starfish. - Ecol.Entomol. 5:293-303.

1981

NOTE. The Proceedings of the 3rd International Symposium on Trichoptera (Perugia), ed.G.P.Moretti, The Hague:Dr.W.Junk publishers, are abbreviated as follows: Protri 3.

Aisa,E., Guerrieri,P., A report on gregarines of Trichoptera in the intestines of Salmo fario L. - Protri 3:1-3.
Alstad,D.N., Nearest-neighbor analysis of microhabitat partitioning in stream insect communities. - Hydrobiologia 79:137-140.
Andruschajtisa,G.P., Katschalova,O.L., Hydrobiological regime of small rivers under conditions of anthropogenic influence. - Acad.Nauk Latv.SSR,Inst.Biol., 166 pp. (Russian)
Badcock,R.M., Pigments and preservation in caddis larvae and pupae. - Protri 3:5-10.
Bird,G.A., Hynes,H.B.N., Movements of adult aquatic insects near streams in southern Ontario. - Hydrobiologia 77:65-69.
Boles,G.L., Macroinvertebrate colonisation of replacement substrate below a hypolimnial release reservoir. - Hydrobiologia 78:133-146.
Botosaneanu,L., Ordo Trichoptera et homo insapiens. - Protri 3:11-19
Botosaneanu,L., Wichard,W., Trichoptera from the Baltic Amber. Protri 3:31-32.
Botosaneanu,L., Giudicelli,J., Observations morphologiques, éthologiques et écologiques sur Hydroptila hirra Mosely. Protri 3: 21-29
Botosaneanu,L., On some Trichoptera collected by Mrs.Drs. A.C.Ellis and Dr.W.N.Ellis on Gomera (Canary Islands). - Ent.Ber.(Amst.) 41:186-190.
Botosaneanu,L., Marlier,G., Les Wormaldia ouest-paléarctiques des collections de l'Institut Royal des Sciences Naturelles de Belgique. Contribution à l'étude de la variabilité des Wormaldia. - Bull.Inst.r.Sci.nat.Belg. 53(19):1-11.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1981

Botosaneanu,L., Giudicelli,J., Les Hydropsyche de Corse. Bull.Zool.Mus.Univ.Amsterdam 8:13-19.
Botosaneanu,L., On a false and a genuine caddis-fly from Burmese Amber (Insecta:Trichoptera,Homoptera). - Bull. Zool.Mus.Univ.Amsterdam 8:73-78.
Bradt,P.T., Wieland,G.E., A comparison of the benthic macroinvertebrate communities in a trout stream: winter and spring 1973 and 1977. - Hydrobiologia 77:31-35.
Bueno-Soria,J., Padilla,J., Rivera,M., Observations on the longitudinal distribution of Trichoptera larvae in a stream at Zempoala Mexico, Maxico. - Protri 3:33-38.
Çakin,F., Ankara ve kismen çevre illerdeki Trichoptera erginlerinin sistekatik yönden incelenmesi. - Thesis, Hacettepe Üniversitesi, Ankara, 263 pp.
Canton,S.P., Ward,J.V., Emergence of Trichoptera from Trout Creek, Colorado, USA. - Protri 3:39-45.
Crichton,M.I., Fisher,D.B., Further observations on Limnephilid life histories, based on the Rothamsted Insect Survey. - Protri 3:47-55.
Dakki,M., Contribution à la connaissance des Leptocerides du Maroc. - Bull.Inst.Sci.Rabat 1979-80, 4:41-52.
Dejoux,C., Elouard,J.M., Forge,P., Jestin,J.M., Mise en évidence de la microdistribution des invertébrés dans les cours d'eau tropicaux. Incidence méthodologique pour la recherche d'une pollution à long terme par insecticide. - Rev.Hydrobiol.trop. 14:253-262.
Denis,C., Action de la photopériode sur la maturation génitale des femelles de quelques limnephilidés. - Protri 3:57-66.
Dieterlen,F., Statzner,B., The African rodent Colomys goslingi Thomas and Wroughton, 1907 (Rodentia:Muridae) - a predator in limnetic ecosystems. - Z.Säugetierkunde 46:369-383.
Döhler,W.: Ein neues Trichopteron aus dem Schwarzwald: Beraeamyia eideli n.sp. - Ent.Z.(Stuttg.)91:34-37.
Dowling,C., O'Connor,J.P., O'Grady,M.F., A baseline survey of the Caragh, an unpolluted river in southwest Ireland: observations on the macroinvertebrates. - J.Life Sci. R.Dubl.Soc. 2:147-159.
Edington,J.M.,Hildrew,A.G., A key to the caseless caddis larvae of the British Isles with notes on their ecology. - Sci.Publ.Freshw.Biol.Ass. 43:1-92.
Elliott,J.M., Drake,C.M., A comparative study of seven grabs used for sampling benthic macroinvertebrates in rivers. - Freshw.Biol.11:99-120.
Elliott,J.M., Drake,C.M., A comparative study of four dredges used for sampling benthic macroinvertebrates in rivers. - Freshw.Biol. 11:245-261.
Elliott,J.M., A quantitative study of the life cycle of the net-spinning caddis Philopotamus montanus in a Lake District stream. - J.Anim.Ecol. 50:867-883.
Flannagan,J.F., Cobb,D.G., Changes in the profundal Trichoptera of Lake Winnipeg 1928-32 to 1969. - Protri 3:67-74.
Flint,O.S.,jr., Studies of Neotropical caddisflies, XXVII: Anomalopsychidae, a new family of Trichoptera. - Protri 3:75-85.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1981

Fuller,R.L., Mackay,R.J., Effects of food quality on the growth of three Hydropsyche species. Can.J.Zool.59:1133-1140.
Geddes,A.J., Observations on the feeding behaviour and diet of the aquatic larva of Allogamus auricollis. Ent.Gaz. 32:271-274.
Georgian,T.J.,jr., Wallace,J.B., A model of seston capture by net-spinning caddisflies. - Oikos 36:147-157.
Gislason,G.M., Predatory exclusion of Apatania zonella by Potamophylax cingulatus in Iceland. Protri 3:93-97.
Gislason, G.M., Distribution and habitat preferences of Icelandic Trichoptera. - Protri 3:99-109.
Gonzalez,M.A., Terra,L.S., Une nueva especie del genero Stactobia en la peninsula Iberica. - Nouv.Rev.Ent.11:203-206.
Gonzalez,M.A.,Otero,J.C., Observaciones sobre los Tricópteros de la Peninsula ibérica: II.- Bol.Asoc.esp.Ent.4:127-129.
Haefner,J.D., Wallace,J.B., Shifts in aquatic insect populations in a first-order southern Appalachian stream following a decade of old field succession. - Can.J.Fish.Aquat.Sci. 38:353-359.
Harrison,J.D., Badcock,R.M., Effects of acclimation to temperature on the respiratory rate of some Limnephilid larvae. - Protri 3:111-116.
Hart,D.D., Foraging and resource patchiness: field experiments with a grazing stream insect. - Oikos 37:46-52.
Hemsworth,R.J., Brookes,M.P., Macroinvertebrate drift in the upper Wye catchment, Wales. - Hydrobiologia 85:145-155.
Higler,L.W.G., The effects of pollution in the drainage area of a Dutch lowland stream on fish and macro-invertebrates. - Verh.Int.Ver.Limnol.21:1077-1082.
Higler,L.W.G., Caddisfly systematics up to 1960 and a review of the genera. - Protri 3:117-126.
Higler,L.W.G., Caddis larvae in a Dutch lowland stream. - Protri 3:127-128.
Higler,L.W.G., The presence of caddis flies in the Netherlands and their role in the aquatic system. - Hydrobiol. Bull.(Amsterdam) 14:209-212.
Hudson,P.L., Morse,J.C., Voshell,J.R.,jr., Larva and pupa of Cernotina spicata. - Ann.Ent.Soc.Amer.74:516-519.
Imhof,J.G.A., Harrison,A.D., Survival of Diplectrona modesta Banks during short periods of desiccation. - Hydrobiologia 77:61-63.
Jacquemart,S., Un Trichoptère Philopotamide nouveau de Nouvelle-Guinée (Irian Jaya): Chimarra leopoldi sp.n. - Bull.Inst.r.Sci.nat.Belg.Ent. 53(5):1-6.
Jacqemart,S., Statzner,B., Trichoptères nouveaux du Zaire. Bull.Inst.r.Sci.nat.Belg.Ent. 53(21):1-25.
de Jalón,D.G., del Tánago,M.G., de Viedma,M.G., Importancia de los insectos en los métodos biológicos para el estudio de la calidad de las aguas: Necesidad de su conocimiento taxonomico. - Graellsia 35-36:143-148.
de Jalon,D.G., Description of Hydropsyche larvae found in the Iberian Peninsula. Protri 3:87-92.
Jude,D.J., Rutecki,T.L., Madenjian,C.P., Noguchi,G.E., Schneeberger,P.P., Klinger,S.A., Godun,G.G., Winnell,M.H., The physical, chemical and biological nature of Pigeon Lake, a Lake Michigan coastal lake. - Spec.Rep.(Great Lakes Res.Div.) 78:1-109.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1981

King,J.M., The distribution of invertebrate communities in a small South African river. - Hydrobiologia 83:43-65.
Kiss,O., Trichoptera in the Ilona stream of the Mátra Mountains, North Hungary. - Protri 3:129-138.
Klausnitzer,B., Kuckelkorn,B., Kuckelkorn,U., Schüler,H., Zur Entomofauna des Tagebaurestsees Olba (Kreis Bautzen) - Abh.Ber.Naturkundemus.Görlitz 54(3):1-16.
Klima,F., Neue Köcherfliegenfunde aus der DDR. - Ent.Nachr. (Dresden) 25:52-56.
Klima,F., Köcherfliegen aus der polnischen Tatra. Ent.Nachr. (Dresden) 25:72-76.
de Kozlowski,S., Bunting,D.L.,II, A laboratory study on the thermal tolerance of four southeastern stream insect species. - Hydrobiologia 79:141-145.
Kumanski,K., Description of two new caddisflies and the probable adults of Philocrena trialetica Lepn. from the Caucasus. - Reichenbachia 19:63-70.
Kumanski,K., Faunistic investigations on Bulgarian Trichoptera to June, 1980 - with a revised check-list. - Protri 3: 139-147.
Ladle,M., Bass,J.A.B., The ecology of a small chalk stream and its responses to drying during drought conditions. - Arch.Hydrobiol. 90:448-466.
Lebrun,P., Vlayen,P., Etude de la bioactivité comparée et des effets secondaires de Bacillus thuringiensis H 14. - Z.angew.Ent. 91:15-25.
Legier,P., Terzian.E., Les milieux aquatiques temporaires de Provence: Resistance des invertébrés à l'assèchement des habitats. - Ecol.Medit. 7:113-129.
Lenat,D.R., Penrose,D.L., Eagleson,K.W., Variable effects of sediment addition on stream benthos.- Hydrobiologia 79:187-194.
Levanidova,I.M., Schmid,F., Considerations on Archithremma ulachensis Martynov. - Aquat.Ins. 3:65-73.
McElravy,E.P., Resh,V.H., Wolda,H., Flint,O.S.,jr., Diversity of adult Trichoptera in a 'Non-seasonal' tropical environment. - Protri 3:149-156.
Malicky,H., Oecismus tjederi n.sp., eine neue Sericostomatide aus Griechenland. - Ent.Z.(Stuttg.) 91:69-70.
Malicky,H., Gonzalez,M.A., Hydroptila vilaverde n.sp., eine neue Köcherfliege von der Iberischen Halbinsel. - Ent.Z.(Stuttg.) 91:151-152.
Malicky,H., Polycentropus pirisinui sp.n. (Trichoptera, Polycentropodidae) from Central Italy.- Aqu.Ins. 3:12.
Malicky,H., Artificial illumination of a mountain stream in Lower Austria: Effect of constant daylength on the phenology of the caddisflies. - Aqu.Ins. 3:25-32.
Malicky,H., Drusus noricus sp.n. from the mountains of Carinthia (Austria). - Aqu.Ins. 3:44.
Malicky,H., The phenology of dispersal of several caddisfly species in the island of Crete. - Protri 3:157-163.
Malicky,H., Eine neue Chimarra aus Neukaledonien. - Rev.suisse Zool. 88:341-342.
Malicky,H., Neues über mediterrane, vorderasiatische und europäische Köcherfliegen.- Entomofauna(Linz) 2:175-188.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

Malicky,H., Weiteres Neues über Köcherfliegen aus dem Mittelmeergebiet. - Entomofauna(Linz) 2:335-355.
Malicky,H., Der Indikatorwert von Köcherfliegen in großen Flüssen. - Mitt.dt.Ges.allg.ang.Ent. 3:135-137.
Manuel,K.L., The caddisfly genus Triaenodes in North America: Phylogeny, Biogeography, Ecology. - Abstr.:N.Am.Benthol. Soc., spring meeting 1981.
Marlier,G., Trichoptères.In:Durand,Lévêque(eds.): Flore et faune aquatiques de l'Afrique Sahelo-Soudanienne 2:521-552. Ed.ORSTOM,Paris.
Marlier,G., Trichoptères. In:Symoens (ed.),Exploration hydrobiologique du bassin du Lac Bangweolo et du Luapula. Vol.14(7):1-70. Bruxelles.
Merritt,R.W., Wallace,J.B., Filter-feeding insects. Sci.Amer. 244(4):132-144.
Messner,B, Taschenberger,D., Zur Funktionsmorphologie des Atembandes von Agriotypus armatus Walk.(Hym.,Agriotypidae). - Dt.Ent.Z., N.F. 28:7-9.
Mey,W., Die von R.Jung und A.Müller in Mittelasien gesammelten Köcherfliegen. - Dt.Ent.Z.N.F. 28:55-66.
Mey,W., Radema curvipennis n.sp., ein neuer Endemit des Baikal-Sees. - Dt.Ent.Z.,N.F. 28:301-303.
Mey,W., Lichtfangergebnisse bei Köcherfliegen im Havelseengebiet. - Beitr.Ent. 31:333-339.
Moretti,G.P., Cianficconi,F., Tucciarelli,F., Ripartizione dei tricotteri nel sistema idrico del Lago di Piediluco e nella Cascata delle Marmore (Umbria-Terni). - Studi Trent. Sci.Nat. 58,Acta Biol.:315-373.
Moretti,G.P., Spinelli Batta,G., Descrizione morfologica e tassonomica della larva di Hydropsyche dissimulata Kum. e Bot. nuova per la scienza. - Riv.Idrobiol. 18:133-171.
Moretti,G.P., Bicchierai,M.C., Struttura androconiale di Hydroptila aegyptia Ulm. - Riv.Idrobiol. 18:173-195.
Moretti,G.P., Bicchierai,M.C., Osservazioni al SEM a TEM del palpo mascellare androconiale nel ♂ di Sericostoma pedemontanum McL. - Riv.Idrobiol. 18:197-213.
Moretti,G.P., Bicchierai,M.C., Placche furcifere della "linea laterale" nella larva di Sericostoma pedemontanum McL. - Riv.Idrobiol. 18:215-228.
Moretti,G.P., Bicchierai,M.C., Immagini al SEM dell'androconia alare di Beraea sp.n. - Riv.Idrobiol.18:229-235.
Moretti,G.P., Spinelli Batta,G., Bilancio sistematico delle rappresentanze del genere Halesus appartenenti al gruppo digitatus, nella peninsola italiana e nella Sardegna. - Riv.Idrobiol. 18:237-265.
Moretti,G.P., Bicchierai,M.C., Androconia alare di Beraeamyia squamosa Mos. - Riv.Idrobiol. 28:329-336.
Moretti,G.P., Cianficconi,F., Biccierai,M.C., Squame androconiali delle ali posteriori di Drusus improvisus McL. Riv.Idrobiol. 28:337-351.
Moretti,G.P., Bicchierai,M.C., Formazioni androconiali dell'ala posteriore di Silo nigricornis Pict. - Riv. Idrobiol. 28:379-390.
Moretti,G.P., New Trichoptera species and subspecies found in Italy. Protri 3: 165-192.
Moretti,G.P., Bicchierai,M.C., Comparative SEM and TEM studies on the androconial structures of Lasiocephala basalis Kol. and other Trichoptera. - Protri 3:193-198.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1981

Moretti,G.P., Cianficconi,F., First list of Italian Trichoptera. - Protri 3:199-211.

Moretti,G.P., Corallini Sorcetti,C., Gregarines in Trichoptera larvae. - Protri 3:213-217

Moretti,G.P., di Giovanni,M.V., The Trichoptera component in the hydrophyton of Lake Chiusi (Tuscany). - Protri 3:219-226.

Moretti,G.P., Mearelli,M., Ecological profiles in three Rhyacophila species. - Protri 3:227-230.

Moretti,G.P., Pirisinu,Q., Morphological characteristics of Leptodrusus budtzi Ulm. in the immature stages. - Protri 3:231-236.

Moretti,G.P., Spinelli Batta,G., Taxonomic and morphological characteristics of the Sericostoma italicum Moret. and Sericostoma pedemontanum McL. larvae. - Protri 3:237-242.

Moretti,G.P., Corallini Sorcetti,C., Gattaponi,P., Trichoptera in the intestinal content of certain fish species. - Protri 3: 243-247.

Moretti,G.P., Gianotti,F.S., Taticchi,M.I., Viganò,A., Trichoptera of the Isle of Elba (Tuscany, Italy). - Protri 3:249-256.

Morse,J.C., A tribute to Herbert H.Ross, 1908-1978. - Protri 3:XVII-XVIII.

Morse,J.C., A phylogeny and classification of family-group taxa of Leptoceridae. Protri 3:257-264.

Nebeker,A.V., Baker,F.D., Weitz,S.L., Survival and adult emergence of aquatic insects in air-supersaturated water. - J.Freshw.Ecol. 1:243-250.

Neboiss,A., Tasmanian Caddis-flies. - Fauna of Tasmania Handbook no. 4, 180 pp.

Neboiss,A., Distribution of Trichoptera Families in Australia with comments on the composition of fauna in the South--West. - Protri 3:265-272.

Nielsen,A., On the evolution of the phallus and other male terminalia in the Hydropsychidae with a proposal for a new generic name. - Protri 3:273-278.

Nilsson,L.M., Otto,C., Active uptake of dissolved organic matter by Potamophylax cingulatus larvae. - Oikos 37:345-348.

Nimmo,A.P., Bibliography of Trichoptera.- Protri 3:279-280.

Novák,K., Trichoptera distribution pattern:differences found by sweeping, beating and light traps at three Southern Bohemian sites. - Protri 3:281-284.

O'Connor,J.P., Barnard,P.C., Limnephilus tauricus Schmid new to Great Britain, with a key to the L.hirsutus (Pictet) group in the British Isles. - Ent.Gaz. 32:115-119.

Olsson,T., Overwintering of benthic macroinvertebrates in ice and frozen sediment in a North Swedish river. - Holarctic Ecol. 4:161-166.

Otto,C., Food related adaptations in stream living caddisfly larvae feeding on leaves. - Oikos 37:117-122.

Otto,C., Why does duration of flight periods differ in caddisflies ? - Oikos 37:383-386.

Otto,C., Svensson,B.S., How do macrophytes growing in or close to water reduce their consumption by aquatic herbivores ? - Hydrobiologia 78:107-112.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

1981

Otto,C., Svensson,B,S., Why do Potamophylax cingulatus larvae aggregate at pupation ? - Protri 3:285-291.
Parker,C.R., Voshell,J.R.,jr., A preliminary checklist of the caddisflies of Virginia. - J.Ga.Ent.Soc. 16:1-7.
Peckarsky, B.L., Reply to comment by Sell. - Limnol.Oceanogr. 26:982-987. (about distr.of benthic invertebrates in stony streams).
Prat,N., The influence of reservoir discharge on benthic fauna in the River Ter, NE Spain.- Protri 3:293-301.
Puig,M.A.,Bautista,I.,Tort,M.J.,Prat,N., Les larves de Trichoptères de la rivière Llobregat (Catalogne,Espagne). Distribution longitudinale et relation avec la qualité de l'eau. - Protri 3:303-309.
Rau,G.H., Anderson,N.H., Use of $^{13}C/^{12}C$ to trace dissolved and particulate organic matter utilization by populations of an aquatic invertebrate. - Oecologia (Berl.) 48:19-21. (Clistoronia magnifica)
Reice,S.R., Interspecific associations in a woodland stream. Can.J.Fish.Aq.Sci. 38:1271-1280.
Resh,V.H., Flynn,T.S., Lamberti,G.A., McElravy,E.P., Sorg,K.L., Wood,J.R., Responses of the Sericostomatid caddisfly Gumaga nigricula (McL.) to environmental disruption. - Protri 3:311-318.
Schembri,S.P., First record of the order Trichoptera from the Maltese Islands. - Bull.Amat.Ent.Soc. 40:25-26.
Schmid,F., Revision des Trichoptères Canadiens. I. La famille des Rhyacophilidae(Annulipalpia). - Mem.Soc. Ent.Can. 116:1-83.
Siegenthaler,C., Trichoptera of western Switzerland. - Protri 3:319-320.
Solem,J.O., Resh,V.H., Larval and pupal description, life cycle, and adult flight behaviour of the sponge-feeding caddisfly, Ceraclea nigronervosa (Retzius), in Central Norway. - Ent.scand. 12:311-319.
Solem,J.O., Systematic position of Agrypnetes crassicornis McLachlan, 1876. - Ent.scand.Suppl.15:398-400.
Solem,J.O., Overwintering strategies in some Norwegian caddisflies. - Protri 3:321-322.
Späh,H., Beitrag zur Kenntnis der Wirbellosen-Fauna einiger Bäche des Wiehengebirges (West-Niedersachsen). - Beitr. Naturk.Niedersachsens 34:77-91.
Standford,J.A., Ward,J.V., Preliminary interpretations of the distribution of Hydropsychidae in a regulated river. - Pritri 3:323-328.
Statzner,B., The relation between "hydraulic stress" and microdistribution of benthic macroinvertebrates in a lowland running water system, the Schierenseebrooks (North Germany). - Arch.Hydrobiol. 91:192-218.
Statzner,B., A progress report on Hydropsychidae from the Ivory Coast: Characters for the specific identification of larvae and population dynamics of four abundant species. - Protri 3:329-335.
Sykora,J.L., Swegman,B.G., Weaver,J.S.,III., Occurrence of the genus Hydropsyche in the North American Great Lakes. - Protri 3:337-345.
Tachet,H., Bournaud,M., Cycles biologiques des Hydropsychidae et d'un Polycentropodidae dans le Rhône en amont de Lyon. - Protri 3:347-364.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

Terra,L.S.W.da, Liste faunística de Tricópteros de Portugal. Bol.Soc.Portug.Ent. 12:1-42.
Tobias,W., Eine neue Apatania-Art aus Norwegen. - Ent.Z. (stuttg) 91:62-65.
Tobias,D., Tobias,W., Verzeichnis der deutschen Köcherfliegen-Arten. - Ent.Z.(Stuttgart) 91:85-90.
Tobias,W., Bemerkungen zur Taxonomie von Hydropsyche exocellata Dufour 1841.- Ent.Z.(Stuttgart) 91:225-230.
Tobias,W., Tobias,D., Trichoptera Germanica. Bestimmungstafeln für die deutschen Köcherfliegen. Teil I: Imagines. - Courier Forschungsinst.Senckenberg 49:1-672.
Tomaszewski,C., The principles of case building behaviour in Trichoptera, - Protri 3:365-373.
Tozer,W.E., Resh,V.H., Solem,J.O., Bionomics and adult behavior of a lentic caddisfly, Nectopsyche albida (Walker). - Amer.Midl.Nat. 106:133-144.
Ujhelyi,S., Über das Vorkommen der Arten der Gattung Rhyacophila in Ungarn. - Fol.Ent.Hung. 42:193-196.
Vinikour,W.S., Aquatic insect drift through a final-cut strip mine pit, with emphasis on drift distances. - Hydrobiologia 77:225-232.
Vinikour,W.S., Anderson,R.V., Diptera larvae (Empididae and Chironomidae) in Trichoptera pupal cases (Glossosomatidae and Limnephilidae). - Ent.News 92:69-74.
Wallace,I.D., A key to larvae of the family Leptoceridae in Great Britain and Ireland. - Freshw.Biol. 11:273-297.
Ward,J.V., Altitudinal distribution and abundance of Trichoptera in a Rocky Mountain stream. - Protri 3:375-381.
Weaver,J.S.III, White,T.R., Larval description of Rhyacophila appalachia Morse and Ross.- J.Ga.Ent.Soc.16:269-271.
Weaver,J.S.III, Wojtowicz,J.A., Etnier,D.A., Larval and pupal descriptions of Dolophilodes (Fumonta) major (Banks). - Ent.News 92:85-90.
Weaver,J.S.III, Sykora,J.L., Larval behavior and dispersion of Pycnopsyche luculenta (Betten) demonstrated by a unique tagging method. - Protri 3:383-393.
Wells,A., The genera Oxyethira Eaton, Gnathotrichia Ulmer and Stenoxyethira Kimmins in Australia. - Aust.J.Zool. 29:103-118.
Welton,J.S., Ladle,M., Bass,J.A.B., Chapman,K., Invertebrate sampling in the substratum of an experimental recirculating stream. - Int.Rev.ges.Hydrobiol. 66:407-414.
Wiberg-Larsen,P., Tricholeiochiton fagesii(Guinard) of Triaenodes simulans Tjeder nye for Danmark. - Ent.Meddr. 49:28-30.
Wichard,W., Köcherfliegen des Dominikanischen Bernsteins. I.Ochrotrichia doehleri sp.nov. - Mitt.Münch.Ent.Ges. 71:161-162.
Wiggins,G.B., Considerations on the relevance of immature stages to the systematics of Trichoptera. - Protri 3:395-407.
Wilkinson,B., Quaternary sub-fossil Trichoptera larvae from a site in the English Lake District. - Protri 3:409-419.
Williams,D.D., The first diets of postemergent brook trout (Salvelinus fontinalis) and Atlantic salmon (Salmo salar) alevins in a Quebec river. - Can.J.Fish.Aq.Sci.38:765-771.
Williams,D., Williams,N.E., Some aspects of the life history and feeding ecology of Dolophilodes distinctus (Walker) in two Ontario streams. - Protri 3:431-440.

© Hans Malicky/Austria; download unter www.biologiezentrum.at

Williams,N.E., Westgate,J.A., Williams,D.D., Morgan,A., Morgan,A.V., Invertebrate fossils (Insecta:Trichoptera, Diptera, Coleoptera) from the Pleistocene Scanborough Formation at Toronto, Ontario, and their paleoenvironmental significance. - Quat.Res. 16:146-166.
Williams,N.E., Wiggins,G.B., A proposed setal nomenclature and homology for larval Trichoptera.- Protri 3:421-429.
Zintl,H., Observations and a theory on the building style change in the Potamophylax latipennis. - Protri 3:441-447.

1982

Botosaneanu,L., Etude de quelques Trichoptères ouest-paléarctiques intéressants appartenant au British Museum (Natural History). - Bull.Zool.Mus.Univ.Amsterdam 8:177-188.
Coutant,C.C., Positive phototaxis in first instar Hydropsyche cockerelli Banks. Aq.Ins. 4:55-59.
Cuffney,T.F., Minshall,G.W., Life history and bionomics of Arctopsyche grandis in a central Idaho stream. - Holarctic Ecol. 4:252-262.
Denning,D.G., A review of the Goeridae.- Can.Ent.114:637-642.
Elliott,J.M., A quantitative study of the life cycle of the case-building caddis Odontocerum albicorne in a Lake District stream. - Freshw.Biol. 12:241-255.
Friedländer,M., Morse,J.C., The aberrant spermatozoa of Hydropsychidae caddisflies: An electron microscope analysis of spermiogenesis. - J.Ultrastruct.Res. 78:84-94.
de Jalón,D.G., Contribución a la zoogeografia de los Tricópteros Espanoles (I). - SHILAP,Revta.lepid., 10:75-78.
de Jalon,D.G., do.(II). SHILAP 10:157-165.
Kondratieff,P.F., Simmons,G.M.,jr., Nutrient retention and macroinvertebrate community structure in a small stream receiving sewage effluent. - Arch.Hydrobiol. 94:83-98.
Kumanski,K., Description of three new caddis-flies from Bulgaria. - Riv.Idrobiol. 19:197-206.
Müller,K., The colonization cycle of freshwater insects. - Oecologia(Berl.) 52:202-207.
Otto,C., Habitat, size and distribution of Scandinavian limnephilid caddisflies. - Oikos 38:355-360.
Resh,Vincent H., Age structure alteration in a caddisfly population after habitat loss and recovery. - Oikos 38: 280-284.
Scheiblreiter,H., Ein Beitrag zur Kenntnis der Köcherfliegenfauna der Großen Tulln (Niederösterreich). - Diss.,Univ. Wien, 309 pp.
Timms,B.V., A study of the benthic communities of twenty lakes in the South Island, New Zealand.- Freshw.Biol. 12:123-138.
Ward,J.V., Stanford,J.A., Thermal responses in the evolutionary ecology of aquatic insects. - Ann.Rev.Ent. 27:97-117.
Williams,D.D., Williams,N.E., Morphological and dietary variations in a riverine population of Pycnopsyche guttifer. - Aquat.Ins. 4:21-27.

(To be continued)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Trichoptera Newsletter

Jahr/Year: 1982

Band/Volume: 09

Autor(en)/Author(s): Anonym

Artikel/Article: Trichopterological Literature 12-26