

TRICHOPTEROLOGICAL LITERATURE

1981

Baumann,R.W., Unzicker,J.D., Preliminary checklist of Utah caddisflies (Trichoptera). - *Encyclia* 58:25-29.

1984

Brittain,J.E., Lillehammer,A., Bildeng,R., The impact of a water transfer scheme on the benthic macroinvertebrates of a Norwegian river. - In: Lillehammer,A., Saltveit,S.J. (eds): Regulated Rivers. Universitetsforlaget Oslo, pp. 189-199.

Décamps,H., Biology of regulated rivers in France. - In:Lillehammer, A., Saltveit,S.J. (eds): Regulated Rivers. Universitetsforlaget Oslo, pp. 495-514.

Henricson,J., Sjöberg,G., Stream zoobenthos below two short-term regulated hydro-power dams in Sweden. - In: Lillehammer,A., Saltveit,S.J. (eds): Regulated Rivers. Universitetsforlaget Oslo, pp.211-221.

Lillehammer,A., Saltveit,S.J., The effect of the regulation on the aquatic macroinvertebrate fauna of the River Suldalslågen, Western Norway. - In: Lillehammer,A., Saltveit,S.J. (eds) Regulated Rivers. Universitetsforlaget Oslo, pp.201-210.)

Millet,X., Prat,N., Las comunidades de macroinvertebrados a lo largo del Rio Llobregat. - *Limnética* 1:222-233.

Puig,M.A., González,G., Soriano,O., Introducción al estudio de las comunidades macrobentónicos de los ríos asturianos: Efemeropteros, Plecopteros, Tricópteros, Similidos y Quironomidos.- *Limnética* 1:187-196.

1986

Baumann,R.W., Dirk Cornelis Geijskes (1907-1985). - *Perla* 7:3.

Cortés,M.V., Rui,Tjarda,K.de, Simoes Graça,M.A., Estudio de un río montaña de una zona granítica del norte de Portugal.- *Limnética* 2:197-204.

Habdija,I., Einfluß der organischen Belastung auf die biozönotische Struktur des Makrozoobenthos in lithorheophilen, psammorheophilen und pelorheophilen Zönosen der Gebirgsbäche. - *Acta hydrochim. hydrobiol.* 14:241-254.

Hallberg,E., Hagberg,M., Ocellar fine structure in *Caenis robusta* (Ephemeroptera), *Trichostegia minor*, *Agrypnia varia*, and *Limnephilus flavicornis* (Trichoptera). - *Protoplasma* 135:12=18.

Holzenthal,R.W., Studies in neotropical Leptoceridae (Trichoptera),VI: Immature stages of *Hudsonema flaminii* (Navas) and the evolution and historical biogeography of Hudsonemini (Triplectidinae).- *Proc.Ent.Soc.Wash.* 88:268-279.

Munoz,I., Prat,N., Millet,X., Martínez Ansemil,E., Heterogeneidad espacial en la distribucion de los macroinvertebrados a lo largo de un transecto en el Rio Llobregat (Barcelona, España).- *Limnética* 2:135-145.

Peterson,R.H., Martin-Robichaud,D.J., Aquatic insect histories and Atlantic salmon fry diets in the St.Croix River, New Brunswick, Canada. - *Can.Tech.Rep.Fish.Aquat.Sci.* 1485:1=30.

1986

- Recasens,L., Murillo,J., Ciclo vital de *Hydropsyche siltalai* Döhler, 1963 (Trichoptera, Hydropsychidae) en la riera de Vallvidrera (Catalunya, Espana). - Limnética 2:157-162.
- Rieradevall,M., Prat,N., Deriva nictemeral de macroinvertebrados en el Rio Llobregat (Barcelona). - Limnética 2:147-156.
- Starmühlner,F., Checklist of the fauna of mountain streams of tropical Indopacific Islands. - Arch.Naturhist.Mus.Wien 88/89B: 457-480.
- Xie,W.-D., Canning,E.V., Pyrotheca hydropsycheae n.sp., a microsporidian parasite of caddis fly larvae, *Hydropsyche siltalai* Doepler, 1963 (Trichoptera, Hydropsychidae). - J.Protazool. 33:462-467.

1987

- Aagaard,K., Hågvar,S., Sjeldne insektarter i Norge. 1.Døgnfluer, steinfluer, øyenstikkere, vannteger, vårflyer, rettvinger, saksedyr, nettvinger, mudderfluer og skorpionflyer. - Økoforsk Utred. 1987,6:1-81.
- Alstad,D.N., Particle size, resource concentration, and the distribution of net-spinning caddisflies. - Oecologia 71:525-531.
- Alstad,D.N., A capture-rate model of net-spinning caddisfly communities. - Oecologia 71:532-536.
- Arens,W., Vergleichende Untersuchungen zur Funktionsmorphologie der Mundwerkzeuge aufwuchsfressender Bergbachtiere. Adaptationen und Konvergenzen. - Diss., Fak.Biol., Univ.Freiburg i.Br., 359pp.
- Badri,A., Giudicelli,J., Prévot,G., Effets d'une crue sur la communauté d'invertébrés benthiques d'une rivière méditerranéenne, le Rdat (Maroc). - Acta Oecol., Oecol.Gener. 8:481-500.
- Flannagan,J.F., MacDonald,S.R., Ephemeroptera and Trichoptera of peatlands and marshes in Canada. - Mem.Ent.Soc.Can. 140:47-56.
- Forsyth,D.J., Lewis,M.H., The Invertebrates. In: Viner,A.B.(ed.): Inland waters of New Zealand, pp. 265-290.
- Gellert,G., Limnologische Untersuchung der Sieg zwischen Auer Mühle und Mündung (Fluß-km 76-146) unter besonderer Berücksichtigung der Gewässergüte. - Decheniana (Bonn):140:148=163.
- Habdić,I., Princ,B., Biocenotical classification of the lithorheophilous communities in the karst running waters according to the macro benthic fauna. - Acta hydrochim.hydrobiol. 15:495-503.
- Lago,P.K., Harris,S.C., An annotated list of the Curvipalpia (Trichoptera) of Alabama. - Ent.News 98:255-262.
- McElhone,M.J., Davies,R.W., Culp,J.M., Factors influencing the abundance of Trichoptera in Hartley Creek, a brownwater stream in northeastern Alberta, Canada. - Arch.Hydrobiol. 109:279-285.
- McElravy,E.P., Resh,V., Diversity, seasonality, and annual variability of caddisfly (Trichoptera) adults from two streams in the California Coast Range. - Pan-Pac.Ent. 63:75-91.
- Marlier,G., Observations sur le labre des larves du genre *Oecetis* MacLachlan (Trichoptera, Leptoceridae). - Bull.Inst.R.Sci. Nat.Belg., Ent. 57:187-190.

1987

- Nógrádi, S.U., Néhány adat Szöcse tegzesfaunájához (Trichoptera). - Praenorica, Fol.Hist.-nat.(Szombathely) 2:113-118.
- Pitsch,T., Martin,K., Settele,J., Zur Köcherfliegenfauna des südlichen Pfälzerwaldes. - In:Roweck,H.,(ed.): Beiträge zur Biologie der Grünlandbrachen im südlichen Pfälzerwald. Pollichia-Buch Nr. 12:383-389.
- Tian Li-xin, Li Yu-wen, Trichoptera. In: Agricultural insects, spiders, plant diseases and weeds of Xizang 1:241-250.(Chinese)
- Williams,D.D., Williams,N.E., Trichoptera from cold freshwater springs in Canada: Records and comments. - Proc.Ent.Soc. Ontario 118:13-23.
- Winterbourn,M.J., Invertebrate communities. - In:Viner,A.B.(ed.): Inland waters of New Zealand,pp. 167-190.

1988

- Anonymous, Donald G.Denning (1911-1988). - Perla 8:5-6.
- Berczik,A., Pham Ngoc,L., Hydrobiologische Zustandsänderung während eines Jahrzehntes in einem Mittelgebirgsbach in Ungarn. - Opusc.Zool.Budapest 23:117-132.
- Camargo,J.A., Garcia de Jalón,D., Principales características morfológicas de los géneros ibéricos de la familia Limnephilidae (Trichoptera), en sus últimos estadios larvarios. - Bol.Asoc. esp.Ent. 12:239-258.
- Colburn, E.A., Factors influencing species diversity in saline waters of Death Valley, USA. - Hydrobiologia 158:215-226.
- Czachorowski,S., Caddis flies (Trichoptera) of the River Pasęka (Northern Poland). - Acta Hydrobiol. 30:393-409.
- Dakki,M., Ecosystèmes d'eau courante du haut Sebou (Moyen Atlas): études typologiques et analyses écologique et biogéographique des principaux peuplements entomologiques. - Trav.Inst.Sci. Rabat, Sér.Zool. 42(1987):1-99.
- Doesburg,P.H.van, In memoriam Dr.D.C.Geijsses (1907-1985), een biografie en een bibliografie. - Ent.Ber.(Amst.) 48:133-139.
- Golembowski,U., Quellbäche des Kottenforstes als Lebensraum gefährdeter Tierarten am Beispiel von Cordulegaster bidentatus Selys (Insecta:Odonata). - Decheniana (Bonn) 141:204-208.
- Görtz,M., Limnologische Untersuchungen des Nitzbaches und seiner Nebenbäche (Rheinland-Pfalz). - Decheniana (Bonn) 141:271=287.
- Greenwood,M.T., Hobday,C., The re-discovery of Tinodes pallidulus (McL.) (Trichopt.,Psychomyiidae) in Britain. - Ent.mon.Mag. 124:99-102.
- Hamilton,S.W., Historical Biogeography of two groups of Caribbean Polycentropus (Trichoptera:Polycentropodidae). - In: Liebherr,J.K. (ed.): Zoogeography of Caribbean Insects. Cornell Univ.Press (Ithaca), pp. 153-182.
- Hemphill,N., Competition between two stream dwelling filter-feeders, Hydropsyche oslari and Simulium virgatum. - Oecologia 77:73=80.
- Herrmann,S.J., New record and range extension for Ceraclea resurgens. (Trichoptera:Leptoceridae) from Colorado, with notes on ecological conditions. - Ent.News 99:253-259.

1988

- Humpesch,U.H., Anderwald,P.H., Beitrag zur Faunistik der österreichischen Donau - das Macrozoobenthos bei Stromkilometer 2005. - Wasser und Abwasser 32:41-55.
- Jackson,J.K., Diel emergence, swarming and longevity of selected adult aquatic insects from a Sonoran Desert stream. - Amer. Midl.Nat. 119:344-352.
- Jalón,D.G.de, Trichoptera. In: Barriatos,J.A.(ed.): Bases para un curso práctica de Entomología. (Asoc.Esp.Ent.,Salamanca), pp.521-531.
- Klima,F., Köcherfliegen-Lichtfänge aus dem Bezirk Frankfurt(Oder) (Insecta,Trichoptera). - Beeskower natwiss.Abh. 2:82-86.
- Ko,Myoung-Kyu, Park,Kyu-Tek, A systematic study of Rhyacophilidae (Trichoptera) in Korea. - Korean J.Ent. 18:7-16.
- Kobayashi,Y., Ando,H., Phylogenetic relationships among the lepidopteran and trichopteran suborders (Insecta) from the embryological standpoint. - Z.Zool.Syst.Evolutionsforsch. 26:186-210.
- Ludwig,M., Marthaler,H., Neugebauer,H., Beitrag zur Fauna des Holderbaches im südlichen Odenwald. - Carolinea 46:43-48.
- Maes,J.-M., Flint,O.S.jr., Catalogo de los Trichoptera de Nicaragua. - Rev.Nicar.Ent. 2:1-11.
- Malicky,H., Zur Insektenfauna des Urserentales, Furkastraße 2000m, Kanton Uri. III.Trichoptera (Köcherfliegen). - Ent.Ber. Luzern 20:113-118.
- Mey,W., Bemerkungen zu einigen interessanten Köcherfliegen (Insecta, Trichoptera) aus der Sammlung des Naturhistorischen Museums Rudolstadt/Thür. - Rudolstädter nat.hist.Schr. 1:88-97.
- Moretti,G.P., Tricotteri del Museo Civico di Scienze Naturali "Enrico Caffi" di Bergamo. - Riv.Mus.civ.Sc.Nat."E.Caffi" Bergamo 13:1-19.
- Neboiss,A., Trichoptera. In: Walton,D.W., Houston,W.W.K.(eds): Zoological Catalogue of Australia 6:177-280. Austral.Governm. Publ.Serv.Canberra.
- Nishimoto,H., A new species of Apatania (Trichoptera, Limnephilidae) from Nepal. - Kontyu 56:723-726.
- Nógrádi,S.U., New data to the caddisfly (Trichoptera) fauna of Hungary,II. - Fol.Ent.Hung. 49:205-210.
- Nógrádi U.,S., Uherkovich,A., Trichopterological results from the Northern Mountains (Hungary). - Fol.Hist.-nat.Mus.Matr. 13:71-90.
- Nógrádi U.,S., Slovakian Trichoptera data from two Hungarian collections. - Fol.Hist.-nat.Mus.Matr. 13:91-94.
- Novak,M.A., Reilly,A.A., Jackling,S.J., Long-term monitoring of polychlorinated biphenyls in the Hudson River (New York) using caddisfly larvae and other macroinvertebrates. - Arch. Environ.Contamin.Toxicol. 17:699-710.
- Outridge,P.M., Seasonal and spatial variations in benthic macro-invertebrate communities of Magela Creek, Northern Territory.- Aust.J.Mar.Freshw.Res. 39:211-223.
- Plachter,H., Plachter,J., Ökologische Studien zur terrestrischen Höhlenfauna Süddeutschlands. - Zoologica 47(139):1-67.

1988

- Resh,V.H., Publication patterns in entomology: An example based on aquatic insects. - Bull.ent.Soc.Amer. 34:145-150.
- Reusch,H., Köcherfliegenfänge in Niedersächsischen Tiefland 1984=85 (Insecta, Trichoptera). - Jb.Naturw.Ver.Fstm.Lüneburg 38:291=304.
- Schauwinhold,U., Limnologische Untersuchungen an Bächen des Angerlandes (Kreis Mettmann). - Decheniana 141:230-240.
- Shields,O., Mesozoic history and neontology of Lepidoptera in relation to Trichoptera, Mecoptera, and angiosperms. - J.Paleontol. 62:251-258.
- Stroot,P., Tachet,H., Dolédec,S., Les larves d'Ecnomus tenellus et d'E.deceptor (Trichoptera, Ecnomidae): Identification, distribution, biologie et écologie. - Bijdr.Dierk. 58:259-269.
- Uherkovich,A., Nógrádi,S.U., The Trichoptera of the Bakony Mountains and environs II. (Hungary). - Fol.Mus.Hist.-Nat.Bakonyiensis 7:35-48.
- Yamamoto,E., Goto,A., Nakaya,K., Amaoka,K., Aquatic insect fauna and feeding habits of two Cottus species in the Daitobetsu River of southern Hokkaido. - Bull.Fac.Fish.,Hokkaido Univ. 39:237-256.

1989

- Arens,W., Comparative functional morphology of the mouthparts of stream animals feeding on epilithic algae. - Arch.Hydrobiol. Suppl. 83:253-354.
- Barmuta,L.A., Habitat patchiness and macrobenthic community structure in an upland stream in temperate Victoria, Australia. - Freshw.Biol. 21:223-236.
- Botosaneanu,L., de Vos,R., van Alewijk,G.L.N., Enkele interessante Trichoptera van voornamelijk bronnen en bronbekken op de Veluwezoom. - Ent.Ber.(Amst.) 49:45-49.
- Botosaneanu,L., Notes on the caddisflies (Insecta, Trichoptera) from Isla Margarita (Venezuela). - Bull.Zool.Mus.Amst. 11:203-208.
- Botosaneanu,L., Seconde contribution à l'étude des trichoptères de la Martinique. - Annls Soc.ent.Fr.(N.S.) 25:95-104.
- Botosaneanu,L., Three western Palaearctic caddisflies (Trichoptera) from the British Museum (Natural History). - Ent.Gaz.40:165=169.
- Botosaneanu,L., Adicella melanella (McLachlan, 1884) (Trichoptera, Leptoceridae) représentée au Maroc par une sous-espèce nouvelle. - Ann.Zool.(Warszawa) 42:331-334.
- Burmeister,E.-G., Der Lectotypus von *Macronema maculatum* (Perty, 1833) (Trichoptera, Hydropsychidae). - Spixiana 11:259-262.
- Casado,C., Garcia de Jalon,D., del Olmo,C.M., Barcelo,E., Menes,F., The effect of an irrigation and hydroelectric reservoir on its downstream communities. - Regulated Rivers 4:275-284.
- Cellot,B., Rhythme nyctéméral et distribution verticale de la dérive des macroinvertébrés benthiques dans une grande rivière européenne. - Arch.Hydrobiol. 115:265-286.

1989

- Chantaramongkol,P., Malicky,H., Some Chimarra (Trichoptera: Philopotamidae) from Thailand (Studies on caddisflies from Thailand, No.2). - *Aquat.Insects* 11:223-240.
- Chen,Y.E., Morse,J.C., A preliminary examination of caddisflies from Taiwan, with special reference to Leptoceridae. - *Techn. Contr.(South Carolina Agric.Exper.Stat., Clemson University)* 3004:1-4.
- Christian,A., Köcherfliegen aus der Stadt Görlitz (Insecta, Trichoptera). *Ent.Nachr.Ber.* 33:225-229.
- Czachorowski,S., Chruściki (Trichoptera) nowe dla Pojezierza Pomorskiego. - *Przegl.Zool.* 33:267-269.
- Czachorowski,S., Differentiation of the habitats of Hydropsychidae larvae (Insecta:Trichoptera) in the Pasłka River as a result of avoidance of trophic competition. - *Pol.Arch.Hydrobiol.* 36:123-132.
- Czachorowski,S., Vertical distribution of Trichoptera in three masurian lakes - results of preliminary studies. - *Pol.Arch. Hydrobiol.* 36:351-358.
- Dall,P.C., The life history and occurrence of *Sialis lutaria* L. (Megaloptera) in the littoral zone of Lake Esrom, Denmark. - *Int.Rev.ges.Hydrobiol.* 74:273-281.
- Doledec,S., Tachet,H., Ecological observations and life histories of five net-spinning caddisflies (Trichoptera) of the lower Ardèche river. - *Aquat.Insects* 11:89-99.
- Erman,N.A., Species composition, emergence, and habitat preferences of Trichoptera of the Sagehen Creek Basin, California, USA. - *Great Basin Naturalist* 49:186-197.
- Fairchild,W.L., Wiggins,G.B., Immature stages and biology of the North American caddisfly genus *Phanocelia* Banks (Trichoptera:Limnephilidae). - *Can.Ent.* 121:515-519.
- Feminella,J.W., Power,M.E., Resh,V.H., Periphyton responses to invertebrate grazing and riparian canopy in three northern California coastal streams. - *Freshw.Biol.* 22:445-457.
- Fjellheim,A., Raddum,G.G., Schnell,Ø.A., Changes in benthic animal production of a weir basin after eight years of succession. - *Regulated Rivers* 3:183-190.
- Flint,O.S.jr., Studies of Neotropical caddisflies, XXXIX: The genus *Smicridea* in the Chilean Subregion (Trichoptera:Hydropsychidae). - *Smiths.Contr.Zool.* 472:1-45.
- Flint,O.S.jr., Denning,D.G., Studies of Neotropical caddisflies, XL: New species of *Smicridea* (*Smicridea*) from Middle America and the West Indies (Trichoptera:Hydropsychidae). - *Proc.Biol. Soc.Wash.* 102:418-433.
- Flint,O.S.jr., Denning,D.G., Studies of Neotropical caddisflies, XL1: New species and records of *Austrotinodes* (Trichoptera: Psychomyiidae). - *Pan-Pac.Ent.* 65:108-122.
- di Giovanni,M.V., Goretti,E., Lorenzoni,M., Macrobenthos ad invertebrati del sistema idrico Chiusi-Montepulciano. - *Riv.Idrobiol.* 27:3-38.

1989

- González, M.A., Iglesias, J.C., Une nouvelle espèce de Trichoptère du Sud de l'Espagne: *Oecetis grazalemae* (Trichoptera:Leptoceridae). - Annls Limnol. 25:69-71.
- Graça, M.A.S., Fonseca, D.M., Castro, S.T., The distribution of macroinvertebrate communities in two Portuguese rivers. - Freshwat.Biol. 22:297-308.
- Green, P.C., The use of Trichoptera as indicators of conservation value: Hertfordshire gravel pits. - J.Environmental Management 29:95-104
- Groom, A.P., Hildrew, A.G., Food quality for detritivores in streams of contrasting pH. - J.Anim.Ecol. 58:863-881.
- Gullefors, B., The egg-laying behaviour of *Glyphotaelius pellucidus* (Retzius) (Trichoptera:Limnephilidae). - Fauna norv.Ser.B 36:59-63.
- Gullefors, B., *Oxyethira falcata* (Trichoptera,Hydroptilidae), en för Sverige ny nattslända. - Ent.Tidsk. 110:119-120.
- Harris, S.C., New Trichoptera from Alabama. - J.New York Ent.Soc. 97:309-316.
- Harris, S.C., Contreras-Ramos, A., *Ithytrichia mexicana* (Trichoptera: Hydroptilidae), a new species of caddisfly from Mexico. - Ent.News 100:176-178.
- Hauer, F.R., Stanford, J.A., Ward, J.V., Serial discontinuities in a Rocky Mountain river. II.Distribution and abundance of Trichoptera. - Regulated Rivers 3:177-182.
- Holzenthal, R.W., Studies in neotropical Leptoceridae (Trichoptera), IX: a new genus and species from southeastern Brazil. - Aquatic Insects 11:29-32.
- Huryn, A.D., Identity of the hydropsychid larva known as "Oropsyche?": the immature stages of *Homoplectra flinti* Weaver. - J.N.Amer. Benthol.Soc. 8:112-116.
- Ito, T., Lepidostomatid caddisflies (Trichoptera) from the Tsushima Islands of Japan, with description of a new species. - Jap.J.Ent. 57:46-60.
- Jackson, J.K., Resh, V.H., Distribution and abundance of adult aquatic insects in the forest adjacent to a northern California stream. - Environ.Ent. 18:278-283.
- Klima, F., Zusammenstellung der bisher im NSG "Lange Dammwiesen und Unteres Annatal" (Kreis Strausberg, Bez.Frankfurt/O.) nachgewiesenen Köcherfliegenarten (Trichoptera). - Novius 8:153-155.
- Klima, F., Qualitative Aspekte der Trichopterenfauna im Berliner Seengebiet und Versuch einer Zonierung urbaner limnischer Habitate. - Verh. 11.SIEEC Gotha 1986:86-90.
- Klima, F., *Hydropsyche dinarica* Marinkovic, 1979 (Insecta, Trichoptera) aus dem Rhithral des westlichen Thüringer Waldes - neu für die Fauna der DDR. - Veröff.Naturhist.Mus.Schleusingen 4:90-92.

1989

- Kočharina,S.L., Growth and production of filter-feeding caddisfly (Trichoptera) larvae in a foothill stream in the Soviet Far East. - *Aquatic Insects* 11:161-179.
- Kofler,A., Malicky,H., Mildner,P., Wieser,C., Faunistische Erhebungen in der Lendorfer Au bei Spittal/Drau. - *Carinthia* II 179/99: 697-713. (Trichoptera:706-707)
- Lawson,D.L., Klug,M.J., Microbial fermentation in the hindgut of two stream detritivores. - *J.N.Am.Benthol.Soc.* 8:85-91.
- Leff,L.G., McArthur,J.V., Microbial colonization of limnephilid caddisfly larvae cases. - *Arch.Hydrobiol.* 116:81-84.
- Lelaud,H.V., Fend,S.V., Dudley,T.L., Carter,J.L., Effects of copper on species composition of benthic insects in a Sierra Nevada, California, stream. - *Freshwater Biol.* 21:163-179.
- Li,J.L., Gregory,S.V., Behavioral changes in the herbivorous caddisfly *Dicosmoecus gilvipes* (Limnephilidae). - *J.N.Am.Benthol.Soc.* 8:250-259.
- Li Youwen, Tian Lixin, Notes on a new subgenus and a new species of *Hydropsyche*. - *J.Nanjing Agric.Univ.* 12:44-45.
- Li Youwen, Tian Lixin, Dudgeon,D., Notes on six new species of *Hydropsychidae* (Trichoptera). - *J.Nanjing Agric.Univ.* 13:37-42.
- McElravy,E.P., Lamberti,G.A., Resh,V.H., Year-to-year variation in the aquatic macro-invertebrate fauna of a northern California stream. - *J.N.Am.Benthol.Soc.* 8:51-63.
- Malicky,H., Köcherfliegen (Trichoptera) von Sumatra und Nias: Die Gattungen *Chimarra* (Philopotamidae) und *Marilia* (Odontoceridae), mit Nachträgen zu *Rhyacophila* (Rhyacophilidae). - *Mitt.Schweiz.Ent.Ges.* 62:131-143.
- Malicky,H., *Chaetopterygopsis siveci* sp.n. from Greece (Trichoptera, Limnephilidae). - *Aquatic Insects* 11:152.
- Malicky,H., Krušnik,C., *Chaetopteryx marinkoviae* sp.n. (Trichoptera, Limnephilidae) from Istria, Yugoslavia. - *Aquat.Insects* 11:180.
- Malicky,H., Ein neuer Drusus aus Nord-Griechenland (Trichoptera: Limnephilidae). - *Ent.Z.(Essen)* 99:303-304.
- Malicky,H., Eine ergänzte Liste der österreichischen Köcherfliegen (Insecta, Trichoptera). - *Z.Arbgem.Öst.Ent.* 41:32-40.
- Malicky,H., Zur Köcherfliegenfauna des Monte Generoso, Kanton Tessin, Südschweiz (Trichoptera). - *Ent.Ber.Luzern* 21:51=60.
- Malicky,H., Chantaramongkol,P. Beschreibung von neuen Köcherfliegen (Trichoptera) aus Thailand und Burma (Arbeiten über thailändische Köcherfliegen Nr. 6). - *Ent.Ber.Luzern* 22:117-126.
- Marchant,R., Changes in the benthic invertebrate communities of the Thomson River, Southeastern Australia, after dam construction. - *Regulated Rivers* 4:71-89.
- Mey,W., Levanidova,I.M., Revision der Gattung *Apataniana* Mosely, 1936 (Trichoptera, Limnephilidae). - *Dt.ent.Z.,N.F.* 36:65=98.
- Mey,W., Taxonomische und faunistische Notizen zu einigen Köcherfliegen (Trichoptera) aus Korea. - *Acta Ent.Bohemoslov.* 86:295=305.

1989

- Mey,W., Setodes fluviavirens sp.n. - a new potamobiontic species from the Mekong River of Cambodia (Trichoptera:Leptoceridae).- Aquatic Insects 11:125-127.
- Mey,W., Jung,R., Neue Köcherfliegen aus dem westlichen Tienschan (UdSSR)(Trichoptera). - Dt.ent.Z.,N.F. 36:283-292.
- Mey,W., Joost,W., Beschreibung der Larve und der Puppe von *Synagapetus dubitans* McLachlan (Insecta, Trichoptera) aus Thüringen. - Rudolstädter nat.hist.Schr. 2:57-62.
- Meyer.E., The relationship between body length parameters and dry mass in running water invertebrates. - Arch.Hydrobiol. 117:191-203.
- Morse,J.C., A redescription of *Symphitoneuria dammermanni* Ulmer, 1951 (Trichoptera:Leptoceridae). - Ent.Mitt.Zool.Mus.Hamburg 9 (135):205-212.
- Morse,J.C., Hamilton,S.W., Hoffman,K.M., Aquatic insects of Lake Jocassee catchment in North and South Carolina, with descriptions of four new species of caddisflies (Trichoptera).- J!Elisha Mitchell Sci.Soc. 105:14-33.
- Neboiss,A., Jackson,J., Walker,K., Caddis-flies (Insecta:Trichoptera) of the World Heritage Area in Tasmania - species composition and distribution. - Occ.Pap.Mus.Victoria 4:1-41.
- Neboiss,A., Walker,K., Types of Trichoptera (Insecta) held in the Museum of Victoria. - Occ.Pap.Mus.Victoria 4:43-56.
- Neboiss,A., Additions and corrections to the Atlas of Trichoptera of the SW Pacific-Australian region. - Occ.Pap.Mus.Victoria 4:63-67.
- Neboiss,A., Caddis-flies (Trichoptera) of the families Polycentropodidae and Hyalopsychidae from Dumoga-Bone National Park, Sulawesi, Indonesia, with comments on identity of *Polycentropus orientalis* McLachlan. - Bull.Zool.Mus.Univ.Amsterdam 12:101=109.
- Nógrádi,S., The trichoptera fauna of Szöce and its environs (West Hungary). - Janus Pannonus Múz.Evkönyve 33:27=33.
- Nógrádi,S.U., Uherkovich,A., On the caddisfly fauna of the Kőszeg mountains, Hungary (Trichoptera). - Janus Pannoniüs Múz. Evkönyve 33:35-42.
- Nógrádi,S.U., Locality data of the Trichoptera collection originating from the Carpathian Basin in the Hungarian Natural History Museum. - Fol.Ent.Hung. 50:147-156.
- Novák,K., Das Vorkommen der Trichopteren in der Moldau in Prag. - Verh.11.SIEEC Gotha (1986):339-341.
- Oláh,J., Thirty-five new hydroptilid species from Vietnam (Trichoptera, Hydroptilidae). - Acta Zool.Hung. 35:255-293.
- Ortlepp,J., Schröder,P., Rey,P., Tomka,I., The longitudinal zonation of macroinvertebrates in the River Rhine. - 24.Congr. Int.Ass.Limnol., Manuscript (13pp.).
- Otto,C., Effects of information asymmetries in contests between net spinning caddis larvae (*Plectrocnemia conspersa*). - Oecologia 81:176-180.
- Otto,C., Wallace,J.B., Life cycle variation and habitat longevity in waterlily leaf beetles. - Holarct.Ecol. 12:144-151.

1989

- Peeters,E.T.H.M., Tachet,H., Comparison of macrobenthos in braided and channelized sectors of the Drôme River, France.- Regulated Rivers 4:317-325.
- Petersen,L.B.-M., Resource utilization of coexisting species of Hydropsychidae (Trichoptera). - Arch.Hydrobiol.Supp.83:83=119.
- Petersson,E., Swarming activity patterns and seasonal decline in adult size in some caddis flies (Trichoptera: Leptoceridae).- Aquatic Insects 11:17-28.
- Petersson,E., Mating in swarming caddis flies (Trichoptera: Leptoceridae). - Acta Univ.Upsaliensis; Compreh.summ.of Uppsala Dissertations from the Faculty of Science, 222.
- Rader,R.B., Ward,J.V., The influence of environmental predictability/ disturbance characteristics on the structure of a guild of mountain stream insects. - Oikos 54:107-116.
- Resh,V.H., Donald G.Denning (1909-1988). - Pan-Pacific Ent.65:97=107.
- Riedel,W., Majecki,J., Postacie doskonałe chrząszcików (Trichoptera) Gór Świętokrzyskich. - Frag.Faun.(Warszawa) 32:227-241.
- Robert,B., Schmidt,C., Retzlaff,H., Schulze,W., Die bisher aus der Senne bekannten Arten der Köcherfliegen (Trichoptera).- Mitt.Arbgem.ostwestfälisch-lippischer Ent. 5(3):89-98.
- Roeding,C.E., Smock,L.A., Ecology of macroinvertebrate shredders in a low-gradient sandy-bottomed stream. - J.N.Am.Benthol. Soc. 8:149-161.
- Rutt,G.P., Weatherley,N.S., Ormerod,S.J., Microhabitat availability in Welsh moorland and forest streams as a determinant of macroinvertebrate distribution. - Freshw.Biol.22:247-261.
- Schöll,F., Zur näheren Kenntnis des Makrozoobenthos der Fließgewässer im Nationalpark Bayerischer Wald. - Ent.Z.(Essen) 99:257-267.
- Schröder,P., Rey,P., Erfassung des biologischen, chemischen und physikalischen Zustandes ausgewählter Zuflüsse des Rheins in der Region Basel. - Zwischenbericht,Zool.Inst.Univ. Fribourg, 158pp.
- Sipahiler,F., Seven new species and a new subspecies of Trichoptera from south western Anatolia. - Aquat.Insects 11:129-140.
- Sipahiler,F., Four new species of Trichoptera from Turkey (Glossosomatidae, Ecnomidae, Lepidostomatidae). - Opusc.zool.flumin. 39:1-7.
- Spuris,Z., Latvijas kukaiņu katalogs. 7.Makstenes (Trichoptera).- Latv.Ent.32:5-42.
- Statzner,B., Morphological adaptation of shape to flow: Micro-currents around lotic macroinvertebrates with known Reynolds numbers at quasi-natural flow conditions. - Oecologia 78:145=157.
- Statzner,B., Holm,T.F., Morphological adaptation of shape to flow: Micro-currents around lotic macroinvertebrates with known Reynolds numbers at quasi-natural flow conditions.- Oecologia 78:145-157.
- Stoltze,M., The Afrotropical caddisfly family Pisuliidae. Systematics, zoogeography, and biology (Trichoptera: Pisuliidae). - Steenstrupia 15(1):1-49.

1989

- Stout,B.M.,III, Stout,J.S., Northern caddisfly (Trichoptera) fauna in a remnant boreal wetlands of West Virginia. - Ent.News 100:37-40.
- Strommer,J.L., Smock,L.A., Vertical distribution and abundance of invertebrates within the sandy substrate of a low-gradient headwater stream. - Freshw.Biol. 22:263-274.
- Stroot,P., The variability of larval coloration patterns of *Agraylea multipunctata* in a population from Belgium (Trichoptera: Hydroptilidae). - Ent.Ber.(Amst.) 49:157-160.
- Stroot,P., Neven,B., A propos de la présence en Belgique de *Molannodes tinctus* (Zetterstedt,1840), Trichoptère nouveau pour la faune belge. - Bull.Inst.R.Sci.Nat.Belg.,Ent.58:179=182.
- Stroot,P., Depiereux,E., Proposition d'une méthodologie pour établir des "listes rouges" d'invertébrés menacés. - Biological Conservation 48:163-179.
- Szczesny,B., The Tatra Mountains. - Leaflet, 7pp.: 6th Int.Symp. Trich.Łódź-Zakopane.
- Tian Lixin, Zheng Jianzhong, Description of two new species of the genus *Stenopsyche* McLachlan from China (Trichoptera).- J.Nanjing Agric.Univ. 12:49-52. (Chinese)
- Tudorancea,C., Baxter,R.M., Fernando,C.H., A comparative limnological study of zoobenthic associations in lakes of the Ethiopian Rift Valley. - Arch.Hydrobiol.Suppl. 83:121-174.
- Usis,J.D., Foote,B.A., New record of caddisflies (Trichoptera) from Ohio, with particular reference to Stillfork Swamp, Carroll County. - Ent.News 100:83-85.
- Usseglio-Polatera,P., Bournaud,M., Trichoptera and Ephemeroptera as indicators of environmental changes of the Rhône river at Lyons over the last twenty-five years. - Regulated Rivers 4:249-262.
- Waringer,J.A., The abundance and temporal distribution of caddis flies (Insecta:Trichoptera) caught by light traps on the Austrian Danube from 1986 to 1987. - Freshw.Biol.21:387=399.
- Waringer,J.A., Resistance of a cased caddis larva to accidental entry into the drift: the contribution of active and passive elements. - Freshwater Biol. 21:411-420.
- Waringer,J.A., Life cycle, horizontal microdistribution and current resistance of *Allogamus auricollis* (Trichoptera:Limnephilidae) in an Austrian mountain brook. - Freshwater Biol. 22:177=188.
- Weaver,John S.,III, Indonesian Lepidostomatidae (Trichoptera) collected by Dr.E.W.Diehl. - Aquatic Insects 11:47-63.
- Weinzierl,A., Dorn,A., Stein- und Köcherfliegen-Nachweise entlang der Münchener Isar. - Lauterbornia 1:6-22.
- Wells,A., Wichard,W., Caddisflies of Dominican amber. VI.Hydroptilidae (Trichoptera). - Stud.Neotrop.Fauna Environm. 24:41=51.
- Wichard,W., Köcherfliegen des Dominikanischen Bernsteins. VII. Fossile Arten der Gattung *Cubanoptila* Sykora,1973. - Mitt. Münch.Ent.Ges. 79:91-100.
- Wichard,W., Anpassungen von Köcherfliegen (Trichoptera) an periodische Gewässer. - Verh.Westd.Ent.Tag 1988:79-88.

1989

- Wiggins,G.B., Wichard,W., Phylogeny of pupation in Trichoptera, with proposals on the origin and higher classification of the order. - J.N.Am.Benthol.Soc. 8:260-276.
- Wiggins,G.B., Larson,D.J., Systematics and biology for a new Nearctic genus in the caddisfly family Phryganeidae (Trichoptera). - Can.J.Zool. 67:1550-1556.
- Wiggins,G.B., Richardson,J.S., Biosystematics of Eocosmoecus, a new Nearctic caddisfly genus (Trichoptera: Limnephilidae, Dicosmoecinae). - J.N.Am.Benthol.Soc. 8:355-369.
- Williams,N.E., Factors affecting the interpretation of caddisfly assemblages from Quaternary sediments. - J.Paleolimnology 1:241-248.
- Wimmer,J., Ein Beitrag zur Kenntnis der Trichopterenfauna von Oberösterreich. - Steyrer Entomologenrunde 23:38-64.
- Winterbourn,M.J., Gregson,K.L.D., Guide to the Aquatic Insects of New Zealand. - Bull.Ent.Soc.NZ 9:1-95.
- Wright,J.F., Armitage,P.D., Furse,M.T., Moss,D., Prediction of invertebrates communities using stream measurements.- Regulated Rivers 4:147-155.
- Yagi,M., Yoshiyasu,Y., Sasakawa,M., A peculiar organ newly found in the Hydropsychidae (Trichoptera). - Jap.J.Ent. 57:176-183.
- Yang Lian-fang, Morse,J.C., Setodini of the People's Republic of China (Trichoptera: Leptoceridae, Leptocerinae). - Contr. Amer.Ent.Inst. 25(4):1-77.
- Yang Liangfang, Tian Lixin, Four new species and two new records of the genus Ceraclea Stephens (Trichoptera: Leptoceridae). - Entomotaxonomia 11:293-297. (Chinese)
- Yoon Il Byong, Ki Hong Kim, A systematic study of the caddisfly larvae in Korea. - Korean J.Ent. 19:25-40.
- Zullini,A., The ecology of the Lambro River. - Riv.Idrobiol. 27:39-58.

1990

- Andersen,T., Hansen,L.O., Johanson,K.A., Solhøy,T., Søli,G.E.E., Faunistical records of Caddis flies (Trichoptera) from Aust-Agder and Vest-Agder, South Norway. - Fauna norv.Ser.B 37:23-32.
- Andersen,T., Hossain,M., Solhøy,T., Søli,G.E.E., Caddisflies (Trichoptera) from Jostedalen, West Norway. - Fauna norv. Ser.B 37:37-41.
- Andersen,T., Søli,G.E.E., Further additions to the caddis fly fauna (Trichoptera) in Vestfold, SE Norway. - Fauna norv. Ser.B 37:46.
- Botosaneanu,L., About Far-eastern Stactobia McLachlan (Trichoptera: Hydroptilidae): a correction. - Aquatic Insects 12:47-48.
- Burmeister,E.-G., Makroinvertebraten der Isar und ihrer Nebengewässer in und südlich von München. - Lauterbornia 4:7=23.

1990

- Füchs,U., Statzner,B., Time scales for the recovery potential of river communities after restoration: Lessons to be learned from smaller streams. - Regulated Rivers 5:77-87.

Gullefors,B., Det sjätte internationella nattländesymposiet, Lodz-Zakopane, Polen 11-16 september 1989. - Ent.Tidskr.111:44=46.

Kobayashi,Y., Ando,H., Early embryonic development and external features of developing embryos of the caddisfly, *Nemotaulius admorsus* (Trichoptera: Limnephilidae). - J.Morphol.203:69=85.

Malicky,H., Revision der Gattung *Melampophylax* Schmid,1955 (Trichoptera, Limnephilidae). - Nachrbl.bayer.Ent. 39:1=13.

Matczak,T.Z., Mackay,R.J., Territoriality in filter-feeding caddisfly larvae: Laboratory experiments. - J.N.Am.Benthol.Soc.9:26=34.

O'Connor,J.P., Good,J.A., Bond,K.G.M., Leptocerus tineiformis Curtis: a caddisfly new to Ireland from Pollardstown Fen, Co.Kildare. - Ent.Rec. 102:115-116.

Robert,B., Schmidt,C., Die Köcherfliegen (Trichoptera) des rheinisch-westfälischen Tieflandes. - Ent.Z.(Essen) 100:98-105.

Waringer-Löschenkohl,A., Waringer,J., Zur Typisierung von Augewässern anhand der Litoralfauna (Evertebraten, Amphibien).- Arch.Hydrobiol.Suppl. 84:73-94.

Wells,A., The Micro-Caddisflies (Trichoptera:Hydroptilidae) of North Sulawesi. - Invetrebr.Taxon. 3:363-406.

X X

LIST OF RESEARCH WORKERS ON TRICHOPTERA

Cynthia R.L.ADLER Dept.of Biological Sciences, Clemson University, Clemson South Carolina 29634-1903, USA
Present interest: Chimarra; ecology of immatures, association of immatures with adults to facilitate specific determination of immatures. Previously studied: Lepidostoma, Pycnopsyche.
Investigation area: Mid-Atlantic and Southeastern North America.
Willing to identify for other workers: Eastern Chimarra immatures.

x x x x x x x x x x

Gabriela S. ROZAS DENNIS, Biologist, Instructor.
Universidad Nacional del Sur, Depart.de Biología, Perú 670,
C.P.8000., B.Blanca, Buenos Aires, Argentina.

Present interest: Hydropsychidae, Leptoceridae, Hydroptilidae. Gut contents of filter-feeding Trichoptera; Trichoptera living in organic polluted streams. Investigation area: Neotropis. Information wanted: Papers on Trichoptera, esp. on ecology, effects of organic pollution on stream caddisflies. Other interests: Function of other macroinvertebrates as bioindicators.

INFORMATION ON TRICHOPTERA FROM SOUTHEASTERN ASIA

Many colleagues have sent their informations. A summary will be given in Newsletter No.18 next year.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Trichoptera Newsletter](#)

Jahr/Year: 1990

Band/Volume: [17](#)

Autor(en)/Author(s): Anonym

Artikel/Article: [Trichopterological Literature 39-51](#)