



The Taxonomic Report

OF THE INTERNATIONAL LEPIDOPTERA SURVEY



ISSN 2643-4776 (print) / ISSN 2643-4806 (online)

Notable Hesperidae collected by Kilian Roever in Arizona, USA

Jing Zhang^{1,2,3}, Qian Cong^{1,3}, Jinhui Shen^{1,2}, Leina Song^{1,2}, and Nick V. Grishin^{1,2*}

Departments of ¹Biophysics, ²Biochemistry, and ³Eugene McDermott Center For Human Growth & Development, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, TX 75390-9050, USA;

*Corresponding author: grishin@chop.swmed.edu

ABSTRACT. Genomic analysis of Hesperidae specimens collected by Kilian Roever in Arizona, USA, and deposited in the McGuire Center for Lepidoptera and Biodiversity (Florida Museum of Natural History, Gainesville, FL, USA) confirms several published state species records, reveals that *Cecropterus* (*Murgaria*) *markwalkeri* Grishin, 2023, *Telegonus* (*Telegonus*) *misitra* (Plötz, 1881), and *Epargyreus cruza* Evans, 1952 have been found in Arizona with sequenced specimens representing first confirmed USA records of these three species, and allows us to discover new taxa *Cecropterus* (*Murgaria*) *roeveri* Grishin, **sp. n.** and *Lerema* (*Lerema*) *ochrius occidus* Grishin, **ssp. n.** This study underscores the value of extensive collecting and careful specimen preservation for research.

Key words: taxonomy, classification, genomics, phylogeny, biodiversity.

ZooBank registration: <http://zoobank.org/4F75FF52-85A5-494B-8757-52A732DCB578>

INTRODUCTION AND METHODS

In contrast to photographs, which are only as good as the pose and our ability to recognize a species from its appearance, collected and carefully preserved specimens enable future studies using various methods. Dissection of genitalia is essential for the identification of many Hesperidae Latreille, 1809 that cannot be confidently told apart by their wing patterns (Evans 1951, 1952, 1953, 1955). Recently, DNA-based approaches, in particular, those deriving power from whole genome sequencing, offer a way of confident identification, species discovery, and even determining collecting localities by genomic comparison (Cong et al. 2021). We use these methods to study butterflies, and this work builds upon our previous publications, following similar principles and employing the same techniques (Cong et al. 2019a, b; Li et al. 2019; Zhang et al. 2019a–d; Cong et al. 2020; Zhang et al. 2020; Cong et al. 2021; Zhang et al. 2021; Robbins et al. 2022; Zhang et al. 2022a, b; 2023a, c–g; 2024a–c).

Here, we apply whole genome sequencing and analysis to specimens of rarely observed species reaching their northern distribution limits in the southern USA. We focus on specimens collected by Kilian Roever in southeastern Arizona and deposited in the McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, Gainesville, FL, USA (MGCL). A leg was sampled from each specimen for DNA extraction and sequencing. The methods were exactly as those described in our latest two publications (Zhang et al. 2024a, b), and the introduction to Zhang et al. (2024b) summarizes general principles we use for species delineation. Here, for brevity, we show only two genomic trees in each figure: one nuclear tree (either constructed from protein-coding genes in autosomes, or the Z chromosome when it reveals better separation between species) and the mitochondrial genome tree.

Sections of this work are arranged in taxonomic order deduced from genome-scale phylogeny complemented by phenotypic considerations. For the new taxa, in addition to brief phenotypic diagnoses frequently accompanied by references that discuss and illustrate morphological characters in greater detail, we provide diagnostic DNA characters in the nuclear genome and in the COI barcode. DNA characters are found in nuclear protein-coding regions using our previously developed procedure (see SI Appendix to Li et al. 2019). The logic behind the character selection was described in Cong et al. (2019b) and is aimed at finding more robust characters likely to stand when additional specimens and species are sequenced. The character states are given in species diagnoses as abbreviations. E.g., aly728.44.1:G672C means position 672 in exon 1 of gene 44 from scaffold 728 of the *Cecropterus lyciades* (Geyer, 1832) (aly, because this species was formerly in the genus *Achalarus* Scudder, 1872) reference genome (Shen et al. 2017) is C, changed from G in the ancestor. When characters are given for the sister clade of the diagnosed taxon, the following notation is used: aly5294.20.2:A548A (not C), which means that position 548 in exon 2 of gene 20 on scaffold 5294 is occupied by the ancestral base pair A, which was changed to C in the sister clade (so it is not C in the diagnosed taxon). The same notation is used for COI barcode characters, but without a prefix ending with ‘:’. The sequences of exons from the reference genome with the positions used as character states highlighted in green are in the supplemental file deposited at < <https://osf.io/sp7fn/> >. This link to the DNA sequences accessible from this publication ensures that DNA characters given in the diagnoses can be readily associated with actual sequences.

Whole genome shotgun datasets we obtained and used in this work are available from the NCBI database < <https://www.ncbi.nlm.nih.gov/> > as BioProject PRJNA1243725 and BioSample entries of the project contain the locality and other collection data of the sequenced specimens shown in the trees. For each specimen in tree figures, the following information is provided (separated by ‘|’): taxon name with comments in square brackets, DNA sample code, type status, general locality, and year of collection (‘old’ if not dated and likely collected 100–150 years ago). Type status abbreviations are: HT holotype, LT lectotype, ST syntype, ?ST possible syntype, NT neotype, PT paratype, PLT paralectotype; and if a synonym name is given (in parenthesis, preceded by ‘=’), type status refers to the synonym. COI barcode sequences reported here have been deposited in GenBank with accessions [PV408305](#) and [PV408306](#). Abbreviations or acronyms for collections are listed in the acknowledgments section.

RESULTS AND DISCUSSION

Specimens of species rarely encountered in the USA were selected for genomic sequencing from the Roever collection, now in the McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, Gainesville, FL, USA. Genomic comparison with other sequenced specimens, including many primary types, provided identification of these species, mostly confirming previously published phenotype-based identifications (Bailowitz and Brock 1991, 2022) corrected for the updates in taxonomy since then (Warren et al. 2024). These advances in our understanding of taxonomy and recent descriptions of new species (Zhang et al. 2023b, f) resulted in three confirmed new US records of butterflies: *Cecropterus (Murgaria) markwalkeri* Grishin, 2023 (type locality in Mexico: Sonora), *Telegonus (Telegonus) misitra* (Plötz, 1881) (type locality in Mexico), and *Epargyreus cruza* Evans, 1952 (type locality in Mexico: Veracruz). Finally, a new species and a new subspecies were discovered and are described here. Details are given for each taxon below.

Cecropterus (Murgaria) markwalkeri Grishin, 2023

As hypothesized in Zhang et al. (2024b), we confirm by genomic analysis that at least some records referred to as *Cecropterus albociliatus* (Mabille, 1877) (type locality in Colombia, Panama, Guatemala) from specimens collected by Kilian Roever in southeastern Arizona (Bailowitz and Brock 1991, 2022) are *Cecropterus (Murgaria) markwalkeri* Grishin, 2023 (type locality in Mexico: Sonora) (Fig. 1 shaded yellow). We sequenced the following three specimens from USA, Arizona, Santa Cruz Co.: NVG-

24074G12 Mt. Hopkins Rd., 8 mi E of Amado, 18-Jun-1978 (Fig. 2a, genitalia Fig. 3); NVG-24074H01 Santa Cruz Mts., Temporal Gulch, 18-Jun-1978 (Fig. 2b); and NVG-24074H02 Sycamore Canyon near Ruby, 26-Aug-1988.




Fig. 1. Phylogenetic trees of selected *Cecropteris* (*Murgaria*) and *Spicauda* species inferred from protein-coding regions in **a)** the Z chromosome, based on 361,947 positions, and **b)** the mitochondrial genome: *C. markwalkeri* (red), *C. albociliatus* (blue), *C. roeveri* sp. n. (magenta), *C. coyote* (green), *C. nigrociliata* (purple), *S. procne* (cyan), and *S. simplicius* (olive). Ultrafast bootstrap (Minh et al. 2013) values are shown at nodes. Labels of specimens from Arizona and Texas are highlighted in yellow and green, respectively. For each specimen, the name adopted in this work is given first, supplemented with the DNA Sample number, type status (see Introduction for abbreviations), general locality, and collection year. Synonyms are given in parentheses preceded by “=” (in subsequent figures). The type status refers to this synonym if the synonym name is provided. The same notations are used throughout this work in other figures showing phylogenetic trees.



Fig. 2. *Cecropterus (Murgaria) markwalkeri* from USA, Arizona in dorsal (left) and ventral (right) views, data in text: **a)** NVG-24074G12 and **b)** NVG-24074H01.



Fig. 3. Male genitalia of *Cecropterus markwalkeri* NVG-24074G12 from AZ: left valva in right lateral view (left) and the rest of the genitalia in left lateral (middle) and dorsal (right) views; unnumbered dissection by Kilian Roever or his helpers.

Cecropterus (Murgaria) roeveri Grishin, new species

<http://zoobank.org/27B71385-5F34-426D-8446-BBBE8549DD27>

(Figs. 1 part, 4, 5a)

Definition and diagnosis. A specimen of *Cecropterus (Murgaria)* E. Watson, 1893 (type species *Telegonus albociliatus* Mabille, 1877) collected by Kilian Roever in southeastern Arizona (Fig. 1 magenta) is sister to both *Cecropterus (Murgaria) coyote* (Skinner, 1892) (type locality in USA: Southern Texas) (Fig. 1 green) and *Cecropterus (Murgaria) nigrociliata* (Mabille & Boulet, 1912) (type locality in Mexico) (Fig. 1 purple) in the Z chromosome and the mitochondrial genome trees, and is genetically differentiated from them at the species level, e.g., their COI barcodes differ by 3.3% (22 bp) from *C. coyote* and by 2.7% (18 bp) from *C. nigrociliata*, and therefore represents a new species. This new species keys to “*Achalarus toxeus*” (C.17.4) in Evans (1952), who misidentified *Aethilla toxeus* Plötz, 1882 (type locality in Mexico), currently treated as a junior subjective synonym of *Cecropterus (Murgaria) albociliatus albociliatus* (Mabille, 1877) (type locality in Colombia, Panama, and Guatemala) (Zhang et al. 2023f). Evans’s “*A. toxeus*” corresponds to the species group with *C. coyote* and *C. nigrociliata*. The new species differs from others by its male genitalia (Fig. 5a) with harpe narrowing in the middle and turning posterodorsad and narrowing to a point directed anterodorsad (harpe of both *C. coyote* and *C. nigrociliata* (Fig. 5b) is blade-like, relatively straight, terminally rounded and dorsally serrated, broader and shorter in *C. coyote* and narrower and longer in *C. nigrociliata*), broader valva, longer tegumen, and thicker and shorter uncus arms. The general shape of the new species’ genitalia is somewhat reminiscent of some *Epargyreus* Hübner, [1819] (type species *Papilio tityrus* Fabricius, 1775, a junior homonym, considered a subjective synonym of *Papilio clarus* Cramer, 1775). While this new species is diagnosed by male genitalia, due to unexplored individual variation and other undescribed species, most reliable identification is achieved by DNA, and a combination of the following base pairs is diagnostic in the nuclear genome: alY10226.35.2:A171G, alY10226.35.2:G180A, alY10226.35.2:G186T, alY6654.9.15:G549A, alY6654.9.15:G558A, alY2633.9.6:C30C (not T), alY173.58.7:A72A (not G), alY1205.1.8:C81C (not T), alY1205.1.8:C93C (not T), alY173.58.7:G63G (not A), and COI barcode: T19T (not C), 205T, T208T (not G), T286C, T472C, C517C (not T).

Barcode sequence of the holotype. Sample NVG-24074H03, GenBank [PV408305](https://www.ncbi.nlm.nih.gov/nuclot/PV408305), 658 base pairs:

```
AAC TT TAT AT TTT AT TTT TTT TGA AT TTG AGC AGG ATTA AT TGG AAC CT CAT TAAG AT TACT TAT TCG AAC TGA AT TAG GA ACC CAG GAT CTT TAA TTG GAG ATG AT CAA AT TTT ATA TACT  
ATT GTA ACAG CT CAT GCT TTT TAT TATA AT TTT TTT TAT AG TTAT ACCT AT TATA AT TGG AGG AT TTT GGA AAT TGA TTA AT TCC TCT TATA CTAG GAG CCCC GAT ATAG CTT TCC CT CGTA  
TAA ATA TATA AG AT TTT GAT TAT TAC CCCC AT CCT TAA CT CT TAA TT TCA AGA AGA AT TGT AGA AAA CCG GTGC AGG TACT GGAT GAAC AGT TTT AT CCCC CTT TAT CAT CCA AT TAT GC  
CC ATCA AGGG CAT CAG TAG AT TTAG CA AT TTT TTT CATT ACAT CTAG CTGG GAT TTT CCT CTAT TCT TGG AGCT ATTA AT TTT TATT ACA ACT AT TATTA AT ATAC GAAT CAATA AT TTTAT CA  
TTT GAC CAA ATAC CAT TAT TTT TTT GAG CTGT CGGA ATTAC AGCT TTT AT TAT TAT TACT TTT CATT ACCT GTTT TAG CAG GAG CTAT TAC CAT AT TAT TAA CTG ATCG AAA TTT TAA TACT T  
CT TTT TTT TAT CAG CAG GTGG GGTG ATCCT AT TTT TAT ATCA AC AT TTT AT T
```



Fig. 4. *Cecropterus (Murgaria) roeveri* sp. n. holotype ♂ NVG-24074H03 in dorsal (left) and ventral (right) views, data in text.




Fig. 5. Male genitalia of *Cecropterus* (*Murgaria*) in left lateral (left) and dorsal (right) views: **a)** *C. roeveri* **sp. n.** holotype NVG-24074H03; **b)** *C. nigrociliata* NVG-24064F03 Mexico: Colima, Comala, 14-Nov-1982, D. W. Jenkins leg. [MGCL].

Type material. Holotype: ♂ deposited in the McGuire Center for Lepidoptera and Biodiversity Collection, Gainesville, FL, USA (MGCL), illustrated in Fig. 4 (genitalia Fig. 5a), bears the following six printed (text in *italics* handwritten) rectangular labels, five white: [Arivaca Creek, 3600' | at Arivaca, Pima Co. | AZ ↑ 7 mi. W. of], [22 Sept.1984 | leg. K. Roever], [MGCL Collection | W. McGuire coll. | Ex. K. Roever], [DNA sample ID: | NVG-24074H03 | c/o Nick V. Grishin], [genitalia | NVG241111-34 | c/o Nick V. Grishin], and one red [HOLOTYPE ♂ | *Cecropterus* (*Murgaria*) | roeveri Grishin].

Type locality. USA: Arizona, Pima Co., 7 mi west of Arivaca, Arivaca Creek, elevation 3600'.

Etymology. The name, a noun in the genitive case, honors Kilian Roever, the collector of the holotype of this new species and numerous other butterfly records. Kilian's deep passion for nature exploration, extensive knowledge, and countless hours in the field were instrumental in discovering all these rarely observed species in Arizona and beyond, significantly enhancing our understanding of them. Without his immense collecting efforts and excellent specimen preservation, which ensured high-quality DNA, this and many other studies would have been impossible. Kilian eagerly shares stories of his field experiences and generously provides details about the localities of his fascinating discoveries, for which we are deeply grateful. His generosity has enriched us all.

Distribution. Currently known only from the holotype collected in southeastern Arizona, expected in western Mexico.

***Spicauda procne* (Plötz, 1881) in Arizona and New Mexico**

Genomic analysis of several *Spicauda* Grishin, 2019 (type species *Goniurus procne* Plötz, 1881) specimens collected in SE Arizona, and SW New Mexico by Kilian Roever confirms their identification as *Spicauda procne* (Plötz, 1881) (type in Brazil) (Fig. 1 cyan, shaded yellow). The following five specimens were sequenced: 1♂ NVG-24074E08 New Mexico, Hidalgo Co., Peloncillo Mts., Cottonwood Canyon, 0-1.5 mi E of AZ state line, 23-Sep-1990 (Fig. 6) and Arizona, Santa Cruz Co.: 2♀♀ NVG-24087B10 and NVG-24087B11 Atascosa Mts., Sycamore Canyon, 27-Sep-1959; 1♂ NVG-24087C01 3 mi SW of Sonoita, Cottonwood Spring, 5-Oct-1969; and 1♂ NVG-24087B12 O'Donnell Canyon at Ewing



Fig. 6. *Spicauda procne* ♂ NVG-24074E08 from USA, New Mexico in dorsal (left) and ventral (right) views, data in text.




Fig. 7. *Telegonus* (*Telegonus*) *misitra* ♂♂ from USA: Arizona in dorsal (left) and ventral (right) views, data in text:
 a) NVG-24087C03 and b) NVG-24087B09.

Ranch, nr. Canelo, 2-Sep-1973. None of the Arizona specimens we sequenced was *Spicauda simplicius* (Stoll, 1790) (type locality in Suriname). However, we confirm *S. simplicius* from Texas, Cameron Co., Brownsville [USNM]: 1♂ NVG-19121D01, USNMENT 01602723 22-Jul-1922, E. M. Smith leg, G. W. Rawson collection and 1♀ NVG-19121C12, USNMENT 01201575 8-Jun-1944 from J. C. Hopfinger collection (Fig. 1 olive, shaded green).

Telegonus (Telegonus) misitra (Plötz, 1881)

Genomic analysis of two males from the *fulgerator* species group of *Telegonus* Hübner, [1819] (type species *Papilio talus* Cramer, 1777) collected by Kilian Roever in Arizona, Santa Cruz Co., Atascosa Mts., Sycamore Canyon (NVG-24087C03 8-May-1960 (Fig. 7a) and NVG-24087B09 27-Aug-1983 (Fig. 7b)) (Fig. 8 shaded yellow) reveals that they are not *Telegonus (Telegonus) tsongae* Grishin, 2023 (type locality in USA: Texas, Starr Co.) (Fig. 8 blue), the only species from the group previously known from the US (Fig. 8 shaded green), but *Telegonus (Telegonus) misitra* (Plötz, 1881) (type locality in Mexico) (Fig. 8 red) instead, because they are in the same clade with its lectotype. Thus, these specimens represent the first US records of *T. misitra*, which was known from Mexico and Costa Rica.




Fig. 8. Phylogenetic trees of selected *Telegonus (Telegonus)* species of the *T. fulgerator* group inferred from protein-coding regions in **a)** the Z chromosome, based on 258,849 positions, and **c)** the mitochondrial genome: *T. tsongae* (blue), *T. misitra* (red), *T. catemacoensis* (purple), *T. azul* (cyan), *T. fulminator* (brown), and *T. fulgerator* (green). Labels of specimens from Arizona and Texas are highlighted in yellow and green, respectively. Note rampant mitochondrial genome introgression that precludes confident species identification using COI barcodes. See Fig. 1 caption for other notations.

Epargyreus cruzi Evans, 1952

Genomic analysis of a male (NVG-24074H05) (Fig. 9a, 10) collected by Kilian Roever in Arizona, Santa Cruz Co., Peña Blanca Canyon on 2-Apr-1960 (Fig. 11 shaded yellow) reveals that it is not *Epargyreus fractigutta* Grishin, 2023 (type locality USA: Texas, Hidalgo County, McAllen) (Fig. 9b), the only representative of Mexican and Central American species of *Epargyreus* Hübner, [1819] confirmed from the US (Fig. 11 blue, shaded green), but *Epargyreus cruzi* Evans, 1952 (type locality Mexico: Veracruz, Cordova) instead, in agreement with its wing patterns (Fig. 9). The differences in wing patterns of these

species are described and illustrated in Zhang et al. (2023b). For instance, the semihyaline spot in the forewing cell $CuA_2-1A+2A$ is offset farther distad from spot in the cell CuA_1-CuA_2 in *E. fractigutta* (Fig. 9b) than in *E. cruza* (Fig. 9a). Whitish lavender overscaling on ventral hindwing in *E. cruza* (Fig. 9a) is better developed between the discal silver spots and the postdiscal white band. However, in *E. fractigutta* (Fig. 9b), the overscaling is better developed between the postdiscal white band and marginal pale-lavender area, which are frequently merged with each other as a result. Due to the potential presence of undescribed species, these phenotypic characters should be used with caution, with genome-based identification providing the highest level of confidence. Thus, this Roever's specimen represents the first




Fig. 9. Specimens of *Epargyreus* from the USA in dorsal (left) and ventral (right) views, data in text: **a)** *E. cruza* ♂ NVG-24074H05 from Arizona; **b)** *E. fractigutta* holotype ♀ NVG-14111F08 from Texas.



Fig. 10. Male genitalia of *Epargyreus cruza* NVG-24074H05 from USA: Arizona in left lateral (left) and dorsal (right) views.




Fig. 11. Phylogenetic trees of selected *Epargyreus* species inferred from protein-coding regions in **a**) the nuclear genome (autosomes), based on 4,239,540 positions, and **b**) the mitochondrial genome: *E. fractigutta* (blue) and *E. cruza* (red). Labels of a specimen from Arizona collected by Roever and other USA specimens are highlighted in yellow and green, respectively. See Fig. 1 caption for other notations.

US record of *E. cruza*, a name previously applied to *E. fractigutta* because of misidentification. We are not aware of *E. cruza* records from Texas.

Ectomis (Asina) mexicanus (H. Freeman, 1969)

Genomic analysis of a male (NVG-24074E05) (Fig. 12) collected by B. Griffin (Bailowitz and Brock 1991, 2022) in Arizona, Pima Co., Box Canyon on 4-Oct-1967 confirms its identification as *Ectomis (Asina) mexicanus* (H. Freeman, 1969) (type locality in Mexico: San Luis Potosi) (Fig. 13 red, shaded yellow). This species has also been recorded from Texas (Fig. 13 red, shaded green).



Fig. 12. *Ectomis (Asina) mexicanus* ♂ NVG-24074E05 from Arizona in dorsal (left) and ventral (right) views, data in text.




Fig. 13. Phylogenetic trees of selected *Ectomis* species inferred from protein-coding regions in **a)** the nuclear genome (autosomes), based on 4,043,244 positions, and **b)** the mitochondrial genome: *E. (Asina) mexicanus* (red) and *Ectomis (Ectomis) octomaculata* (purple). Labels of specimens from Arizona and Texas are highlighted in yellow and green, respectively. The clade corresponding to the subgenus *Asina* Grishin, 2019 (type species *Eudamus asine* Hewitson, 1867) is labeled. See Fig. 1 caption for other notations.

Ectomis (Ectomis) octomaculata (Sepp, [1844])

Genomic analysis of a male (NVG-24074E06) (Fig. 14) collected by Kilian Roever in Arizona, Santa Cruz Co., Santa Rita Mts., Temporal Gulch on 29-Sep-1990 confirms its identification as *Ectomis (Ectomis) octomaculata* (Sepp, [1844]) (type locality in Suriname) (Fig. 13 purple, shaded yellow). This species is also known from Texas (Fig. 13 purple, shaded green).



Fig. 14. *Ectomis (Ectomis) octomaculata* ♂ NVG-24074E06 from AZ in dorsal (left) and ventral (right) views, data in text.

Celaenorrhinus fritzgaertneri (Bailey, 1880)

Genomic analysis of a specimen (NVG-24074F03) (Fig. 15) collected by Kilian Roever in Arizona, Cochise Co., Peloncillo Mts., Cottonwood Canyon, 4800-5200' on 23-Aug-1983 confirms its

identification as *Celaenorrhinus fritzgaertneri* (Bailey, 1880) (type locality in El Salvador) (Fig. 16 shaded yellow). Both *C. fritzgaertneri* and *Celaenorrhinus stallingsi* H. Freeman, 1946 (type locality Mexico: Nuevo Leon, Monterrey) are confirmed from Texas by genomic sequencing (Fig. 16 shaded green).




Fig. 15. *Celaenorrhinus fritzgaertneri* NVG-24074F03 from USA: Arizona in dorsal (left) and ventral (right) views, data in text.




Fig. 16. Phylogenetic trees of selected *Celaenorrhinus* species inferred from protein-coding regions in **a**) the nuclear genome (autosomes), based on 2,246,907 positions, and **b**) the mitochondrial genome: *C. fritzgaertneri* (red) and *C. stallingsi* (blue). Labels of specimens from Arizona and Texas are highlighted in yellow and green, respectively. See Fig. 1 caption for other notations.

***Lerema (Lerema) ochrius occidus* Grishin, new subspecies**

<http://zoobank.org/E9489DAF-BB97-4F6A-9372-68B9B2CA4D91>

(Figs. 17 part, 18)

Definition and diagnosis. Closely related to *Lerema (Lerema) ochrius* Grishin, 2023 (type locality in USA: Texas, Hidalgo Co.) this new subspecies is genetically differentiated from it and corresponds to a separate clade in the Z chromosome and mitochondrial genome trees (Fig. 17), while differing by 1.5% (10 bp) in the COI barcode. Due to somewhat limited genetic differentiation and phenotypic similarity, this taxon is proposed as a subspecies. This new subspecies keys to *Lerema accius accius* (J. E. Smith, 1797) (type locality in USA: Georgia) (J.39.2(a)) in Evans (1955), and is more similar to *L. ochrius* in less reddish-brown and more ochreous-brown tones of ventral side of wings, but differs from it by less olive and more brown tones; being paler and more washed out on the ventral surface with larger and blotchier hindwing pale spots and patches (especially the discal area basad of the postdiscal band); and




Fig. 17. Phylogenetic trees of selected *Lerema* (*Lerema*) species inferred from protein-coding regions in **a**) the Z chromosome, based on 187,050 positions, and **b**) the mitochondrial genome: *Lerema* (*Lerema*) *accius* (blue), *Lerema* (*Lerema*) *ochrius* (purple), and *Lerema* (*Lerema*) *ochrius occidus* ssp. n. (red, labels of AZ specimens highlighted in yellow). See Fig. 1 caption for other notations.

females with larger forewing semihyaline hyaline spots. Due to the cryptic nature of this subspecies, most reliable identification is achieved by DNA, and a combination of the following base pairs is diagnostic in the nuclear genome: *aly276558.4.10*:G749A, *aly276558.4.10*:G844A, *aly276558.4.10*:T897C, *aly276558.4.10*:C936T, *aly276558.4.10*:C1458T, and COI barcode: A100G, A214G, T247T (not C), C274C (not T), T340C, T644C.

Barcode sequence of the holotype. Sample NVG-24087E12, GenBank [PV408306](#), 658 base pairs:

AACCTTTATATTTTATTTTGGAAATTTGAGCAGGAATATTAGGAACCTCTTTAAGCTTATTAATTCGAACAGAAATAGGAAATCCAGGATCTTAAATGGGGATGATCAAAATTTATAATCT
ATTGTTACAGCTCATGCTTTTATTAATTTTATAGTTATACCTATTATAATTTGGAGGATTTGGTAATTGATTAGTCCATTAATACCTGGGAGCTCCTGATATGATATTCACAGAA
TAAATAACATAAGATTTTGAATATTACCTCCCTCATTAACTATTAAATTTCAAGTAGAATTTAGAGAAATGGTGCAGGAACAGGATGAACAGTTTACCCACCTTTATCTCTAATATTGC
CCATCAAGGAGCATCAGTTGATTAGCAATTTTCTCTTCATTTAGCAGGATTTCTTCAATTTTAGGAGCTATTAATTTTATTACTACAATTATTAATACGAGATAGAAATTTATCT
TTTGATCAAAATACCTTTATTTGTTGATCTGTGCGAATTACAGGATTTATTTATTTACTCTCTACTACCTGTATTAGCTGGAGCTATTACTATCTTTTAACTGATCGAAATCTTAATCTCT
CTTTTTTGTCTGAGGAGGTGGTGATCTCTTTTATACCAACATTTATTT

Type material. Holotype: ♂ deposited in the McGuire Center for Lepidoptera and Biodiversity Collection, Gainesville, FL, USA (MGCL), illustrated in Fig. 18a, bears the following five printed (text in italics handwritten) rectangular labels, four white: [AZ: Santa Cruz Co., | Lake Patagonia S.P.], [8 Sept.1990 | leg. Kilian Roever], [MGCL collection | W. McGuire Coll. | Ex K. Roever], [DNA sample ID: | NVG-24087E12 | c/o Nick V. Grishin], and one red [HOLOTYPE ♂ | *Lerema* (*Lerema*) *ochrius* | *occidus* Grishin]. **Paratypes:** 9♂♂ and 5♀♀: 1♂ NVG-24074E09 USA, Arizona, Santa Cruz Co., O'Donnell Canyon at Ewing Ranch nr. Canelo, 23-Aug-1998, Kilian Roever leg. [MGCL] and from Mexico: 1♂ NVG-23076D06 no location details, from Herrich-Schäffer collection, before 1874 [MFNB]; 1♂ NVG-22101G06, CASENT8566924 Baja California Sur, San Jose del Cabo, Pescadero 22-Mar-1944 [CAS]; 1♂ NVG-24075C06 Sonora, Santa Rosa-Yécora Road, 1–2 mi SE of Santa Rosa, 13-Apr-1987, Jim P. Brock [MGCL]; Sinaloa: 1♀ NVG-22056C03 Hwy 40 El Palmito, 1970 m, 12-Oct-1988, John Kemner leg. [TMMC] and Mazatlan, Dec-1916, J. August Kusche leg. [USNM]: 1♂ NVG-7734, USNMMENT 01321574, genitalia NVG170205-19 and 1♀ NVG-7735, USNMMENT 01321575, genitalia




Fig. 18. Type specimens of *Lerema (Lerema) ochrius occidus* in dorsal (left) and ventral (right) views, data in text: **a)** holotype ♂ NVG-24087E12 from USA: Arizona and **b)** paratype ♀ NVG-7735 from Mexico: Sinaloa.

NVG170205-20 (Fig. 18b); 1♂ NVG-5020 Durango, 8 km W Los Mimbres Creek, 16-Sep-1977, W. W. McGuire leg., genitalia NVG151101-71 [USNM]; Jalisco: 1♀ NVG-24075B12 La Garita, 20-Nov-1991, D. L. & J. Lindsley leg. [MGCL] and 1♀ NVG-8664 Hesp-EB 02 424 San Juan de la Montana, 1820 m, GPS 19.67, -103.07, 6-Aug-2010, G. Nogueira leg. [Ernst Brockmann collection]; 1♂ NVG-24075C02 Guanajuato, San Miguel de Allende, 25-Dec-1994, D. L. Lindsley leg. [MGCL]; Oaxaca, Hwy 175, ca. 5 mi N of Oaxaca, J. Kemner leg.: 1♂ NVG-24105C06 11-Aug-1988 [TMMC] and 1♀ NVG-24076D03 22-Jul-1988 [MGCL]; and 1♂ NVG-23085D01 “Mexico, Orizaba, ex coll. Neuburger” old, likely mislabeled [ZSMC].

Type locality. USA: Arizona, Santa Cruz Co., Patagonia Lake State Park.

Etymology. In Latin, *occidens* means west or western, forming the basis of the name, which is treated as a noun in apposition.

Distribution. From the southwestern US through western to southern Mexico.

Comment. While conservatively proposed here as a subspecies pending further research, it may be a species-level taxon.

ACKNOWLEDGMENTS

We acknowledge Ping Chen and Ming Tang for their excellent technical assistance. We are grateful to David Grimaldi and Courtney Richenbacher (AMNH: American Museum of Natural History, New York, NY, USA), Blanca Huertas, David Lees, and Geoff Martin (BMNH: Natural History Museum, London, UK), Chris Grinter, Denise Montelongo, David Bettman, Vince Lee, and the late Norm Penny (CAS:

California Academy of Sciences, San Francisco, CA, USA), Jim Fetzner, Bob Androw, Vanessa Verdecia, Cat Giles, and the late John Rawlins (CMNH: Carnegie Museum of Natural History, Pittsburgh, PA, USA), the late Paul Opler, Chuck Harp, and the late Boris Kondratieff (CSUC: Colorado State University Collection, Fort Collins, CO, USA), Crystal Maier and Rebekah Baquiran (FMNH: Field Museum of Natural History, Chicago, IL, USA), Weiping Xie and Giar-Ann Kung (LACM: Los Angeles County Museum of Natural History, Los Angeles, CA, USA), Théo Léger, Christoph L. Häuser, Viola Richter, and Wolfram Mey (MFNB: Museum für Naturkunde, Berlin, Germany), Andrei Sourakov, Andrew D. Warren, Debbie Matthews-Lott, Riley J. Gott, and Keith R. Willmott (MGCL: McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, Gainesville, FL, USA), Rodolphe Rougerie (MNHP: Muséum National d'Histoire Naturelle, Paris, France), Matthias Nuss and Manuela Bartel (MTD: Museum für Tierkunde, Dresden, Germany), Mario Cupello, Edward G. Riley, Karen Wright, and John Oswald (TAMU: Texas A&M University Insect Collection, College Station, TX, USA), Alex Wild (TMMC: University of Texas Biodiversity Center, Austin, TX, USA), Robert K. Robbins, John M. Burns, and Brian Harris (USNM: National Museum of Natural History, Smithsonian Institution, Washington, DC, USA), Andreas Werno and Ernst Brockmann (ZfBS: Zentrum für Biodokumentation des Saarlandes, Schiffweiler, Germany), and Axel Hausmann, Andreas Segerer, and Ulf Buchsbaum (ZSMC: Zoologische Staatssammlung München, Germany), for granting access to the collections under their care, sampling specimens, and stimulating discussions; to Ernst Brockmann (Lich, Germany), Bernard Hermier (Rémire-Montjoly, French Guiana, France), and Mark Walker (Oceanside, CA, USA) for specimens and leg samples, and to Bernard Hermier for critical review of the manuscript and many helpful comments and corrections. We are indebted to Texas Parks and Wildlife Department (Natural Resources Program Director David H. Riskind) for the research permit 08-02Rev, to U. S. National Park Service for the research permits: Big Bend (Raymond Skiles) for BIBE-2004-SCI-0011 and Yellowstone (Erik Oberg and Annie Carlson) for YELL-2017-SCI-7076. We acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing HPC resources. This study was supported in part by the HHMI Investigator funds and by grants from the National Institutes of Health GM127390 and the Welch Foundation I-1505 to N.V.G.

LITERATURE CITED

- Bailowitz, R. A., and J. P. Brock. 1991.** Butterflies of southeastern Arizona. Sonoran Arthropod Studies, Inc.; Tucson. ix + 342 pp.
- Bailowitz, R. A., and J. P. Brock. 2022.** Southeastern Arizona Butterflies. Wheatmark Publishing Company; Tucson. x + 356 pp.
- Cong, Q., J. Shen, J. Zhang, W. Li, L. N. Kinch, J. V. Calhoun, A. D. Warren, and N. V. Grishin. 2021.** Genomics reveals the origins of historical specimens. *Molecular Biology and Evolution* 38(5): 2166–2176.
- Cong, Q., J. Zhang, and N. V. Grishin. 2019a.** Genomic determinants of speciation. *bioRxiv* BIORXIV/2019/837666.
- Cong, Q., J. Zhang, J. Shen, X. Cao, C. Brevignon, and N. V. Grishin. 2020.** Speciation in North American *Junonia* from a genomic perspective. *Systematic Entomology* 45(4): 803–837.
- Cong, Q., J. Zhang, J. Shen, and N. V. Grishin. 2019b.** Fifty new genera of HesperIIDae (Lepidoptera). *Insecta Mundi* 0731: 1–56.
- Evans, W. H. 1951.** A catalogue of the American HesperIIDae indicating the classification and nomenclature adopted in the British Museum (Natural History). Part I. Introduction and Group A Pyrrhopyginae. The Trustees of the British Museum (Natural History); London. x + 92 pp., pls. 1–9.
- Evans, W. H. 1952.** A catalogue of the American HesperIIDae indicating the classification and nomenclature adopted in the British Museum (Natural History). Part II. Pyrginae. Section I. The Trustees of the British Museum (Natural History); London. v + 178 pp., pls. 10–25.

- Evans, W. H. 1953. A catalogue of the American Hesperiidæ indicating the classification and nomenclature adopted in the British Museum (Natural History). Part III. Pyrginae. Section 2. The Trustees of the British Museum (Natural History); London. v + 246 pp., pls. 26–53.
- Evans, W. H. 1955. A catalogue of the American Hesperiidæ indicating the classification and nomenclature adopted in the British Museum (Natural History). Part IV. Hesperinae and Megathyminae. The Trustees of the British Museum (Natural History); London. v + 499 pp., pls. 54–88.
- Li, W., Q. Cong, J. Shen, J. Zhang, W. Hallwachs, D. H. Janzen, and N. V. Grishin. 2019. Genomes of skipper butterflies reveal extensive convergence of wing patterns. *Proceedings of the National Academy of Sciences of the United States of America* 116(13): 6232–6237.
- Minh, B. Q., M. A. Nguyen, and A. von Haeseler. 2013. Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195.
- Robbins, R. K., Q. Cong, J. Zhang, J. Shen, R. C. Busby, C. Faynel, M. Duarte, A. R. P. Martins, C. Prieto, G. Lamas, and N. V. Grishin. 2022. Genomics-based higher classification of the species-rich hairstreaks (Lepidoptera: Lycaenidae: Eumaeini). *Systematic Entomology* 47(3): 445–469.
- Shen, J., Q. Cong, D. Borek, Z. Otwinowski, and N. V. Grishin. 2017. Complete genome of *Achalarus lyciades*, the first representative of the Eudaminae subfamily of skippers. *Current Genomics* 18(4): 366–374.
- Warren, A. D., K. J. Davis, E. M. Stangeland, J. P. Pelham, K. R. Willmott, and N. V. Grishin. 2024. Illustrated Lists of American Butterflies. [9-III-2024] <<https://www.butterfliesofamerica.com/>>.
- Zhang, J., Q. Cong, and N. V. Grishin. 2023a. Descriptions of one hundred new species of Hesperiidæ. *Insecta Mundi* 1026: 1–115.
- Zhang, J., Q. Cong, and N. V. Grishin. 2023b. Thirteen new species of butterflies (Lepidoptera: Hesperiidæ) from Texas. *Insecta Mundi* 0969: 1–58.
- Zhang, J., Q. Cong, J. Shen, E. Brockmann, and N. V. Grishin. 2019a. Genomes reveal drastic and recurrent phenotypic divergence in firetip skipper butterflies (Hesperiidæ: Pyrrhopyginae). *Proceedings of the Royal Society B: Biological Sciences* 286(1903): 1–6.
- Zhang, J., Q. Cong, J. Shen, and N. V. Grishin. 2022a. Taxonomic changes suggested by the genomic analysis of Hesperiidæ (Lepidoptera). *Insecta Mundi* 0921: 1–135.
- Zhang, J., Q. Cong, J. Shen, P. A. Opler, and N. V. Grishin. 2019b. Changes to North American butterfly names. *The Taxonomic Report of the International Lepidoptera Survey* 8(2): 1–11.
- Zhang, J., Q. Cong, J. Shen, P. A. Opler, and N. V. Grishin. 2019c. Genomics of a complete butterfly continent. *bioRxiv* BIORXIV/2019/829887.
- Zhang, J., Q. Cong, J. Shen, P. A. Opler, and N. V. Grishin. 2020. Genomic evidence suggests further changes of butterfly names. *The Taxonomic Report of the International Lepidoptera Survey* 8(7): 1–40.
- Zhang, J., Q. Cong, J. Shen, P. A. Opler, and N. V. Grishin. 2021. Genomics-guided refinement of butterfly taxonomy. *The Taxonomic Report of the International Lepidoptera Survey* 9(3): 1–54.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2022b. Genomic DNA sequencing reveals two new North American species of *Staphylus* (Hesperiidæ: Pyrginae: Carcharodini). *The Taxonomic Report of the International Lepidoptera Survey* 10(4): 1–13.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2023c. Butterfly classification and species discovery using genomics. *The Taxonomic Report of the International Lepidoptera Survey* 11(3): 1–93.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2023d. Genomic analysis reveals new species and subspecies of butterflies. *The Taxonomic Report of the International Lepidoptera Survey* 11(6): 1–62.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2023e. Genomics-based taxonomic rearrangement of Achlyodini and Carcharodini (Lepidoptera: Hesperiidæ: Pyrginae). *Insecta Mundi* 1016: 1–33.

- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2024a.** Genomic analysis reveals hidden species diversity in *Emesis* Fabricius (Lepidoptera: Riodinidae). *Insecta Mundi* 1082: 1–48.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2024b.** New taxa of butterflies supported by genomic analysis. *The Taxonomic Report of the International Lepidoptera Survey* 12(3): 1–62.
- Zhang, J., Q. Cong, J. Shen, L. Song, and N. V. Grishin. 2024c.** Taxonomic advances driven by the genomic analysis of butterflies. *The Taxonomic Report of the International Lepidoptera Survey* 11(7): 1–42.
- Zhang, J., Q. Cong, J. Shen, L. Song, P. A. Opler, and N. V. Grishin. 2023f.** Additional taxonomic refinements suggested by genomic analysis of butterflies. *The Taxonomic Report of the International Lepidoptera Survey* 11(1): 1–25.
- Zhang, J., D. R. Dolibaina, Q. Cong, J. Shen, L. Song, C. G. C. Mielke, M. M. Casagrade, O. H. H. Mielke, and N. V. Grishin. 2023g.** Taxonomic notes on Neotropical HesperIIDae (Lepidoptera). *Zootaxa* 5271(1): 91–114.
- Zhang, J., J. Shen, Q. Cong, and N. V. Grishin. 2019d.** Genomic analysis of the tribe Emesidini (Lepidoptera: Riodinidae). *Zootaxa* 4668(4): 475–488.

The Taxonomic Report

is a platinum open access peer-reviewed publication of

The International Lepidoptera Survey (TILS)

The International Lepidoptera Survey is registered as a non-profit Limited Liability Company (LLC) in the state of Virginia, U.S.A. The Taxonomic Report (TTR), ISSN 2643-4776 (print) / ISSN 2643-4806 (online) is published for the purpose of providing a public and permanent scientific record. Articles are peer-reviewed but not necessarily through the anonymous editor-mediated review process. Typically, the authors are encouraged to solicit reviews of their manuscripts from knowledgeable lepidopterists before submission. TTR appears in digital, open-access form, is disseminated as a hardcopy to select institutional repositories, and is available as printed copy upon request at the discretion of authors and/or the editor. Printing and postage charges may apply. An initial run of 25 copies is printed on paper to meet ICZN Code recommendation 8B. All published TTR articles are freely available at the archival TTR website (<http://lepsurvey.carolinanature.com/report.html>) and via the following digital repositories:

Internet Archive (<https://archive.org/>)
Biodiversity Heritage Library (<https://www.biodiversitylibrary.org>)
Zobodat (<https://www.zobodat.at/>)
Zenodo (<https://zenodo.org>)
Digital Commons (<https://digitalcommons.unl.edu>)

TILS Purpose

TILS is devoted to the worldwide collection of Lepidoptera for the purpose of scientific discovery, determination, and documentation, without which there can be no preservation.

TILS Motto

“As a world community, we cannot protect that which we do not know”

Articles for publication are sought

Manuscripts may deal with any area of research on Lepidoptera, including faunal surveys, conservation topics, life histories and foodplant records, matters of nomenclature, descriptions of new taxa, methods, etc. Taxonomic papers are particularly welcome. There are no publication charges for authors. Before submitting a manuscript, email **TTR editor, Harry Pavulaan, 606 Hunton Place NE, Leesburg, VA, 20176, USA** at intlepsurvey@gmail.com (cc: to harrypav@hotmail.com if you do not receive a reply within one week) to initiate discussion on how to best handle your material for publication, and to discuss peer review options.

Visit *The International Lepidoptera Survey* on the World Wide Web at:

<http://lepsurvey.carolinanature.com>

&

Join the discussion at our list serve on Groups.io at:

<https://groups.io/g/TILS>

You can subscribe by sending an email to: TILS+subscribe@groups.io

&

Join The International Lepidoptera Survey on Facebook at:

<https://www.facebook.com/groups/1072292259768446>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [The Taxonomic Report](#)

Jahr/Year: 2025

Band/Volume: [12-4](#)

Autor(en)/Author(s): diverse

Artikel/Article: [The Taxonomic Report 12/4 1-18](#)